

## REVIEW ARTICLE

## Advancements in APOE and dementia research: Highlights from the 2023 AAIC Advancements: APOE conference

Courtney M. Kloske<sup>1</sup> | Michael E. Belloy<sup>2,3,4</sup> | Elizabeth E. Blue<sup>5,6</sup> |  
 Gregory R. Bowman<sup>7</sup> | Maria C. Carrillo<sup>1</sup> | Xiaoying Chen<sup>8</sup> | Ornit Chiba-Falek<sup>9</sup> |  
 Albert A. Davis<sup>10</sup> | Gilbert Di Paolo<sup>11</sup> | Francesca Garrett<sup>12,13</sup> | David Gate<sup>14</sup> |  
 Lesley R. Golden<sup>15,16</sup> | Jay W. Heinecke<sup>17</sup> | Joachim Herz<sup>18</sup> | Yadong Huang<sup>19,20</sup> |  
 Costantino Iadecola<sup>21</sup> | Lance A. Johnson<sup>15,16</sup> | Takahisa Kanekiyo<sup>22</sup> |  
 Celeste M. Karch<sup>23</sup> | Anastasia Khvorova<sup>24</sup> | Sascha J. Koppes-den Hertog<sup>25,26</sup> |  
 Bruce T. Lamb<sup>27</sup> | Paige E. Lawler<sup>4,28</sup> | Yann Le Guen<sup>29,30</sup> | Alexandra Litvinchuk<sup>8</sup> |  
 Chia-Chen Liu<sup>22</sup> | Simin Mahinrad<sup>1</sup> | Edoardo Marcora<sup>31</sup> | Claudia Marino<sup>32</sup> |  
 Danny M. Michaelson<sup>33</sup> | Justin J. Miller<sup>7,34</sup> | Josh M. Morganti<sup>16,35</sup> |  
 Priyanka S. Narayan<sup>36</sup> | Michel S. Naslavsky<sup>37,38</sup> | Marlies Oosthoek<sup>39</sup> |  
 Kapil V. Ramachandran<sup>40,41,42</sup> | Abhirami Ramakrishnan<sup>14</sup> | Ana-Caroline Raulin<sup>43</sup> |  
 Aiko Robert<sup>25,26</sup> | Rasha N. M. Saleh<sup>44,45</sup> | Claire Sexton<sup>1</sup> | Nilomi Shah<sup>46</sup> |  
 Francis Shue<sup>43</sup> | Isabel J. Sible<sup>47</sup> | Andrea Soranno<sup>48</sup> | Michael R. Strickland<sup>4</sup> |  
 Julia TCW<sup>49,50</sup> | Manon Thierry<sup>51</sup> | Li-Huei Tsai<sup>52</sup> | Ryan A. Tuckey<sup>53</sup> |  
 Jason D. Ulrich<sup>8</sup> | Rik van der Kant<sup>25,26</sup> | Na Wang<sup>54</sup> | Cheryl L. Wellington<sup>55</sup> |  
 Stacie C. Weninger<sup>56</sup> | Hussein N. Yassine<sup>57</sup> | Na Zhao<sup>22</sup> | Guojun Bu<sup>58</sup> |  
 Alison M. Goate<sup>59</sup> | David M. Holtzman<sup>8</sup>

**Funding information:** NIH, Grant/Award Numbers: K99AG075238, U19AG069701, RF1AG067194, K08NS101118, W81XWH2010934, RF1AG083753, F32AG08483, T32 AG078110, R56AG081417, T32AG05851804, 5R01AG070830, 3RF1NS118558, K24AG053435, 7DP5OD028133, T32AG058518, R01AG082362, R01AG083941, K01AG062683, R56AG078733, R01AG081228, R01AG068395, T32 GM008361, U19AG069701, RF1AG046205, R01AG66395, U54NS110435, U01AG058635, P30EY003790; Alzheimer's Association, Grant/Award Numbers: AARF-20-683984, 23AARG-1026607, FAPESP13/08028-1; NIH/NIA, Grant/Award Numbers: R01AG059737, U01AG058589, R41AG077992, R01AG057522, P01AG073082, U19AG069701, R01AG062837, P30AG066512, P01AG060882; TRIAD, Grant/Award Numbers: R01HL149685, R01AG061186, P01AG073082; BrightFocus Foundation, Grant/Award Numbers: A2022010F, A20174585, A2021015F; NS126467 ApoE4, neurovascular injury and cognitive impairment; National Institute on Aging (NIA), Grant/Award Numbers: R01AG060056, R01AG062550, R01AG080589, R01AG081421, R01AG054671, RF1AG077627, U19, RF1AG076124, RF1AG078362, R01AG067063, R01AG054434, R01AG055770, R21AG056518, P30AG066530; National Institute of General Medical Sciences (NIGMS) Center of Biomedical Research Excellence in CNS Metabolism, Grant/Award Number: P20GM148326; Biology and pathobiology of apoE in aging and Alzheimer's Disease, Grant/Award Numbers: GM131839-03, U54 AG065181, U54AG054345, RF1AG074566; Cure Alzheimer's Fund; Marie Skłodowska-Curie Actions, Grant/Award Number: 890650; Edward N. and Della L. Thome Memorial Foundation; Robert J. Kleberg, Jr. and Helen C. Kleberg Foundation; Remondi Family Foundation; US National Institute of Neurological Disorders and Stroke and National Institute on Aging, Grant/Award Numbers: UH3 NS100121, RF1 NS110048; Good Ventures and Open Philanthropy; NIH Office of the Director, Grant/Award Number: DP5 OD019833; Massachusetts General Hospital Executive Committee (ECOR) on Research; MGH Research Scholar Award; Eichenbaum Foundation; NIH-NIDDK Intramural Research Program; CNPq, Grant/Award Number: 304746/2022-3; Department of Defense, Grant/Award Number: W81XWH-21-1-0093; Fidelity Biomedical Research Initiative; Medical Research Council (MRC, UK); NuBrain Consortium, Grant/Award Number: MR/T001852/1; NSF GRFP Fellowship, Grant/Award Number: DGE-1745038; Alzheimer's Drug Discovery Foundation, Grant/Award Number: GC-201711-2014197; Cure Alzheimer Fund and BrightFocus Foundation; Kenneth and Bette Volk Endowed Chair of Neurology; Vranos and Tiny Foundations; Ms. Lynne Nauss; JPB Foundation

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). *Alzheimer's & Dementia* published by Wiley Periodicals LLC on behalf of Alzheimer's Association.

**Correspondence**

Courtney M. Kloske, Alzheimer's Association,  
225 N Michigan Ave, Floor 17, Chicago, IL  
60601, USA.  
Email: [cmkloske@alz.org](mailto:cmkloske@alz.org)

**Abstract**

**INTRODUCTION:** The apolipoprotein E gene (*APOE*) is an established central player in the pathogenesis of Alzheimer's disease (AD), with distinct apoE isoforms exerting diverse effects. apoE influences not only amyloid-beta and tau pathologies but also lipid and energy metabolism, neuroinflammation, cerebral vascular health, and sex-dependent disease manifestations. Furthermore, ancestral background may significantly impact the link between *APOE* and AD, underscoring the need for more inclusive research.

**METHODS:** In 2023, the Alzheimer's Association convened multidisciplinary researchers at the "AAIC Advancements: APOE" conference to discuss various topics, including apoE isoforms and their roles in AD pathogenesis, progress in apoE-targeted therapeutic strategies, updates on disease models and interventions that modulate apoE expression and function.

**RESULTS:** This manuscript presents highlights from the conference and provides an overview of opportunities for further research in the field.

**DISCUSSION:** Understanding apoE's multifaceted roles in AD pathogenesis will help develop targeted interventions for AD and advance the field of AD precision medicine.

**KEYWORDS**

Alzheimer's disease, APOE, apolipoprotein E, conference proceedings, dementia, lipids, microglia, neuroinflammation, risk factor, therapeutics, vasculature

**Highlights**

- APOE is a central player in the pathogenesis of Alzheimer's disease.
- APOE exerts a numerous effects throughout the brain on amyloid-beta, tau, and other pathways.
- The AAIC Advancements: APOE conference encouraged discussions and collaborations on understanding the role of APOE.

**1 | INTRODUCTION**

In 1973, researchers identified an arginine-rich protein in the very low-density lipoprotein (VLDL) of patients with a rare lipid metabolism disorder called familial hypercholesterolemia type III.<sup>1</sup> This protein was further characterized and named apolipoprotein E, or apoE,<sup>2</sup> which is encoded by the apolipoprotein E gene (*APOE*) and has three main alleles:  $\epsilon 2$ ,  $\epsilon 3$ , and  $\epsilon 4$  that translate to three isoforms: apoE2, apoE3, and apoE4, respectively.<sup>3</sup> The apoE protein was later found to play a role in lipid metabolism, cardiovascular disease (CVD), and Alzheimer's disease (AD).<sup>4-8</sup>

Over the past decades, apoE has attracted greater attention in AD research. The *APOE*  $\epsilon 2$  allele has been determined to be protective against AD, while the  $\epsilon 4$  allele was found to be associated with increased AD risk.<sup>7-9</sup> Research has also found an important correlation between the frequency of *APOE* variants across different

populations and their associated risk for AD pathology.<sup>10</sup> This landscape of findings has opened an important avenue for a more precise mechanistic understanding of AD etiology and pointed toward numerous potential treatment targets. However, it is important to note that most research in the area of apoE and AD has focused on non-Hispanic White populations, leading to a lack of diversity in the field and undermining the breadth and impact of apoE and AD research on diverse populations.

The Alzheimer's Association convened the Alzheimer's Association International Conference (AAIC) Advancements: APOE conference on March 6-7, 2023, to foster new collaborations and develop novel research directions with the potential to break down multidisciplinary barriers and drive transformative neuroscience research. This manuscript provides an overview of the discussions from this conference while highlighting knowledge gaps in the field that need to be addressed by future research.

## 2 | THE APOE GENE AND NEURODEGENERATIVE DISEASE RISK

The *APOE*  $\epsilon 4$  has been identified as the most significant genetic risk factor associated with AD<sup>11</sup> and other neurodegenerative diseases, including frontotemporal lobar dementia (FTLD), Lewy body dementia (LBD), and other amyloid-beta ( $A\beta$ ) and tau pathologies.<sup>12</sup> While *APOE*  $\epsilon 2$  is established as protective against AD, *APOE*  $\epsilon 4$  is generally associated with greater AD risk if using *APOE*  $\epsilon 3$  as a reference allele. In addition, *APOE*  $\epsilon 4$  is associated with decreased age of AD onset and promotion of  $A\beta$  and tau pathology, inflammation, and neurodegeneration.<sup>13</sup> Pathways associated with the risk of AD and other dementia, such as lipid metabolism, cardiovascular and cerebrovascular diseases, altered efferocytosis, inflammation, trafficking, and integrated stress response, are also associated with *APOE*.<sup>13-18</sup> By interrogating the relationship between *APOE* and various disease states and pathways, researchers have learned that *APOE* can influence AD risk through a variety of pathways.

Several rare *APOE* variants have also been identified. For example, the rare *APOE*-Ch (R136S) mutation, located within the overlapping N-terminal heparan sulfate proteoglycan (HSPG) and receptor binding domain of apoE,<sup>4,19</sup> was identified in a Colombian kindred with a dominant *PSEN1* E280A mutation.<sup>20</sup> While *PSEN1* mutation carriers often exhibit significant  $A\beta$  and tau and burden, hippocampal atrophy, and brain hypometabolism,<sup>21,22</sup> *APOE*-Ch carriers with *PSEN1* mutation have been shown to be protective against AD, with one homozygous individual exhibiting minimal tau pathology, hippocampal atrophy, and hypometabolism despite the presence of a *PSEN1* mutation.<sup>20,21</sup> Studies using the Alzheimer's Disease Sequencing Project (ADSP) data have identified several other rare *APOE* variants associated with AD, including the R145C commonly found in the African American AD population.<sup>23</sup> The R145C variant causes a heterozygotic effect, associated with an increased risk of AD and reduced age of onset only in the presence of an  $\epsilon 4$  allele.<sup>24</sup> Insights from the conference highlight *APOE* variants identified in populations of European ancestry including L28P, V236E (also known as the Jacksonville variant), and R251G. Studies have shown that the V236E and R251G variants are associated with decreased AD risk, whereas the L28P variant has not been associated with AD risk.<sup>23,24</sup>

## 3 | RACIAL AND ETHNIC DIFFERENCES IN APOE AND AD RISK

Although significant progress has been made in understanding the link between *APOE* and AD, most studies have focused on non-Hispanic White populations and do not adequately represent human diversity.<sup>25</sup> The AAIC *advancements: APOE conference* emphasized the need to explore and address population-level distinctions in *APOE* and AD, particularly through research involving diverse populations. For example, several presentations highlighted that the frequency of *APOE*  $\epsilon 4$  varies substantially across different populations, with central Africa and northern Europe displaying the highest *APOE*  $\epsilon 4$  frequencies.<sup>25</sup>

### RESEARCH IN CONTEXT

- 1. Systematic review:** The role of apolipoprotein E (*APOE*) in neurodegenerative diseases, including Alzheimer's and other dementia, is an active and growing area of research. The authors of this manuscript report updates and advances in research presented at the 2023 AAIC Advancements: *APOE Conference*, held in March of 2023.
- 2. Interpretation:** There have been strides in research identifying the role of *APOE* in dementia research. This manuscript highlights the research presented at the 2023 AAIC Advancements *APOE Conference* including the role of apoE isoforms and their roles in AD pathogenesis, progress in apoE-targeted therapeutic strategies, and updates on disease models and interventions that modulate apoE expression and function. This manuscript also highlights apoE influences not only amyloid-beta and tau pathologies but also lipid and energy metabolism, neuroinflammation, cerebral vascular health, and sex-dependent disease manifestations. Furthermore, ancestral background may significantly impact the link between *APOE* and AD, underscoring the need for more inclusive research.
- 3. Future directions:** Understanding apoE's multifaceted role in AD pathogenesis will help develop targeted interventions for AD and advance the field of AD precision medicine.

Among European, African American, Hispanic, and Japanese populations, *APOE*  $\epsilon 4/\epsilon 4$  homozygotes are most common in Japanese populations.<sup>26</sup>

The three most common *APOE* alleles appear within the general global population at varying frequencies. The *APOE*  $\epsilon 3$  allele is the most common across all human populations, with frequencies ranging from 85% in Asia to 69% in Africa. The *APOE*  $\epsilon 4$  allele is enriched in indigenous populations of Central Africa, Oceania, and Australia, with frequencies ranging from 26% to 40%, while the Mediterranean area or south China has a low frequency of <10%. The *APOE*  $\epsilon 2$  allele is the least common, with a worldwide frequency of about 7% and no apparent geographical trend.<sup>13</sup> Overall, accumulating evidence suggests that the frequencies of the major *APOE* alleles differ dramatically across geographical, racial, and ethnic groups, the genetic-environmental interplay heterogeneously impacts disease risk within those populations,<sup>27-29</sup> and *APOE*-related risk of AD may be sex-dependent.<sup>29,30</sup>

Furthermore, research has shown that the impact of *APOE* genotype on AD risk varies across populations with diverse ancestral backgrounds. For example, in Caribbean Hispanic individuals, African-derived ancestry of *APOE* genotype is associated with a lower risk of

AD compared to individuals with European-derived *APOE* genotype.<sup>31</sup> These results are consistent with studies in African American and Puerto Rican populations.<sup>32</sup> Therefore, a commitment to the diversification of *APOE* and AD research is needed to expand the understanding of the role of *APOE* in AD for diverse populations affected by the disease.

## 4 | STRUCTURE OF THE APOE PROTEIN AND ITS BIOCHEMICAL PROPERTIES

apoE is the most abundant apolipoprotein in the brain<sup>33</sup> and, depending on the isoform, can either contribute to AD pathologies or protect against them. Initially studied in hyperlipidemia, *APOE* encodes a 317-amino-acid protein, apoE, that includes an 18-residue signal peptide. Cleavage of this signal peptide, followed by glycosylation at one of the several glycosylation sites within the sequence, results in the mature apoE protein, 299 amino acids in length and approximately 34 kDa. This protein's tertiary structure comprises two independently folded domains separated by a hinge region. The N-terminal domain includes the receptor binding domain and one of two heparin-binding regions. The C-terminal domain includes the lipid-binding domain and the other heparin-binding region. The work presented at the conference and through this section suggests these domains enable the spectrum of interactions and functions that make apoE a critical component of various neurodegenerative diseases.<sup>34</sup>

### 4.1 | Conformational heterogeneity in apoE protein

While apoE plays a significant role in AD pathology, the structural determinants of apoE that contribute to this pathogenicity remain unclear. In vivo, apoE is secreted primarily by astrocytes in discoidal HDL-like lipoproteins.<sup>35–37</sup> Under disease conditions, microglia upregulate apoE expression.<sup>38,39</sup> Research suggests that apoE can exist in various conformational states in lipid-free and lipid-bound forms, which may have important implications for its biological functions.<sup>40</sup> apoE undergoes extensive conformational changes upon binding to lipids. The four-helix bundle in nonlipidated apoE unfolds, and the hydrophobic portions of the amphipathic  $\alpha$ -helix associate with lipid.<sup>41,42</sup> This conformational change is critical to apoE's function as nonlipidated apoE is unable to bind low-density lipoprotein receptor (LDLR).<sup>43,44</sup>

The structural changes accompanying lipidation are poorly understood. The monomeric form of the nonlipidated apoE has been proposed as the competent form for lipid binding.<sup>45</sup> However, structural characterization has been elusive because of the strong propensity of apoE for oligomerization, with dimers already forming at a nanomolar concentration.<sup>45–47</sup> As a result, the structural features of nonlipidated apoE have been determined only for N-terminal domain fragments that lack the C-terminal domain<sup>46,47</sup> and for a “monomeric” mutant of apoE  $\epsilon$ 3 that contains several mutations in the C-terminal domain.<sup>48</sup> To close this knowledge gap and overcome these

experimental challenges, single-molecule Förster resonance energy transfer (smFRET) was implemented to isolate apoE4 monomers and measure distances between fluorescently labeled positions of apoE4.<sup>40</sup> smFRET revealed three major conformational ensembles of apoE, which differ largely for the conformations of the C-terminal domain with respect to the closed, open, and extended N-terminal domains. Furthermore, smFRET experiments using DMPC liposomes to analyze lipidated apoE4 revealed both compact and expanded conformations.<sup>40</sup>

### 4.2 | Biochemical and structural changes in rare APOE variants

Recombinant forms of apoE  $\epsilon$ 3 variants—apoE3-Christchurch (apoE3-Ch) and *APOE**E*poE3-R145C, but not apoE3-Jacksonville (apoE3-Jac)—show reduced heparin-binding affinity and mixed LDLR binding affinity.<sup>20,49,50</sup>

Molecular dynamics simulations allow for comparing conformational and structural characteristics of common and rare apoE isoforms when in a closed conformation. Hydrogen bond occupancy analyses between common apoE isoforms, apoE  $\epsilon$ 3, apoE  $\epsilon$ 2, and apoE  $\epsilon$ 4, support prior findings that apoE  $\epsilon$ 2 has altered hydrogen bond occupancy between residues in the receptor binding domain.<sup>51</sup> In addition, all apoE isoforms show similar conformational flexibility measured via root mean square fluctuation; however, the C112R substitution present in apoE  $\epsilon$ 4 may increase conformational motion in specific functional domains of apoE  $\epsilon$ 4 compared to apoE  $\epsilon$ 3 and apoE  $\epsilon$ 2. While the rare variant apoE4-R251G has the same C112R substitution, it shows a distinct pattern of conformational motion in these same regions compared to apoE  $\epsilon$ 4.

### 4.3 | Post-translational modifications in apoE

Post-translational modifications (PTMs) are chemical transformations that proteins undergo after translation, resulting in a functional alteration for the protein. Many proteins associated with AD undergo PTMs, including apoE, APP, tau, and  $A\beta$ . In most cases, PTMs are imperative to the proper functioning of a protein. Such PTMs can directly affect apoE structure and function. However, some PTMs may impede protein function and result in downstream consequences. One particularly critical PTM in AD is glycosylation, a process required to produce mature apoE.<sup>52</sup> Immunoprecipitation and mass spectrometry assessing O-glycosylated apoE in plasma and cerebrospinal fluid (CSF) samples of older adults showed that CSF-derived apoE has a higher proportion of total glycosylation than plasma-derived apoE.<sup>52</sup> In addition, plasma-derived apoE O-glycosylation levels differ significantly by *APOE* genotype and CSF amyloid status, with *APOE*  $\epsilon$ 4/ $\epsilon$ 4 carriers showing low apoE O-glycosylation. These findings presented at the conference suggest that O-glycosylation of apoE could be a potential CSF or blood-based biomarker for brain amyloidosis and AD diagnosis.

#### 4.4 | Role of apoE and ABCA1 in lipid transport and metabolism

In addition to modifications to APOE, APOE also plays an important role in interacting with other proteins. The adenosine triphosphate (ATP)-binding cassette transporters A1 (ABCA1) and apoE are both critical players in lipid metabolism, particularly in the context of lipid transport in the central nervous system (CNS).<sup>53</sup> In high-density lipoprotein (HDL) particles, apoA1 binds ABCA1, leading to the translocation of cholesterol across the phospholipid bilayer membrane, thereby regulating cholesterol efflux. ABCA1 is a critical element to consider in AD studies because of its potential relevance to apoE metabolism and supporting apoE protein levels in the CNS. Given that research suggests a link between ABCA1, apoE levels in the CNS, and AD pathology, proper ABCA1 functioning is important to understanding apoE metabolism and AD progression. While the structure of APOE is necessary to consider, the additional functions of APOE were also discussed at the AAIC Advancements conference.

### 5 | APOE, MICROGLIAL FUNCTION, IMMUNOMODULATION, AND CHANGES TO GLIAL LIPID METABOLISM IN AD

The mechanisms underlying the pathogenic effects of apoE in AD are complex and involve multiple pathways.<sup>54</sup> apoE is involved in several potentially pathologic processes that may lead to the development of dementia. For example, apoE isoforms are linked to differential levels of A $\beta$ -accumulation in the brain, with apoE  $\epsilon$ 4 associated with the highest accumulation levels and apoE  $\epsilon$ 2 associated with the lowest.<sup>55</sup> In addition, apoE plays a role in blood-brain barrier (BBB) integrity and induction of microglia-driven phagocytosis and inflammation. apoE  $\epsilon$ 4 is associated with BBB dysfunction, increased phagocytosis, and proinflammatory activity.<sup>56-58</sup> A growing number of studies suggest that apoE mediates the induction of disease-associated microglia (DAM).<sup>38</sup> apoE has also been implicated in several metabolic abnormalities, including abnormal glucose metabolism, altered lipidome and metabolome, mitochondrial dysfunction, and decreased oxygen consumption, all of which are involved in the pathology of dementias.<sup>59-63</sup> The AAIC Advancements: APOE conference highlighted new data to help delineate the mechanisms underlying apoE's contribution to AD-associated pathologies.

Evidence suggests that inflammation and immunomodulation pathways play a role in the pathogenesis of AD. Pathways that rely on or are modulated by specific AD risk factors, including triggering receptors expressed on myeloid cells 2 (TREM2) and specific cell types, such as glial cells, have been implicated in AD. The combination of these risk factors, cellular changes, and individual APOE genotypes results in the heterogeneous immunophenotypes observed in AD.<sup>64</sup>

#### 5.1 | apoE in microglial functions and AD

Research from mouse models of AD pathology suggests that while the microglial response to A $\beta$  is generally protective against axonal injury, microglial responses to tau pathology promote neurodegeneration. In A $\beta$  and tau pathology, microglia exhibit TREM2 and apoE-dependent upregulation of genes, colloquially referred to as DAM. apoE isoforms may differentially impact microglial activation in a cell-autonomous manner.<sup>64</sup> One study presented at the conference suggested microglial apoE  $\epsilon$ 3 expression improves cognitive behavior, synaptic function, and microglial responses to injury in a tamoxifen-inducible transgenic animal model, iE3/Cx3cr1-Cre<sup>ER</sup> mice, while microglial from iE4/Cx3cr1-Cre<sup>ER</sup> elicits minimal or an opposite effect. Furthermore, microglial apoE  $\epsilon$ 3 expression increases plaque-associated microgliosis and reduces A $\beta$  deposition and associated neuronal toxicity, whereas microglial apoE  $\epsilon$ 4 expression either compromises or has no effects on these outcomes by impairing lipid metabolism.<sup>65</sup>

In addition, research has shown that TREM2 can interact with apoE and its isoforms to induce microgliosis.<sup>66</sup> Research presented at the conference showed that TREM2-independent microgliosis promotes tau-mediated neurodegeneration in the presence of APOE4, increasing neutral lipid accumulation in microglial phagolysosomes.<sup>67</sup> TREM2 deletion does not counteract the detrimental effect of apoE  $\epsilon$ 4 on tau-mediated neurodegeneration and synaptic loss, nor does it lower tau pathology or attenuate microglial lysosomal burden. Homeostatic microglial markers are preserved despite neurodegeneration in Tau-APOE $\epsilon$ 4 (TE4)-TREM2 knockout (KO) mice. However, some reactive microglial markers, such as Clec7a, are reduced by TREM2 KO in TE4 mice. TE4 and TE4-TREM2 KO mice display increased lysosomal enzymes; thus, TREM2 KO increases phagolysosome volume in TE4 mice.<sup>67</sup>

##### 5.1.1 | apoE-modulated microglial response to myelin damage

Many of the identified 75 genomic loci that impact the risk of AD regulate lipid metabolism.<sup>11</sup> Recently, it has been shown that apoE  $\epsilon$ 4 is associated with gene expression changes in cholesterol- and lipid-related pathways across all human brain cell types.<sup>10,68</sup> This dysregulated cholesterol activity is associated with reduced myelination in apoE  $\epsilon$ 4 carriers. Solubilizing cholesterol relieves the cholesterol burden, and treatment with cyclodextrin increases myelination in apoE  $\epsilon$ 4 culture and restores axonal myelination in apoE  $\epsilon$ 4 KI mice. Notably, cyclodextrin-treated mice exhibit greater interhemispheric fast gamma phase-locking, associated with improved cognition.<sup>68</sup>

In response to myelin damage, microglia clear myelin debris, secrete regenerative factors, and modulate the extracellular matrix. Microglial response to myelin damage is a common pathological feature in neurodegenerative disorders. To determine how apoE isoforms affect microglial responses to myelin damage, a recent study by Wang et al.

used the apoE-targeted replacement (TR) mouse models fed with normal diet or cuprizone to induce demyelination. Significant isoform-dependent differences in microglial activation and function were observed. apoE  $\epsilon$ 4-TR mice had more myelin debris accumulation than apoE  $\epsilon$ 2-TR mice upon cuprizone treatment. The microglia in apoE  $\epsilon$ 2-TR mice proliferated more and demonstrated hyperactivity compared to apoE  $\epsilon$ 4-TR mice. Genes related to the immune response, inflammatory signaling, and lipid metabolism were upregulated in apoE  $\epsilon$ 2-TR mice and downregulated in apoE  $\epsilon$ 4-TR mice. Moreover, apoE  $\epsilon$ 4 microglia had a reduced ability to clear myelin debris due to a weaker phagocytosis ability and enhanced lipid droplet accumulation than apoE  $\epsilon$ 2 microglia.<sup>69</sup> Thus, apoE isoforms may differentially regulate microglia activation and lipid metabolism in the context of myelin damage.

## 5.2 | apoE, immunomodulation, and immunometabolism in AD

A study presented at the conference demonstrated that mice with tauopathy, but not A $\beta$ , developed a unique innate and adaptive immune response and that the depletion of microglia or T-cells or inhibition of interferon-gamma (IFN- $\gamma$ ) signaling can significantly ameliorate brain atrophy and improve cognitive behavior. Consistent with the finding that apoE  $\epsilon$ 4 exacerbates tau-mediated neurodegeneration, T cell infiltration increases in mice with apoE  $\epsilon$ 4 but did not present in the tau mice lacking apoE. These data suggest apoE is important in immunomodulation and may link innate and adaptive immunity.<sup>70,71</sup>

Spatial transcriptomics in 5XFAD mice expressing apoE  $\epsilon$ 4 revealed a unique cortical transcriptomic signature characterized by increased expression of DAM/MGnD microglia genes and biomarkers related to microglial activation, lipid metabolism, complement activation, and synapse pruning.<sup>72</sup> Furthermore, in the presence of both advanced age and A $\beta$  overexpression, apoE  $\epsilon$ 4 exacerbated expression of plaque-induced genes (PIGs), decreased expression of myelin and oligodendrocyte genes (OLIGs), and was associated with transcriptomic signatures related to microglial activation and lipid metabolism.<sup>73</sup> Finally, when analyzing the brain in a spot-by-spot manner, increases in local plaque load were highly correlated with changes in genes related to lipid metabolism in apoE  $\epsilon$ 4 but not apoE  $\epsilon$ 3, expressing 5XFAD mice. These data suggest that apoE  $\epsilon$ 4 negatively impacts microglial function and alters lipid metabolism, which in concert may lead to the dysregulated immunometabolism characteristic of AD.<sup>72</sup>

## 5.3 | apoE and glial lipid metabolism

Lipids play a major role in the processes that are associated with AD pathogenesis, including synaptic plasticity, inflammation, and oxidative stress. Glial lipid metabolism is one of the several pathways implicated in AD pathology.<sup>74</sup> Researchers have interrogated the relationships between AD pathologies and lipid metabolism, such as the association of specific apoE isoforms and calcium-dependent phospholipase A2 (cPLA2) activation.<sup>75</sup> For example, apoE  $\epsilon$ 4 increases calcium-

dependent cPLA2 activation, leading to accentuated eicosanoid lipid metabolism and neuroinflammation. Compared to individuals with the apoE  $\epsilon$ 3/ $\epsilon$ 3 genotype, astrocytes from apoE  $\epsilon$ 4 carriers exhibit increased cPLA2 expression, particularly surrounding A $\beta$  plaques. Notably, cPLA2 reduction ameliorates cognitive deficits in an AD mouse model and resolves chronic neuroinflammation from low docosahexaenoic acid DHA diets in apoE  $\epsilon$ 4-TR mice and increases brain DHA and eicosapentaenoic acid (EPA) levels.<sup>76</sup> cPLA2 inhibition also reverses acute LPS-induced inflammation. No cPLA2 inhibitor has progressed to human studies; future research will involve examining small molecules that inhibit cPLA2 for mitigating neuroinflammation.

Another major piece of lipid metabolism involves cholesterol.<sup>77,78</sup> Cholesterol metabolism is dysregulated in both *post mortem* AD brains and animal AD models, leading to cholesterol ester accumulation in the brain and glial cells in a manner consistent with defects in cholesterol efflux. apoE  $\epsilon$ 4 microglia display cholesterol dysregulation consistent with a cholesterol ester storage disorder.<sup>79–83</sup> A recent study presented at the conference suggests the potential mechanism underlying dysregulated cholesterol metabolism in apoE  $\epsilon$ 4/ $\epsilon$ 4 astrocytes. A change in intracellular cholesterol distribution causes cells to upregulate *de novo* cholesterol synthesis and increase cell surface lipid receptors to facilitate cholesterol uptake.<sup>84</sup> Simultaneously, reduced phagocytic capacity, apoE levels, and lipid transporter levels lead to decreased cholesterol secretion and efflux. apoE  $\epsilon$ 4/ $\epsilon$ 4 astrocytes exhibit an overall decrease in genes associated with the autophagy pathway and network, especially lysosomal acidification, membrane, biogenesis, and hydrolases. Compared to apoE  $\epsilon$ 3/ $\epsilon$ 3 astrocytes, apoE  $\epsilon$ 4/ $\epsilon$ 4 astrocytes exhibit more LDL binding but reduced uptake of lipids, exposing lipids on cell surfaces and leading to reduction of adhesion to the actin cytoskeleton by apoE  $\epsilon$ 4/ $\epsilon$ 4 astrocytes. Furthermore, enriched matrisome pathways associated with upregulated chemotaxis, glial activation, and lipid biosynthesis originate from astrocytes in the presence of neurons in the AD brain.<sup>84</sup>

Another way to interrogate the relationship between AD and lipid metabolism is through unbiased lipidomics analysis of induced pluripotent stem cells (iPSC)-derived brain cells. Lipidomics data demonstrates that APOE genotypes uniquely affect lipid metabolism in iPSC-microglia, neurons, and astrocytes. Compared to apoE  $\epsilon$ 3, apoE  $\epsilon$ 4 microglia and astrocytes display increased triglycerides<sup>85</sup> and cholesterol ester levels.<sup>86,87</sup> Proteomic analyses showed an upregulation of mitochondrial pathways in apoE  $\epsilon$ 4 and apoE KO astrocytes, whereas the same pathways are downregulated in the apoE  $\epsilon$ 4 and apoE KO microglia. Additionally, apoE  $\epsilon$ 4 glia increased cholesterol synthesis and altered immune signaling, indicating tight coupling between intracellular cholesterol distribution and immune activity of glia.<sup>86,87</sup>

Using a novel conditional mouse model, researchers have identified that a “midlife midlife switch” involving a full-body transition from the expression of apoE  $\epsilon$ 4 to apoE  $\epsilon$ 2 impacts the cerebral transcriptome and lipidome. Single-cell RNA sequencing identified astrocytes, oligodendrocytes, endothelial cells, and microglia as the cell types most impacted by the apoE  $\epsilon$ 4 to apoE  $\epsilon$ 2 “switch.” Each of these cell types has differentially expressed genes that have been associated with AD-related pathways, particularly pathways involved with lipid and

carbohydrate metabolism. Lipidomic analyses further showed that several glycerophospholipid species, including phosphatidylcholines, were impacted by the midlife apoE  $\epsilon$ 4:apoE  $\epsilon$ 2 allele switch, which is consistent with previous *post mortem* human AD studies. These findings suggest that apoE  $\epsilon$ 4's transcriptional and lipidomic signatures are not set in stone during development or early life. Rather, these signatures appear acutely malleable within the specific parameters tested here when mice are switched at a "mid-lifemidlife" (6 months) and assessed 1 month post apoE  $\epsilon$ 4:apoE  $\epsilon$ 2 switch.<sup>88</sup>

Using unbiased lipidomics coupled with immunostaining, a recent study demonstrated that apoE  $\epsilon$ 4 promotes cholesterol ester accumulation in the endolysosomal compartment of microglia of aged P301S tauopathy mice when compared to P301S mice expressing apoE  $\epsilon$ 3 or apoE KO animals.<sup>83</sup> Increasing cholesterol efflux via LXR agonist diet or by ABCA1 overexpression significantly decreased lipid droplet accumulation *in vitro* and dramatically reduced tau pathology and associated neurodegeneration, neuroinflammation, and microglial cholesterol ester accumulation *in vivo*. These data demonstrate that promoting cholesterol efflux could be a beneficial therapeutic approach to reduce tau pathology and apoE  $\epsilon$ 4-linked neurodegeneration.

## 6 | APOE AND NEUROVASCULAR DYSFUNCTION IN AD

### 6.1 | Role of apoE receptors in cerebrovascular integrity

apoE  $\epsilon$ 4 has been previously shown to impair neurovascular integrity, which may in turn enhance neurodegeneration.<sup>89</sup> At the conference, research presented highlighted that deleting the major apoE receptor LRP1 affects the cerebrovascular system and cognitive performance in a genotype-dependent manner. Specifically, impaired spatial memory, excessive perivascular glial activation, reduced cerebrovascular collagen IV, and disrupted BBB integrity were observed in apoE  $\epsilon$ 4 mice, but not in apoE  $\epsilon$ 3 mice with smLrp1 removed. This suggests that LRP1 in vascular mural cells maintains cerebrovascular integrity and functions in an apoE genotype-dependent manner.<sup>90</sup>

One mechanism underlying the link between apoE  $\epsilon$ 4 and impaired neurovascular integrity could be attributed to BBB leakage. Human AD brains show widespread decreases in tight junction proteins, which are critical for maintaining BBB integrity.<sup>10</sup> APOE  $\epsilon$ 4 carriers display BBB breakdown in the hippocampus and medial temporal lobe, while non-carriers do not. This BBB breakdown increases with impaired cognition and is independent of CSF A $\beta$  and CSF tau presence.<sup>57</sup>

### 6.2 | Role of border-associated macrophages

Recent findings also suggest that apoE  $\epsilon$ 4-induced neurovascular dysfunction may be mediated by border-associated macrophages, specifically perivascular macrophages (PVM). A recent study presented at the conference showed that apoE  $\epsilon$ 4, but not apoE  $\epsilon$ 3, induces reactive oxy-

gen species (ROS) production in PVM *in vivo* and *ex vivo*. Conditional deletion of apoE  $\epsilon$ 4 in PVM rescues neurovascular dysfunction, indicating that PVM cells are both the source and target of apoE  $\epsilon$ 4 and drive neurovascular dysfunction.<sup>91</sup>

### 6.3 | apoE and blood pressure variability in dementia

Blood pressure variability (BPV) has also emerged as a risk factor for dementia, particularly in individuals with apoE  $\epsilon$ 4. Elevated BPV, independent of mean BP, has been associated with cognitive decline, AD progression, and cerebrovascular disease burden. Interestingly, a recent observational study found that only people with apoE  $\epsilon$ 4 isoform displayed an association between high BPV and cognitive decline.<sup>92</sup> The accelerated cognitive decline among apoE  $\epsilon$ 4 carriers may be due to the "tsunami effect," in which large fluctuations of BP exacerbates apoE  $\epsilon$ 4-associated vulnerability, such as a leaky BBB. Such findings provide opportunities to develop targeted interventions for APOE  $\epsilon$ 4 carriers with high BPV.<sup>92</sup>

## 7 | MODELS OF APOE PHENOTYPES

Both animal and iPSC models are important not only for understanding the pathology and biological mechanisms of apoE but also for developing therapeutics. These models provide opportunities for preclinical therapeutic testing once researchers have identified mechanisms for modifying apoE function. Although iPSC lines enable researchers to study molecular mechanisms, the AD research community has primarily used animal models to characterize apoE's broad physiological effects. Recently, researchers have developed humanized mouse models by using knock-in (KI) techniques to integrate human APOE into the mouse genome.<sup>93</sup>

### 7.1 | Mouse models

The first KI mouse model of human APOE was developed by researchers at Duke University, which is now distributed through Taconic Biosciences. The Jackson Laboratory (Jax) Model-AD Consortium also developed KI mouse models of apoE  $\epsilon$ 3 and apoE  $\epsilon$ 4 on the C57BL6/J background. Jax is currently developing models of the apoE  $\epsilon$ 3-Ch and apoE  $\epsilon$ 3-Jac variants, as well as inducible apoE reporters and flips of apoE  $\epsilon$ 4 to  $\epsilon$ 3, apoE  $\epsilon$ 3 to  $\epsilon$ 4, and apoE  $\epsilon$ 3 to  $\epsilon$ 2. The Cure Alzheimer's Fund also has a KI model with floxed alleles that allow for conditional deletion.<sup>93</sup> Understanding the limitations of each model is important in deciphering the results from studies and furthering the understanding of the field.

Most phenotypes of Jax and Taconic mouse models are reproducible, but some researchers have not observed increases in SERPINA3 as previously described in the Taconic model. In addition, apoE expression levels are typically higher in Jax mice compared to the

Taconic mice, which may be due to a leftover neomycin cassette in the Taconic model. Other phenotypic changes are limited by age, as these mouse models do not show behavioral differences until 16 months old. However, Jax has observed sex-dependent changes in vascular coupling from fluorodeoxyglucose (FDG) positron emission tomography (PET) by 12 months old.

## 7.2 | Other animal models

Using rats as an apoE model can potentially facilitate greater characterizations in vascular changes and CSF flow. In addition, the Jax Marmo-AD program aims to develop an early-onset AD marmoset model harboring a *PSEN1* mutation and a late-onset AD marmoset model harboring an *ABCA7* mutation. However, nonhuman primate (NHP) models can limit study sample sizes and increase study timelines due to challenges in producing large, aged cohorts of NHPs.<sup>93</sup>

## 8 | THERAPEUTIC STRATEGIES

A range of therapeutic strategies targeting apoE or apoE signaling mechanisms and building on recent technological advances are in various stages of development. These approaches include genetic therapies such as an adeno-associated virus (AAV), antisense oligonucleotides (ASOs), and RNA interference (RNAi); antibodies; and small-molecule drugs. Incorporating refined trial designs and biomarker developments may facilitate meaningful mechanistic insight regardless of trial outcomes.<sup>94</sup>

### 8.1 | RNAi modulation of apoE expression

Using RNAi to silence disease-associated genes on demand is a particularly attractive strategy in the context of neurodegeneration. The entrapment of siRNA compounds in the endolysosomal system creates a slowly released intracellular depot of drug that enables multimonth efficacy. In addition, optimization of chemical scaffolds for DNA delivery (i.e., dianophores) enables the selective delivery of siRNAs to the CNS or other target tissues.<sup>95</sup> The recent development of a novel divalent siRNA dianophore enables potent and sustained modulation of gene expression throughout the CNS of mice as well as larger NHPs, specifically reducing gene expression in the CNS by more than 99% for up to 12 months.<sup>96</sup>

RNAi for apoE  $\epsilon$ 4 is currently under development as a novel therapeutic approach for treating late-onset AD (LOAD). Genetic evidence suggests that in the CNS, apoE  $\epsilon$ 4 promotes neurodegeneration through a toxic gain-of-function mechanism<sup>55,77,97</sup> and that removing apoE completely in a mouse model of tauopathy blocks most tau-mediated neurodegeneration<sup>98</sup> and decreasing apoE  $\epsilon$ 4 in the brain by ~50% using antisense oligonucleotides also significantly reduces tau-mediated neurodegeneration.<sup>99</sup> However, because apoE is critical for systemic cholesterol metabolism, maintaining the systemic function of

apoE is essential.<sup>100-102</sup> These findings constrain the development of RNAi to target apoE  $\epsilon$ 4.

### 8.2 | Correcting endolysosomal dysfunction mediated by apoE $\epsilon$ 4

Converging lines of research implicate dysfunction of the endolysosomal system in aging and AD and suggest that apoE  $\epsilon$ 4 exacerbates these deficits.<sup>14,15,103-111</sup> At a fundamental level, differences between the net charge of the apoE  $\epsilon$ 2, apoE  $\epsilon$ 3, and apoE  $\epsilon$ 4 isoforms confer differences in the isoelectric point (IEP) (i.e., the point at which the surface charge is neutral), which leads to differences in how the isoforms are processed and trafficked by the endolysosomal system.<sup>111</sup>

The match between the IEP of apoE  $\epsilon$ 4 and the pH of early endosomes suggests that apoE  $\epsilon$ 4 may lose solubility in the early endosome leading to increased self-interaction of apoE particles. This, in turn, would be predicted to increase their apparent affinity to clustered apoE receptors within the endosome, hindering the dissociation of the particles from their receptors and thereby impairing normal endosome maturation, recycling, retrograde sorting, and lysosomal degradation. In support of this idea, it has been shown that biochemical and genetic inhibition of a proton leak channel in the early endosome, the Na<sup>+</sup>/H<sup>+</sup> exchanger 6 (NHE6), restores normal apoE and glutamate receptor trafficking through the endosomal compartment.<sup>111</sup> Furthermore, NHE6 disruption equalizes the phenotypes between apoE isoforms and prevents A $\beta$  plaque accumulation in apoE  $\epsilon$ 4 KI mice. Targeting endolysosomal dysfunction thus presents a pharmacologically tractable mechanism that may be fundamental to AD risk. Furthermore, taken together these findings suggest that balancing endolysosomal pH homeostasis may be a possible pathway for AD prevention, not only for the increased risk imposed by apoE  $\epsilon$ 4 but also for other forms of AD.<sup>109</sup>

### 8.3 | Targeting microglial pathways downstream of apoE/TREM2

The Target Enablement to Accelerate Therapy Development for AD (TREAT-AD) initiative has developed a strategy for small-molecule targeting of pathways downstream of apoE and TREM2. Analysis of apoE/TREM2 signaling networks identified inositol polyphosphate-5-phosphatase (INPP5D), also known as SHIP1, to have a significant genetic association with LOAD. INPP5D/SHIP1 is co-expressed with 73 other genes in an AD immune response module, selectively expressed in microglia, and postulated to limit TREM2-mediated microglia activation. Furthermore, studies using the 5xFAD mouse model show that INPP5D/SHIP1 is upregulated as A $\beta$  pathology progresses and that INPP5D/SHIP1 haploinsufficiency normalizes AD-related neuropathology and behavioral deficits, which suggests that inhibition of INPP5D/SHIP1 early in AD would increase TREM2 signaling and microglial protective functions, resulting in reduced rate of disease progression and cognitive decline in AD.<sup>112</sup>



A high-throughput small-molecule screen (HTS) for INPP5D/SHIP1 inhibitors using co-purified INPP5D/SHIP1 identified a lead candidate chemical probe, called compound 23. In vitro studies in primary mouse microglia demonstrated concentration-dependent effects of compound 23 on INPP5D/SHIP1 signaling, a lack of toxicity, and good exposure in brain and plasma. Next steps include additional structure-activity studies and development of a lead clinical compound. Studies are also underway to identify inhibitors of additional neuroinflammation targets and, in parallel, to develop relevant biomarkers for INPP5D/SHIP1 and other identified targets.<sup>113</sup> In addition, a “Target Enablement Package,” available on the [AD Knowledge Portal](#) has been recently released that includes data, methods, and research tools to catalyze further drug discovery efforts by academic and industry researchers.<sup>114</sup>

## 8.4 | Novel interventions targeting apoE-heparan sulfate proteoglycan interactions

Compared to apoE  $\epsilon$ 3, apoE  $\epsilon$ 4 displays a higher affinity to heparan sulfate proteoglycans (HSPGs), whereas apoE  $\epsilon$ 2 and apoE3-Ch display lower binding affinities when used heparin-affinity chromatography to model this interaction.<sup>20</sup> This observation suggests that the protective effect of the apoE3-Ch variant in the Colombian PSEN1 FAD kindred may be mediated by diminished interactions between apoE-Ch and HSPGs. Marino and colleagues generated anti-apoE-HSPG antibodies to mimic the reduced apoE-Ch-HSPG interaction, and characterized them using both biomolecular approaches. Among the antibodies screened for inhibition of binding, the top candidate, 7C11, was tested in vitro and in vivo. Notably, 7C11 reduced the cytotoxicity of apoE4 in vitro and rescued apoE4-induced tau pathology in vivo. Recently, a study using isogenic hiPSC-derived neurons with apoE4 or apoE4-Ch showed that apoE4-Ch reduced HSPG-mediated tau uptake by neurons in culture.<sup>115</sup> Together, these findings suggest that developing novel inhibitors of apoE-HSPGs interactions might lead to effective disease-modifying therapies for AD.<sup>116</sup>

## 8.5 | APOE-targeted epigenome therapy

A global reduction in overall brain APOE levels may have beneficial effects on AD pathogenesis.<sup>99,117-120</sup> Consistently, integrative single nucleus multiomic analysis revealed an overexpression of APOE  $\epsilon$ 3 in specific cellular subtypes in AD brains versus control, as well as more open chromatin in several genomic sites linked to the promoter of the APOE gene, which implies disease-associated cis-regulatory elements.<sup>121</sup> In addition, a new study suggested that apoE  $\epsilon$ 4 drives AD risk through a gain of abnormal function (contrary to the earlier mentioned loss), providing further support that reducing apoE  $\epsilon$ 4 levels is a promising therapeutic strategy.<sup>97</sup> These findings informed the development of an epigenome therapeutic approach for treating AD based on a modification of the CRISPR/Cas9 technology

and delivered by an all-in-one viral vector platform that specifically recognizes the APOE  $\epsilon$ 4 allele and represses and fine-tunes APOE  $\epsilon$ 4 expression. Homozygous APOE  $\epsilon$ 4 and APOE  $\epsilon$ 3 hiPSC-derived cholinergic neurons and organoids showed that expression of the APOE  $\epsilon$ 4 allele, but not the APOE  $\epsilon$ 3 allele, was specifically repressed. In vivo studies following injection into mouse hippocampus found that apoE protein expression was reduced by approximately 70% and that the repression effect exhibited sex-specific differences. This novel epigenome therapy platform offers the opportunity for refinement to develop gene-, allele-, cell type-, and population-specific therapies, thereby advancing strategies for precision medicine in LOAD.<sup>122</sup>

## 8.6 | Phase 1 AAV gene therapy in patients with APOE $\epsilon$ 4 homozygote AD

While APOE  $\epsilon$ 4 is associated with great AD risk and earlier age of onset, emerging evidence suggests that APOE  $\epsilon$ 2 has protective effects against AD. Lexeo Therapeutics has a clinical program that is investigating their AAVrh10 gene therapy candidate, LX1001, in patients with homozygous APOE  $\epsilon$ 4-associated AD. LX1001 is designed to deliver into the CSF and express the protective APOE  $\epsilon$ 2 gene. Administering the APOE  $\epsilon$ 2 gene to APOE  $\epsilon$ 4 homozygous individuals has the potential to address several pathways that are involved in the progression of AD disease.<sup>101,123</sup>

LX1001 is being evaluated in an open-label, dose-escalation Phase 1/2 clinical trial in APOE  $\epsilon$ 4 homozygous individuals who are 50 years of age or older and have mild cognitive impairment (MCI) to moderate dementia and biomarkers consistent with AD. The primary endpoint is safety; additional measures include expression of APOE  $\epsilon$ 2 in the brain, CSF biomarkers (A $\beta$ 42, t-tau, p-tau), A $\beta$  and tau PET scans, quantitative MRI, and cognitive testing. In the first dose cohort in the trial, the study investigators observed a consistent trend toward improvement in AD CSF biomarkers, such as total tau and phosphorylated tau. Investigators have also observed expression of the protective apoE  $\epsilon$ 2 protein in all patients in the first dose cohort with follow-up data. Among all patients, treatment with LX1001 has been well-tolerated with no serious related adverse events reported as of July 2023. Although these initial data are promising, evaluation of the dose-response relationship, effects of higher doses, and inclusion of more patients will provide additional insight into this therapeutic approach and its potential clinical impact (Clinical Trial NCT03634007).

### 8.6.1 | Hormone replacement therapy in women with APOE $\epsilon$ 4

Estrogen decline during the menopausal transition is emerging as a key factor enhancing AD risk in women.<sup>124</sup> Estrogen regulates multiple neurophysiological processes, including cerebrovascular function, BBB integrity, synaptic plasticity, neuroinflammation, and brain energy metabolism. Furthermore, estrogen and progesterone

receptors are expressed in multiple brain regions relevant to cognitive function and AD.<sup>125</sup> Hormone replacement therapy (HRT) has thus been of interest as a strategy to reduce cognitive decline in women, although clinical trial results have been inconsistent thus far.<sup>126-128</sup> A recently published study by Saleh and colleagues investigated whether HRT would have greater cognitive benefits in APOE ε4 compared to non-APOE ε4 women, particularly when introduced early during the menopausal transition. The study found that APOE ε4 women on HRT compared to non-HRT scored higher on delayed memory tests and had larger entorhinal and amygdala volumes. Interestingly, earlier HRT initiation was associated with larger hippocampal volume. Findings thus far emphasize the importance of personalized medicine in AD prevention.<sup>129</sup>

## 9 | CONCLUSION

The neurodegenerative disease and biomedical research communities have made significant progress in understanding the structure, function, associated pathologies, and clinical impact of apoE. The "AAIC advancements: APOE conference," which assembled over 850 participants from 54 countries, helped to facilitate discussions on emerging APOE research, formed collaborative efforts, and provided a forward perspective on the field. The conversations clarified areas in which further study is necessary and worthwhile, including studies on the mechanisms underlying sex-dependent impacts of apoE isoforms, development of improved animal and cell models of AD and other apoE-related diseases, clinical impact of modified apoE proteins (e.g., lipidated), and assessment of therapeutics targeting apoE-related physiological aberrations.

Another major priority for the entire research community is to diversify the patient populations within studies and clinical trials to be representative of the population impacted by APOE-related conditions, particularly AD. To better understand the neural-genetic-environmental interplay underlying apoE-pathology and improve the livelihood of all individuals, research studies must prioritize understanding the intricacies of APOE in various geographical, racial, and ethnic groups that are impacted. These approaches can improve the many ongoing and future APOE-focused research studies and therapeutic trials and enable new discoveries that ultimately help prevent, diagnose, and treat-AD.

## AFFILIATIONS

<sup>1</sup>Alzheimer's Association, Chicago, Illinois, USA

<sup>2</sup>Department of Neurology and Neurological Sciences, Stanford University, Stanford, Palo Alto, California, USA

<sup>3</sup>NeuroGenomics and Informatics Center, Washington University School of Medicine, St. Louis, Missouri, USA

<sup>4</sup>Department of Neurology, Washington University School of Medicine, St. Louis, Missouri, St. Louis, Missouri, USA

<sup>5</sup>Division of Medical Genetics, Department of Medicine, University of Washington, Seattle, Washington, USA

<sup>6</sup>Institute for Public Health Genetics, University of Washington, Seattle, Washington, USA

<sup>7</sup>Departments of Biochemistry & Biophysics and Bioengineering, University of Pennsylvania, Philadelphia, Pennsylvania, USA

<sup>8</sup>Department of Neurology, Hope Center for Neurological Disorders, Knight Alzheimer's Disease Research Center, Washington University School of Medicine, St. Louis, Missouri, USA

<sup>9</sup>Division of Translational Brain Sciences, Department of Neurology, Duke University School of Medicine, Durham, North Carolina, USA

<sup>10</sup>Department of Neurology Hope Center for Neurological Disorders Washington University School of Medicine, St. Louis, Missouri, USA

<sup>11</sup>Denali Therapeutics Inc., San Francisco, California, USA

<sup>12</sup>Ronald M. Loeb Center for Alzheimer's Disease, New York, New York, USA

<sup>13</sup>Department of Genetics & Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, New York, USA

<sup>14</sup>The Ken & Ruth Davee Department of Neurology, Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA

<sup>15</sup>Department of Physiology, University of Kentucky, Lexington, Kentucky, USA

<sup>16</sup>Sanders-Brown Center on Aging, University of Kentucky, Lexington, Kentucky, USA

<sup>17</sup>Department of Medicine, University of Washington, UV Medicine, Seattle, Washington, USA

<sup>18</sup>Center for Translational Neurodegeneration Research, University of Texas Southwestern Medical Center, Dallas, Texas, USA

<sup>19</sup>Gladstone Institute of Neurological Disease, Gladstone Institutes, San Francisco, California, USA

<sup>20</sup>Department of Neurology, University of California, San Francisco, California, USA

<sup>21</sup>Feil Family Brain and Mind Research Institute, Weill Cornell Medicine, New York, New York, USA

<sup>22</sup>Department of Neuroscience, Mayo Clinic Jacksonville, Jacksonville, Florida, USA

<sup>23</sup>Department of Psychiatry, Washington University in St. Louis, St. Louis, Missouri, USA

<sup>24</sup>RNA Therapeutic Institute, UMass Chan Medical School, Worcester, Massachusetts, USA

<sup>25</sup>Department of Functional Genomics, Center for Neurogenomics and Cognitive Research (CNCR), VU University Amsterdam, Amsterdam, USA

<sup>26</sup>Alzheimer Center Amsterdam, Department of Neurology, Amsterdam Neuroscience, Amsterdam University Medical Center, Amsterdam, USA

<sup>27</sup>Stark Neurosciences Research Institute Indiana University School of Medicine, Indianapolis, Indiana, USA

<sup>28</sup>The Tracy Family SILQ Center, Washington University School of Medicine, Indianapolis, Indiana, USA

<sup>29</sup>Department of Neurology and Neurological Sciences, Stanford University, Palo Alto, California, USA

<sup>30</sup>Institut du Cerveau-Paris Brain Institute-ICM, Paris, France

<sup>31</sup>Department of Genetics and Genomic Sciences, Nash Family Department of Neuroscience, Icahn Genomics Institute; Icahn School of Medicine at Mount Sinai, New York, New York, USA

<sup>32</sup>Schepens Eye Research Institute of Mass Eye and Ear and Department of Ophthalmology at Harvard Medical School, Boston, Massachusetts, USA

<sup>33</sup>Sagol School of Neuroscience Tel Aviv University, Tel Aviv, Israel

<sup>34</sup>Department of Biochemistry and Molecular Biophysics, Washington University School of Medicine, St. Louis, Missouri, USA

<sup>35</sup>Department of Neuroscience, University of Kentucky, Lexington, Kentucky, USA

<sup>36</sup>Genetics and Biochemistry Branch, National Institute of Diabetes and Digestive and Kidney Diseases, National Institute of Neurological Disorders and

Stroke, Center for Alzheimer's and Related Dementias (CARD), National Institutes of Health, Maryland, USA

<sup>37</sup>Human Genome and Stem-cell Research Center, Biosciences Institute, University of São Paulo, Rua do Matao, São Paulo, Brazil

<sup>38</sup>Hospital Israelita Albert Einstein, Avenida Albert Einstein, São Paulo, Brazil

<sup>39</sup>Neurochemistry Laboratory, Department of Laboratory Medicine, Vrije Universiteit Amsterdam, Amsterdam UMC, Amsterdam, Netherlands

<sup>40</sup>Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Columbia University Vagelos College of Physicians and Surgeons, New York, New York, USA

<sup>41</sup>Department of Neurology, Columbia University Irving Medical Center, New York, New York, USA

<sup>42</sup>Department of Neuroscience, Columbia University Vagelos College of Physicians and Surgeons, New York, USA

<sup>43</sup>Mayo Clinic Florida, Jacksonville, Florida, USA

<sup>44</sup>Norwich Medical School, University of East Anglia, UK Clinical and Chemical Pathology, Norfolk, UK

<sup>45</sup>Faculty of Medicine, Alexandria University, Alexandria Governorate, Egypt

<sup>46</sup>Lexeo Therapeutics, New York, New York, USA

<sup>47</sup>University of Southern California, Los Angeles, California, USA

<sup>48</sup>Washington University in Saint Louis, St. Louis, Missouri, USA, St. Louis, Missouri, USA

<sup>49</sup>Department of Pharmacology, Physiology & Biophysics, Chobanian and Avedisian School of Medicine, Boston University, Boston, Massachusetts, USA

<sup>50</sup>Bioinformatics Program, Faculty of Computing & Data Sciences, Boston University, Boston, Massachusetts, USA

<sup>51</sup>Center for Cognitive Neurology, Department of Neurology, New York University Grossman School of Medicine, New York, New York, USA

<sup>52</sup>Picower Institute for Learning and Memory, Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

<sup>53</sup>Department of Neurology, Center for Neurodegeneration and Experimental Therapeutics, Medical Scientist Training Program, University of Alabama at Birmingham, Birmingham, Alabama, USA

<sup>54</sup>Mayo Clinic Rochester, Rochester, Minnesota, USA

<sup>55</sup>Djavad Mowafaghian Centre for Brain Health Department of Pathology and Laboratory Medicine International Collaboration on Repair Discoveries School of Biomedical Engineering University of British Columbia, Vancouver, Canada

<sup>56</sup>FBRI, Massachusetts, USA

<sup>57</sup>Department of Neurology, Keck School of Medicine, University of Southern California, Los Angeles, California, USA

<sup>58</sup>Division of Life Science, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong

<sup>59</sup>Department of Genetics & Genomic Sciences, Ronald M. Loeb Center for Alzheimer's disease, Icahn School of Medicine at Mount Sinai, New York, New York, USA

## ACKNOWLEDGMENTS

We thank and acknowledge all conference organizers, speakers and session chairs for your tremendous contributions to this conference. M. E. Belloy received funding for this work from the NIH (K99AG075238) and Alzheimer's Association (AARF-20-683984). E. E. Blue was funded by the National Institutes of Health/National Institute on Aging grant number R01AG059737 and is related to her work funded by grant number U01AG058589. G. R. Bowman was funded by NIH grants U19AG069701 and RF1AG067194. X.

Chen received the 2023 Early Career Achievement Award from Alzheimer's Association. O. Chiba-Falek was funded in part by the National Institutes of Health/National Institute on Aging (NIH/NIA) [R41 AG077992, R01 AG057522 to OC-F]. A. A. Davis has received research funding that is focused on APOE but which did not directly support this manuscript includes: NIH K08NS101118 DoD W81XWH2010934 NIH RF1AG083753. F. Garretti is funded by F32AG08483. D. Gate is funded by Alzheimer's Association 23AARG-1026607. L. R. Golden is funded by NIH T32 AG078110 "Training in Translational Research in Alzheimer's and Related Dementias (TRIAD)". J. Heinecke is funded by R01HL149685, R01AG061186. J. Herz acknowledges funding by the NIH, the Brightfocus Foundation and the Alzheimer's Association. Y. Huang is funded by NIH/NIA P01AG073082. C. Iadecola is partially supported by NS126467 ApoE4, neurovascular injury and cognitive impairment. L. A. Johnson is supported by the National Institute on Aging (R01AG060056, R01AG062550, R01AG080589, and R01AG081421), the National Institute of General Medical Sciences (NIGMS) Center of Biomedical Research Excellence in CNS Metabolism P20GM148326, and the Alzheimer's Association. T. Kanekiyo is supported by NIH/NIA U19AG069701. C. M. Karch is funded by NIH-NIA U19 AG069701 Title: Biology and pathobiology of apoE in aging and Alzheimer's Disease. A. Khvorova is funded by GM131839-03. B. T. Lamb is funded by U54 AG065181, U54AG054345, RF1 AG074566. P. E. Lawler is funded by Cure Alzheimer's Fund. Y. Le Guen is supported by the European Union's Horizon 2020 Research and Innovation Program under the Marie Skłodowska-Curie Actions grant 890650 (Dr Le Guen). A. Litvinchuk is funded by BrightFocus Foundation Postdoctoral Fellowship A2022010F (A.L.). E. Marcora is funded by NIH R56AG081417 NIH U19AG069701 BrightFocus Foundation A2017458S. C. Marino is funded by the Edward N. & Della L. Thome Memorial Foundation, from the Robert J. Kleberg, Jr. and Helen C. Kleberg Foundation, from the Remondi Family Foundation, from the US National Institute of Neurological Disorders and Stroke and National Institute on Aging co-funded grants UH3 NS100121 and RF1 NS110048 and from Good Ventures and Open Philanthropy to J.F.A.-V. Grants from the US National Institutes of Health (NIH) Office of the Director grant DP5 OD019833 and US National Institute on Aging grants R01 AG054671, and RF1AG077627, the Massachusetts General Hospital Executive Committee (ECOR) on Research (MGH Research Scholar Award), and grants from the Alzheimer's Association to Y.T.Q. NEI-NIH P30EY003790. D. M. Michaelson received a donation from the Eichenbaum Foundation. J. J. Miller is funded by NIH training grant T32AG05851804. J. M. Morganti is funded by 5R01AG070830, 3RF1NS118558. P. S. Narayan is supported by NIH-NIDDK Intramural Research Program. M. S. Naslavsky is supported by FAPESP13/08028-1 Alzheimer's Association/NIH K24AG053435 CNPq 304746/2022-3. K. V. Ramachandran is supported by NIH Director's Early Independence Award 7DP5OD028133, Department of Defense CDMRP award W81XWH-21-1-0093, Fidelity Biomedical Research Initiative. A. C. Raulin is funded by BrightFocus foundation postdoctoral fellowship A2021015F. R. N. M. Saleh is funded by Medical Research Council (MRC, UK), NuBrain Consortium (MR/T001852/1). F. Shue

is supported by U19AG069701. A. Soranno is funded by NIH NIA U19AG069701 (Project 1, A.S.) NIH NIA R01AG062837 (to A.S.). M. R. Strickland has funding provided by the NIH T32 Fellowship (T32AG058518, MRS) and NSF GRFP Fellowship (DGE-1745038, MRS). J. TCW is supported by NIH R01AG082362, R01AG083941, K01AG062683, U19AG069701, R56AG078733. M. Thierry is supported by NIH/NIA grants: P30AG066512 and P01AG060882.R. Tuckey is supported by Alzheimer's Drug Discovery Foundation and NIH grants R01 AG081228, R01 AG068395, and T32 GM008361. J. Ulrich is funded by 1U19AG069701. N. Wang is funded by a NIA U19 grant. C. L. Wellington is supported by Cure Alzheimer Fund and BrightFocus Foundation. H. N. Yassine holds the Kenneth and Bette Volk Endowed Chair of Neurology. HNY is supported by RF1AG076124, RF1AG078362, R01AG067063, R01AG054434, R01AG055770, R21AG056518, and P30AG066530 from the National Institute on Aging, GC-201711-2014197 from the Alzheimer's Drug Discovery Foundation (ADDF), and generous donations from the Vranos and Tiny Foundations and from Ms. Lynne Nauss. N. Zhao is supported by NIH grants U19AG069701, RF1AG046205, R01AG66395, and U54NS110435; a grant from BrightFocus Foundation; and a grant from Cure Alzheimer's Fund. G Bu is supported by Cure Alzheimer's Fund. A. M. Goate receives funding from NIH (U19AG069701, U01AG058635) and the JPB FOundation. D. M. Holtzman is funded by NIH 1U19AG069701 (DMH).

#### CONFLICT OF INTEREST STATEMENT

C. M. Kloske, M. C. Carrillo, S. Mahinrad, and C. E. Sexton are all full time employees of the Alzheimer's Association. M. Belloy has nothing to disclose. E. E. Blue was funded by the National Institutes of Health/National Institute on Aging grant number R01AG059737 and is related to her work funded by grant number U01AG058589. Dr. Blue is a member of the Board of Directors for the International Genetic Epidemiology Society. G. R. Bowman holds equity in Decrypt Biomedicine. X. Chen has nothing to disclose. O. Chiba-Falek is a co-inventor of the related IP. Duke University filed patent applications for the technology developed in this study. CLAIRGene has an exclusive, worldwide option agreement from Duke for the related patent portfolio for all fields of use. Dr. Chiba-Falek is a Co-Founder at CLAIRGene, LLC. A. Davis has nothing to disclose. G. Di Paolo is a full-time employee and shareholder of Denali Therapeutics Inc. F. Garretti has nothing to disclose. D. Gate has nothing to disclose. L. M. Golden has nothing to disclose. J. Heinecke has nothing to disclose. J. Herz is a cofounder of Reelin Therapeutics. Y. Huang has nothing to disclose. C. Iadecola has nothing to disclose. L. A. Johnson has nothing to disclose. T. Kanekiyo has nothing to disclose. C. Karch has nothing to disclose. A. Khvorova is a Co-Founder of Atalanta. S. Koppes-den Hertog has nothing to disclose. B. Lamb Co-Founded of Monument Biosciences. P. E. Lawler has nothing to disclose. Y. Le Guen has nothing to disclose. A. Litvinchuk has nothing to disclose. C. Liu has nothing to disclose. E. Marcora has nothing to disclose. C. Marino has nothing to disclose. D. M. Michaelson has nothing to disclose. J. J. Miller has nothing to disclose. J. M. Morganti has nothing to disclose. P. S. Narayan has nothing to disclose. M. S. Naslavsky has nothing to disclose. M. Oosthoek has nothing to

disclose. K. V. Ramachandran has nothing to disclose. A Ramakrishnan has nothing to disclose. A. Raulin has nothing to disclose. A. Robert has nothing to disclose. R. N. M. Saleh has nothing to disclose. N. Shah is an employee of Lexeo Therapeutics. F. Shue has nothing to disclose. I. J. Sible has nothing to disclose. A. Soranno has nothing to disclose. M. R. Strickland has nothing to disclose. J. TCW has nothing to disclose. M. Thierry has nothing to disclose. L. Tsai has nothing to disclose. R. A. Tuckey has nothing to disclose. J. D. Ulrich has nothing to disclose. R. van der Kant has nothing to disclose. N. Wang has nothing to disclose. C. L. Wellington has nothing to disclose. S. C. Weninger has nothing to disclose. H. N. Yassine has nothing to disclose. N. Zhao has nothing to disclose. G. Bu is the Editor-in-Chief of Molecular Neurodegeneration and a consultant for SciNeuro Pharmaceutical. A. M. Goate serves on the scientific advisory boards for Genentech and Muna Therapeutics. D. M. Holtzman is an inventor on a patent licensed by Washington University to NextCure on the therapeutic use of anti-APOE antibodies. Pub. No. US 2022/0411485 A1 Pub date Dec 29, 2022. Title of patent: ANTI-APOE ANTIBODIES. D.M.H. co-founded and is on the scientific advisory board of C2N Diagnostics. D.M.H. is on the scientific advisory board of Denali, Genentech, and Cajal Neuroscience and consults for Asteroid. Author disclosures are available in the [supporting information](#).

#### REFERENCES

- Havel RJ, Kane JP, Kashyap ML. Interchange of apolipoproteins between chylomicrons and high density lipoproteins during alimentary lipemia in man. *J Clin Invest*. 1973;52:32-38. doi:10.1172/JCI107171
- Utermann G, Jaeschke M, Menzel J. Familial hyperlipoproteinemia type III: deficiency of a specific apolipoprotein (APO E-III) in the very-low-density lipoproteins. *FEBS Lett*. 1975;56:352-355. doi:10.1016/0014-5793(75)81125-2
- Zannis VI, Breslow JL, Utermann G, et al. Proposed nomenclature of apoE isoproteins, apoE genotypes, and phenotypes. *J Lipid Res*. 1982;23:911-914.
- Mahley RW, Rall SC. Apolipoprotein E: far more than a lipid transport protein. *Annu Rev Genomics Hum Genet*. 2000;1:507-537. doi:10.1146/annurev.genom.1.1.507
- Saunders AM, Strittmatter WJ, Schmechel D, et al. Association of apolipoprotein E allele  $\epsilon 4$  with late-onset familial and sporadic Alzheimer's disease. *Neurology*. 1993;43:1467-1467. doi:10.1212/WNL.43.8.1467
- Kim J, Basak JM, Holtzman DM. The role of apolipoprotein E in Alzheimer's disease. *Neuron*. 2009;63:287-303. doi:10.1016/j.neuron.2009.06.026
- Corder EH, Saunders AM, Strittmatter WJ, et al. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science*. 1993;261:921-923. doi:10.1126/science.8346443
- Strittmatter WJ, Saunders AM, Schmechel D, et al. Apolipoprotein E: high-avidity binding to beta-amyloid and increased frequency of type 4 allele in late-onset familial Alzheimer disease. *Proc Natl Acad Sci*. 1993;90:1977-1981. doi:10.1073/pnas.90.5.1977
- Corder EH, Saunders AM, Risch NJ, et al. Protective effect of apolipoprotein E type 2 allele for late onset Alzheimer disease. *Nat Genet*. 1994;7:180-184. doi:10.1038/ng0694-180
- Yamazaki Y, Zhao N, Caulfield TR, Liu C-C, Bu G. Apolipoprotein E and Alzheimer disease: pathobiology and targeting strategies. *Nat Rev Neurol*. 2019;15:501-518. doi:10.1038/s41582-019-0228-7

11. Bellenguez C, Küçükali F, Jansen IE, et al. New insights into the genetic etiology of Alzheimer's disease and related dementias. *Nat Genet.* 2022;54:412-436. doi:10.1038/s41588-022-01024-z
12. Deming Y, Li Z, Kapoor M, Harari O, et al. Genome-wide association study identifies four novel loci associated with Alzheimer's endophenotypes and disease modifiers. *Acta Neuropathol (Berl).* 2017;133:839-856. doi:10.1007/s00401-017-1685-y
13. Belloy ME, Napolioni V, Greicius MD. A quarter century of APOE and Alzheimer's disease: progress to date and the path forward. *Neuron.* 2019;101:820-838. doi:10.1016/j.neuron.2019.01.056
14. Cataldo AM, Peterhoff CM, Troncoso JC, Gomez-Isla T, Hyman BT, Nixon RA. Endocytic pathway abnormalities precede amyloid beta deposition in sporadic Alzheimer's disease and Down syndrome: differential effects of APOE genotype and presenilin mutations. *Am J Pathol.* 2000;157:277-286. doi:10.1016/s0002-9440(10)64538-5
15. Narayan P, Sienski G, Bonner JM, et al. PICALM rescues endocytic defects caused by the Alzheimer's disease risk factor APOE4. *Cell Rep.* 2020;33:108224. doi:10.1016/j.celrep.2020.108224
16. Miyashita A, Koike A, Jun G, et al. SORL1 is genetically associated with late-onset Alzheimer's disease in Japanese, Koreans and Caucasians. *PLoS ONE.* 2013;8:e58618. doi:10.1371/journal.pone.0058618
17. Romero-Molina C, Garrett F, Andrews SJ, Marcora E, Goate AM. Microglial efferocytosis: diving into the Alzheimer's disease gene pool. *Neuron.* 2022;110:3513-3533. doi:10.1016/j.neuron.2022.10.015
18. Wu HM, Goate AM, O'Reilly PF. Heterogeneous effects of genetic risk for Alzheimer's disease on the phenome. *Transl Psychiatry.* 2021;11:406. doi:10.1038/s41398-021-01518-0
19. Futamura M, Dhanasekaran P, Handa T, Phillips MC, Lund-Katz S, Saito H. Two-step mechanism of binding of apolipoprotein E to heparin. *J Biol Chem.* 2005;280:5414-5422. doi:10.1074/jbc.M411719200
20. Arboleda-Velasquez JF, Lopera F, O'Hare M, et al. Resistance to autosomal dominant Alzheimer's disease in an APOE3 Christchurch homozygote: a case report. *Nat Med.* 2019;25:1680-1683. doi:10.1038/s41591-019-0611-3
21. Zalocusky KA, Nelson MR, Huang Y. An Alzheimer's-disease-protective APOE mutation. *Nat Med.* 2019;25:1648-1649. doi:10.1038/s41591-019-0634-9
22. Acosta-Baena N, Sepulveda-Falla D, Lopera-Gómez CM, et al. Pre-dementia clinical stages in presenilin 1 E280A familial early-onset Alzheimer's disease: a retrospective cohort study. *Lancet Neurol.* 2011;10:213-220. doi:10.1016/S1474-4422(10)70323-9
23. Le Guen Y, Raulin A-C, Logue MW, et al. Association of African Ancestry-Specific APOE missense variant R145C with risk of Alzheimer disease. *JAMA.* 2023;329:551. doi:10.1001/jama.2023.0268
24. Le Guen Y, Belloy ME, Grenier-Boley B, et al. Association of rare APOE missense variants V236E and R251G with risk of Alzheimer disease. *JAMA Neurol.* 2022;79:652. doi:10.1001/jamaneurol.2022.1166
25. Karczewski KJ, Francioli LC, Tiao G, et al. The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature.* 2020;581:434-443. doi:10.1038/s41586-020-2308-7
26. Kobayashi S, Tateno M, Park TW, et al. Apolipoprotein E4 frequencies in a Japanese population with Alzheimer's disease and dementia with Lewy bodies. *PLoS One.* 2011;6:e18569. doi:10.1371/journal.pone.0018569
27. Belloy ME, Andrews SJ, Le Guen Y, et al. APOE genotype and Alzheimer disease risk across age, sex, and population ancestry. *JAMA Neurol.* 2023;80:1284. doi:10.1001/jamaneurol.2023.3599
28. Corbo RM, Scacchi R. Apolipoprotein E (APOE) allele distribution in the world. Is APOE\*4 a "thrifty" allele? *Ann Hum Genet.* 1999;63:301-310. doi:10.1046/j.1469-1809.1999.6340301.x
29. Farrer LA, Cupples LA, Haines JL, et al. Effects of age, sex, and ethnicity on the association between apolipoprotein E genotype and Alzheimer disease. A meta-analysis. APOE and Alzheimer Disease Meta Analysis Consortium. *JAMA.* 1997;278:1349-1356.
30. Dumitrescu L, Mayeda ER, Sharman K, Moore AM, Hohman TJ. Sex differences in the genetic architecture of Alzheimer's disease. *Curr Genet Med Rep.* 2019;7:13-21. doi:10.1007/s40142-019-0157-1
31. Blue EE, Horimoto ARVR, Mukherjee S, Wijsman EM, Thornton TA. Local ancestry at APOE modifies Alzheimer's disease risk in Caribbean Hispanics. *Alzheimers Dement J Alzheimers Assoc.* 2019;15:1524-1532. doi:10.1016/j.jalz.2019.07.016
32. Rajabli F, Feliciano BE, Celis K, et al. Ancestral origin of ApoE ε4 Alzheimer disease risk in Puerto Rican and African American populations. *PLoS Genet.* 2018;14:e1007791. doi:10.1371/journal.pgen.1007791
33. Raulin A-C, Martens YA, Bu G. Lipoproteins in the central nervous system: from biology to pathobiology. *Annu Rev Biochem.* 2022;91:731-759. doi:10.1146/annurev-biochem-032620-104801
34. Mahley RW, Weisgraber KH, Huang Y. Apolipoprotein E: structure determines function, from atherosclerosis to Alzheimer's disease to AIDS. *J Lipid Res.* 2009;50:S183-188. doi:10.1194/jlr.R800069-JLR200
35. Fagan AM, Holtzman DM, Munson G, et al. Unique lipoproteins secreted by primary astrocytes from wild type, apoE (-/-), and human apoE transgenic mice. *J Biol Chem.* 1999;274:30001-30007. doi:10.1074/jbc.274.42.30001
36. LaDu MJ, Gilligan SM, Lukens JR, et al. Nascent astrocyte particles differ from lipoproteins in CSF. *J Neurochem.* 1998;70:2070-2081. doi:10.1046/j.1471-4159.1998.70052070.x
37. DeMattos RB, Brendza RP, Heuser JE, et al. Purification and characterization of astrocyte-secreted apolipoprotein E and J-containing lipoproteins from wild-type and human apoE transgenic mice. *Neurochem Int.* 2001;39:415-425. doi:10.1016/s0197-0186(01)00049-3
38. Krasemann S, Madore C, Cialic R, et al. The TREM2-APOE pathway drives the transcriptional phenotype of dysfunctional microglia in neurodegenerative diseases. *Immunity.* 2017;47:566-581. doi:10.1016/j.immuni.2017.08.008.e9.
39. Keren-Shaul H, Spinrad A, Weiner A, et al. A unique microglia type associated with restricting development of Alzheimer's disease. *Cell.* 2017;169:1276-1290. doi:10.1016/j.cell.2017.05.018.e17.
40. Stuchell-Brereton MD, Zimmerman MI, Miller JJ, et al. Apolipoprotein E4 has extensive conformational heterogeneity in lipid-free and lipid-bound forms. *Proc Natl Acad Sci.* 2023;120:e2215371120. doi:10.1073/pnas.2215371120
41. Lu B, Morrow JA, Weisgraber KH. Conformational reorganization of the four-helix bundle of human apolipoprotein E in binding to phospholipid. *J Biol Chem.* 2000;275:20775-20781. doi:10.1074/jbc.M003508200
42. Fisher CA, Ryan RO. Lipid binding-induced conformational changes in the N-terminal domain of human apolipoprotein E. *J Lipid Res.* 1999;40:93-99.
43. Gupta V, Narayanaswami V, Budamagunta MS, Yamamoto T, Voss JC, Ryan RO. Lipid-induced extension of apolipoprotein E helix 4 correlates with low density lipoprotein receptor binding ability. *J Biol Chem.* 2006;281:39294-39299. doi:10.1074/jbc.M608085200
44. Yamamoto T, Choi HW, Ryan RO. Apolipoprotein E isoform-specific binding to the low-density lipoprotein receptor. *Anal Biochem.* 2008;372:222-226. doi:10.1016/j.ab.2007.09.005
45. Garai K, Mustafi SM, Baban B, Frieden C. Structural differences between apolipoprotein E3 and E4 as measured by <sup>19</sup>F NMR. *Protein Sci.* 2010;19:66-74. doi:10.1002/pro.283
46. Dong LM, Wilson C, Wardell MR, et al. Human apolipoprotein E. Role of arginine 61 in mediating the lipoprotein preferences of the E3 and E4 isoforms. *J Biol Chem.* 1994;269:22358-22365.

47. Wilson C, Mau T, Weisgraber KH, Wardell MR, Mahley RW, Agard DA. Salt bridge relay triggers defective LDL receptor binding by a mutant apolipoprotein. *Structure*. 1994;2:713-718. doi:10.1016/S0969-2126(00)00072-1
48. Chen J, Li Q, Wang J. Topology of human apolipoprotein E3 uniquely regulates its diverse biological functions. *Proc Natl Acad Sci U S A*. 2011;108:14813-14818. doi:10.1073/pnas.1106420108
49. Liu C-C, Murray ME, Li X, et al. APOE3 -Jacksonville (V236E) variant reduces self-aggregation and risk of dementia. *Sci Transl Med*. 2021;13:eabc9375. doi:10.1126/scitranslmed.abc9375
50. Ji ZS, Fazio S, Mahley RW. Variable heparan sulfate proteoglycan binding of apolipoprotein E variants may modulate the expression of type III hyperlipoproteinemia. *J Biol Chem*. 1994;269:13421-13428.
51. Dong L-M, Parkin S, Trakhanov SD, et al. Novel mechanism for defective receptor binding of apolipoprotein E2 in type III hyperlipoproteinemia. *Nat Struct Mol Biol*. 1996;3:718-722. doi:10.1038/nsb0896-718
52. Lawler PE, Bollinger JG, Schindler SE, et al. Apolipoprotein E O-glycosylation is associated with amyloid plaques and APOE genotype. *Anal Biochem*. 2023;672:115156. doi:10.1016/j.ab.2023.115156
53. Koldamova R, Fitz NF, Lefterov I. ATP-binding cassette transporter A1: from metabolism to neurodegeneration. *Neurobiol Dis*. 2014;72:13-21. doi:10.1016/j.nbd.2014.05.007. Pt A.
54. Martens YA, Zhao N, Liu C-C, et al. ApoE Cascade Hypothesis in the pathogenesis of Alzheimer's disease and related dementias. *Neuron*. 2022;110:1304-1317. doi:10.1016/j.neuron.2022.03.004
55. Liu C-C, Kanekiyo T, Xu H, Bu G. Apolipoprotein E and Alzheimer disease: risk, mechanisms and therapy. *Nat Rev Neurol*. 2013;9:106-118. doi:10.1038/nrneurol.2012.263
56. Arnold M, Nho K, Kueider-Paisley A, et al. Sex and APOE  $\epsilon$ 4 genotype modify the Alzheimer's disease serum metabolome. *Nat Commun*. 2020;11:1148. doi:10.1038/s41467-020-14959-w
57. Montagne A, Nation DA, Sagare AP, et al. APOE4 leads to blood-brain barrier dysfunction predicting cognitive decline. *Nature*. 2020;581:71-76. doi:10.1038/s41586-020-2247-3
58. Serrano-Pozo A, Li Z, Noori A, et al. Effect of APOE alleles on the glial transcriptome in normal aging and Alzheimer's disease. *Nat Aging*. 2021;1:919-931. doi:10.1038/s43587-021-00123-6
59. Area-Gomez E, Larrea D, Pera M, et al. APOE4 is associated with differential regional vulnerability to bioenergetic deficits in aged APOE mice. *Sci Rep*. 2020;10:4277. doi:10.1038/s41598-020-61142-8
60. Farmer BC, Williams HC, Devanney NA, et al. APOE4 lowers energy expenditure in females and impairs glucose oxidation by increasing flux through aerobic glycolysis. *Mol Neurodegener*. 2021;16:62. doi:10.1186/s13024-021-00483-y
61. Keeney JT-R, Ibrahim S, Zhao L. Human ApoE isoforms differentially modulate glucose and amyloid metabolic pathways in female brain: evidence of the mechanism of neuroprotection by apoE2 and implications for Alzheimer's disease prevention and early intervention. *J Alzheimers Dis*. 2015;48:411-424. doi:10.3233/JAD-150348
62. Qi G, Mi Y, Shi X, Gu H, Brinton RD, Yin F. ApoE4 impairs neuron-astrocyte coupling of fatty acid metabolism. *Cell Rep*. 2021;34:108572. doi:10.1016/j.celrep.2020.108572
63. Reiman EM, Caselli RJ, Yun LS, et al. Preclinical evidence of Alzheimer's disease in persons homozygous for the  $\epsilon$ 4 allele for apolipoprotein E. *N Engl J Med*. 1996;334:752-758. doi:10.1056/NEJM199603213341202
64. Kloske CM, Wilcock DM. The important interface between apolipoprotein E and neuroinflammation in Alzheimer's disease. *Front Immunol*. 2020;11:754. doi:10.3389/fimmu.2020.00754
65. Liu C-C, Wang N, Chen Y, et al. Cell-autonomous effects of APOE4 in restricting microglial response in brain homeostasis and Alzheimer's disease. *Nat Immunol*. 2023;24:1854-1866. doi:10.1038/s41590-023-01640-9
66. Wolfe CM, Fitz NF, Nam KN, Lefterov I, Koldamova R. The role of APOE and TREM2 in Alzheimer's disease-current understanding and perspectives. *Int J Mol Sci*. 2018;20:81. doi:10.3390/ijms20010081
67. Gratuze M, Schlachetzki JCM, D'Oliveira Albanus R, et al. TREM2-independent microgliosis promotes tau-mediated neurodegeneration in the presence of ApoE4. *Neuron*. 2023;111:202-219. doi:10.1016/j.neuron.2022.10.022. e7.
68. Blanchard JW, Akay LA, Davila-Velderrain J, et al. APOE4 impairs myelination via cholesterol dysregulation in oligodendrocytes. *Nature*. 2022;611:769-779. doi:10.1038/s41586-022-05439-w
69. Wang N, Wang M, Jeevaratnam S, et al. Opposing effects of apoE2 and apoE4 on microglial activation and lipid metabolism in response to demyelination. *Mol Neurodegener*. 2022;17:75. doi:10.1186/s13024-022-00577-1
70. Chen X, Firulyova M, Manis M, et al. Microglia-mediated T cell infiltration drives neurodegeneration in tauopathy. *Nature*. 2023;615:668-677. doi:10.1038/s41586-023-05788-0
71. Chen X, Holtzman DM. Emerging roles of innate and adaptive immunity in Alzheimer's disease. *Immunity*. 2022;55:2236-2254. doi:10.1016/j.immuni.2022.10.016
72. Lee S, Devanney NA, Golden LR, et al. APOE modulates microglial immunometabolism in response to age, amyloid pathology, and inflammatory challenge. *Cell Rep*. 2023;42:112196. doi:10.1016/j.celrep.2023.112196
73. Chen W-T, Lu A, Craessaerts K, et al. Spatial transcriptomics and in situ sequencing to study Alzheimer's disease. *Cell*. 2020;182:976-991. doi:10.1016/j.cell.2020.06.038. e19.
74. Yang LG, March ZM, Stephenson RA, Narayan PS. Apolipoprotein E in lipid metabolism and neurodegenerative disease. *Trends Endocrinol Metab TEM*. 2023;34:430-445. doi:10.1016/j.tem.2023.05.002
75. Wang S, Li B, Solomon V, et al. Calcium-dependent cytosolic phospholipase A2 activation is implicated in neuroinflammation and oxidative stress associated with ApoE4. *Mol Neurodegener*. 2022;17:42. doi:10.1186/s13024-022-00549-5
76. Sanchez-Mejia RO, Newman JW, Toh S, et al. Phospholipase A2 reduction ameliorates cognitive deficits in a mouse model of Alzheimer's disease. *Nat Neurosci*. 2008;11:1311-1318. doi:10.1038/nn.2213
77. Kim J, Jiang H, Park S, et al. Haploinsufficiency of human APOE reduces amyloid deposition in a mouse model of amyloid- $\beta$  amyloidosis. *J Neurosci*. 2011;31:18007-18012. doi:10.1523/JNEUROSCI.3773-11.2011
78. Feringa FM, van der Kant R. Cholesterol and Alzheimer's disease: from risk genes to pathological effects. *Front Aging Neurosci*. 2021;13:690372. doi:10.3389/fnagi.2021.690372
79. Andreone BJ, Przybyla L, Llapashtica C, et al. Alzheimer's-associated PLC $\gamma$ 2 is a signaling node required for both TREM2 function and the inflammatory response in human microglia. *Nat Neurosci*. 2020;23:927-938. doi:10.1038/s41593-020-0650-6
80. Chan RB, Oliveira TG, Cortes EP, et al. Comparative lipidomic analysis of mouse and human brain with Alzheimer disease. *J Biol Chem*. 2012;287:2678-2688. doi:10.1074/jbc.M111.274142
81. Nugent AA, Lin K, Van Lengerich B, et al. TREM2 regulates microglial cholesterol metabolism upon chronic phagocytic challenge. *Neuron*. 2020;105:837-854. doi:10.1016/j.neuron.2019.12.007. e9.
82. Van Der Kant R, Langness VF, Herrera CM, et al. Cholesterol metabolism is a druggable axis that independently regulates tau and amyloid- $\beta$  in iPSC-derived Alzheimer's disease neurons. *Cell Stem Cell*. 2019;24:363-375. doi:10.1016/j.stem.2018.12.013. e9.
83. Litvinchuk A, Suh JH, Guo JL, et al. Amelioration of Tau and ApoE4-linked glial lipid accumulation and neurodegeneration with an LXR agonist. *Neuron*. 2024;112(3):384-403.e8. doi:10.1016/j.neuron.2023.10.023

84. Tcw J, Qian L, Pipalia NH, et al. Cholesterol and matrisome pathways dysregulated in astrocytes and microglia. *Cell*. 2022;185:2213-2233. doi:10.1016/j.cell.2022.05.017. e25.
85. Sienski G, Narayan P, Bonner JM, et al. APOE4 disrupts intracellular lipid homeostasis in human iPSC-derived glia. *Sci Transl Med*. 2021;13:eaa4564. doi:10.1126/scitranslmed.aaz4564
86. Victor MB, Leary N, Luna X, et al. Lipid accumulation induced by APOE4 impairs microglial surveillance of neuronal-network activity. *Cell Stem Cell*. 2022;29:1197-1212. doi:10.1016/j.stem.2022.07.005
87. Marschallinger J, Iram T, Zardeneta M, et al. Lipid-droplet-accumulating microglia represent a dysfunctional and proinflammatory state in the aging brain. *Nat Neurosci*. 2020;23:194-208. doi:10.1038/s41593-019-0566-1
88. Lefterov I, Wolfe CM, Fitz NF, et al. APOE2 orchestrated differences in transcriptomic and lipidomic profiles of postmortem AD brain. *Alzheimers Res Ther*. 2019;11:113. doi:10.1186/s13195-019-0558-0
89. Koizumi K, Hattori Y, Ahn SJ, et al. Apoε4 disrupts neurovascular regulation and undermines white matter integrity and cognitive function. *Nat Commun*. 2018;9:3816. doi:10.1038/s41467-018-06301-2
90. Oue H, Yamazaki Y, Qiao W, et al. LRP1 in vascular mural cells modulates cerebrovascular integrity and function in the presence of APOE4. *JCI Insight*. 2023;8:e163822. doi:10.1172/jci.insight.163822
91. Iadecola C, Anfray A, Schaeffer S, et al. Cell autonomous role of border associated macrophages in ApoE4 neurovascular dysfunction and susceptibility to white matter injury. *Res Sq*. 2023. doi:10.21203/rs.3.rs-3222611/v1
92. Sibley IJ, Nation DA. Blood pressure variability and cognitive decline: a post hoc analysis of the SPRINT MIND trial. *Am J Hypertens*. 2023;36:168-175. doi:10.1093/ajh/hpac128
93. Oblak AL, Forner S, Territo PR, et al. Model organism development and evaluation for late-onset Alzheimer's disease: mODEL-AD. *Alzheimers Dement Transl Res Clin Interv*. 2020;6:e12110. doi:10.1002/trc2.12110
94. Cummings J, Zhou Y, Lee G, Zhong K, Fonseca J, Cheng F. Alzheimer's disease drug development pipeline: 2023. *Alzheimers Dement Transl Res Clin Interv*. 2023;9:e12385. doi:10.1002/trc2.12385
95. Khvorova A, Watts JK. The chemical evolution of oligonucleotide therapies of clinical utility. *Nat Biotechnol*. 2017;35:238-248. doi:10.1038/nbt.3765
96. Alterman JF, Godinho BMDC, Hassler MR, et al. A divalent siRNA chemical scaffold for potent and sustained modulation of gene expression throughout the central nervous system. *Nat Biotechnol*. 2019;37:884-894. doi:10.1038/s41587-019-0205-0
97. Chemparathy A, Guen YL, Chen S, et al. APOE loss-of-function variants: compatible with longevity and associated with resistance to Alzheimer's Disease pathology. *medRxiv*. 2023;2023.07.20.23292771. doi:10.1101/2023.07.20.23292771 medRxiv
98. Shi Y, Yamada K, Liddelow SA, et al. ApoE4 markedly exacerbates tau-mediated neurodegeneration in a mouse model of tauopathy. *Nature*. 2017;549:523-527. doi:10.1038/nature24016
99. Litvinchuk A, Huynh T-PV, Shi Y, et al. Apolipoprotein E4 reduction with antisense oligonucleotides decreases neurodegeneration in a tauopathy model. *Ann Neurol*. 2021;89:952-966. doi:10.1002/ana.26043
100. Xu Q, Bernardo A, Walker D, Kanegawa T, Mahley RW, Huang Y. Profile and regulation of apolipoprotein E (ApoE) expression in the CNS in mice with targeting of green fluorescent protein gene to the ApoE locus. *J Neurosci Off J Soc Neurosci*. 2006;26:4985-4994. doi:10.1523/JNEUROSCI.5476-05.2006
101. Zhao L, Gottesdiener AJ, Parmar M, et al. Intracerebral adeno-associated virus gene delivery of apolipoprotein E2 markedly reduces brain amyloid pathology in Alzheimer's disease mouse models. *Neurobiol Aging*. 2016;44:159-172. doi:10.1016/j.neurobiolaging.2016.04.020
102. Rebeck GW. The role of APOE on lipid homeostasis and inflammation in normal brains. *J Lipid Res*. 2017;58:1493-1499. doi:10.1194/jlr.R075408
103. Chen Y, Durakoglugil MS, Xian X, Herz J. ApoE4 reduces glutamate receptor function and synaptic plasticity by selectively impairing ApoE receptor recycling. *Proc Natl Acad Sci*. 2010;107:12011-12016. doi:10.1073/pnas.0914984107
104. Fote GM, Geller NR, Efstathiou NE, et al. Isoform-dependent lysosomal degradation and internalization of apolipoprotein E requires autophagy proteins. *J Cell Sci*. 2022;135:jcs258687. doi:10.1242/jcs.258687
105. Heeren J, Grewal T, Laatsch A, et al. Impaired recycling of apolipoprotein E4 is associated with intracellular cholesterol accumulation. *J Biol Chem*. 2004;279:55483-55492. doi:10.1074/jbc.M409324200
106. Lee J-H, Yang D-S, Goulbourne CN, et al. Faulty autolysosome acidification in Alzheimer's disease mouse models induces autophagic build-up of Aβ in neurons, yielding senile plaques. *Nat Neurosci*. 2022;25:688-701. doi:10.1038/s41593-022-01084-8
107. Nixon RA, Wegiel J, Kumar A, et al. Extensive involvement of autophagy in Alzheimer disease: an immuno-electron microscopy study. *J Neuropathol Exp Neurol*. 2005;64:113-122. doi:10.1093/jnen/64.2.113
108. Nuriel T, Peng KY, Ashok A, et al. The endosomal-lysosomal pathway is dysregulated by APOE4 expression in vivo. *Front Neurosci*. 2017;11:702. doi:10.3389/fnins.2017.00702
109. Pohlkamp T, Xian X, Wong CH, et al. NHE6 depletion corrects ApoE4-mediated synaptic impairments and reduces amyloid plaque load. *eLife*. 2021;10:e72034. doi:10.7554/eLife.72034
110. Van Acker ZP, Bretou M, Annaert W. Endo-lysosomal dysregulations and late-onset Alzheimer's disease: impact of genetic risk factors. *Mol Neurodegener*. 2019;14:20. doi:10.1186/s13024-019-0323-7
111. Xian X, Pohlkamp T, Durakoglugil MS, et al. Reversal of ApoE4-induced recycling block as a novel prevention approach for Alzheimer's disease. *eLife*. 2018;7:e40048. doi:10.7554/eLife.40048
112. Lin PB, Tsai AP, Soni D, et al. INPP5D deficiency attenuates amyloid pathology in a mouse model of Alzheimer's disease. *Alzheimers Dement*. 2023;19:2528-2537. doi:10.1002/alz.12849
113. Rahmoune H, Guest PC. Studies of isolated peripheral blood cells as a model of immune dysfunction. In: Guest PC, ed. *Investig. Early Nutr. Eff. Long-Term Health*. Springer New York; 2018:221-229. doi:10.1007/978-1-4939-7614-0\_12
114. Jesudason CD, Mason ER, Chu S, et al. SHIP1 therapeutic target enablement: identification and evaluation of inhibitors for the treatment of late-onset Alzheimer's disease. *Alzheimers Dement N Y N*. 2023;9:e12429. doi:10.1002/trc2.12429
115. Nelson MR, Liu P, Agrawal A, et al. The APOE-R136S mutation protects against APOE4-driven Tau pathology, neurodegeneration and neuroinflammation. *Nat Neurosci*. 2023;26:2104-2121. doi:10.1038/s41593-023-01480-8
116. Marino C, Perez-Corredor P, O'Hare M, et al. APOE Christchurch-mimetic therapeutic antibody reduces APOE-mediated toxicity and tau phosphorylation. *Alzheimers Dement*. 2023;20(2):819-836. doi:10.1002/alz.13436
117. Yang A, Kantor B, Chiba-Falek O. APOE: the new frontier in the development of a therapeutic target towards precision medicine in late-onset Alzheimer's. *Int J Mol Sci*. 2021;22:1244. doi:10.3390/ijms22031244
118. Huynh T-PV, Liao F, Francis CM, et al. Age-dependent effects of apoE reduction using antisense oligonucleotides in a model of β-amyloidosis. *Neuron*. 2017;96:1013-1023. doi:10.1016/j.neuron.2017.11.014. e4.

119. Gottschalk KW, Mihovilovic M. The role of upregulated APOE in Alzheimer's disease etiology. *J Alzheimers Dis Park*. 2016;06. doi:10.4172/2161-0460.1000209
120. Linnertz C, Anderson L, Gottschalk W, et al. The cis -regulatory effect of an Alzheimer's disease-associated poly-T locus on expression of TOMM40 and apolipoprotein E genes. *Alzheimers Dement*. 2014;10:541-551. doi:10.1016/j.jalz.2013.08.280
121. Gamache J, Gingerich D, Shwab EK, et al. Integrative single-nucleus multi-omics analysis prioritizes candidate cis and trans regulatory networks and their target genes in Alzheimer's disease brains. *Cell Biosci*. 2023;13:185. doi:10.1186/s13578-023-01120-5
122. Kantor B, Odonovan B, Rittiner J, et al. All-in-one AAV-delivered epigenome-editing platform: *proof-of-concept* and therapeutic implications for neurodegenerative disorders. *Bioengineering*; 2023. doi:10.1101/2023.04.14.536951. bioRxiv
123. Rosenberg JB, Kaplitt MG, De BP, et al. AAVrh.10-mediated APOE2 Central Nervous System Gene Therapy for APOE4-associated Alzheimer's Disease. *Hum Gene Ther Clin Dev*. 2018;29:24-47. doi:10.1089/humc.2017.231
124. Nichols E, Steinmetz JD, Vollset SE, et al. Estimation of the global prevalence of dementia in 2019 and forecasted prevalence in 2050: an analysis for the Global Burden of Disease Study 2019. *Lancet Public Health*. 2022;7:e105-125. doi:10.1016/S2468-2667(21)00249-8
125. Boyle CP, Raji CA, Erickson KI, et al. Estrogen, brain structure, and cognition in p OSTMENOPAUSAL women. *Hum Brain Mapp*. 2021;42:24-35. doi:10.1002/hbm.25200
126. Kantarci K, Tosakulwong N, Lesnick TG, et al. Brain structure and cognition 3 years after the end of an early menopausal hormone therapy trial. *Neurology*. 2018;90(16):e1404-e1412. doi:10.1212/WNL.0000000000005325
127. Russell JK, Jones CK, Newhouse PA. The role of estrogen in brain and cognitive aging. *Neurotherapeutics*. 2019;16:649-665. doi:10.1007/s13311-019-00766-9
128. Shumaker SA. Conjugated equine estrogens and incidence of probable dementia and mild cognitive impairment in postmenopausal women's health initiative memory study. *JAMA*. 2004;291:2947. doi:10.1001/jama.291.24.2947
129. Saleh RNM, Hornberger M, Ritchie CW, Minihane AM. Hormone replacement therapy is associated with improved cognition and larger brain volumes in at-risk APOE4 women: results from the European Prevention of Alzheimer's Disease (EPAD) cohort. *Alzheimers Res Ther*. 2023;15:10. doi:10.1186/s13195-022-01121-5

#### SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Kloske CM, Belloy ME, Blue EE, et al. Advancements in APOE and dementia research: Highlights from the 2023 AAIC Advancements: APOE conference. *Alzheimer's Dement*. 2024;20:6590–6605. <https://doi.org/10.1002/alz.13877>