

The Role of STING-Mediated Activation of Dendritic Cells in Cancer Immunotherapy

Ana RS Ribeiro ^{1,2}, Theresa Neuper¹⁻³, Jutta Horejs-Hoek¹⁻³

¹Department of Biosciences and Medical Biology, Paris Lodron University of Salzburg, Salzburg, Austria; ²Cancer Cluster Salzburg (CCS), Salzburg, 5020, Austria; ³Center for Tumor biology and Immunology (CTBI), Salzburg, 5020, Austria

Correspondence: Jutta Horejs-Hoek, Department of Biosciences and Medical Biology, Paris-Lodron University Salzburg, Hellbrunner Strasse 34, Salzburg, 5020, Austria, Tel +43(0)662 8044 5709, Email jutta.horejs-hoek@plus.ac.at

Abstract: The signaling pathway that comprises cyclic guanosine monophosphate–adenosine monophosphate (cGAMP or GMP-AMP) synthase (cGAS) and Stimulator of Interferon Genes (STING) is emerging as a druggable target for immunotherapy, with tumor-resident dendritic cells (DC) playing a critical role in mediating its effects. The STING receptor is part of the DNA-sensing cellular machinery, that can trigger the secretion of pro-inflammatory mediators, priming effector T cells and initiating specific antitumor responses. Yet, recent studies have highlighted the dual role of STING activation in the context of cancer: STING can either promote antitumor responses or enhance tumor progression. This dichotomy often depends on the cell type in which cGAS-STING signaling is induced and the activation mode, namely acute versus chronic. Of note, STING activation at the DC level appears to be particularly important for tumor eradication. This review outlines the contribution of the different conventional and plasmacytoid DC subsets and describes the mechanisms underlying STING-mediated activation of DCs in cancer. We further highlight how the STING pathway plays an intricate role in modulating the function of DCs embedded in tumor tissue. Additionally, we discuss the strategies being employed to harness STING activation for cancer treatment, such as the development of synthetic agonists and nano-based delivery systems, spotlighting the current techniques used to prompt STING engagement specifically in DCs.

Keywords: dendritic cell, cGAS-STING pathway, cancer immunotherapy, nanoparticle

Introduction

The paradigm of natural antitumor immunity involves the recognition, processing and presentation of cancer-associated antigens by dendritic cells (DCs) on major histocompatibility (MHC) complexes, triggering effective T cell-mediated antitumor immunity that results in killing of the cancer cells.¹⁻⁴ Accordingly, DCs are highly specialized tissue-scanning cells equipped with a plethora of receptors to detect damage- or pathogen-associated molecular patterns (DAMPs or PAMPs). Although the immune system's task of filtering and responding to onco-signals emanating from cancerous tissue is an exceedingly complex one, the wide range of innate receptors expressed by DCs enables them to effectively recognize several by-products associated with tumor growth, such as tumor-derived DNA present in the tumor micro-environment (TME).⁵⁻⁷ The cGAS-STING pathway is an innate DNA-sensing mechanism which was originally described to mediate protective immune responses against infections but was recently found to play a crucial role in driving antitumor immunity. In particular, activation of STING signaling in DCs has the potential to generate spontaneous as well as long-lasting and highly specific antitumor immunity.⁸⁻¹⁰ This review focuses on the contribution of individual DC subsets to cancer defense and explores emerging developments in STING-mediated activation of DCs for novel cancer therapeutics.

The cGAS-STING Pathway and Its Effects in Cancer Cells

The cGAS-cGAMP-STING signaling pathway, discovered in 2013, is a promising therapeutic target to enhance antitumor immunity. Its activation induces an inflammatory response that converts “cold” tumors into “hot” tumors, ie, cancers that have higher infiltration of immune cells and consequently are more likely to exhibit higher response rates to currently used

immunotherapies such as immune checkpoint inhibitors (ICI).^{8,9,11} STING is a ubiquitous intracellular receptor that is located in the endoplasmic reticulum membrane and activated by binding of the agonist cGAMP. cGAMP is a naturally occurring cyclic dinucleotide (CDN) produced when the enzyme cGAS detects intracellular nucleic acids. Upon binding cGAMP, STING dimerizes and undergoes vesicular trafficking to the Golgi. This triggers downstream signaling which involves activation of TANK binding kinase 1 (TBK1), phosphorylation of interferon (IFN) regulatory factor 3 (IRF3), and induction of nuclear factor kappa B (NF- κ B).^{12,13} As a result, type I IFNs (IFN I) are upregulated, leading to expression and secretion of pro-inflammatory factors that can affect the TME and influence disease progression.^{14–16} In cancer cells, the cGAS-STING pathway is frequently activated due to genomic instability caused by higher occurrence of mitotic errors compared to healthy cells. Consequently, tumors are rich in micronuclei, and chromatin bridges. Micronuclei consist of residual chromosomes or chromosome fragments that are surrounded by a nuclear envelope that persists after mitosis. These structures are prone to rupture, and by releasing nucleic acids into the cytosol, they have been reported to trigger STING activation.^{17,18} Interestingly, Flynn et al report that in spite of the ability of these structures to efficiently recruit cGAS, they fail to activate the cGAS-STING pathway.¹⁹ Instead, the authors convincingly show that chromatin bridges, single-handedly trigger cGAS activation, and subsequent IFN signaling.¹⁹ Similarly to micronuclei, these structures are also byproducts of mitotic processes which persist throughout interphase and can be strongly stretched, until they break due to tension caused by cell migration.¹⁷ Self-chromatin can induce STING activation, that in turn, has the potential to impede early tumor progression by upregulating inflammatory genes and by inducing apoptosis as well as cell cycle arrest, in a process mediated by IFN I and NF- κ B.^{20,21} In melanoma, for example, restoration of STING signaling in tumor cells increases their immunogenicity and thus their recognition by T cells.²² Species-specific molecular determinants of self-DNA reactivity of cGAS were also uncovered using a melanoma model, which contributed to a better understanding of innate immunity involved in self-nonsel self discrimination. By engineering melanoma cells to express several species-specific cGAS transgenes with a doxycycline-inducible system, the authors showed that the key factor that protected mice from tumor-related mortality was the self-DNA reactive cGAS present in melanoma cells.²³ In fact, induction of STING in tumor cells can induce cell death, increasing the availability of tumor antigens to be recognized by antigen-presenting cells, such as DCs. Accordingly, besides the self-sustained protection mechanisms supported by STING activation in cancer cells, the continuous engagement of this receptor recruits various immune cells to the TME; therefore, enabling the cGAS-STING pathway in the immune compartment is of critical importance to overcome the generation of tumor-suppressive immune cells by the TME.^{24,25}

STING-Mediated Activation in DCs

Tumor-resident DCs are attractive targets for cancer immunotherapy because potent activation of DCs can initiate effective antitumor immune responses by priming CD4⁺ and CD8⁺ T cells.^{1,26,27} Upon recognition of PAMPs or DAMPs, DCs undergo a severe phenotypic and functional transformation, characterized by the secretion of a wide variety of inflammatory cytokines and the expression of maturation and activation markers such as MHC II and CD86. During this process, DCs become competent in cross-talking with lymphocytes, sustaining the expansion of specific effector cell subsets. As activation of the cGAS-STING pathway is well-known to promote DC activation, efforts have been made to understand the impact of STING activation in DCs within the tumor context. Accordingly, while it is widely accepted that DCs undergo a metabolic shift from oxidative phosphorylation to glycolysis upon activation, Hu et al recently found that increased ATP production via glycolysis facilitates STING-mediated antitumor functions in tumor-resident DCs.^{28,29} In an attempt to unravel the discrepancies observed with respect to the effects of STING activation in the tumor context in murine models and the human setting, Pang et al discovered that STING stimulation activated plasmacytoid (pDC) and conventional (cDC) DC subsets in both species, resulting in increased production of IFN I.³⁰ The authors also reported differential cell death responses upon STING activation, with rapid ablation occurring particularly for pDCs, highlighting the importance of understanding the impact of STING activation on different DC subsets.

cDC1s

cDC1s, identified by CD141⁺ in humans and CD103⁺ or CD8 α ⁺ markers in murine models, play a vital role in antitumor immunity by efficiently presenting tumor antigens on MHC I and thereby activating cytotoxic CD8⁺ T cells.^{1,31} Using

mice with deficiencies in different subsets of antigen-presenting cells, Wang et al demonstrated that specific STING licensing of cDC1s using cGAMP-loaded micelle nanoparticles (NPs) is essential for tumor rejection in mice. These NPs not only inhibited tumor growth *in vivo* but also induced a STING-activated cDC1 signature that correlated with higher survival rates among patients with non-small cell lung cancer (NSCLC).³² Another recent study found that the antitumor effects induced by cGAMP encapsulated in virus-like NPs also required STING activation in resident cDC1s (XCR1⁺ CD172⁻) but not in cDC2s. The choice of delivery vehicle, route of administration and STING agonist are all crucial factors, as the same study reported that administration of the synthetic CDN ADU-S100 induced cDC1 ablation.³³ In fact, ADU-S100 activates STING in cDC1s and not only fails to engage cDC2s but also induces upregulation of T-cell immunoglobulin and mucin-domain containing-3 (TIM-3), a traditional exhaustion marker.³⁴ Of note, another study reported that inhibiting TIM-3 also supports activation of the cGAS-STING pathway and increases the uptake of extracellular DNA in intratumoral cDC1s, emphasizing the role of TIM-3 as a regulator of DNA-sensing mechanisms.³⁵ Interestingly, cDC1 activation can also be prompted by IFN I secreted from STING-activated CD4⁺ T cells, thus establishing a mechanism that further supports antitumor immunity.³⁶

cDC2s

cDC2s, defined as CD1c⁺ and CD11b⁺ cells in humans and in murine animals, respectively, play a crucial role in halting cancer progression by directly promoting CD4⁺ T helper responses that support cytotoxic responses induced by CD8⁺ T cells.³⁷ Ulrich-Lewis et al convincingly demonstrated that upon treatment with a DNA vaccine, STING^{-/-} and cDC STING conditional knockout mice show lower Th1 responses compared to wild-type animals.³⁸ In line with previous findings, blocking of TIM-3 has been reported to enhance antitumor immunity mediated by STING agonist in 4T1 tumor models, by regulating cDC2s and improving the release of CD4⁺ T cells.³⁴

pDCs

Plasmacytoid DCs (pDCs) have a lower ability to act as antigen-presenting cells compared to cDCs. However, they are major producers of IFN and are key players in innate immunity, particularly in mounting responses against viral infections.^{39,40} In 2016, work by Bode et al demonstrated that human pDCs detect cytosolic DNA and CDNs via the cGAS-STING pathway, leading to robust activation and identifying pDCs as a potential target for new antitumor strategies.⁴¹ However, while stimulation of human pDCs with cGAMP or double-stranded immunostimulatory DNA (dsDNA) did induce nuclear translocation of IRF3, NF-κB was not activated.⁴² Interestingly, STING signaling in pDCs has been shown to be susceptible to TME cues, as it is impaired in melanomas.⁴³ Furthermore, Brewitz et al reported that pDC-secreted IFN can support maturation and cross presentation of cDCs in close proximity, enhancing CD8⁺ T cell-mediated immunity, and thus pinpointing pDCs as potential targets of interest for novel cancer therapies.⁴⁴

Tumor-Associated DCs

In the context of cancer, all DC subsets may be present within the TME, albeit with different phenotypes and functions compared to DCs present in healthy tissues. Initially identified in NSCLC patients by Maier et al, tumor-associated DCs are characterized by high expression of activation markers such as MHC and CD80/86, as well as markers associated with immune tolerance such as indoleamine 2,3-dioxygenase (IDO) and programmed death-ligand 1 (PD-L1).^{27,45-47} Thus, tumor-associated DCs are often referred to as mature-regulatory DCs (mregDCs) and both type I and type II conventional DCs, cDC1s and cDC2s respectively, can acquire that phenotype. mregDCs are metabolically very active and play a critical role in modulating the TME by influencing the recruitment and function of other immune cells, exerting significant control over tumor progression.²⁷ By increasing the expression of regulatory genes, mregDCs can interfere with T-cell responses by favoring regulatory T-cell polarization and decreasing the number of antitumor effector T cells.^{48,49} Accordingly, while several studies describe tumor-suppressing effects of STING signaling, STING activation in the TME has also been reported to induce IDO expression in tumor-associated DCs in a murine model of lung carcinoma. Moreover, the authors showed that only in tumors with low antigenicity did DNA-sensing by STING promote a suppressive TME and thus tumor growth.⁴⁹ In contrast, activation of STING signaling in tumor-associated DCs in hepatocellular carcinoma models increased the secretion of CXCL9 and IL-12, leading to infiltration and activation of

natural killer (NK) cells.⁵⁰ The STING-mediated antitumor function of DCs can also be induced by oxidative stress in the TME. In fact, Hu and co-authors showed that oxidative stress is required for DC antitumor functions mediated by STING, as it promotes the accumulation of the redox sensor SUMO Specific Peptidase 3 (SEN3) which detects reactive oxygen species (ROS) and enables STING-mediated antitumor responses.⁵¹ Oxidative stress, mediated by ROS, is a feature of the TME that can activate immunosuppressive mechanisms and also induce DNA oxidation, which enhances immune recognition by DCs.^{33,34} Due to the rapid growth of cells during neoplasm formation, tumors are often deprived of oxygen, and the hypoxic TME can intensify the oxidative stress. This leads to a metabolic shift of tumor-resident DCs from oxidative phosphorylation to glycolysis, thus supporting downstream antitumor responses mediated by STING. As a result of STING activation induced by glycolysis, there is upregulation of hypoxia-inducible factor 1 α (HIF-1 α) that can sustain a self-perpetuating HIF-1 α -mediated glycolysis, prompting DC activation.²⁹ Engagement of STING receptor is essential to start the activation and maturation of antigen-bearing tumor-resident DCs, supporting adaptive immunity by effector T cells (Figure 1).^{8,29,52}

STING Activation Strategies and Application in Cancer Therapies

Due to the strong immunostimulatory effect of STING signaling in immune oncology, much effort has focused on identifying the optimal agonist. Recent reports show that not only the induction of STING in immune cells can reduce TME-induced immunosuppression but also highlight that some agonists have the potential to enhance T cell infiltration at the tumor site, addressing a critical limitation of the success of adoptive cell therapy (ACT).^{53,54} Both Benoit-Lizon and




Figure 1 Illustration of the dual effects of STING activation in tumor-associated DCs. Tumor-associated DCs often display a mature-regulatory phenotype, characterized by the upregulation of both co-stimulatory as well as co-inhibitory molecules. In this context, it has been reported that the activation of STING in tumor-associated DCs may also play a dual role. On the one hand, engagement of the STING receptor can sustain antitumor effects by inducing HIF-1 α , which supports glycolysis in DCs, associated with an active and mature phenotype of these cells. This induces an IFN I-mediated response with secretion of pro-inflammatory cytokines that trigger effector T cell responses fighting the tumor (right panel). Conversely, activation of the STING pathway has also been reported to support pro-tumorigenesis, by inducing regulation of IDO expression and consequently enhancing regulatory T cell polarization (left panel). Created with BioRender.com.

Li have reported that STING agonists can directly activate T cells with no significant associated toxicity, improving effector responses mediated by CD8⁺ T cells, and Th1 and Th9, respectively.^{55,56} As STING induction may enhance anti-tumor immunity triggered by ICI and ACT, and as these therapies can be combined, it is possible that a suitable agonist may be the missing link for improved clinical success.^{56,57} Furthermore, Xu et al have shown that CAR T cells stimulated with STING agonists cGAMP or 5,6-dimethylxanthine-4-acetic acid (DMXAA) improved the control of tumor development as it enhanced the permanence of CAR T cells at the tumor site. Of note, the authors also show that sustained anti-tumor responses required co-treatment with ICI or anti-GR-1 Ab, which deplete myeloid-derived suppressor cells.⁵⁴ The relevance of the cGAS-STING pathway as a druggable target in immune-oncology is largely acknowledged for its classical inflammatory response, but also for its ability to induce anti-cancer immunity via IFN I-independent mechanisms (described in¹⁰). Hines et al have nicely reviewed candidate agonist molecules currently being evaluated clinically.⁵⁸ In broad terms, STING-activating agents can be categorized into CDNs and non-CDNs of natural or of synthetic origin.

Natural Agonists in the Tumor Context

cGAMP is a CDN that is expressed in both physiological and pathological conditions.⁵⁹ It can activate STING directly in the cell of origin or it can act on surrounding cells.^{60,61} However, due to its hydrophilic nature, it cannot passively diffuse through the cell membrane, requiring protein transporters for intercellular access. Solute Carrier Family 19 Member 1 (SLC19A1) was the first reported cGAMP importer, but several others have been identified subsequently and were thoroughly reviewed by Blest et al.⁶⁰ Nevertheless, none of them is particularly efficient in mediating cGAMP internalization in primary cells, leading to very low accumulation of this agonist in DCs.^{60,62,63} Furthermore, cGAMP is rapidly destroyed by nucleases in the plasma, which limits its uptake by antigen-presenting cells under physiological conditions, thereby preventing constitutive STING activation and collateral chronic inflammation.^{64,65} In cancer, DCs are more exposed to STING agonists secreted by cancer cells in the TME because of chromosomal instability and DNA leakage. This is particularly notable in primary tumors, where cancer cell DNA-sensing mechanisms can autonomously trigger antitumor immunity. Consequently, byproducts of the cGAS-STING pathway such as cGAMP or dsDNA are fed into the TME through necrosis or cell leakage, boosting immunity via DC activation.^{7,66} In fact, recent studies have shown that the success rate of radiation as an anticancer therapy is dependent on STING signaling. Ionizing radiation may promote the activation of exosomes loaded with tumor-derived dsDNA that mediate the cargo transfer to DCs, initiating pro-inflammatory cascades.⁶⁷ This hypothesis is supported by another study where cancer cells were treated with another DNA damaging agent, a topoisomerase I inhibitor. This promoted the activation of DCs through delivery of tumor-associated DNA by exosomes.⁶⁸ Furthermore, ionizing radiation induces genotoxicity and immunogenic cell death not only in irradiated cells but also in neighboring cells, increasing the exposure of DCs to natural tumor-derived STING agonists.^{69–71}

Synthetic Agonists

Synthetic agonists comprise CDNs and small molecules, often developed to improve the pharmacokinetics of STING agonists or to maximize STING signaling. The use of natural or synthetic CDNs as a standalone or combined cancer therapy is limited not only by their hydrophilicity, but also by their small size which causes rapid diffusion out of the tumors even when administered intratumorally.⁷² CDNs can be isolated from bacteria, such as cyclic di-GMP or cyclic di-AMP, or synthesized and modified to enhance their stability and immunogenicity for systemic administration.^{73,74} Despite achieving robust CD8⁺ T cell-mediated specific responses in preclinical models, prominent examples of synthetic CDNs like ADU-S100 did not perform well in clinical trials.^{75–78} Other CDNs have since been tested in clinical trials but all produced underwhelming results, as reviewed by others.^{79,80} To overcome the limitations of CDNs, other small molecules that sustain STING-mediated antitumor immune responses have been developed. For example, DMXAA was developed as an angiogenic disrupting agent in 2002, but it gained distinction for its potent immunomodulatory effects on the STING pathway, leading to unprecedented success in inducing tumor regression in pre-clinical models.^{81–83} However, DMXAA failed to replicate the results obtained in preclinical studies when used to treat NSCLC patients in combination with chemotherapy in Phase III trials. This disappointing result was traced to the drug's high specificity

exclusively for mouse STING.^{84–86} Several other non-CDN small molecules, most of which share structural similarities to DMXAA, are currently under development.^{80,87} For instance, dimeric amidobenzimidazole ligand (diABZI) is a synthetic non-CDN small molecule that demonstrated a 400-fold higher potency over cGAMP in activating peripheral blood mononuclear cells.⁶⁵ Nonetheless, small molecules lack strong cell selectivity across different subsets of immune cells, which leads to their uptake by lymphocytes and subsequent cell death. This highlights the need for drug delivery systems able to translate STING agonists into clinical applications.^{78,88–90}

STING Agonist Delivery Systems

Nano-based drug delivery systems offer promising advantages in modulating cargo pharmacokinetics and enabling targeted delivery, which can reduce drug-related side effects and enhance therapeutic effects.^{91,92} Liposomes were the first platforms investigated for encapsulating CDNs, due to their cationic lipid structure and aqueous core. In fact, encapsulation of cyclic di-GMP resulted in a STING-dependent enhanced antitumor response in metastatic melanoma mouse models.⁹³ Similarly, Doshi et al demonstrated that intravenous administration of ADU-S100 loaded into liposomes improved the therapeutic efficacy of that STING agonist and the responses to an immune checkpoint inhibitor (ICI) in both immunogenic and non-immunogenic *in vivo* tumor models.⁹⁴ CDNs were also successfully delivered using pegylated (PEG)-lipid discoid NPs, promoting efficient penetration into tumors after intravenous injection, and thus exposing most tumor cells to the loaded STING agonist in mice bearing colon adenocarcinomas.⁹⁵ The recently FDA-approved solid lipid nanoparticles (LNP) have also been exploited to this end, as they are known to successfully encapsulate nucleic acids and have intrinsic adjuvant properties. In line with this, Zhivaki et al have shown *in vivo* that LNPs loaded with mRNA encoding a self-DNA reactive variant of cGAMP had stronger immunostimulatory activity than other mRNA-encoded antigens also delivered by LNPs.⁹⁶ Mesoporous silica NPs also have been shown to be suitable for CDN delivery, achieving remarkable loading capacities and sustained release, resulting in a robust immune response in melanoma models.⁹⁷ Furthermore, several other polymeric-based NPs have successfully improved the delivery of CDNs, boosting the natural potency of cGAMP and inducing a “hot” immune TME in different tumor models.^{98,99} Accordingly, chitosan-based NPs, for instance, have been explored for their ability to induce cGAS-STING signaling, making them a promising adjuvant for use in novel formulations.^{100,101} Moreover, different ions, including Cu^{2+} , Zn^{2+} , Mn^{2+} , have been reported to support cGAS-STING signaling, but their exact mechanism has not yet been unveiled.^{102–105} In line with these findings, manganese has been deemed as essential not only in inducing cGAS-STING as a defense mechanism against dsDNA, but also in initiating innate immune responses to tumors by promoting maturation of DCs and macrophages and also supporting downstream effector T-cell responses.¹⁰⁶ The transporters for many of ions mentioned earlier are dysregulated in cancer (reviewed in¹⁰⁷), indicating that altered ion transport may be linked to cGAS-STING-mediated tumor immune evasion.

Nano-Based Platforms for Efficient Activation of STING in DCs

Several preclinical studies have explored different strategies to effectively deliver STING agonists to the TME and support STING-mediated tumor cell clearance. The interplay between cGAS-STING signaling and ions in the cancer context motivated the development of new strategies to tackle DCs with manganese nano-based systems.^{108–110} Using hybrid exosomes formulated with mannose ligands to target DCs, and protein nano-based structures without specific DC targeting, the effect of Mn^{2+} ions on STING activation of cDCs and bone marrow-derived DCs, respectively, was assessed in breast cancer tumor models. In both approaches, the authors included a DNA-damaging agent in their formulation that induces immunogenic cell death (ICD), thereby increasing the prevalence of dsDNA and conferring synergistic effects on STING engagement and further supporting antitumor responses mediated by the engagement of DCs with natural killer (NK) cells or with T-cells.^{111,112} Inducing tumor cell ICD to engage the cGAS-STING pathway has also been explored by Gu et al, in a study where colorectal cancer mouse models were co-treated with oxaliplatin-loaded liposomes and with ADU-S100.¹¹³ The authors observed that liposome-based immunotherapy not only induced cDC maturation, but also enhanced infiltration by CD4^{+} and CD8^{+} T cells. The effect of ADU-S100-loaded liposomes in DCs has also been assessed in colorectal and melanoma murine models. In this case, DCs were directly targeted by incorporating WH (a peptide with high affinity to Clec9a) into liposomes, and this resulted in enhanced potency of the

synthetic STING agonist.⁹⁴ The same DC-targeting strategy was used in another study, where poly(lactide-co-glycolide) (PLGA)-NPs encapsulating 2'3'-cGAMP were coated with another ligand of Clec9a, peptide CBP-12, and administered to animals with melanoma or breast cancer.¹¹⁴ The natural STING agonist 2'3'-cGAMP has also been encapsulated in virus-like particles and in liposomes and used to treat melanoma models. While there was no DC-targeting strategy for liposomes, in the virus-like particles a fusion glycoprotein was used for this purpose.^{33,115} To treat MC38-OVA tumor-bearing mice, 2'3'-cGAMP was loaded into polymeric STING-activating NPs (PolySTING) and the antitumor effects observed were attributed specifically to cDC1 activation.³² A formulation with the same name, consisting of a prodrug platform that was used to deliver diABZI to DCs via mannose ligands, resulted in selective targeting of DCs in the TME and boosted not only cDC1/CD8⁺ T-cell crosstalk and lymphocyte infiltration, but also supported maturation of other DC subsets in tumor-draining lymph nodes.¹¹⁶ The recent observations made by Dosta et al, in which CDN-NP-mediated antitumor effects were assessed in melanoma models, further support the observations reported above. CDN-NPs behaved as a potent immune agonist, sustaining cDC activation either through direct uptake or through posterior transfer of NPs internalized by cancer cells.¹¹⁷ By exploiting natural intercellular mechanisms, Jang et al showed that DC targeting can be achieved to some extent by using CDN-loaded HEK293-derived extracellular vesicles (ExoSTING) and that this formulation can induce maturation of all DC subsets.¹¹⁸ To provide clarity on the efforts made to directly activate DCs in order to drive antitumor immunity, a summary of the effects of nano-based systems for activating the STING signaling pathway in DCs is provided in Table 1.

STING Controversy – the Tumor-Promoting Effect

Recent studies have highlighted the dual role of STING-induced responses in immune oncology. While activation of the STING pathway is crucial for natural antitumor immune responses in low immunogenic primary tumors, overactivation can lead to unproductive nonimmunogenic cell ablation, rendering patients more vulnerable to relapse or metastasis.^{33,78} In metastatic tumors with high chromosomal instability, targeting STING may further support tumor immune evasion.^{119,120} In fact, chronic activation of the cGAS-STING pathway, even at low levels, can sustain immunoregulatory gene expression such as IDO in DCs, thereby supporting tumor outgrowth and metastasis.^{49,121,122} Moreover, non-canonical activation of the STING pathway can also induce tumor-promoting responses, as such activation has been reported to support breast cancer cell survival in patient-derived xenograft models.¹²³ Persistent inflammation following STING engagement in phagocytic cells engulfing dead cells may be tumor-promoting, which is consistent with the theory that prolonged, unresolved inflammation contributes to oncogenesis. Thus, targeting the cGAS-STING pathway in DCs is particularly challenging, as DCs appear to waver between pro- and antitumor T-cell responses.

Concluding Remarks

The net effect of STING activation on cancer is heavily influenced by several factors, including the timing and magnitude of STING stimulation and the STING-responsive target cells. As this receptor is ubiquitously expressed in different cells and independent of pathological states, identification of an optimal STING agonist candidate has proven to be difficult, based on the results of clinical trials.^{80,90,124} Nevertheless, it is uniformly recognized that DCs play a crucial role in stimulating antitumor immune responses, with STING engagement being essential to prompt this process through IFN I secretion.¹⁰ However, because the development of DC-targeted vaccines for cancer therapy remains challenging, no standalone STING-based therapies have been successful to date.^{77,125} Instead, current approaches combine the administration of STING agonists with ICIs such as PD-L1 or CTLA-4, because at optimal doses, STING induction prompts antitumor immunity mediated by IFN I signaling and it facilitates tumor infiltration by T cells, thereby converting “cold” tumors into “hot” tumors that are more prone to respond to ICI therapies.^{8,9,126–129} However, the underwhelming results of this approach underline the challenge of applying STING agonists in the clinic. The major hurdles hindering the clinical success of STING engagement in cancer immunotherapy are not only the poor pharmacokinetic and physico-chemical properties of CDNs that limit their systemic administration, but also the frequent impairment of STING signaling in multiple cancers, although its restoration may be achieved by epigenetic reprogramming.^{80,130–132} To overcome these difficulties, small molecule STING agonists have been tested in combination with immune checkpoint

Table 1 Strategies for Activation of the STING Pathway in DCs

Delivery System	Cargo	Size (nm)	DC-targeting Approach	DC Subset Analyzed	Model	Route of Administration	Ref
Hybrid exosomes	SN38 and MnO ₂	149	Unspecific (ICD of cancer cells and stimulation of STING)	cDC (CD11c ⁺)	4T1 tumor model	i.v.	[112]
Protein-based nanostructure (Man@Mn ²⁺ -Ft@Lap)	Mn ²⁺ and β-lapachone	100	Mannose ligands	BMDC	4T1 tumor model	i.v.	[111]
Liposomes	Oxaliplatin	122	Unspecific (ICD of cancer cells and stimulation of STING through co-administration of ADU-S100)	cDC1 and cDC2 (CD11c ⁺)	CT26 murine model	Liposomes: i.v. ADU-S100: i.t.	[113]
Liposomes	ADU-S100	121.3	WH peptide (Clec9a targeting)	cDC1 (CD103 ⁺)	MC38 and B16F10 murine tumor	i.v.	[94]
PLGA NPs	2'3'-cGAMP	157.5	CBP-12 EPBM NP coating (Clec9a targeting)	cDC1 (CD8α ⁺)	B16-OVA melanoma and 4T1 breast cancer	s.c.	[114]
Liposomes	2'3'-cGAMP	160	Unspecific	BMDC	B16-F10 melanoma model	in vitro stimulation	[115]
Virus-like particles	2'3'-cGAMP	158	Viral fusion glycoprotein	cDC1 and cDC2 (CD11c ⁺)	B16-F10 melanoma model	i.t.	[33]
PolySTING	2'3'-cGAMP	25	Not specified	cDC1 (CD8α ⁺ and CD103 ⁺)	MC38-OVA tumor-bearing	i.t.	[32]
PolySTING	diABZI	-	Mannose ligands	cDC1 (CD8α ⁺ and CD103 ⁺)	B16F10 and 4T1 tumor models	i.v.	[116]
CDN-NPs	Synthetic CDN	41	Unspecific	cDC (CD11c ⁺)	B16-F10 melanoma model	i.v.	[117]
ExoSTING	MR SS-2 CDA or cAIM(PS)2 Difluor (Rp/Sp)	50–200	EV import	pDC (CD123 ⁺), cDC (CD11c ⁺)	B16F10, EG7.OVA and CT26 models	i.t.	[118]

Abbreviations: BMDC, bone marrow-derived dendritic cell; i.v., intravenously; i.t., intratumorally; s.c., subcutaneous.

blockade, but these attempts were reported to induce toxicity in lymphocytes, highlighting that antitumor immunity requires STING licensing specifically in myeloid cells.^{88,133}

In summary, the cGAS-STING pathway is an undisputed target for novel developments in cancer immunotherapy, mediating the orchestration of critical antitumor responses by tumor-resident DCs. However, although STING signaling is essential to combat cancer progression, recent evidence has revealed its dual role, where chronic activation can paradoxically support tumor growth. Thus, the future of immuno-oncology relies on a deeper understanding of the effects of STING activation on different DC subsets in the tumor context, as well as on the development of efficient, tailored STING agonist delivery systems that will allow for optimal antitumor immune responses while mitigating potential side effects.

Acknowledgment

The image was created with Biorender. This work was supported by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie Action – Innovative Training Network DIRNANO (grant agreement No. 956544), the County of Salzburg, Cancer Cluster Salzburg [grant number 20102-P1601064-FPR01-2017], and by the Priority program ACBN, University of Salzburg.

Disclosure

The author(s) report no conflicts of interest in this work.

References

1. Broz ML, Binnewies M, Boldajipour B, et al. Dissecting the tumor myeloid compartment reveals rare activating antigen-presenting cells critical for T cell immunity. *Cancer Cell*. 2014;26(5):638–652. doi:10.1016/j.ccell.2014.09.007
2. Binnewies M, Mujal AM, Pollack JL, et al. Unleashing type-2 dendritic cells to drive protective antitumor CD4+ T cell immunity. *Cell*. 2019;177(3):556–571.e16. doi:10.1016/j.ccell.2019.02.005
3. Chiang MC, Tullett KM, Lee YS, et al. Differential uptake and cross-presentation of soluble and necrotic cell antigen by human DC subsets. *Eur J Immunol*. 2016;46(2):329–339. doi:10.1002/eji.201546023
4. Fu C, Jiang A. Dendritic cells and CD8 T cell immunity in tumor microenvironment. *Front Immunol*. 2018;9:423775. doi:10.3389/fimmu.2018.03059
5. Samson N, Ablasser A. The cGAS–STING pathway and cancer. *Nat Cancer*. 2022;3(12):1452–1463. doi:10.1038/s43018-022-00468-w
6. Kang TH, Mao C-P, Kim YS, et al. TLR9 acts as a sensor for tumor-released DNA to modulate anti-tumor immunity after chemotherapy. *J Immunol Cancer*. 2019;7(1):1–8. doi:10.1186/s40425-019-0738-2
7. Woo S-R, Fuertes MB, Corrales L, et al. STING-dependent cytosolic DNA sensing mediates innate immune recognition of immunogenic tumors. *Immunity*. 2014;41(5):830–842. doi:10.1016/j.immuni.2014.10.017
8. Wang H, Hu S, Chen X, et al. cGAS is essential for the antitumor effect of immune checkpoint blockade. *Proc Natl Acad Sci*. 2017;114(7):1637–1642. doi:10.1073/pnas.1621363114
9. Deng L, Liang H, Xu M, et al. STING-dependent cytosolic DNA sensing promotes radiation-induced type I interferon-dependent antitumor immunity in immunogenic tumors. *Immunity*. 2014;41(5):843–852. doi:10.1016/j.immuni.2014.10.019
10. Li G, Zhao X, Zheng Z, et al. cGAS-STING pathway mediates activation of dendritic cell sensing of immunogenic tumors. *Cell Mol Life Sci*. 2024;81(1):149. doi:10.1007/s00018-024-05191-6
11. X-D L, Wu J, Gao D, Wang H, Sun L, Chen ZJ. Pivotal roles of cGAS-cGAMP signaling in antiviral defense and immune adjuvant effects. *Science*. 2013;341(6152):1390–1394. doi:10.1126/science.1244040
12. Jeltama D, Abbott K, Yan N. STING trafficking as a new dimension of immune signaling. *J Exp Med*. 2023;220(3):e20220990. doi:10.1084/jem.20220990
13. Gonugunta VK, Sakai T, Pokatayev V, et al. Trafficking-mediated STING degradation requires sorting to acidified endolysosomes and can be targeted to enhance anti-tumor response. *Cell Rep*. 2017;21(11):3234–3242. doi:10.1016/j.celrep.2017.11.061
14. Decout A, Katz JD, Venkatraman S, Ablasser A. The cGAS–STING pathway as a therapeutic target in inflammatory diseases. *Nat Rev Immunol*. 2021;21(9):548–569. doi:10.1038/s41577-021-00524-z
15. Demaria O, De Gassart A, Coso S, et al. STING activation of tumor endothelial cells initiates spontaneous and therapeutic antitumor immunity. *Proc Natl Acad Sci*. 2015;112(50):15408–15413. doi:10.1073/pnas.1512832112
16. Arwert EN, Milford EL, Rullan A, et al. STING and IRF3 in stromal fibroblasts enable sensing of genomic stress in cancer cells to undermine oncolytic viral therapy. *Nat Cell Biol*. 2020;22(7):758–766. doi:10.1038/s41556-020-0527-7
17. Harding SM, Benci JL, Irianto J, Discher DE, Minn AJ, Greenberg RA. Mitotic progression following DNA damage enables pattern recognition within micronuclei. *Nature*. 2017;548(7668):466–470. doi:10.1038/nature23470
18. Mackenzie KJ, Carroll P, Martin C-A, et al. cGAS surveillance of micronuclei links genome instability to innate immunity. *Nature*. 2017;548(7668):461–465. doi:10.1038/nature23449
19. Flynn PJ, Koch PD, Mitchison TJ. Chromatin bridges, not micronuclei, activate cGAS after drug-induced mitotic errors in human cells. *Proc Natl Acad Sci*. 2021;118(48):e2103585118. doi:10.1073/pnas.2103585118

20. Ranoa DRE, Widau RC, Mallon S, et al. STING promotes homeostasis via regulation of cell proliferation and chromosomal stability. *Cancer Res.* 2019;79(7):1465–1479. doi:10.1158/0008-5472.CAN-18-1972
21. Guo Q, Chen X, Chen J, et al. STING promotes senescence, apoptosis, and extracellular matrix degradation in osteoarthritis via the NF- κ B signaling pathway. *Cell Death Dis* 2021;12(1):13. doi:10.1038/s41419-020-03341-9
22. Falahat R, Berglund A, Putney RM, et al. Epigenetic reprogramming of tumor cell–intrinsic STING function sculpts antigenicity and T cell recognition of melanoma. *Proc Natl Acad Sci.* 2021;118(15):e2013598118. doi:10.1073/pnas.2013598118
23. Mosallanejad K, Kennedy SN, Bahleda KM, et al. Species-specific self-DNA detection mechanisms by mammalian cyclic GMP-AMP synthases. *Sci Immunol.* 2023;8(79):eabp9765. doi:10.1126/sciimmunol.abp9765
24. Kim JC, Liu X, Fitzgerald K, et al. Brief report: STING expressed in tumor and non-tumor compartments has distinct roles in regulating anti-tumor immunity. *Cancer Immunol Immunother.* 2023;72(5):1327–1335. doi:10.1007/s00262-022-03327-w
25. Li S, Mirlekar B, Johnson BM, et al. STING-induced regulatory B cells compromise NK function in cancer immunity. *Nature.* 2022;610(7931):373–380. doi:10.1038/s41586-022-05254-3
26. Wculek SK, Cueto FJ, Mujal AM, Melero I, Krummel MF, Sancho D. Dendritic cells in cancer immunology and immunotherapy. *Nat Rev Immunol.* 2020;20(1):7–24. doi:10.1038/s41577-019-0210-z
27. Li J, Zhou J, Huang H, Jiang J, Zhang T, Ni C. Mature dendritic cells enriched in immunoregulatory molecules (mregDCs): a novel population in the tumour microenvironment and immunotherapy target. *Clin J Transl Med.* 2023;13(2):e1199. doi:10.1002/ctm2.1199
28. Møller SH, Wang L, P-C H. Metabolic programming in dendritic cells tailors immune responses and homeostasis. *Cell Mol Immunol* 2022;19(3):370–383. doi:10.1038/s41423-021-00753-1
29. Hu Z, Yu X, Ding R, et al. Glycolysis drives STING signaling to facilitate dendritic cell antitumor function. *J Clin Invest.* 2023;133(7). doi:10.1172/JCI166031
30. Pang ES, Daraj G, Balka KR, et al. Discordance in STING-induced activation and cell death between mouse and human dendritic cell populations. *Front Immunol.* 2022;13:794776. doi:10.3389/fimmu.2022.794776
31. Böttcher JP, e Sousa CR. The role of type 1 conventional dendritic cells in cancer immunity. *Trends Cancer.* 2018;4(11):784–792. doi:10.1016/j.trecan.2018.09.001
32. Wang J, Li S, Wang M, et al. STING licensing of type I dendritic cells potentiates antitumor immunity. *Sci Immunol.* 2024;9(92):eadj3945. doi:10.1126/sciimmunol.adj3945
33. Jneid B, Bochnakian A, Hoffmann C, et al. Selective STING stimulation in dendritic cells primes antitumor T cell responses. *Sci Immunol.* 2023;8(79):eabn6612. doi:10.1126/sciimmunol.abn6612
34. Luo J, Pang S, Hui Z, et al. Blocking Tim-3 enhances the anti-tumor immunity of STING agonist ADU-S100 by unleashing CD4+ T cells through regulating type 2 conventional dendritic cells. *Theranostics.* 2023;13(14):4836. doi:10.7150/thno.86792
35. De mingo Pulido Á, Hänggi K, Celiás DP, et al. The inhibitory receptor TIM-3 limits activation of the cGAS-STING pathway in intra-tumoral dendritic cells by suppressing extracellular DNA uptake. *Immunity.* 2021;54(6):1154–1167.e7. doi:10.1016/j.immuni.2021.04.019
36. Lei X, de Groot DC, Welters MJ, et al. CD4+ T cells produce IFN- γ to license cDC1s for induction of cytotoxic T-cell activity in human tumors. *Cell Mol Immunol* 2024;21:1–19.
37. Lei X, Khatri I, de Wit T, et al. CD4+ helper T cells endow cDC1 with cancer-impeding functions in the human tumor micro-environment. *Nat Commun.* 2023;14(1):217. doi:10.1038/s41467-022-35615-5
38. Ulrich-Lewis JT, Draves KE, Roe K, O'Connor MA, Clark EA, Fuller DH. STING is required in conventional dendritic cells for DNA vaccine induction of type I T helper cell-dependent antibody responses. *Front Immunol.* 2022;13:861710. doi:10.3389/fimmu.2022.861710
39. Villadangos JA, Young L. Antigen-presentation properties of plasmacytoid dendritic cells. *Immunity.* 2008;29(3):352–361. doi:10.1016/j.immuni.2008.09.002
40. Ziegler-Heitbrock L, Ohteki T, Ginhoux F, Shortman K, Spits H. Reclassifying plasmacytoid dendritic cells as innate lymphocytes. *Nat Rev Immunol.* 2023;23(1):1–2. doi:10.1038/s41577-022-00806-0
41. Bode C, Fox M, Tewary P, et al. Human plasmacytoid dendritic cells elicit a type I interferon response by sensing DNA via the cGAS-STING signaling pathway. *Eur J Immunol.* 2016;46(7):1615–1621. doi:10.1002/eji.201546113
42. Deb P, Dai J, Singh S, Kalyoussef E, Fitzgerald-Bocarsly P. Triggering of the cGAS–STING pathway in human plasmacytoid dendritic cells inhibits TLR9-mediated IFN production. *J Immunol.* 2020;205(1):223–236. doi:10.4049/jimmunol.1800933
43. Monti M, Ferrari G, Grosso V, et al. Impaired activation of plasmacytoid dendritic cells via toll-like receptor 7/9 and STING is mediated by melanoma-derived immunosuppressive cytokines and metabolic drift. *Front Immunol.* 2024;14:1227648. doi:10.3389/fimmu.2023.1227648
44. Brewitz A, Eickhoff S, Dähling S, et al. CD8+ T cells orchestrate pDC-XCR1+ dendritic cell spatial and functional cooperativity to optimize priming. *Immunity.* 2017;46(2):205–219. doi:10.1016/j.immuni.2017.01.003
45. Maier B, Leader AM, Chen ST, et al. A conserved dendritic-cell regulatory program limits antitumor immunity. *Nature.* 2020;580(7802):257–262. doi:10.1038/s41586-020-2134-y
46. Zhang Y, Chen H, Mo H, et al. Single-cell analyses reveal key immune cell subsets associated with response to PD-L1 blockade in triple-negative breast cancer. *Cancer Cell.* 2021;39(12):1578–1593.e8. doi:10.1016/j.ccell.2021.09.010
47. Smalley I, Chen Z, Phadke M, et al. Single-cell characterization of the immune microenvironment of melanoma brain and leptomeningeal metastases. *Clin Cancer Res.* 2021;27(14):4109–4125. doi:10.1158/1078-0432.CCR-21-1694
48. Munn DH, Mellor AL. IDO in the tumor microenvironment: inflammation, counter-regulation, and tolerance. *Trends Immunol.* 2016;37(3):193–207. doi:10.1016/j.it.2016.01.002
49. Lemos H, Mohamed E, Huang L, et al. STING promotes the growth of tumors characterized by low antigenicity via IDO activation. *Cancer Res.* 2016;76(8):2076–2081. doi:10.1158/0008-5472.CAN-15-1456
50. Wang S, Wu Q, Chen T, et al. Blocking CD47 promotes antitumor immunity through CD103+ dendritic cell–NK cell axis in murine hepatocellular carcinoma model. *J Hepatol.* 2022;77(2):467–478. doi:10.1016/j.jhep.2022.03.011
51. Hu Z, Teng X-L, Zhang T, et al. SENP3 senses oxidative stress to facilitate STING-dependent dendritic cell antitumor function. *Molecular Cell.* 2021;81(5):940–952.e5. doi:10.1016/j.molcel.2020.12.024
52. Andzinski L, Spanier J, Kasnitz N, et al. Growing tumors induce a local STING dependent type I IFN response in dendritic cells. *Int J Cancer.* 2016;139(6):1350–1357. doi:10.1002/ijc.30159

53. Conde E, Vercher E, Soria-Castellano M, et al. Epitope spreading driven by the joint action of CART cells and pharmacological STING stimulation counteracts tumor escape via antigen-loss variants. *J Immun Cancer*. 2021;9(11):e003351. doi:10.1136/jitc-2021-003351
54. Xu N, Palmer DC, Robeson AC, et al. STING agonist promotes CAR T cell trafficking and persistence in breast cancer. *J Exp Med*. 2021;218(2). doi:10.1084/jem.20200844
55. Li W, Lu L, Lu J, et al. cGAS-STING-mediated DNA sensing maintains CD8⁺ T cell stemness and promotes antitumor T cell therapy. *Sci, trans med*. 2020;12(549):eaay9013. doi:10.1126/scitranslmed.aay9013
56. Benoit-Lizon I, Jacquin E, Vargas TR, et al. CD4 T cell-intrinsic STING signaling controls the differentiation and effector functions of T H 1 and T H 9 cells. *J Immun Cancer*. 2022;10(1):e003459. doi:10.1136/jitc-2021-003459
57. Mullinax JE, Hall M, Prabhakaran S, et al. Combination of ipilimumab and adoptive cell therapy with tumor-infiltrating lymphocytes for patients with metastatic melanoma. *Front Oncol*. 2018;8:44. doi:10.3389/fonc.2018.00044
58. Hines JB, Kacew AJ, Sweis RF. The development of STING agonists and emerging results as a cancer immunotherapy. *Curr Oncol Rep*. 2023;25(3):189–199. doi:10.1007/s11912-023-01361-0
59. Carozza JA, Böhnert V, Nguyen KC, et al. Extracellular cGAMP is a cancer-cell-produced immunotransmitter involved in radiation-induced anticancer immunity. *Nat Cancer*. 2020;1(2):184–196. doi:10.1038/s43018-020-0028-4
60. Blest HT, Chauveau L. cGAMP the travelling messenger. *Front Immunol*. 2023;14:1150705. doi:10.3389/fimmu.2023.1150705
61. Ablasser A, Schmid-Burgk JL, Hemmerling I, et al. Cell intrinsic immunity spreads to bystander cells via the intercellular transfer of cGAMP. *Nature*. 2013;503(7477):530–534. doi:10.1038/nature12640
62. Ritchie C, Cordova AF, Hess GT, Bassik MC, Li L. SLC19A1 is an importer of the immunotransmitter cGAMP. *Molecular Cell*. 2019;75(2):372–381.e5. doi:10.1016/j.molcel.2019.05.006
63. Luteijn RD, Zaver SA, Gowen BG, et al. SLC19A1 transports immunoreactive cyclic dinucleotides. *Nature*. 2019;573(7774):434–438. doi:10.1038/s41586-019-1553-0
64. Carozza JA, Cordova AF, Brown JA, et al. ENPP1's regulation of extracellular cGAMP is a ubiquitous mechanism of attenuating STING signaling. *Proc Natl Acad Sci*. 2022;119(21):e2119189119. doi:10.1073/pnas.2119189119
65. Garland KM, Sheehy TL, Wilson JT. Chemical and biomolecular strategies for STING pathway activation in cancer immunotherapy. *Chem Rev*. 2022;122(6):5977–6039. doi:10.1021/acs.chemrev.1c00750
66. Fuertes MB, Woo S-R, Burnett B, Fu Y-X, Gajewski TF. Type I interferon response and innate immune sensing of cancer. *Trends Immunol*. 2013;34(2):67–73. doi:10.1016/j.it.2012.10.004
67. Diamond JM, Vanpouille-Box C, Spada S, et al. Exosomes shuttle TREX1-sensitive IFN-stimulatory dsDNA from irradiated cancer cells to DCs. *Cancer Immunol Res*. 2018;6(8):910–920. doi:10.1158/2326-6066.CIR-17-0581
68. Kitai Y, Kawasaki T, Sueyoshi T, et al. DNA-containing exosomes derived from cancer cells treated with topotecan activate a STING-dependent pathway and reinforce antitumor immunity. *J Immunol*. 2017;198(4):1649–1659. doi:10.4049/jimmunol.1601694
69. Azzam EI, De Toledo SM, Spitz DR, Little JB. Oxidative metabolism modulates signal transduction and micronucleus formation in bystander cells from α -particle-irradiated normal human fibroblast cultures. *Cancer Res*. 2002;62(19):5436–5442.
70. Demaria S, Formenti SC. Radiotherapy effects on anti-tumor immunity: implications for cancer treatment. *Frontiers Media S A*. 2013;128.
71. Zhou H, Randers-Pehrson G, Waldren CA, Vannais D, Hall EJ, Hei TK. Induction of a bystander mutagenic effect of alpha particles in mammalian cells. *Proc Natl Acad Sci*. 2000;97(5):2099–2104. doi:10.1073/pnas.030420797
72. Wang J, Meng F, Yao Y. Delivery of STING agonists for cancer immunotherapy. *Curr Opin Biotechnol*. 2024;87:103105. doi:10.1016/j.copbio.2024.103105
73. Woodward JJ, Iavarone AT, Portnoy DA. c-di-AMP secreted by intracellular *Listeria monocytogenes* activates a host type I interferon response. *Science*. 2010;328(5986):1703–1705. doi:10.1126/science.1189801
74. Burdette DL, Monroe KM, Sotelo-Troha K, et al. STING is a direct innate immune sensor of cyclic di-GMP. *Nature*. 2011;478(7370):515–518. doi:10.1038/nature10429
75. Knelson EH, Ivanova EV, Tarannum M, et al. Activation of tumor-cell STING primes NK-cell therapy. *Cancer Immunol Res*. 2022;10(8):947–961. doi:10.1158/2326-6066.CIR-22-0017
76. Zaidi AH, Kelly RJ, Gorbunova A, et al. Intratumoral immunotherapy with STING agonist, ADU-S100, induces CD8⁺ T-cell mediated anti-tumor immunity in an esophageal adenocarcinoma model. *Oncotarget*. 2021;12(4):292. doi:10.18632/oncotarget.27886
77. Meric-Bernstam F, Sweis RF, Hodi FS, et al. Phase I dose-escalation trial of MIW815 (ADU-S100), an intratumoral STING agonist, in patients with advanced/metastatic solid tumors or lymphomas. *Clin Cancer Res*. 2022;28(4):677–688. doi:10.1158/1078-0432.CCR-21-1963
78. Sivick KE, Desbrien AL, Glickman LH, et al. Magnitude of therapeutic STING activation determines CD8⁺ T cell-mediated anti-tumor immunity. *Cell Rep*. 2018;25(11):3074–3085.e5. doi:10.1016/j.celrep.2018.11.047
79. Amouzegar A, Chelvanambi M, Filderman JN, Storkus WJ, Luke JJ. STING agonists as cancer therapeutics. *Cancers*. 2021;13(11):2695. doi:10.3390/cancers13112695
80. Motedayen Aval L, Pease JE, Sharma R, Pinato DJ. Challenges and opportunities in the clinical development of STING agonists for cancer immunotherapy. *J Clin Med*. 2020;9(10):3323. doi:10.3390/jcm9103323
81. Prantner D, Perkins DJ, Lai W, et al. 5, 6-Dimethylxanthenone-4-acetic acid (DMXAA) activates stimulator of interferon gene (STING)-dependent innate immune pathways and is regulated by mitochondrial membrane potential. *J Biol Chem*. 2012;287(47):39776–39788. doi:10.1074/jbc.M112.382986
82. Corrales L, Glickman LH, McWhirter SM, et al. Direct activation of STING in the tumor microenvironment leads to potent and systemic tumor regression and immunity. *Cell Rep*. 2015;11(7):1018–1030. doi:10.1016/j.celrep.2015.04.031
83. Ching L, Cao Z, Kieda C, Zwain S, Jameson M, Baguley B. Induction of endothelial cell apoptosis by the antivascular agent 5, 6-dimethylxanthenone-4-acetic acid. *Br J Cancer*. 2002;86(12):1937–1942. doi:10.1038/sj.bjc.6600368
84. Lara PN Jr, Douillard J-Y, Nakagawa K, et al. Randomized phase III placebo-controlled trial of carboplatin and paclitaxel with or without the vascular disrupting agent vadimezan (ASA404) in advanced non-small-cell lung cancer. *J Clin Oncol*. 2011;29(22):2965–2971. doi:10.1200/JCO.2011.35.0660
85. Conlon J, Burdette DL, Sharma S, et al. Mouse, but not human STING, binds and signals in response to the vascular disrupting agent 5, 6-dimethylxanthenone-4-acetic acid. *J Immunol*. 2013;190(10):5216–5225. doi:10.4049/jimmunol.1300097

86. Gao P, Ascano M, Zillinger T, et al. Structure-function analysis of STING activation by c [G (2', 5') pA (3', 5') p] and targeting by antiviral DMXAA. *Cell*. 2013;154(4):748–762. doi:10.1016/j.cell.2013.07.023
87. Hao X, Chen Z, Li H, Wei M, Zuo Z, Su Q. Small-molecule drugs in immunotherapy. *Mini Reviews in Med Chem*. 2023;23(13):1341–1359. doi:10.2174/1389557522666220930154527
88. Gulen MF, Koch U, Haag SM, et al. Signalling strength determines proapoptotic functions of STING. *Nat Commun*. 2017;8(1):427. doi:10.1038/s41467-017-00573-w
89. Wu J, Dobbs N, Yang K, Yan N. Interferon-independent activities of mammalian STING mediate antiviral response and tumor immune evasion. *Immunity*. 2020;53(1):115–126.e5. doi:10.1016/j.immuni.2020.06.009
90. Zou Y, Zhang M, Zhou J. Recent trends in STING modulators: structures, mechanisms, and therapeutic potential. *Drug Discovery Today*. 2023;28(9):103694. doi:10.1016/j.drudis.2023.103694
91. Huang P, Wang X, Liang X, et al. Nano-, micro-, and macroscale drug delivery systems for cancer immunotherapy. *Acta Biomater*. 2019;85:1–26. doi:10.1016/j.actbio.2018.12.028
92. Shao K, Singha S, Clemente-Casares X, Tsai S, Yang Y, Santamaria P. Nanoparticle-based immunotherapy for cancer. *ACS nano*. 2015;9(1):16–30. doi:10.1021/nn5062029
93. Nakamura T, Miyabe H, Hyodo M, Sato Y, Hayakawa Y, Harashima H. Liposomes loaded with a STING pathway ligand, cyclic di-GMP, enhance cancer immunotherapy against metastatic melanoma. *J Control Release*. 2015;216:149–157. doi:10.1016/j.jconrel.2015.08.026
94. Doshi AS, Cantin S, Prickett LB, Mele DA, Amiji M. Systemic nano-delivery of low-dose STING agonist targeted to CD103+ dendritic cells for cancer immunotherapy. *J Control Release*. 2022;345:721–733.
95. Dane EL, Belessiotis-Richards A, Backlund C, et al. STING agonist delivery by tumour-penetrating PEG-lipid nanodiscs primes robust anticancer immunity. *Nature Mater*. 2022;21(6):710–720. doi:10.1038/s41563-022-01251-z
96. Zhivaki D, Gosselin EA, Sengupta D, et al. mRNAs encoding self-DNA reactive cGAS enhance the immunogenicity of lipid nanoparticle vaccines. *Mbio*. 2023;14(6):e02506–23. doi:10.1128/mbio.02506-23
97. Park KS, Xu C, Sun X, Louttit C, Moon JJ. Improving STING agonist delivery for cancer immunotherapy using biodegradable mesoporous silica nanoparticles. *Adv Ther*. 2020;3(10):2000130. doi:10.1002/adtp.202000130
98. Shae D, Becker KW, Christov P, et al. Endosomolytic polymersomes increase the activity of cyclic dinucleotide STING agonists to enhance cancer immunotherapy. *Nature Nanotechnol*. 2019;14(3):269–278. doi:10.1038/s41565-018-0342-5
99. Wang-Bishop L, Kimmel BR, Ngwa VM, et al. STING-activating nanoparticles normalize the vascular-immune interface to potentiate cancer immunotherapy. *Sci Immunol*. 2023;8(83):eadd1153. doi:10.1126/sciimmunol.add1153
100. Carroll EC, Jin L, Mori A, et al. The vaccine adjuvant chitosan promotes cellular immunity via DNA sensor cGAS-STING-dependent induction of type I interferons. *Immunity*. 2016;44(3):597–608. doi:10.1016/j.immuni.2016.02.004
101. Zhang S, Zeng Y, Wang K, et al. Chitosan-based nano-micelles for potential anti-tumor immunotherapy: synergistic effect of cGAS-STING signaling pathway activation and tumor antigen absorption. *Carbohydr Polym*. 2023;321:121346. doi:10.1016/j.carbpol.2023.121346
102. Wang P, Wang Y, Li H, et al. A homologous-targeting cGAS-STING agonist multimodally activates dendritic cells for enhanced cancer immunotherapy. *Acta Biomater*. 2024;177:400–413—.
103. Wang Q, Gao Y, Li Q, He A, Xu Q, Mou Y. Enhancing dendritic cell activation through manganese-coated nanovaccine targeting the cGAS-STING pathway. *Int J Nanomed*. 2024;Volume 19:263–280. doi:10.2147/IJN.S438359
104. Molinaro C, Martoriati A, Pelinski L, Cailliau K. Copper complexes as anticancer agents targeting topoisomerases I and II. *Cancers*. 2020;12(10):2863. doi:10.3390/cancers12102863
105. Du M, Chen ZJ. DNA-induced liquid phase condensation of cGAS activates innate immune signaling. *Science*. 2018;361(6403):704–709. doi:10.1126/science.aat1022
106. Lv M, Chen M, Zhang R, et al. Manganese is critical for antitumor immune responses via cGAS-STING and improves the efficacy of clinical immunotherapy. *Cell Res*. 2020;30(11):966–979. doi:10.1038/s41422-020-00395-4
107. Yu L, Liu P. Cytosolic DNA sensing by cGAS: regulation, function, and human diseases. *Signal Transduct Target Ther*. 2021;6(1):170. doi:10.1038/s41392-021-00554-y
108. Gao M, Xie YQ, Lei K, et al. A manganese phosphate nanocluster activates the cGAS-STING pathway for enhanced cancer immunotherapy. *Adv Ther*. 2021;4(8):2100065. doi:10.1002/adtp.202100065
109. Sun X, Zhang Y, Li J, et al. Amplifying STING activation by cyclic dinucleotide–manganese particles for local and systemic cancer metalloimmunotherapy. *Nature Nanotechnol*. 2021;16(11):1260–1270. doi:10.1038/s41565-021-00962-9
110. Deng Z, Xi M, Zhang C, et al. Biomimetic MnO₂ nanoplateforms mediated delivery of immune checkpoint inhibitors with STING pathway activation to potentiate cancer radio-immunotherapy. *ACS nano*. 2023;17(5):4495–4506. doi:10.1021/acsnano.2c10352
111. Wang X, Liu Y, Xue C, et al. A protein-based cGAS-STING nanoagonist enhances T cell-mediated anti-tumor immune responses. *Nat Commun*. 2022;13(1):5685. doi:10.1038/s41467-022-33301-0
112. Cheng L, Zhang P, Liu Y, et al. Multifunctional hybrid exosomes enhanced cancer chemo-immunotherapy by activating the STING pathway. *Biomaterials*. 2023;301:122259. doi:10.1016/j.biomaterials.2023.122259
113. Gu Z, Hao Y, Schomann T, Ossendorp F, Ten Dijke P, Cruz LJ. Enhancing anti-tumor immunity through liposomal oxaliplatin and localized immunotherapy via STING activation. *J Control Release*. 2023;357:531–544. doi:10.1016/j.jconrel.2023.04.011
114. Gou S, Liu W, Wang S, et al. Engineered nanovaccine targeting Clec9a+ dendritic cells remarkably enhances the cancer immunotherapy effects of STING agonist. *Nano Lett*. 2021;21(23):9939–9950. doi:10.1021/acs.nanolett.1c03243
115. Koshy ST, Cheung AS, Gu L, Graveline AR, Mooney DJ. Liposomal delivery enhances immune activation by STING agonists for cancer immunotherapy. *Adv Biosyst*. 2017;1(1–2):1600013. doi:10.1002/adbi.201600013
116. Nguyen DC, Song K, Jokonya S, et al. Mannosylated STING agonist drugamers for dendritic cell-mediated cancer immunotherapy. *ACS Cent Sci* 2024;10:666–75.
117. Dosta P, Cryer AM, Dion MZ, et al. Investigation of the enhanced antitumor potency of STING agonist after conjugation to polymer nanoparticles. *Nature Nanotechnol*. 2023;18(11):1351–1363. doi:10.1038/s41565-023-01447-7
118. Jang SC, Economides KD, Moniz RJ, et al. ExoSTING, an extracellular vesicle loaded with STING agonists, promotes tumor immune surveillance. *Commun Biol*. 2021;4(1):497. doi:10.1038/s42003-021-02004-5

119. Motwani M, Pesiridis S, Fitzgerald KA. DNA sensing by the cGAS–STING pathway in health and disease. *Nat Rev Genet.* 2019;20(11):657–674. doi:10.1038/s41576-019-0151-1
120. Bakhoun SF, Ngo B, Laughney AM, et al. Chromosomal instability drives metastasis through a cytosolic DNA response. *Nature.* 2018;553(7689):467–472. doi:10.1038/nature25432
121. Sharma S, Campbell AM, Chan J, et al. Suppression of systemic autoimmunity by the innate immune adaptor STING. *Proc Natl Acad Sci.* 2015;112(7):E710–E717. doi:10.1073/pnas.1420217112
122. Lemos H, Huang L, Chandler PR, et al. Activation of the STING adaptor attenuates experimental autoimmune encephalitis. *J Immunol.* 2014;192(12):5571–5578. doi:10.4049/jimmunol.1303258
123. Cheradame L, Guerrero IC, Gaston J, et al. STING protects breast cancer cells from intrinsic and genotoxic-induced DNA instability via a non-canonical, cell-autonomous pathway. *Oncogene.* 2021;40(49):6627–6640. doi:10.1038/s41388-021-02037-4
124. Kong X, Zuo H, Huang H-D, et al. STING as an emerging therapeutic target for drug discovery: perspectives from the global patent landscape. *J Adv Res.* 2023;44:119–133. doi:10.1016/j.jare.2022.05.006
125. Harrington K, Brody J, Ingham M, et al. Preliminary results of the first-in-human (FIH) study of MK-1454, an agonist of stimulator of interferon genes (STING), as monotherapy or in combination with pembrolizumab (pembro) in patients with advanced solid tumors or lymphomas. *Ann Oncol.* 2018;29:viii712. doi:10.1093/annonc/mdy424.015
126. Marloye M, Lawler SE, Berger G. Current patent and clinical status of stimulator of interferon genes (STING) agonists for cancer immunotherapy. *Future Sci.* 2019;8:87–90.
127. Li A, Yi M, Qin S, Song Y, Chu Q, Wu K. Activating cGAS-STING pathway for the optimal effect of cancer immunotherapy. *J Hematol Oncol.* 2019;12(1):1–12. doi:10.1186/s13045-019-0721-x
128. Sistigu A, Yamazaki T, Vacchelli E, et al. Cancer cell–autonomous contribution of type I interferon signaling to the efficacy of chemotherapy. *Nature Med.* 2014;20(11):1301–1309. doi:10.1038/nm.3708
129. Ghaffari A, Peterson N, Khalaj K, et al. STING agonist therapy in combination with PD-1 immune checkpoint blockade enhances response to carboplatin chemotherapy in high-grade serous ovarian cancer. *Br J Cancer.* 2018;119(4):440–449. doi:10.1038/s41416-018-0188-5
130. Song S, Peng P, Tang Z, et al. Decreased expression of STING predicts poor prognosis in patients with gastric cancer. *Sci Rep.* 2017;7(1):39858. doi:10.1038/srep39858
131. Falahat R, Berglund A, Perez-Villarreal P, et al. Epigenetic state determines the in vivo efficacy of STING agonist therapy. *Nat Commun.* 2023;14(1):1573. doi:10.1038/s41467-023-37217-1
132. Lin Z, Liu Y, Lin P, Li J, Gan J. Clinical significance of STING expression and methylation in lung adenocarcinoma based on bioinformatics analysis. *Sci Rep.* 2022;12(1):13951. doi:10.1038/s41598-022-18278-6
133. Larkin B, Ilyukha V, Sorokin M, Buzdin A, Vannier E, Poltorak A. Cutting edge: activation of STING in T cells induces type I IFN responses and cell death. *J Immunol.* 2017;199(2):397–402. doi:10.4049/jimmunol.1601999

International Journal of Nanomedicine

Dovepress

Publish your work in this journal

The International Journal of Nanomedicine is an international, peer-reviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch®, Current Contents®/Clinical Medicine, Journal Citation Reports/Science Edition, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-nanomedicine-journal>