

Review Article

Transcriptional regulation of insect steroid hormone biosynthesis and its role in controlling timing of molting and metamorphosis

Yuko S. Niwa^{1*} and Ryusuke Niwa^{2,3*}

¹Life Science Center, Tsukuba Advanced Research Alliance, University of Tsukuba; ²Faculty of Life and Environmental Sciences, University of Tsukuba, Tennoudai 1-1-1, Tsukuba, Ibaraki 305-8572, Japan; and ³PRESTO, Japan Science and Technology Agency, Honcho 4-1-8, Kawaguchi 332-0012, Saitama, Japan

The developmental transition from juvenile to adult is often accompanied by many systemic changes in morphology, metabolism, and reproduction. Curiously, both mammalian puberty and insect metamorphosis are triggered by a pulse of steroid hormones, which can harmonize gene expression profiles in the body and thus orchestrate drastic biological changes. However, understanding of how the timing of steroid hormone biosynthesis is regulated at the molecular level is poor. The principal insect steroid hormone, ecdysteroid, is biosynthesized from dietary cholesterol in the specialized endocrine organ called the prothoracic gland. The periodic pulses of ecdysteroid titers determine the timing of molting and metamorphosis. To date, at least nine families of ecdysteroidogenic enzyme genes have been identified. Expression levels of these genes correlate well with ecdysteroid titers, indicating that the transcriptional regulatory network plays a critical role in regulating the ecdysteroid biosynthesis pathway. In this article, we summarize the transcriptional regulation of ecdysteroid biosynthesis. We first describe the development of prothoracic gland cells during *Drosophila* embryogenesis, and then provide an overview of the transcription factors that act in ecdysteroid biosynthesis and signaling. We also discuss the external signaling pathways that target these transcriptional regulators. Furthermore, we describe conserved and/or diverse aspects of steroid hormone biosynthesis in insect species as well as vertebrates.

Key words: ecdysteroid, insect, metamorphosis, prothoracic gland, transcription.

Introduction

Temporal coordination of organismal development, simply called developmental timing, is one of the fundamental aspects in developmental biology (Ambros 2000; Thummel 2001; Banerjee & Slack 2002; Rougvie 2005). In various multicellular organisms, appropriate regulation of developmental timing allows

organisms to be sexually mature adults from the juvenile stage. For example, in mammals including humans, puberty is one of the major temporal changes during development, which initiates a series of drastic morphological and physiological changes that make organisms reproductive. Such a drastic developmental transition to transform sexually immature individuals to fecund adults, known as metamorphosis, is also found in evolutionarily distant animals such as insects.

Both mammalian puberty and insect metamorphosis are triggered by steroid hormones, which are small fat-soluble bioactive molecules that can pass through the cell membrane into the cytoplasm (Miller & Auchus 2011; Yamanaka *et al.* 2013a; Niwa & Niwa 2014b). Steroid hormones can systemically harmonize changes of gene expression in the body and thus orchestrate the drastic biological changes. One of the crucial keys to determine the timing of both puberty and metamorphosis is temporal regulation of steroid hormone biosynthesis *in vivo*, which generates temporally specific

*Author to whom all correspondence should be addressed.

Emails: shimada.yuko.gn@u.tsukuba.ac.jp; ryusuke-niwa@u.tsukuba.ac.jp

Received 2 September 2015; revised 11 October 2015; accepted 11 October 2015.

© 2015 The Authors

Development, Growth & Differentiation published by John Wiley & Sons Australia, Ltd on behalf of Japanese Society of Developmental Biologists.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.


peaks of hemolymph steroid hormone titers to trigger developmental transitions (Hariharan 2012). Therefore, understanding the mechanisms that modulate the timing of biosynthesis is important to comprehend developmental timing at the molecular level.

An initial step towards the elucidation of steroid hormone biosynthesis is identification and characterization of steroidogenic enzymes responsible for converting precursor sterols to active steroid hormones. Studies on mammalian steroidogenic enzymes have been conducted since the 1980s, and a number of essential steroidogenic enzymes have been identified (Miller 1988; Hanukoglu 1992). By contrast, insect steroidogenic enzymes have been reported only since 2000. The previous 15 years, however, have been a fruitful period in terms of the elucidation of a number of insect steroidogenic enzymes (Niwa & Niwa 2014a). The principal insect steroid hormones are ecdysteroids, including ecdysone and its active derivative 20-hydroxyecdysone (20E) that trigger metamorphosis as well as molting. Molecular genetics and biochemical studies using the fruit fly *Drosophila melanogaster* and the silkworm *Bombyx mori* have revealed that for ecdysteroid biosynthesis in the ecdysone-producing organ, called the prothoracic gland (PG, Fig. 1), at least nine families of enzymes are required: *noppera-bo* (*nobo*) (Enya *et al.* 2014, 2015),

neverland (*nvd*) (Yoshiyama *et al.* 2006; Yoshiyama-Yanagawa *et al.* 2011), *non-molting glossy/shroud* (*sro*) (Niwa *et al.* 2010), *Cyp307a1/spook* (*spo*) (Niwa *et al.* 2005; Ono *et al.* 2006), *Cyp307a2/spookier* (*spok*) (Ono *et al.* 2006), *Cyp6t3* (Ou *et al.* 2011), *Cyp306a1/phantom* (*p hm*) (Niwa *et al.* 2004; Warren *et al.* 2004), *Cyp302a1/disembodied* (*dib*) (Chávez *et al.* 2000; Warren *et al.* 2002), and *Cyp315a1/shadow* (*sad*) (Warren *et al.* 2002). After release from the PG to the hemolymph, ecdysone is converted to 20E by another enzyme *Cyp314a1/shade* (*shd*) in the peripheral tissues (Petryk *et al.* 2003). All of these enzymes (except *nvd*, *spok*, and *Cyp6t3*) are collectively referred to as the Halloween genes (Rewitz *et al.* 2007; Niwa & Niwa 2014a).

After the discovery of the ecdysteroidogenic enzymes, researchers promptly realized that the expression levels of these biosynthesis genes in the PG correlate very well with the levels of hemolymph ecdysteroid titers (Warren *et al.* 2002, 2006; Niwa *et al.* 2005; Parvy *et al.* 2005). This is reminiscent of mammalian steroidogenic gene expression, which well reflects steroid hormone production (Mizutani *et al.* 2015). Therefore, it turns out that the timing of steroid hormone biosynthesis depends on transcriptional regulation of the steroidogenic enzyme genes. Indeed, transcriptional regulation of mammalian steroidogenic

Fig. 1. Overview of the ecdysteroid biosynthesis pathway and the regulatory mechanisms through transcriptional network in the prothoracic gland (PG) of *Drosophila melanogaster*. Several distinct signaling pathways regulate ecdysteroid biosynthesis and some pathways potentially target transcription factors (TFs) to regulate ecdysteroidogenic enzyme gene expressions. The ecdysteroid biosynthesis pathway is colored in blue. It starts with dietary cholesterol. Ecdysteroid biosynthesis enzymes are highlighted in green. Hexagons represent ecdysteroidogenic TFs that are listed in Table 1. Among them, nuclear receptors are colored in sky blue. In each signaling pathway, only the key components are depicted. Yellow “P” means phosphorylation. DILPS, *Drosophila* insulin-like peptides; ERK, Extracellular signal-regulated kinase; InR, Insulin receptor; NO, nitric oxide; PI3K, Phosphoinositide 3-kinase; PG, prothoracic gland; PTTH, Prothoracicotropic hormone; TOR, Target of rapamycin.



genes has been an important research issue for a long time, and the key steroidogenic transcriptional factors (TFs) have been identified. The nuclear receptor NR5A1, also known as Ad4BP/Steroidogenic Factor 1 (SF-1), is recognized as a master regulator for steroidogenic gene expression, as this TF controls the transcription of almost all steroidogenic genes in steroidogenic tissues (Parker *et al.* 2002; Morohashi *et al.* 2013). By contrast, elucidation of ecdysteroidogenic TFs and their regulatory roles in insects has lagged far behind that of mammalian steroidogenic transcriptional regulation.

In this review, we will examine the recent progress of the spatio-temporal transcriptional regulatory mechanisms underlying ecdysteroid biosynthesis in the PG. Based on studies published in the previous 15 years, we will illustrate that multiple transcription factors are cooperatively working in a network to achieve the differentiation and morphogenesis of the PG cells, and the appropriate control of ecdysteroid biosynthesis during larval development. In addition, we will also discuss how TFs are regulated by extracellular stimuli, affecting ecdysteroid biosynthesis by controlling ecdysteroidogenic gene expressions.

Transcription factors specify the origin of the prothoracic gland cells

In mammals, Ad4BP/SF-1 serves as a master TF to induce differentiation of the cell into a steroidogenic cell lineage (Parker *et al.* 2002; Morohashi *et al.* 2013; Mizutani *et al.* 2015). In insects, much less is known about molecular mechanisms to regulate the development of the PG itself during embryogenesis. Very recently, Sánchez-Higuera *et al.* (2014) reported that differentiation and morphogenesis of the PG requires a proper combination of TFs during *Drosophila* embryogenesis (Fig. 2). The study demonstrates that the PG has a homologous origin with the respiratory tracheal system in *Drosophila*. The PG and the corpora allata (CA), which biosynthesizes juvenile hormone, are derived from identical primordia in successive segments of the head and trunk of the embryo: the PG arises in the labial (Lb) segment, the CA arises in the maxillary (Mx) segment, and the trachea arises in thoracic T2 to abdominal A8 segments. In each segment, the identity of the PG, CA, and tracheas is initially specified by a unique Homeotic TF with the Signal Transducers and Activator of Transcription (STAT) via inducing expression of the POU-domain TF gene *ventral veins lacking* (also known as *ventral veinless*; *vvl*). In detail, Deformed (Dfd) and Sex comb reduced (Scr) control *vvl* expressions in the Mx and Lb patches, respectively. Subsequently, a subgroup of *vvl*-expressing




Fig. 2. Schematic of ectodermal endocrine and respiratory primordia in embryos and their specification gene regulatory network. A part of this cartoon is adapted from figure 4 in Sánchez-Higuera *et al.* (2014). In each segment, Homeotic genes and Signal Transducers and Activators of Transcription (STAT) induce expression of the early transcription factors. Deformed (Dfd)-STAT in the maxillary primordium (Mx) and Sex combs reduced (Scr)-STAT in the labial primordium (Lb) induce *ventral veins lacking* (*vvl*) and *snail* (*sna*) expression. Spalt (Sal) represses trunk Homeotic gene expression in these primordia, preventing tracheless (*trh*) expression. The *sna*-expressing cells undergo the epithelial-mesenchymal transition (EMT) and migrate to form the corpora allata (CA) and the prothoracic gland (PG) cells. The corpora cardiaca (CC) cells are separately derived from anterior mesodermal cells. Representative lineage markers are Glass (Gl) in the CC, Seven-up (Svp) in the CA, and Sal in the PG. These cells make a composite endocrine organ called the ring gland (RG).

cells activates the Zn-finger gene *snail*, a key regulator of the epithelial-mesenchymal transition. The *snail*-expressing cells then migrate dorsally and merge into the corpora cardiaca (CC), which originates from anterior mesodermal cells. In cyclorrhaphous Diptera, including *Drosophila*, the PG, CA, and CC form a composite endocrine organ “the ring gland (RG)”. The differentiated PG and CA cells eventually express the specific marker TF genes, *spalt* and *seven-up*, respectively (Sánchez-Higuera *et al.* 2014). In summary, the specification of the embryonic PG primordia requires a specific combination of TFs, at least including Scr, STAT, Vvl, and Snail (Fig. 2). These TF codes might be essential for establishing the cellular status of the PG cells expressing the special set of ecdysteroidogenic genes.

Transcription factors are required for expression of ecdysteroidogenic genes in the prothoracic gland

After the PG is specified, a number of ecdysteroidogenic enzyme genes and ecdysteroidogenic regulators begin to be expressed, which is an important characteristic of this endocrine organ. Such gene expressions are induced and/or maintained by a number of TFs, hereinafter referred to as “ecdysteroidogenic TFs”. Some but not all ecdysteroidogenic TFs directly bind to the promoters of ecdysteroidogenic genes, confirmed by an electrophoresis mobility shift assay or chromatin immunoprecipitation assay (Xiang *et al.* 2010; Deng & Kerppola 2013; Danielsen *et al.* 2014; Meng *et al.* 2015). The known validated and putative ecdysteroidogenic TFs required in the PG are listed in Table 1.

The first identified TF that influences the expression levels of validated ecdysteroidogenic genes was β Ftz-f1, a critical regulator of insect metamorphosis in many

tissues (Parvy *et al.* 2005). β Ftz-f1 is a homologue of Ad4BP/SF-1, suggesting a conserved role of steroidogenic TFs. Studies on *Drosophila* clearly show that the protein levels of Phm and Dib are significantly reduced with the loss of β Ftz-f1 function in PG cells (Parvy *et al.* 2005).

Most of the ecdysteroidogenic TFs, listed in Table 1, have originally been characterized as TFs for their non-steroidogenic functions: β Ftz-f1, Ultraspiracle (USP) (Koyama *et al.* 2014), Broad (Br) (Xiang *et al.* 2010; Moeller *et al.* 2013), DHR3 (Parvy *et al.* 2014), and DHR4 (Ou *et al.* 2011) are encoded by the well-known ecdysteroid-inducible genes that are expressed in many types of cells (Thummel 2001; Ou & King-Jones 2013). Other TFs are known for their roles in spatial pattern formation. For example, Vvl described above is involved in cellular differentiation of several types of cells including the PG (Cheng *et al.* 2014; Danielsen *et al.* 2014; Sánchez-Higueras *et al.* 2014). *Knirps* is quite well known as a gap gene during embryogenesis (Danielsen *et al.* 2014). Moreover, TFs involved in

Table 1. A list of ecdysteroidogenic transcription factors

TF names	Protein family	Reported ecdysteroidogenic genes whose expression are affected†	Organisms analyzed in published studies	References
Antp	homeotic	<u>phm</u>	<i>Bombyx mori</i>	Meng <i>et al.</i> (2015)
Br	C ₂ H ₂ zinc finger	<u>phm</u> , <u>dib</u> , <u>sad</u> , <u>npc1</u>	<i>Drosophila melanogaster</i>	Xiang <i>et al.</i> (2010); Moeller <i>et al.</i> (2013)
CncC	basic leucine zipper	<u>nvd</u> , <u>spok</u> , <u>dib</u> , <u>sad</u>	<i>D. melanogaster</i>	Deng & Kerppola (2013)
DHR3	nuclear receptor	<u>phm</u> , <u>dib</u> , <u>sad</u>	<i>D. melanogaster</i>	Parvy <i>et al.</i> (2014)
DHR4	nuclear receptor	<u>Cyp6t3</u>	<i>D. melanogaster</i>	Ou <i>et al.</i> (2011)
dKeap1	BTB	<u>nvd</u> , <u>spok</u> , <u>phm</u> , <u>dib</u> , <u>sad</u>	<i>D. melanogaster</i>	Deng & Kerppola (2013)
E75	nuclear receptor	<u>phm</u>	<i>D. melanogaster</i>	Bialecki <i>et al.</i> (2002); Cáceres <i>et al.</i> (2011); Parvy <i>et al.</i> (2014)
EcR	nuclear receptor	<u>phm</u> , <u>dib</u> , <u>sad</u>	<i>D. melanogaster</i>	Moeller <i>et al.</i> (2013); Parvy <i>et al.</i> (2014)
FOXO	Forkhead	<u>phm</u> , <u>dib</u>	<i>D. melanogaster</i>	Koyama <i>et al.</i> (2014)
β Ftz-f1	nuclear receptor	<u>phm</u> , <u>dib</u> , <u>sad</u>	<i>D. melanogaster</i>	Parvy <i>et al.</i> (2005, 2014)
Kni	C ₂ C ₂ zinc finger	<u>phm</u> , <u>dib</u> , <u>sad</u>	<i>D. melanogaster</i>	Danielsen <i>et al.</i> (2014)
Mld‡	C ₂ H ₂ zinc finger	<u>nvd</u> , <u>spok</u> , <u>sro</u>	<i>D. melanogaster</i>	Ono <i>et al.</i> (2006); Danielsen <i>et al.</i> (2014)
Ouib‡	C ₂ H ₂ zinc finger	<u>spok</u>	<i>D. melanogaster</i>	Komura-Kawa <i>et al.</i> (2015).
USP	C ₂ C ₂ zinc finger, ligand binding	<u>phm</u> , <u>dib</u>	<i>D. melanogaster</i>	Koyama <i>et al.</i> (2014)
Vvl/POU-M2	POU	<u>spok</u> , <u>sro</u> , <u>phm</u> , <u>dib</u> , <u>sad</u> , <u>torso</u>	<i>D. melanogaster</i> <i>Tribolium castaneum</i> <i>B. mori</i>	Cheng <i>et al.</i> (2014); Danielsen <i>et al.</i> (2014); Meng <i>et al.</i> (2015)

†Underlines indicate genes whose promoter sequences can be physically associated with TFs, which are shown by electrophoresis mobility shift assay and/or chromatin immunoprecipitation analyses. ‡It must be noted that these TF genes are found only in genomes of Drosophilidae species. Abbreviations: Antp, Antennapedia; Br, Broad; CncC, Cap'n'collar; DHR3, Drosophila Hormone Receptor 3; DHR4, Drosophila Hormone Receptor 4; E75, Ecdysone-induced protein 75; EcR, Ecdysone receptor; FOXO, Forkhead box, sub-group O; β Ftz-F1, β -fushi tarazu transcription factor 1; Kni, Knirps; Mld, Molting defective; Ouib, Ouija board; USP, Ultraspiracle; Vvl, Ventral veins lacking

metabolic responses have also been identified, such as the Cap'n'collar (CncC)-dKeap1 complex and Forkhead box, sub-group O (FOXO) that are mediators of the xenobiotic metabolism signaling pathway and of the insulin/insulin-like peptide signaling pathway, respectively (Deng & Kerppola 2013; Koyama *et al.* 2014).

In addition to typical TFs, chromatin remodeling factors influence ecdysteroid biosynthesis by affecting expression of many but not all ecdysteroidogenic genes in the PG, as evidenced by genetic analyses on the dATAC histone acetylase complex (Pankotai *et al.* 2010; Borsos *et al.* 2015) and the insulator protein CTCF (Fresán *et al.* 2015). It should be noted that these chromatin remodeling factors are also known as crucial proteins for many biological processes other than ecdysteroid biosynthesis, indicating that these 'generalist'-type transcription regulators act on a certain group of ecdysteroidogenic genes in the PG.

The evolutionarily conserved function of steroidogenic TFs

Many of the ecdysteroidogenic TFs are evolutionarily conserved across a wide variety of animal species from insects to mammals. The most epitomized example is β Ftz-f1, an insect orthologue of vertebrate Ad4BP/SF1. As described above, Ad4BP/SF1 is the key regulator of vertebrate steroidogenic organ specifications and steroidogenic gene expressions (Parker *et al.* 2002; Morohashi *et al.* 2013; Mizutani *et al.* 2015). It is also interesting to note that a chromosomal deletion of a human homologue of *vvl*, known as *POU3F2*, is associated with hypogonadotropic hypogonadism and adrenal insufficiency (Bonaglia *et al.* 2008; Izumi *et al.* 2013). Insect orthologues of *vvl* are also involved in ecdysteroid biosynthesis not only in *D. melanogaster* but also in other insects such as the red flour beetle *Tribolium castaneum* (Cheng *et al.* 2014) and *B. mori* (referred as *POU-M2*) (Meng *et al.* 2015). Therefore, the β Ftz-f1/SF-1 and Vvl/POU3F2 families emphasize the conserved regulatory mechanisms of steroidogenesis between insects and vertebrates.

On the other hand, the ecdysteroidogenic C_2H_2 zinc finger TF Molting defective (Mld) is very unique because its orthologues are found only in genomes of Drosophilidae species (Neubueser *et al.* 2005; Ono *et al.* 2006). The less conservative nature of Mld does not mean a lesser importance of the TF; Mld plays an essential role in ecdysteroid biosynthesis via inducing *nvd* and *spok* (Ono *et al.* 2006; Danielsen *et al.* 2014). It is also noteworthy that the classical temperature-sensitive dominant mutant *lethal(3)dts3*, which shows

ecdysteroid deficiency (Walker *et al.* 1987), has been recognized and used as an allele of *mld* (Simon *et al.* 2003; Ishimoto *et al.* 2013). Besides *mld*, we have also recently identified another novel ecdysteroidogenic C_2H_2 zinc finger TF gene designated *ouija board* (*ouib*), which is required for expression of *spok* but whose orthologues are found only in Drosophilidae genomes (Komura-Kawa *et al.* 2015). These data imply that some essential ecdysteroidogenic TFs might have rapidly evolved only in very small insect clade(s), and thus future studies using a variety of insects would be valuable to unravel ecdysteroidogenic TFs that are not well conserved in Drosophilidae.

PTTH signaling regulates transcription of ecdysteroidogenic genes in the prothoracic glands

In general, expression levels of ecdysteroidogenic genes correlate well with temporal fluctuation of the ecdysteroid titer during development. Thus, regarding the issue of "Time in development", an important question to be addressed is how the ecdysteroidogenic TFs described above contribute to timing of ecdysteroid biosynthesis in the PG.

Previous studies have demonstrated that timing of ecdysteroid biosynthesis in the PG is influenced by multiple extracellular stimuli (Niwa & Niwa 2014b). The most important and classical humoral factor is the neuropeptide Prothoracicotropic hormone (PTTH), which stimulates the biosynthesis and secretion of ecdysteroids in the PG (Tanaka 2011). The temporal coordination of PTTH secretion regulates timing of molting and metamorphosis in many insects (Mizoguchi *et al.* 2001, 2015; Halme *et al.* 2010; Yamanaka *et al.* 2013b). While PTTH has been extensively studied for its short-term prothoracicotropic activity, which requires the *de novo* translation of proteins (Gilbert *et al.* 2002), PTTH also influences transcription in the PG, which might have a long-term effect on ecdysteroid biosynthesis (Ou & King-Jones 2013). For example, *in vitro* experimental assays have demonstrated that a recombinant PTTH protein stimulates transcription of *spo*, *dib*, and *phm* in the PG of the silkworm *B. mori* (Namiki *et al.* 2005; Niwa *et al.* 2005; Yamanaka *et al.* 2007). Conversely, PTTH neuron-ablated animals or loss-of-function animals of *torso*, which encodes a PTTH receptor (Rewitz *et al.* 2009b) exhibit drastic reduction of many ecdysteroidogenic enzyme genes in *D. melanogaster* (McBrayer *et al.* 2007; Niwa *et al.* 2010; Enya *et al.* 2014). Therefore, the PTTH signaling pathway in the PG, which consists of Torso, Ras small GTPase, Raf kinase, and Extracellular signal-related kinase (ERK) (Rewitz *et al.* 2009b), should

control the activities of some ecdysteroidogenic TFs (Fig. 1).

Although ecdysteroidogenic TFs acting downstream of the Torso-Ras-ERK pathway are not fully understood, one striking example is the nuclear receptor DHR4 (Ou *et al.* 2011). The activity of DHR4 is regulated by its subcellular localization between the nucleus and cytoplasm of the PG cells. Furthermore, DHR4 protein accumulates in the PG nuclei at developmental times when the ecdysteroid titer is high. A crucial function of DHR4 is to negatively regulate expression of the ecdysteroidogenic P450 gene *Cyp6t3*, which functions in the 'Black Box' in the biosynthesis pathway but whose substrate is still unknown (Ou *et al.* 2011).

Currently, DHR4 is the only good example of ecdysteroidogenic TFs acting downstream of PTTH signaling. Nevertheless, DHR4 is not extensively involved in regulating expression of any other characterized ecdysteroidogenic enzyme genes (Ou *et al.* 2011). Therefore, PTTH signaling must target other ecdysteroidogenic TFs in the PG to control ecdysteroidogenic gene expressions. Recently, it was shown that loss of *cncC* function suppresses a developmental acceleration phenotype of the activated *Ras* overexpression (Deng & Kerppola 2013), suggesting that the CncC-dKeap1 complex would be a candidate acting downstream of the PTTH signaling pathway.

Regulation of ecdysteroidogenic gene expression in the prothoracic gland by 20-hydroxyecdysone and the ecdysteroid-signaling cascade

In addition to PTTH, other humoral factors that influence ecdysteroidogenic gene expressions in the PG are ecdysteroids *per se*, particularly 20E, the most biologically active form of ecdysteroids, and its downstream signaling. 20E has a large impact on ecdysteroid biosynthesis in the PG (Gilbert *et al.* 2002). For example, in cultured *B. mori* PGs, both ecdysteroid biosynthesis activity and responsiveness to PTTH are inhibited by 20E administration (Takaki & Sakurai 2003). Curiously, the inhibitory effect of 20E on the PG is dependent on the larval developmental stage of *B. mori*, implying that 20E seems to have a feedback effect on the PG to generate a peak of ecdysteroid level from juvenile to adult transition.

An *in vivo* biological significance of the feedback effect of 20E has recently been demonstrated by genetics of *D. melanogaster* (Fig. 3). During the larval stages, reduced ecdysteroid-EcR signaling in the PG decreases ecdysteroidogenic gene expressions, leading to a delay in the larva-to-pupa transition (Moeller

et al. 2013). This indicates that ecdysteroid has a positive-feedback effect on the PG, rapidly amplifying its own synthesis to trigger pupariation. Indeed, the isoforms of the ecdysteroid-regulated factor Br, Br-Z1, and Br-Z4 bind directly to the *phm* and *dib* promoters/enhancers to regulate their transcriptions. By contrast, after pupariation, reduced ecdysteroid-EcR signaling increases ecdysteroidogenic gene expressions, leading to incomplete metamorphosis. This means that a negative-feedback signal ensures the decline in ecdysteroid levels in the prepupa-to-pupa transition. Notably, these opposing signals depend on the different responses of Br isoforms to ecdysteroid levels: low levels of ecdysteroid should quickly induce Br-Z4, whereas high levels of ecdysteroid should induce Br-Z1 that transcriptionally silences the ecdysteroidogenic genes (Moeller *et al.* 2013). The inhibitory effect of Br is reminiscent of the fact that overexpression of *Br* in the PG blocks molting, and this larval arrest phenotype can be rescued by feeding 20E (Zhou *et al.* 2004).

Other ecdysteroid-response genes modulate timing of ecdysteroid biosynthesis in the PG. Similar to the repressive function of Br-Z1 described above, the ecdysteroid-inducible nuclear receptor DHR3 represses ecdysteroidogenic enzyme genes through EcR function during the prepupal-to-pupal transition (Parvy *et al.* 2014). Conversely, before the transition, the other ecdysteroid-inducible nuclear receptor E75 positively regulates ecdysteroid biosynthesis (Bialecki *et al.* 2002). Indeed, E75 and β *Ftz-f1* counteracts DHR3 to avoid premature repression of ecdysteroid biosynthesis (Cáceres *et al.* 2011; Parvy *et al.* 2014). These data demonstrate that the metamorphic 20E peak relies on ecdysteroid-mediated feedback control of PG activity through transcriptional regulatory networks.

E75 contains a heme moiety and thus can bind to nitric oxide (NO), an important secondary messenger acting as a short-range signaling molecule in a vast array of important physiological processes (Reinking *et al.* 2005). NO is produced by NO synthase (NOS) in the PG and blocks the function of E75 (Wildemann & Bicker 1999; Cáceres *et al.* 2011). While it is unclear whether and how NO synthesis is temporally regulated in the PG, the NO signaling pathway possibly modulates ecdysteroid biosynthesis in the PG (Jaszczak *et al.* 2015).

Other extracellular signals that influence ecdysteroidogenic gene expression

Recent studies have accumulated evidence that ecdysteroidogenic gene expression in the PGs is influenced not only by PTTH and 20E, but also by




Fig. 3. Feedback control of ecdysteroids affects ecdysteroidogenic gene expressions in the larva-to-pupa transition in *Drosophila*. The names of TFs are listed in Table 1. In the late 3rd instar stage, ecdysteroid biosynthesis is amplified by the positive feedback of 20-hydroxyecdysone (20E)-EcR signaling and nitric oxide (NO)-E75 signaling to trigger pupariation. Br-B4 upregulates the expressions of ecdysteroidogenic enzyme genes such as *phm* and *dib*. After pupariation, 20E-EcR signaling negatively regulates ecdysteroid biosynthesis by Br-Z1, which suppresses the expressions of ecdysteroidogenic enzyme genes. Moreover, DHR3 also represses ecdysteroidogenic enzyme gene expressions. The positive and negative feedback circuits accomplish the temporal peak of ecdysteroid titer in the developmental transition. NOS, nitric oxide synthase.

other extracellular signals and their signaling pathways (Fig. 1). These pathways include the insulin like peptides-TOR pathway (Colombani *et al.* 2005; Koyama *et al.* 2013), the TGF β /Activin-Smad pathway (Gibbens *et al.* 2011), serotonin and its receptor 5HT-7 (Shimada-Niwa & Niwa 2014), octopamine and its receptor Oct β 3R (Ohhara *et al.* 2012, 2015), and Bommo-FMRamide that is a prothoracicostatic neuropeptide (Yamanaka *et al.* 2007), all of which are required for determining the proper timing of ecdysteroid biosynthesis in the PG during development (Niwa & Niwa 2014b). It will be interesting to determine which ecdysteroidogenic TFs act downstream of any of these signaling cascades, and whether any ecdysteroidogenic TFs are regulated by those multiple signaling inputs.

Some ecdysteroidogenic TFs, however, control expression of genes encoding essential components of these signaling pathways. For example, Vvl and Kni are required for expression of *torso*, *Insulin receptor*, *akt*, *4E-BP*, and *S6 kinase* (Danielsen *et al.* 2014), indicating that some ecdysteroidogenic TFs directly and indirectly regulate ecdysteroid biosynthesis by transcriptional control. Related to this point, ecdysteroidogenic TFs also regulate transcription of genes that are involved in uptake and transport of extracellular cholesterol or plant sterols, which are the precursors of ecdysteroids. For example, Br plays an indispens-

able role in controlling expression of not only ecdysteroidogenic genes (Danielsen *et al.* 2014) but also the *Niemann-Pick type C1* gene, which encodes the evolutionarily conserved cholesterol transporter (Xiang *et al.* 2010). β Ftz-f1 is also required for ecdysteroid biosynthesis in the PG via regulating expression of a scavenger receptor gene *Snmp1*, which appears to be involved in lipid uptake in the PG cells (Talamillo *et al.* 2008, 2013).

It must be noted that regulatory mechanisms to control ecdysteroidogenic gene expression might be diversified among insect species. For example, several studies have reported that juvenile hormones (JHs) have significant effects on some ecdysteroidogenic enzyme genes and *torso* in the PG of *B. mori* (Yamanaka *et al.* 2007; Young *et al.* 2012; Ogihara *et al.* 2015). This is less likely the case with the PG in *D. melanogaster*, considering that JHs appear not to have the typical "status quo" effect on larval development in *D. melanogaster* (Niwa *et al.* 2008; Liu *et al.* 2009; Riddiford *et al.* 2010; Ono 2014; Wen *et al.* 2015). Furthermore, some prothoracicotropic factors might primarily control translation, but not transcription, in the PG. A recent example is Pigment dispersing factor (PDF), as PDF stimulates ecdysteroid biosynthesis in the PG of *B. mori* but does not influence any known ecdysteroidogenic enzyme genes (Iga *et al.* 2014).

Protein modifications and protein–protein interactions to modulate the activity of ecdysteroidogenic TFs

Downstream of the signaling pathways, the activity of ecdysteroidogenic TFs must be modulated at the protein level. The striking example is the protein SUMOylation. A recent study has demonstrated that the activity of β Ftz-f1 in ecdysteroidogenic cells, including PG and ovarian follicle cells, is modulated by its SUMOylation (Talamillo *et al.* 2008, 2013). Curiously, mammalian Ad4BP/SF-1 is also SUMOylated. Moreover, the disruption of Ad4BP/SF-1 SUMOylation in mice exhibits the inappropriate activation of target genes, leading to abnormalities of endocrine development (Lee *et al.* 2011), suggesting that a part of post-translational modification of ecdysteroidogenic TFs is evolutionarily conserved between insects and vertebrates. β Ftz-f1 also appears to be controlled by its acetylation at least in *D. melanogaster* cultured cells, resulting in its protein stabilization (Borsos *et al.* 2015). In addition, phosphorylation might be one of the essential post-translational regulations in the PG, as PTTTH signaling involves the Ras-ERK pathway (Rewitz *et al.* 2009b). However, currently, there are no reports demonstrating the phosphorylation of any validated ecdysteroidogenic TFs including DHR4 (Rewitz *et al.* 2009a).

Physical interactions between ecdysteroidogenic TFs and other regulator proteins have also been recognized as another layer of mechanisms regulating ecdysteroid biosynthesis in the PG, as shown by the interaction between USP and FOXO (Koyama *et al.* 2014) and among CDK8, CyclinC, EcR, and USP (Xie *et al.* 2015). Further protein interactome analyses would be important in the future.

Outlook

In the previous 15 years, significant progress has been made that allows for a better understanding of transcriptional regulation of ecdysteroid biosynthesis, particularly in the fruit fly *D. melanogaster*. It is now obvious that ecdysteroidogenic genes are regulated by multiple numbers of TFs in the PG. However, the current list of ecdysteroidogenic TFs is incomplete, as candidate ecdysteroidogenic TFs are still present as previously discussed (Ou & King-Jones 2013). For example, mutant animals of *without children* (*woc*), encoding a C₂H₂-type zinc finger protein, is a larval lethal with ecdysteroid deficiency (Wismar *et al.* 2000). Because the mutant can be partially rescued by feeding 7-dehydrocholesterol (Warren *et al.* 2001), *Woc* has been hypothesized to activate transcription

of a gene involved in the cholesterol 7,8-dehydrogenation to produce 7dC. However, it is still unclear whether *Woc* regulates the expression of *nvd*, encoding cholesterol 7,8-dehydrogenase (Yoshiyama *et al.* 2006), or any other downstream targets. Another example is a basic-Helix-Loop-Helix TF gene *HLH54F* that is predominantly expressed in the PG of both *D. melanogaster* and *B. mori* (Namiki *et al.* 2009). No validated targets of HLH54F in the PG have been reported. More interestingly, the central circadian clock genes *period* and *timeless*, which do not encode actual TFs but regulatory proteins modulating transcription, are rhythmically expressed in the pupal PG and play a role in controlling eclosion rhythms (Myers *et al.* 2003; Morioka *et al.* 2012). Therefore, gene expression levels of a certain set of genes in the PG might be transcriptionally oscillated under the control of the central clock network. Identification and characterization of the oscillatory genes may elucidate a connection between clock and metamorphosis.

Undoubtedly, the current studies on ecdysteroidogenic TFs are almost only focusing on their functions in the PG cells, while ecdysteroid biosynthesis occurs in other types of cells during embryonic and adult stages. In the embryo, ecdysteroid biosynthesis is activated during mid-embryogenesis before the development of PG primordial cells. At this stage, a subset of epidermal cells and the amnioserosa cells appears to be responsible for ecdysteroid biosynthesis, as the Halloween genes are strongly expressed in these cells (Chávez *et al.* 2000; Warren *et al.* 2002, 2004; Petryk *et al.* 2003; Niwa *et al.* 2004, 2010; Namiki *et al.* 2005; Ono *et al.* 2006; Yoshiyama *et al.* 2006; Enya *et al.* 2014). Importantly, the temporal fluctuation of the Halloween gene expressions during mid-embryogenesis correlates very well with that of the embryonic ecdysteroid titer (Niwa *et al.* 2010; Enya *et al.* 2014). In the case of the adults, female ovarian follicle cells are the classically famous sites of ecdysteroid biosynthesis and indeed require ecdysteroidogenic enzymes for biosynthesis (Ono *et al.* 2006; Domanitskaya *et al.* 2014; Sieber & Spradling 2015) (T. Ameku and R.N., unpublished data). However, it is unclear which TFs regulate such embryonic and ovarian ecdysteroidogenic gene expressions, except for β Ftz-f1 that appears to regulate *dib* expression in the follicle cells (Talamillo *et al.* 2013).

We must now unravel the higher regulatory mechanisms that control ecdysteroid biosynthesis through transcription. As described above, the most important issue to draw a signaling network for controlling ecdysteroid biosynthesis is to understand which ecdysteroidogenic TFs act downstream of which extracellular

stimulus-triggered signaling pathway. As PTTH signaling promotes activated ERK phosphorylating nuclear target proteins such as TFs, it is feasible to hypothesize that PTTH signaling regulates the activity of several ecdysteroidogenic TFs including DHR4 (Ou *et al.* 2011). A proteomic approach would be helpful to identify PTTH-stimulated phosphorylated proteins in the PG in future, while a previous trial using the tobacco hornworm *Manduca sexta* did not identify any known ecdysteroidogenic TFs described above (Rewitz *et al.* 2009a).

>All of the evidence clearly demonstrates the necessity of ecdysteroidogenic TFs to control ecdysteroidogenic gene expression in the PG. By contrast, there is no reported study examining whether any ecdysteroidogenic TFs are sufficient to induce ecdysteroidogenic gene expression in non-steroidogenic cells. In the case of mammals, Ad4BP/SF-1 is both necessary and sufficient for the induction and maintenance of steroidogenic genes. For example, overexpression of Ad4BP/SF-1 differentiates cultured stem cells into steroidogenic cell lineages with the expression of various steroidogenesis-related genes (Miyamoto *et al.* 2011). Moreover, transgenic expression of Ad4BP/SF-1 in mice leads to ectopic adrenal formation (Zubair *et al.* 2009). By contrast, it is unlikely that β Ftz-f1, the insect homologue of Ad4BP/SF-1, acts as a master regulator to induce ecdysteroidogenic gene expression in the PG because β Ftz-f1 plays a crucial role in ecdysteroid-dependent transcriptional cascades not only in the PG, but also in many other tissues (Thummel 2001). Besides β Ftz-f1, other ecdysteroidogenic TFs identified to date are also highly expressed in non-ecdysteroidogenic cells and have important functions other than ecdysteroid biosynthesis. Therefore, it is an interesting open question to examine whether and how forced expression of one or more TFs can differentiate the PG cells and/or induce ecdysteroidogenic gene expression. This point might be important to comprehensively understand the evolutionary commonality of steroidogenic TF function during animal evolution.

Finally, we would like to point out that the current published studies on ecdysteroidogenic TFs have focused on their roles in regulating just a handful of identified ecdysteroidogenic enzyme genes and other known regulatory protein genes, but no studies have examined the entire transcriptome in the PG. It is possible that the set of known genes is just the tip of the iceberg of ecdysteroidogenic TF-regulating genes. Interestingly, a recent study using mice reveals that Ad4BP/SF-1 governs the coordinated regulation of not only typical steroidogenic genes, but also essential genes within a glycolytic pathway (Baba *et al.* 2014). In the future, next-generation sequencing approaches

could help to employ RNA-sequencing to comprehensively understand the ecdysteroidogenic TF-dependent regulation of gene expression profiles at the transcriptome system level.

Acknowledgments

We thank Tomotsune Ameku for valuable comments on the manuscript. This work was supported by grants to R.N. from JST/PRESTO and Japan Society for the Promotion of Science KAKENHI Grant Number 25712010.

References

- Ambros, V. 2000. Control of developmental timing in *Caenorhabditis elegans*. *Curr. Opin. Genet. Dev.* **10**, 428–433.
- Baba, T., Otake, H., Sato, T., Miyabayashi, K., Shishido, Y., Wang, C.-Y., Shima, Y., Kimura, H., Yagi, M., Ishihara, Y., Hino, S., Ogawa, H., Nakao, M., Yamazaki, T., Kang, D., Ohkawa, Y., Chung, B.-C. & Morohashi, K. 2014. Glycolytic genes are targets of the nuclear receptor Ad4BP/SF-1. *Nat. Commun.* **5**, 3634.
- Banerjee, D. & Slack, F. 2002. Control of developmental timing by small temporal RNAs: A paradigm for rna-mediated regulation of gene expression. *BioEssays* **24**, 119–129.
- Bialecki, M., Shilton, A., Fichtenberg, C., Segraves, W. A. & Thummel, C. S. 2002. Loss of the ecdysteroid-inducible E75A orphan nuclear receptor uncouples molting from metamorphosis in *Drosophila*. *Dev. Cell* **3**, 209–220.
- Bonaglia, M. C., Ciccone, R., Gimelli, G., Gimelli, S., Marelli, S., Verheij, J., Giorda, R., Grasso, R., Borgatti, R., Pagone, F., Rodríguez, L., Martínez-Frias, M.-L., van Ravenswaaij, C. & Zuffardi, O. 2008. Detailed phenotype–genotype study in five patients with chromosome 6q16 deletion: narrowing the critical region for Prader-Willi-like phenotype. *Eur. J. Hum. Genet.* **16**, 1443–1449.
- Borsos, B. N., Pankotai, T., Kovács, D., Popescu, C., Páhi, Z. & Boros, I. M. 2015. Acetylations of Ftz-F1 and histone H4K5 are required for the fine-tuning of ecdysone biosynthesis during *Drosophila* metamorphosis. *Dev. Biol.* **404**, 80–87.
- Cáceres, L., Necakov, A. S., Schwartz, C., Kimber, S., Roberts, I. J. & Krause, H. M. 2011. Nitric oxide coordinates metabolism, growth, and development via the nuclear receptor E75. *Genes Dev.* **25**, 1476–1485.
- Chávez, V. M., Marqués, G., Delbecque, J. P., Kobayashi, K., Hollingsworth, M., Burr, J., Natzle, J. E. & O'Connor, M. B. 2000. The *Drosophila disembodied* gene controls late embryonic morphogenesis and codes for a cytochrome P450 enzyme that regulates embryonic ecdysone levels. *Development* **127**, 4115–4126.
- Cheng, C., Ko, A., Chaieb, L., Koyama, T., Sarwar, P., Mirth, C. K., Smith, W. A. & Suzuki, Y. 2014. The POU factor ventral veins lacking/Drifter directs the timing of metamorphosis through ecdysteroid and juvenile hormone signaling. *PLoS Genet.* **10**, e1004425.
- Colombani, J., Bianchini, L., Layalle, S., Pondeville, E., Dauphin-Villemant, C., Antoniewski, C., Carré, C., Noselli, S. & Léopold, P. 2005. Antagonistic actions of ecdysone and insulins determine final size in *Drosophila*. *Science* **310**, 667–670.

- Danielsen, T. E., Moeller, M. E., Dorry, E., Komura-Kawa, T., Fujimoto, Y., Troelsen, J. T., Herder, R., O'Connor, M. B., Niwa, R. & Rewitz, K. F. 2014. Transcriptional control of steroid biosynthesis genes in the *Drosophila* prothoracic gland by Ventral veins lacking and Knirps. *PLoS Genet.* **10**, e1004343.
- Deng, H. & Kerppola, T. K. 2013. Regulation of *Drosophila* metamorphosis by xenobiotic response regulators. *PLoS Genet.* **9**, e1003263.
- Domanitskaya, E., Anllo, L. & Schüpbach, T. 2014. Phantom, a cytochrome P450 enzyme essential for ecdysone biosynthesis, plays a critical role in the control of border cell migration in *Drosophila*. *Dev. Biol.* **386**, 408–418.
- Enya, S., Ameku, T., Igarashi, F., Iga, M., Kataoka, H., Shinoda, T. & Niwa, R. 2014. A Halloween gene *noppera-bo* encodes a glutathione S-transferase essential for ecdysteroid biosynthesis via regulating the behaviour of cholesterol in *Drosophila*. *Sci. Rep.* **4**, 6586.
- Enya, S., Daimon, T., Igarashi, F., Kataoka, H., Uchibori, M., Sezutsu, H., Shinoda, T. & Niwa, R. 2015. The silkworm glutathione S-transferase gene *noppera-bo* is required for ecdysteroid biosynthesis and larval development. *Insect Biochem. Mol. Biol.* **61**, 1–7.
- Fresán, U., Cuartero, S., O'Connor, M. B. & Espinàs, M. L. 2015. The insulator protein CTCF regulates *Drosophila* steroidogenesis. *Biol. Open* **4**, 852–857.
- Gibbens, Y. Y., Warren, J. T., Gilbert, L. I. & O'Connor, M. B. 2011. Neuroendocrine regulation of *Drosophila* metamorphosis requires TGF β /Activin signaling. *Development* **138**, 2693–2703.
- Gilbert, L. I., Rybczynski, R. & Warren, J. T. 2002. Control and biochemical nature of the ecdysteroidogenic pathway. *Annu. Rev. Entomol.* **47**, 883–916.
- Halme, A., Cheng, M. & Hariharan, I. K. 2010. Retinoids regulate a developmental checkpoint for tissue regeneration in *Drosophila*. *Curr. Biol.* **20**, 458–463.
- Hanukoglu, I. 1992. Steroidogenic enzymes: structure, function, and role in regulation of steroid hormone biosynthesis. *J. Steroid Biochem. Mol. Biol.* **43**, 779–804.
- Hariharan, I. K. 2012. How growth abnormalities delay “puberty” in *Drosophila*. *Sci. Signal.* **5**, pe27.
- Iga, M., Nakaoka, T., Suzuki, Y. & Kataoka, H. 2014. Pigment dispersing factor regulates ecdysone biosynthesis via *Bombyx* neuropeptide G protein coupled receptor-B2 in the prothoracic glands of *Bombyx mori*. *PLoS ONE* **9**, e103239.
- Ishimoto, H., Wang, Z., Rao, Y., Wu, C.-F. & Kitamoto, T. 2013. A novel role for ecdysone in *Drosophila* conditioned behavior: Linking GPCR-mediated non-canonical steroid action to cAMP signaling in the adult brain. *PLoS Genet.* **9**, e1003843.
- Izumi, K., Housam, R., Kapadia, C., Stallings, V. A., Medne, L., Shaikh, T. H., Kublaoui, B. M., Zackai, E. H. & Grimberg, A. 2013. Endocrine phenotype of 6q16.1-q21 deletion involving SIM1 and Prader-Willi syndrome-like features. *Am. J. Med. Genet. A* **161**, 3137–3143.
- Jaszczak, J. S., Wolpe, J. B., Dao, A. Q. & Halme, A. 2015. Nitric oxide synthase regulates growth coordination during *Drosophila melanogaster* imaginal disc regeneration. *Genetics* **200**, 1219–1228.
- Komura-Kawa, T., Hirota, K., Shimada-Niwa, Y., Yamauchi, R., Shimell, M., Shinoda, T., Fukamizu, A., O'Connor, M. B. & Niwa, R. 2015. The *Drosophila* zinc finger transcription factor Ouija board controls ecdysteroid biosynthesis through specific regulation of *spookier*. *PLoS Genet.* DOI: 10.1371/journal.pgen.1005712.
- Koyama, T., Mendes, C. C. & Mirth, C. K. 2013. Mechanisms regulating nutrition-dependent developmental plasticity through organ-specific effects in insects. *Front. Physiol.* **4**, 263.
- Koyama, T., Rodrigues, M. A., Athanasiadis, A., Shingleton, A. W. & Mirth, C. K. 2014. Nutritional control of body size through FoxO-Ultraspicle mediated ecdysone biosynthesis. *Elife* **3**, e03091.
- Lee, F. Y., Faivre, E. J., Suzawa, M., Lontok, E., Ebert, D., Cai, F., Belsham, D. D. & Ingraham, H. A. 2011. Eliminating SF-1 (NR5A1) Sumoylation in vivo results in ectopic Hedgehog signaling and disruption of endocrine development. *Dev. Cell* **21**, 315–327.
- Liu, Y., Sheng, Z., Liu, H., Wen, D., He, Q., Wang, S., Shao, W., Jiang, R.-J., An, S., Sun, Y., Bendena, W. G., Wang, J., Gilbert, L. I., Wilson, T. G., Song, Q. & Li, S. 2009. Juvenile hormone counteracts the bHLH-PAS transcription factors MET and GCE to prevent caspase-dependent programmed cell death in *Drosophila*. *Development* **136**, 2015–2025.
- McBrayer, Z., Ono, H., Shimell, M., Parvy, J.-P., Beckstead, R. B., Warren, J. T., Thummel, C. S., Dauphin-Villemant, C., Gilbert, L. I. & O'Connor, M. B. 2007. Prothoracicotropic hormone regulates developmental timing and body size in *Drosophila*. *Dev. Cell* **13**, 857–871.
- Meng, M., Cheng, D., Peng, J., Qian, W., Li, J., Dai, D., Zhang, T. & Xia, Q. 2015. The homeodomain transcription factors Antennapedia and POU-M2 regulate the transcription of the steroidogenic enzyme gene Phantom in the silkworm. *J. Biol. Chem.* **290**, 24438–24452.
- Miller, W. L. 1988. Molecular biology of steroid hormone synthesis. *Endocr. Rev.* **9**, 295–318.
- Miller, W. L. & Auchus, R. J. 2011. The molecular biology, biochemistry, and physiology of human steroidogenesis and its disorders. *Endocr. Rev.* **32**, 81–151.
- Miyamoto, K., Yazawa, T., Mizutani, T., Imamichi, Y., Kawabe, S., Kanno, M., Matsumura, T., Ju, Y. & Umezawa, A. 2011. Stem cell differentiation into steroidogenic cell lineages by NR5A family. *Mol. Cell. Endocrinol.* **336**, 123–126.
- Mizoguchi, A., Ohashi, Y., Hosoda, K., Ishibashi, J. & Kataoka, H. 2001. Developmental profile of the changes in the prothoracicotropic hormone titer in hemolymph of the silkworm *Bombyx mori*: Correlation with ecdysteroid secretion. *Insect Biochem. Mol. Biol.* **31**, 349–358.
- Mizoguchi, A., Kamimura, M., Kiuchi, M. & Kataoka, H. 2015. Positive feedback regulation of prothoracicotropic hormone secretion by ecdysteroid—a mechanism that determines the timing of metamorphosis. *Insect Biochem. Mol. Biol.* **58**, 39–45.
- Mizutani, T., Ishikane, S., Kawabe, S., Umezawa, A. & Miyamoto, K. 2015. Transcriptional regulation of genes related to progesterone production. *Endocr. J.* **62**, 757–763.
- Moeller, M. E., Danielsen, E. T., Herder, R., O'Connor, M. B. & Rewitz, K. F. 2013. Dynamic feedback circuits function as a switch for shaping a maturation-inducing steroid pulse in *Drosophila*. *Development* **140**, 4730–4739.
- Morioka, E., Matsumoto, A. & Ikeda, M. 2012. Neuronal influence on peripheral circadian oscillators in pupal *Drosophila* prothoracic glands. *Nat. Commun.* **3**, 909.
- Morohashi, K., Baba, T. & Tanaka, M. 2013. Steroid hormones and the development of reproductive organs. *Sex Dev.* **7**, 61–79.
- Myers, E. M., Yu, J. & Sehgal, A. 2003. Circadian control of eclosion: Interaction between a central and peripheral clock in *Drosophila melanogaster*. *Curr. Biol.* **13**, 526–533.
- Namiki, T., Niwa, R., Sakudoh, T., Shirai, K.-I., Takeuchi, H. & Kataoka, H. 2005. Cytochrome P450 CYP307A1/Spook: a

- regulator for ecdysone synthesis in insects. *Biochem. Biophys. Res. Commun.* **337**, 367–374.
- Namiki, T., Niwa, R., Higuchi, A., Yoshiyama, T., Mita, K. & Kataoka, H. 2009. A basic-HLH transcription factor, *HLH54F*, is highly expressed in the prothoracic gland in the silkworm *Bombyx mori* and the fruit fly *Drosophila melanogaster*. *Biosci. Biotechnol. Biochem.* **73**, 762–765.
- Neubueser, D., Warren, J. T., Gilbert, L. I. & Cohen, S. M. 2005. *molting defective* is required for ecdysone biosynthesis. *Dev. Biol.* **280**, 362–372.
- Niwa, R. & Niwa, Y. S. 2014a. Enzymes for ecdysteroid biosynthesis: their biological functions in insects and beyond. *Biosci. Biotechnol. Biochem.* **78**, 1283–1292.
- Niwa, Y. S. & Niwa, R. 2014b. Neural control of steroid hormone biosynthesis during development in the fruit fly *Drosophila melanogaster*. *Genes Genet. Syst.* **89**, 27–34.
- Niwa, R., Matsuda, T., Yoshiyama, T., Namiki, T., Mita, K., Fujimoto, Y. & Kataoka, H. 2004. CYP306A1, a cytochrome P450 enzyme, is essential for ecdysteroid biosynthesis in the prothoracic glands of *Bombyx* and *Drosophila*. *J. Biol. Chem.* **279**, 35942–35949.
- Niwa, R., Sakudoh, T., Namiki, T., Saida, K., Fujimoto, Y. & Kataoka, H. 2005. The ecdysteroidogenic P450 *Cyp302a1/disembodied* from the silkworm, *Bombyx mori*, is transcriptionally regulated by prothoracicotrophic hormone. *Insect Mol. Biol.* **14**, 563–571.
- Niwa, R., Niimi, T., Honda, N., Yoshiyama, M., Itoyama, K., Kataoka, H. & Shinoda, T. 2008. Juvenile hormone acid O-methyltransferase in *Drosophila melanogaster*. *Insect Biochem. Mol. Biol.* **38**, 714–720.
- Niwa, R., Namiki, T., Ito, K., Shimada-Niwa, Y., Kiuchi, M., Kawaoka, S., Kayukawa, T., Banno, Y., Fujimoto, Y., Shigenobu, S., Kobayashi, S., Shimada, T., Katsuma, S. & Shinoda, T. 2010. *Non-molting glossy/shroud* encodes a short-chain dehydrogenase/reductase that functions in the “Black Box” of the ecdysteroid biosynthesis pathway. *Development* **137**, 1991–1999.
- Ogihara, M. H., Hikiba, J., Iga, M. & Kataoka, H. 2015. Negative regulation of juvenile hormone analog for ecdysteroidogenic enzymes. *J. Insect Physiol.* **80**, 42–47.
- Ohara, Y., Kayashima, Y., Hayashi, Y., Kobayashi, S. & Yamakawa-Kobayashi, K. 2012. Expression of β -adrenergic-like octopamine receptors during *Drosophila* development. *Zoolog. Sci.* **29**, 83–89.
- Ohara, Y., Shimada-Niwa, Y., Niwa, R., Kayashima, Y., Hayashi, Y., Akagi, K., Ueda, H., Yamakawa-Kobayashi, K. & Kobayashi, S. 2015. Autocrine regulation of ecdysone synthesis by β 3-octopamine receptor in the prothoracic gland is essential for *Drosophila* metamorphosis. *Proc. Natl Acad. Sci. USA* **112**, 1452–1457.
- Ono, H. 2014. Ecdysone differentially regulates metamorphic timing relative to 20-hydroxyecdysone by antagonizing juvenile hormone in *Drosophila melanogaster*. *Dev. Biol.* **391**, 32–42.
- Ono, H., Rewitz, K. F., Shinoda, T., Itoyama, K., Petryk, A., Rybczynski, R., Jarcho, M., Warren, J. T., Marqués, G., Shimmell, M. J., Gilbert, L. I. & O'Connor, M. B. 2006. *Spook* and *Spookier* code for stage-specific components of the ecdysone biosynthetic pathway in Diptera. *Dev. Biol.* **298**, 555–570.
- Ou, Q. & King-Jones, K. 2013. What goes up must come down. Transcription factors have their say in making ecdysone pulses. *Curr. Top. Dev. Biol.* **103**, 35–71.
- Ou, Q., Magico, A. & King-Jones, K. 2011. Nuclear receptor DHR4 controls the timing of steroid hormone pulses during *Drosophila* development. *PLoS Biol.* **9**, e1001160.
- Pankotai, T., Popescu, C., Martín, D., Grau, B., Zsindely, N., Bodai, L., Tora, L., Ferrús, A. & Boros, I. 2010. Genes of the ecdysone biosynthesis pathway are regulated by the dATAC histone acetyltransferase complex in *Drosophila*. *Mol. Cell. Biol.* **30**, 4254–4266.
- Parker, K. L., Rice, D. A., Lala, D. S., Ikeda, Y., Luo, X., Wong, M., Bakke, M., Zhao, L., Frigeri, C., Hanley, N. A., Stallings, N. & Schimmer, B. P. 2002. Steroidogenic factor 1: an essential mediator of endocrine development. *Recent Prog. Horm. Res.* **57**, 19–36.
- Parvy, J. P., Blais, C., Bernard, F., Warren, J. T., Petryk, A., Gilbert, L. I., O'Connor, M. B. & Dauphin-Villemant, C. 2005. A role for β FTZ-F1 in regulating ecdysteroid titers during post-embryonic development in *Drosophila melanogaster*. *Dev. Biol.* **282**, 84–94.
- Parvy, J.-P., Wang, P., Garrido, D., Maria, A., Blais, C., Poidevin, M. & Montagne, J. 2014. Forward and feedback regulation of cyclic steroid production in *Drosophila melanogaster*. *Development* **141**, 3955–3965.
- Petryk, A., Warren, J. T., Marqués, G., Jarcho, M. P., Gilbert, L. I., Kahler, J., Parvy, J.-P., Li, Y., Dauphin-Villemant, C. & O'Connor, M. B. 2003. Shade is the *Drosophila* P450 enzyme that mediates the hydroxylation of ecdysone to the steroid insect molting hormone 20-hydroxyecdysone. *Proc. Natl Acad. Sci. USA* **100**, 13773–13778.
- Reinking, J., Lam, M. M. S., Pardee, K., Sampson, H. M., Liu, S., Yang, P., Williams, S., White, W., Lajoie, G., Edwards, A. & Krause, H. M. 2005. The *Drosophila* nuclear receptor E75 contains heme and is gas responsive. *Cell* **122**, 195–207.
- Rewitz, K. F., O'Connor, M. B. & Gilbert, L. I. 2007. Molecular evolution of the insect Halloween family of cytochrome P450s: Phylogeny, gene organization and functional conservation. *Insect Biochem. Mol. Biol.* **37**, 741–753.
- Rewitz, K. F., Larsen, M. R., Lobner-Olesen, A., Rybczynski, R., O'Connor, M. B. & Gilbert, L. I. 2009a. A phosphoproteomics approach to elucidate neuropeptide signal transduction controlling insect metamorphosis. *Insect Biochem. Mol. Biol.* **39**, 475–483.
- Rewitz, K. F., Yamanaka, N., Gilbert, L. I. & O'Connor, M. B. 2009b. The insect neuropeptide PTTH activates receptor tyrosine kinase Torso to initiate metamorphosis. *Science* **326**, 1403–1405.
- Riddiford, L. M., Truman, J. W., Mirth, C. K. & Shen, Y.-C. 2010. A role for juvenile hormone in the prepupal development of *Drosophila melanogaster*. *Development* **137**, 1117–1126.
- Rougvie, A. E. 2005. Intrinsic and extrinsic regulators of developmental timing: from miRNAs to nutritional cues. *Development* **132**, 3787–3798.
- Sánchez-Higueras, C., Sotillos, S. & Castelli-Gair Hombría, J. 2014. Common origin of insect trachea and endocrine organs from a segmentally repeated precursor. *Curr. Biol.* **24**, 76–81.
- Shimada-Niwa, Y. & Niwa, R. 2014. Serotonergic neurons respond to nutrients and regulate the timing of steroid hormone biosynthesis in *Drosophila*. *Nat. Commun.* **5**, 5778.
- Sieber, M. H. & Spradling, A. C. 2015. Steroid signaling establishes a female metabolic state and regulates SREBP to control oocyte lipid accumulation. *Curr. Biol.* **25**, 1–12.
- Simon, A. F., Shih, C., Mack, A. & Benzer, S. 2003. Steroid control of longevity in *Drosophila melanogaster*. *Science* **299**, 1407–1410.
- Takaki, K. & Sakurai, S. 2003. Regulation of prothoracic gland ecdysteroidogenic activity leading to pupal metamorphosis. *Insect Biochem. Mol. Biol.* **33**, 1189–1199.

- Talamillo, A., Sánchez, J., Cantera, R., Pérez, C., Martín, D., Caminero, E. & Barrio, R. 2008. Smt3 is required for *Drosophila melanogaster* metamorphosis. *Development* **135**, 1659–1668.
- Talamillo, A., Herboso, L., Pirone, L., Pérez, C., González, M., Sánchez, J., Mayor, U., Lopitz-Otsoa, F., Rodríguez, M. S., Sutherland, J. D. & Barrio, R. 2013. Scavenger receptors mediate the role of SUMO and Ftz-f1 in *Drosophila* steroidogenesis. *PLoS Genet.* **9**, e1003473.
- Tanaka, Y. 2011. Recent topics on the regulatory mechanism of ecdysteroidogenesis by the prothoracic glands in insects. *Front. Endocrinol.* **2**, 107.
- Thummel, C. S. 2001. Molecular mechanisms of developmental timing in *C. elegans* and *Drosophila*. *Dev. Cell* **1**, 453–465.
- Walker, V. K., Watson, K. L., Holden, J. J. & Steel, C. G. H. 1987. Vitellogenesis and fertility in *Drosophila* females with low ecdysteroid titres; the *l(3)3DTS* mutation. *J. Insect Physiol.* **33**, 137–142.
- Warren, J. T., Wismar, J., Subrahmanyam, B. & Gilbert, L. I. 2001. *Woc (without children)* gene control of ecdysone biosynthesis in *Drosophila melanogaster*. *Mol. Cell. Endocrinol.* **181**, 1–14.
- Warren, J. T., Petryk, A., Marque, G., Jarcho, M., Parvy, J.-P., Dauphin-villemant, C., O'Connor, M. B. & Gilbert, L. I. 2002. Molecular and biochemical characterization of two P450 enzymes in the ecdysteroidogenic pathway of *Drosophila melanogaster*. *Proc. Natl Acad. Sci. USA* **99**, 11043–11048.
- Warren, J. T., Petryk, A., Marqués, G., Parvy, J.-P., Shinoda, T., Itoyama, K., Kobayashi, J., Jarcho, M., Li, Y., O'Connor, M. B., Dauphin-Villemant, C. & Gilbert, L. I. 2004. Phantom encodes the 25-hydroxylase of *Drosophila melanogaster* and *Bombyx mori*: a P450 enzyme critical in ecdysone biosynthesis. *Insect Biochem. Mol. Biol.* **34**, 991–1010.
- Warren, J. T., Yerushalmi, Y., Shimell, M. J., O'Connor, M. B., Restifo, L. L. & Gilbert, L. I. 2006. Discrete pulses of molting hormone, 20-hydroxyecdysone, during late larval development of *Drosophila melanogaster*: Correlations with changes in gene activity. *Dev. Dyn.* **235**, 315–326.
- Wen, D., Rivera-Perez, C., Abdou, M., Jia, Q., He, Q., Liu, X., Zyaan, O., Xu, J., Bendena, W. G., Tobe, S. S., Noriega, F. G., Palli, S. R., Wang, J. & Li, S. 2015. Methyl farnesoate plays a dual role in regulating *Drosophila* metamorphosis. *PLoS Genet.* **11**, e1005038.
- Wildemann, B. & Bicker, G. 1999. Developmental expression of nitric oxide/cyclic GMP synthesizing cells in the nervous system of *Drosophila melanogaster*. *J. Neurobiol.* **38**, 1–15.
- Wismar, J., Habtemichael, N., Warren, J. T., Dai, J. D., Gilbert, L. I. & Gateff, E. 2000. The mutation *without children^{9f}* causes ecdysteroid deficiency in third-instar larvae of *Drosophila melanogaster*. *Dev. Biol.* **226**, 1–17.
- Xiang, Y., Liu, Z. & Huang, X. 2010. *br* regulates the expression of the ecdysone biosynthesis gene *npc1*. *Dev. Biol.* **344**, 800–808.
- Xie, X.-J., Hsu, F.-N., Gao, X., Xu, W., Ni, J.-Q., Xing, Y., Huang, L., Hsiao, H.-C., Zheng, H., Wang, C., Zheng, Y., Xiaoli, A. M., Yang, F., Bondos, S. E. & Ji, J.-Y. 2015. CDK8-Cyclin C mediates nutritional regulation of developmental transitions through the ecdysone receptor in *Drosophila*. *PLoS Biol.* **13**, e1002207.
- Yamanaka, N., Honda, N., Osato, N., Niwa, R., Mizoguchi, A. & Kataoka, H. 2007. Differential regulation of ecdysteroidogenic P450 gene expression in the silkworm, *Bombyx mori*. *Biosci. Biotechnol. Biochem.* **71**, 2808–2814.
- Yamanaka, N., Rewitz, K. F. & O'Connor, M. B. 2013a. Ecdysone control of developmental transitions: lessons from *Drosophila* research. *Annu. Rev. Entomol.* **58**, 497–516.
- Yamanaka, N., Romero, N. M., Martin, F. A., Rewitz, K. F., Sun, M., O'Connor, M. B. & Léopold, P. 2013b. Neuroendocrine control of *Drosophila* larval light preference. *Science* **341**, 1113–1116.
- Yoshiyama, T., Namiki, T., Mita, K., Kataoka, H. & Niwa, R. 2006. Neverland is an evolutionally conserved Rieske-domain protein that is essential for ecdysone synthesis and insect growth. *Development* **133**, 2565–2574.
- Yoshiyama-Yanagawa, T., Enya, S., Shimada-Niwa, Y., Yaguchi, S., Haramoto, Y., Matsuya, T., Shiomi, K., Sasakura, Y., Takahashi, S., Asashima, M., Kataoka, H. & Niwa, R. 2011. The conserved Rieske oxygenase DAF-36/Neverland is a novel cholesterol-metabolizing enzyme. *J. Biol. Chem.* **286**, 25756–25762.
- Young, S. C., Yeh, W. L. & Gu, S. H. 2012. Transcriptional regulation of the PTTH receptor in prothoracic glands of the silkworm, *Bombyx mori*. *J. Insect Physiol.* **58**, 102–109.
- Zhou, X., Zhou, B., Truman, J. W. & Riddiford, L. M. 2004. Overexpression of *broad*: a new insight into its role in the *Drosophila* prothoracic gland cells. *J. Exp. Biol.* **207**, 1151–1161.
- Zubair, M., Oka, S., Parker, K. & Morohashi, K. 2009. Transgenic expression of Ad4BP/SF-1 in fetal adrenal progenitor cells leads to ectopic adrenal formation. *Mol. Endocrinol.* **23**, 1657–1667.