CORRECTION Open Access

Correction: Three-dimensional visual technique based on CT lymphography data combined with methylene blue in endoscopic sentinel lymph node biopsy for breast cancer

Baiye Wang^{1†}, Caifeng Ou^{2,3†}, Jingang Yu⁴, Jianping Ye⁵, Yunfeng Luo³, Yu Wang⁶ and Pusheng Zhang^{3*}

Correction: European Journal of Medical Research (2024) 27:274

https://doi.org/10.1186/s40001-022-00909-3

In the original article [1], the image of the patient in Fig. 1 has the patient's face visible. It requires explicit informed consent for identifiable details or images of research participants to be published and not satisfied that this has been met for this case as the eye bar provides sufficient anonymisation to the patient.

The authors have provided revised image (Fig. 1).

[†]Baiye Wang and Caifeng Ou contributed equally to this work and should be considered co-first authors

The original article can be found online at https://doi.org/10.1186/s40001-022-00909-3.

*Correspondence:

Pusheng Zhang

drpushengz@163.com

¹ Department of Radiology, Zhujiang Hospital, Southern Medical University, Guangzhou, Guangdong, China

² Present Address: Department of Breast Care Surgery, The First Affiliated Hospital of Guangdong Pharmaceutical University, Guangzhou 510080, Guangdong, China

³ Department of Breast Surgery, Zhujiang Hospital, Southern Medical University, 253, Gongye Dadao Zhong, Haizhu District, Guangzhou 510282, Guangdong, China

⁴ School of Automation Science and Engineering, South China University of Technology, Guangzhou, Guangdong, China

Shenzhen Smart Vision Co. LTD., Shenzhen, Guangdong, China

⁶ Department of Pathology, Zhujiang Hospital, Southern Medical University, Guangzhou, Guangdong, China



Fig. 1 Injection of the contrast agent. 4 mL of lymphatic contrast agent was intradermally injected into the periareolar region in four (clock-wise) quadrants of the breast before CT lymphography

The original article has been corrected.



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/joublicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Published: 5 November 2024

Reference

 Wang B, Ou C, Yu J, Ye J, Luo Y, Wang Y, Zhang P. Three-dimensional visual technique based on CT lymphography data combined with methylene blue in endoscopic sentinel lymph node biopsy for breast cancer. Eur J Med Res. 2022;27:274.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.