



Published in final edited form as:

*J Food Prot.* 2016 November ; 79(11): 1891–1897. doi:10.4315/0362-028X.JFP-16-195.

## Molecular Subtyping and Source Attribution of *Campylobacter* Isolated from Food Animals

GREGORY H. TYSON<sup>1,\*</sup>, HEATHER P. TATE<sup>1</sup>, JASON ABBOTT<sup>1</sup>, THU-THUY TRAN<sup>1</sup>, CLAUDINE KABERA<sup>1</sup>, EMILY CRAREY<sup>1</sup>, SHENIA YOUNG<sup>1</sup>, PATRICK F. MCDERMOTT<sup>1</sup>, GRISELLE SPRAGUE<sup>2</sup>, MARK CAMPBELL<sup>2</sup>, OYEWOLE ADEYEMO<sup>2</sup>, JOHNETTE BROWNE-SILVA<sup>2</sup>, MICHAEL MYERS<sup>2</sup>, SUTAWEE THITARAM<sup>2</sup>, SHAOHUA ZHAO<sup>1</sup>

<sup>1</sup>U.S. Food and Drug Administration, Center for Veterinary Medicine, 8401 Muirkirk Road, Laurel, Maryland 20708;

<sup>2</sup>U.S. Department of Agriculture, Food Safety and Inspection Service, 950 College Station Road, Athens, Georgia 30601, USA

### Abstract

*Campylobacter* spp. commonly cause gastrointestinal illness in humans. Poultry meats have long been considered the predominant source of these infections, but few in-depth *Campylobacter* source attribution studies have been completed. We analyzed more than 1,300 *Campylobacter* isolates recovered from a number of animal and food sources, including dairy and beef cattle, pigs, poultry, and retail poultry meat, and compared them with *Campylobacter* isolates recovered from human clinical samples. Each isolate was subtyped using pulsed-field gel electrophoresis (PFGE) with *Sma*I and queried against the Centers for Disease Control and Prevention PulseNet database to identify human isolates with indistinguishable patterns. Half (49.5%) of the PFGE patterns from poultry animal and retail meat isolates were indistinguishable from patterns of at least one human isolate. Among the isolates from beef and dairy cows, 56.6 and 65.0%, respectively, of their PFGE patterns were indistinguishable from those of human isolates. Only a small portion of the PFGE patterns of *Campylobacter* isolated from pigs (9.5%) were found to have PFGE patterns in common with human isolates. These data imply that cattle may be larger contributors to *Campylobacter* infections than previously recognized and help further our understanding of potential sources of human campylobacteriosis.

### Keywords

*Campylobacter*; Food animals; Pulsed-field gel electrophoresis; Source attribution

---

*Campylobacter* spp. are among the most common bacterial causes of foodborne illness in the United States, causing approximately 1 million illnesses each year (21), with approximately 90% of human infections attributed to *Campylobacter jejuni* (12). Most infections result in self-limiting gastroenteritis; however, some cases can progress to chronic sequelae such as Guillain-Barré syndrome, irritable bowel syndrome, and reactive arthritis (9). Therefore,

---

\* Author for correspondence. Fax: 301-210-4685; Gregory.Tyson@fda.hhs.gov.

identifying sources of *Campylobacter* is key to preventing its spread and ensuring a safe food supply.

Source attribution studies are critical to understanding the predominant sources of human illnesses and to determining where to direct resources to prevent the consumption of contaminated food. To perform source attribution studies, researchers need to discriminate and subtype bacterial strains to allow the direct comparison of isolates from human cases and source materials. Multilocus sequence typing has been used previously to differentiate *Campylobacter* isolates based on sequence polymorphisms in discrete genes (6). Although multilocus sequence typing data generally correlate well with data acquired using pulsed-field gel electrophoresis (PFGE) (29), PFGE has been found to have superior discriminatory power (20).

Food animals are significant reservoirs of *Campylobacter*. These bacteria have been found associated with pig, cattle, and poultry sources (3); however, most attribution studies have identified chickens as the major source of infection. In the United States, *Campylobacter* contaminates approximately 50% of retail chicken meat, with significantly less found in pork, ground turkey, and ground beef retail products (32). In Europe, as many as 80% of human *Campylobacter* infections have been thought to be associated with the consumption of chicken products (7), with some case-control studies in the United States supporting this conclusion (8). However, PFGE profile comparisons of *Campylobacter* isolates from retail poultry meats and humans have revealed significant differences (26), indicating the likelihood of additional *Campylobacter* reservoirs that contribute to human infection. In support of this, some studies have suggested that a large number of animal, food, and environmental reservoirs are important contributors to *Campylobacter* infection (30).

To better evaluate potential sources of human *Campylobacter* infections in the United States, we analyzed more than 1,300 *Campylobacter* isolates recovered from cecal samples of dairy and beef cattle, pigs, chickens, and turkeys, in addition to isolates recovered from retail poultry meats. Using PFGE to discriminate strains, we compared patterns from these isolates with those from human patients.

## MATERIALS AND METHODS

### Cecal sampling and *Campylobacter* isolation.

Cecal contents from pigs, cattle, chicken, and turkey animals were collected by U.S. Department of Agriculture, Food Safety and Inspection Service (USDA-FSIS) personnel from animals presented for slaughter at USDA-FSIS-inspected facilities throughout the United States in 2013 (28). Cecal samples were shipped to the Eastern USDA-FSIS laboratory, where samples were cultured for *Campylobacter* using published methods (15). Briefly, cecal samples suspended in buffered peptone water were enriched with Bolton enrichment broth for 48 h before samples were streaked onto *Campylobacter*-modified charcoal-cefoperazone-deoxycho-late agar plates for isolation. An attempt was made to obtain at least 300 isolates each from pig, dairy cattle, beef cattle, and chicken and turkey sources. All the pig and poultry isolates were analyzed, but the beef and dairy cattle isolates that were analyzed were randomly selected from a larger number of isolates. In the absence

of the requisite number of poultry isolates, we added poultry retail meat isolates obtained in 2013 by the National Antimicrobial Resistance Monitoring System retail meat surveillance (28) to the analysis. *Campylobacter* speciation was performed using a PCR-based assay, as described in a previous article (31).

### PFGE.

PFGE was performed per standardized protocols, in accordance with Centers for Disease Control and Prevention (CDC) PulseNet procedures (26). Briefly, *Campylobacter* genomic DNA was subjected to digestion by *Sma*I, electrophoresis, and GelRed 3X stain (Phenix Research, Candler, NC). The band patterns were analyzed using BioNumerics version 6.6 (Applied Maths, Austin, TX). We performed the analysis using 1.5% optimization and 1.5% band matching tolerance, with an unweighted pair group method with arithmetic mean-based cluster analysis. We compared the PFGE patterns of *Campylobacter* isolates to those from the CDC PulseNet database (24), which contains isolates obtained nationwide from human patients (10,320 isolates for *C. jejuni* and 413 for *Campylobacter coli* at time of analysis). Patterns were identified as indistinguishable using PFGE if they were identical according to this analysis. The same analysis criteria were also used in comparing the patterns from this study with each other. We used the program JVenn to display the number of PFGE patterns shared among sources (Fig. 1) (2).

### Statistical analysis.

We analyzed the results using a chi-square test of independence (or Fisher's exact test, where appropriate) to determine whether the differences were significant among sources in the percentage of indistinguishable patterns or isolates and in the percentage of isolates that were *C. jejuni*. We made additional pairwise comparisons using the Bonferroni correction to see which source accounted for more of the chi-square statistic. *P* values of less than 0.05 were denoted as significant.

## RESULTS

To better understand the sources of the *Campylobacter* causing human infections, we selected 1,363 isolates from various sources for study. The goal was to compare at least 300 isolates each from the ceca of poultry, beef cattle, dairy cattle, and pigs with isolates from humans. All the poultry and pig isolates were selected, but the isolates we analyzed from beef and dairy cattle were randomly selected from different isolation dates to represent samples isolated from throughout the year because a larger number was available. The frequency of *Campylobacter* isolation was highest among the cecal samples collected from dairy cattle (42.6%), beef cattle (41.5%), and pigs (31.7%). In contrast, *Campylobacter* prevalence was lower among poultry, with only 21.8% of chickens and 9.5% of turkeys found to be positive. As a result, we obtained only 79 poultry animal isolates, so we collected an additional 230 isolates from poultry retail meats during the same time period to include in the analysis. Among the isolates selected for study, 721 (52.8%) were *C. jejuni* and 636 (46.7%) were *C. coli*. Five isolates were *Campylobacter lari*, and one was *Campylobacter fetus*. The frequency of isolation of each *Campylobacter* species differed greatly by source (Table 1). For instance, *C. jejuni* predominated among beef cattle (76.4%),

dairy cattle (74.9%), and to a lesser extent, poultry animals and retail meat (62.8%). In contrast, *C. coli* constituted the vast majority of the isolates from pigs (97.8%), with only 1.6% of pig isolates being *C. jejuni*. The differences in frequency of *C. jejuni* isolated from the sources were significant for all sources ( $P < 0.05$ ), except for beef and dairy cattle.

To subtype strains and perform source attribution analysis, we performed PFGE on each isolate, comparing the patterns with each other as well as with those from human isolates in the CDC PulseNet database. There was considerable diversity in the PFGE patterns isolated from each source. For instance, among the 309 poultry animal and retail meat isolates, there were 196 distinct PFGE patterns (Table 2). Less diversity was observed among the isolates obtained from cattle, with a total of 129 and 157 PFGE patterns corresponding to the 318 beef and 371 dairy cattle isolates, respectively. The pig isolates had the greatest diversity of PFGE patterns, with 283 total patterns among 365 isolates.

Despite the strain differences in *Campylobacter* from different sources, there were some PFGE patterns shared by multiple sources (Fig. 1). In total, 69 of 679 total patterns (10.2%) were shared among multiple sources, with just two patterns shared among all four sources (Fig. 1). Many patterns were shared between the isolates from beef and dairy cattle, with a total of 52 PFGE patterns in common. This contrasted with the poultry and pig isolates, which had relatively few patterns shared with either each other or with the cattle isolates (Fig. 1). Many isolates clustered into groups by isolation source; the representative PFGE patterns depicting these differences are shown in Figure 2.

Each of the PFGE patterns from the animal and poultry meat sources was compared with those present in the CDC PulseNet database, which includes patterns from human isolates obtained from both sporadic and outbreak infections. This analysis showed that 83.9% of *C. jejuni* and 32.1% of *C. coli* isolates had PFGE patterns that were indistinguishable from patterns from human isolates (Table 1). The difference between the two species was expected because *C. coli* is less common as a cause of human illness than *C. jejuni* (28). This is also reflected by the fact that, at the time of analysis, CDC PulseNet contained only about 400 *C. coli* human isolates with PFGE patterns, compared with more than 10,000 *C. jejuni* human isolates.

The different animal sources had substantial disparities when their PFGE patterns were compared to those from human isolates. Among the 196 PFGE patterns from poultry animal and retail meat isolates, 97 (49.5%) were indistinguishable from those of human isolates in the CDC PulseNet database (Table 2). Each of the 19 most common poultry patterns, containing at least three isolates per cluster, had corresponding patterns among human isolates. As a result, most *Campylobacter* isolates from poultry animals and retail meat (64.1%) had PFGE patterns indistinguishable from those of bacteria isolated from human patients (Table 1). In contrast, among the pig isolates only 9.5% of the PFGE patterns and 13.7% of all isolates had patterns indistinguishable from those of human isolates (Tables 1 and 2). Only 5 of the 14 common patterns (present in at least three pig strains) were indistinguishable from those of human isolates in the CDC PulseNet database. This suggests that pigs may be a less common source of *Campylobacter* causing human infections than poultry sources; however, it may also be attributable to the fact that nearly all the isolates

from pigs were *C. coli*. For the cattle isolates, the majority of patterns were indistinguishable from those of the human isolates, comprising 56.6% of beef cattle and 65.0% of dairy cattle patterns (Table 2). Moreover, all the most common patterns from the isolates from each cattle source (23 patterns from beef cattle and 34 from dairy cattle sources) had patterns indistinguishable from those of human isolates. As a result, the beef and dairy cattle isolates had the highest correspondence with the human isolate PFGE profiles (80.5 and 83.6%, respectively) (Table 1). These percentages of isolates with indistinguishable patterns for the cattle isolates were significantly greater than those from any of the other sources ( $P < 0.0001$ ), regardless of the *Campylobacter* species (Table 1).

One recent public health concern is the emergence of a *Campylobacter* clone associated with higher virulence, called clone SA (19). This clone, designated DBRS16.0008 (18), causes abortions in sheep as well as increased pathogenesis in mammalian hosts (4). The PFGE pattern corresponding to this clone was present in the CDC PulseNet database and was one of the most common PFGE patterns found in our collection, encompassing 37 total isolates: 1 from retail poultry meat, 2 from pigs, 14 from beef cattle, and 20 from dairy cattle (Fig. 3). Although these isolates are from only a single clonal group, these results suggest that some cattle isolates may be associated with greater pathogenesis in human hosts.

## DISCUSSION

Very few large-scale animal sampling studies of *Campylobacter* have been conducted for source attribution. We found that chickens, thought to be a major source of *Campylobacter*, had a relatively low prevalence of *Campylobacter* in their ceca despite the high contamination rate of poultry retail meats. The low cecal colonization rates were consistent with some previous studies (23), suggesting that cross-contamination from processing facilities may be responsible for the higher colonization in retail meats (13). In contrast to the low *Campylobacter* prevalence in poultry animals, we isolated much more *Campylobacter* from both cattle and pigs. Nevertheless, retail meat surveillance of ground beef and pork chops has found less than 1% of each meat type is contaminated with *Campylobacter* (27); therefore, these meat types presumably pose a lower risk for *Campylobacter* exposure. This could account for some of our results because only a minority of *Campylobacter* from pig ceca had patterns indistinguishable from those of human patients (9.5% of the patterns and 13.7% of the isolates). Over 80% of isolates from beef cattle ceca, however, had PFGE patterns indistinguishable from those of human patient isolates, despite the rare *Campylobacter* contamination of retail beef. This high frequency of indistinguishable patterns may be explained by the fact that beef cattle isolate PFGE patterns often clustered with those from dairy cattle, which may be sources of *Campylobacter* associated with the consumption of raw milk or dairy products (16).

Although chicken meat has long been considered the principal source of *Campylobacter* infections, other important sources have been identified based on outbreak investigations. In a joint U.S. Food and Drug Administration, CDC, and USDA retrospective epidemiological study (17), approximately 67% of *Campylobacter* outbreak infections were attributed to milk and dairy products. A more recent report (11) based on *Campylobacter* outbreaks from 1998 to 2012 estimated that 66% of outbreaks were attributable to the consumption

of dairy products (90% credibility interval, 57 to 74%). The large proportion of cases associated with dairy products is mostly attributable to the consumption of raw milk because *Campylobacter* is effectively killed by pasteurization (1). Additionally, cattle may also be a source through water and environmental dissemination (5, 22). Despite the fact that these reports focused solely on *Campylobacter* outbreaks and not sporadic infections, these data indicate the importance of raw milk and other nonpoultry sources as significant contributors to *Campylobacter* infection.

The results of this study emphasize the continued relevance of PFGE in performing source attribution studies. Although there is some disagreement about whether PFGE is more discriminatory than multilocus sequence typing (20, 25), the vast amount of *Campylobacter* PFGE data from human clinical samples has proven valuable (14). Whole-genome sequencing is in the process of transforming outbreak detection and source attribution, although most outbreaks are still initially detected using PFGE, with subsequent whole-genome sequencing helping with trace-back analyses (10).

Despite the significant results of this study, there are some limitations to our analysis. For instance, the data are based solely on the presence of PFGE patterns from animal sources that are indistinguishable from human sources. This does not unambiguously assert that a given infection was caused by the consumption of a certain product because consumption data from the human disease cases was not available. Also, the poultry and pig isolates had a greater PFGE pattern diversity, making it potentially less likely for human sampling to result in as many pattern matches. In addition, for patterns shared by multiple sources, it is unclear which source was predominantly responsible for the human illnesses. The implementation of newer technologies, such as whole-genome sequencing, may provide sufficient resolution to perform these analyses. Additional case control studies focusing on exposures associated with human illnesses caused by specific PFGE patterns are necessary to strengthen the conclusions made in this study.

Several features of *Campylobacter* have impeded our ability to conduct thorough source attribution studies to date. Because most infections are sporadic, it is often not possible to identify the food vehicles causing the illness. The fastidious growth requirements of the genus also make it challenging to recover sufficient isolate numbers from food sources to make adequate comparisons. Applying common cecal culture methods resulted in high recovery rates and provided the number of isolates we needed to conduct the study reported here. As described previously, PFGE has been helpful for determining source attribution and investigating common source outbreaks of illness, and this study demonstrates that increased surveillance of *Campylobacter* from different sources can lead to new insights into the origins of human infections.

## ACKNOWLEDGMENTS

We thank the CDC PulseNet Task Force for curating the PulseNet database. The views expressed in this article are those of the author(s) and do not necessarily reflect the official policy of the Department of Health and Human Services, the U.S. Food and Drug Administration, or the U.S. Government. Reference to any commercial materials, equipment, or process does not in any way constitute approval, endorsement, or recommendation by the Food and Drug Administration.

## REFERENCES


1. American Academy of Pediatrics, Committee on Infectious Diseases and Committee on Nutrition. 2014. Consumption of raw or unpasteurized milk and milk products by pregnant women and children. *Pediatrics* 133:175–179. [PubMed: 24344105]
2. Bardou P, Mariette J, Escudie F, Djemiel C, and Klopp C. 2014. jvenn: an interactive Venn diagram viewer. *BMC Bioinformatics* 15:293. [PubMed: 25176396]
3. Boes J, Nersting L, Nielsen EM, Kranker S, Enoe C, Wachmann HC, and Baggesen DL. 2005. Prevalence and diversity of *Campylobacter jejuni* in pig herds on farms with and without cattle or poultry. *J. Food Prot* 68:722–727. [PubMed: 15830662]
4. Burrough ER, Sahin O, Plummer PJ, Zhang Q, and Yaeger MJ. 2009. Pathogenicity of an emergent, ovine abortifacient *Campylobacter jejuni* clone orally inoculated into pregnant guinea pigs. *Am. J. Vet. Res* 70:1269–1276. [PubMed: 19795942]
5. Clark CG, Price L, Ahmed R, Woodward DL, Melito PL, Rodgers FG, Jamieson F, Ciebin B, Li A, and Ellis A. 2003. Characterization of waterborne outbreak-associated *Campylobacter jejuni*, Walkerton, Ontario. *Emerg. Infect. Dis* 9:1232–1241. [PubMed: 14609457]
6. Dingle KE, Colles FM, Wareing DR, Ure R, Fox AJ, Bolton FE, Bootsma HJ, Willems RJ, Urwin R, and Maiden MC. 2001. Multilocus sequence typing system for *Campylobacter jejuni*. *J. Clin. Microbiol* 39:14–23. [PubMed: 11136741]
7. European Food Safety Authority (EFSA) Panel on Biological Hazards. 2010. Scientific opinion on quantification of the risk posed by broiler meat to human campylobacteriosis in the EU. *EFSA J.* 8:1437. doi:10.2903/j.efsa.2010.1437.
8. Friedman CR, Hoekstra RM, Samuel M, Marcus R, Bender J, Shiferaw B, Reddy S, Ahuja SD, Helfrick DL, Hardnett F, Carter M, Anderson B, Tauxe RV, and for the Emerging Infections Program FoodNet Working Group. 2004. Risk factors for sporadic *Campylobacter* infection in the United States: a case-control study in FoodNet sites. *Clin. Infect. Dis* 38(Suppl. 3):S285–S296. [PubMed: 15095201]
9. Gibney KB, O'Toole J, Sinclair M, and Leder K. 2014. Disease burden of selected gastrointestinal pathogens in Australia, 2010. *Int. J. Infect. Dis* 28:176–185. [PubMed: 25281904]
10. Hoffmann M, Luo Y, Monday SR, Gonzalez-Escalona N, Ottesen AR, Muruvanda T, Wang C, Kastanis G, Keys C, Janies D, Senturk IF, Catalyurek UV, Wang H, Hammack TS, Wolfgang WJ, Schoonmaker-Bopp D, Chu A, Myers R, Haendiges J, Evans PS, Meng J, Strain EA, Allard MW, and Brown EW. 2016. Tracing origins of the *Salmonella* Bareilly strain causing a food-borne outbreak in the United States. *J. Infect. Dis* 213(4):502–508. [PubMed: 25995194]
11. Interagency Food Safety Analytics Collaboration (IFSAC). 2015. Foodborne illness source attribution estimates for *Salmonella*, *Escherichia coli* O157 (*E. coli* O157), *Listeria monocytogenes* (Lm), and *Campylobacter* using outbreak surveillance data. Available at: <https://www.cdc.gov/foodsafety/pdfs/ifsac-project-report-508c.pdf>.
12. Janssen R, Krogfelt KA, Cawthraw SA, van Pelt W, Wagenaar JA, and Owen RJ. 2008. Host-pathogen interactions in *Campylobacter* infections: the host perspective. *Clin. Microbiol. Rev* 21:505–518. [PubMed: 18625685]
13. Keener KM, Bashor MP, Curtis PA, Sheldon BW, and Kathariou S. 2004. Comprehensive review of *Campylobacter* and poultry processing. *Compr. Rev. Food Sci. Food Saf* 3:105–116. [PubMed: 33430546]
14. Lehner A, Schneck C, Feierl G, Pless P, Deutz A, Brandl E, and Wagner M. 2000. Epidemiologic application of pulsed-field gel electrophoresis to an outbreak of *Campylobacter jejuni* in an Austrian youth centre. *Epidemiol. Infect* 125:13–16. [PubMed: 11057954]
15. National Antimicrobial Resistance Monitoring System (NARMS). 2015. Manual of laboratory methods, 2nd ed. U.S. Food and Drug Administration, Washington, DC.
16. Orr KE, Lightfoot NF, Sisson PR, Harkis BA, Tweddle JL, Boyd P, Carroll A, Jackson CJ, Wareing DR, and Freeman R. 1995. Direct milk excretion of *Campylobacter jejuni* in a dairy cow causing cases of human enteritis. *Epidemiol. Infect* 114:15–24. [PubMed: 7867733]

17. Painter JA, Hoekstra RM, Ayers T, Tauxe RV, Braden CR, Angulo FJ, and Griffin PM. 2013. Attribution of foodborne illnesses, hospitalizations, and deaths to food commodities by using outbreak data, United States, 1998–2008. *Emerg. Infect. Dis* 19:407–415. [PubMed: 23622497]
18. Sahin O, Fitzgerald C, Stroika S, Zhao S, Sippy RJ, Kwan P, Plummer PJ, Han J, Yaeger MJ, and Zhang Q. 2012. Molecular evidence for zoonotic transmission of an emergent, highly pathogenic *Campylobacter jejuni* clone in the United States. *J. Clin. Microbiol* 50:680–687. [PubMed: 22189122]
19. Sahin O, Plummer PJ, Jordan DM, Sulaj K, Pereira S, Robbe-Austerman S, Wang L, Yaeger MJ, Hoffman LJ, and Zhang Q. 2008. Emergence of a tetracycline-resistant *Campylobacter jejuni* clone associated with outbreaks of ovine abortion in the United States. *J. Clin. Microbiol* 46:1663–1671. [PubMed: 18322054]
20. Sails AD, Swaminathan B, and Fields PI. 2003. Utility of multilocus sequence typing as an epidemiological tool for investigation of outbreaks of gastroenteritis caused by *Campylobacter jejuni*. *J. Clin. Microbiol* 41:4733–4739. [PubMed: 14532212]
21. Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, Jones JL, and Griffin PM. 2011. Foodborne illness acquired in the United States—major pathogens. *Emerg. Infect. Dis* 17:7–15. [PubMed: 21192848]
22. Smith KE, Stenzel SA, Bender JB, Wagstrom E, Soderlund D, Leano FT, Taylor CM, Belle-Isle PA, and Danila R. 2004. Outbreaks of enteric infections caused by multiple pathogens associated with calves at a farm day camp. *Pediatr. Infect. Dis. J* 23:1098–1104. [PubMed: 15626945]
23. Stern NJ, Reiersen J, Lowman R, Bisailon JR, Fridriksdottir V, Gunnarsson E, Hiatt KL, and Campy-on-Ice C. 2005. Occurrence of *Campylobacter* spp. in cecal contents among commercial broilers in Iceland. *Foodborne Pathog. Dis* 2:82–89. [PubMed: 15992302]
24. Swaminathan B, Barrett TJ, Hunter SB, and Tauxe RV. 2001. PulseNet: the molecular subtyping network for foodborne bacterial disease surveillance, United States. *Emerg. Infect. Dis* 7:382–389. [PubMed: 11384513]
25. Thakur S, White DG, McDermott PF, Zhao S, Kroft B, Gebreyes W, Abbott J, Cullen P, English L, Carter P, and Harbottle H. 2009. Genotyping of *Campylobacter coli* isolated from humans and retail meats using multilocus sequence typing and pulsed-field gel electrophoresis. *J. Appl. Microbiol* 106:1722–1733. [PubMed: 19226383]
26. Thakur S, Zhao S, McDermott PF, Harbottle H, Abbott J, English L, Gebreyes WA, and White DG. 2010. Antimicrobial resistance, virulence, and genotypic profile comparison of *Campylobacter jejuni* and *Campylobacter coli* isolated from humans and retail meats. *Foodborne Pathog. Dis* 7:835–844. [PubMed: 20367499]
27. U.S. Food and Drug Administration. 2013. National Antimicrobial Resistance Monitoring System—Enteric Bacteria (NARMS): 2011 executive report. U.S. Department of Health and Human Services, Food and Drug Administration, Rockville, MD.
28. U.S. Food and Drug Administration. 2015. National Antimicrobial Resistance Monitoring System (NARMS): 2012–2013 Integrated Report. U.S. Department of Health and Human Services, Food and Drug Administration, Rockville, MD.
29. Wang X, Zhao S, Harbottle H, Tran T, Blickenstaff K, Abbott J, and Meng J. 2011. Antimicrobial resistance and molecular subtyping of *Campylobacter jejuni* and *Campylobacter coli* from retail meats. *J. Food Prot* 74:616–621. [PubMed: 21477476]
30. Whiley H, van den Akker B, Giglio S, and Bentham R. 2013. The role of environmental reservoirs in human campylobacteriosis. *Int. J. Environ. Res. Public Health* 10:5886–5907. [PubMed: 24217177]
31. Zhao C, Ge B, De Villena J, Sudler R, Yeh E, Zhao S, White DG, Wagner D, and Meng J. 2001. Prevalence of *Campylobacter* spp., *Escherichia coli*, and *Salmonella* serovars in retail chicken, turkey, pork, and beef from the greater Washington, D.C., area. *Appl. Environ. Microbiol* 67:5431–5436. [PubMed: 11722889]
32. Zhao S, Young SR, Tong E, Abbott JW, Womack N, Friedman SL, and McDermott PF. 2010. Antimicrobial resistance of *Campylobacter* isolates from retail meat in the United States between 2002 and 2007. *Appl. Environ. Microbiol* 76:7949–7956. [PubMed: 20971875]






**FIGURE 1.** Diversity of PFGE patterns by source. The Venn diagram depicts the number of PFGE patterns that are unique and shared among sources. Poultry refers to patterns of isolates from both poultry animals and poultry retail meat.



**FIGURE 2.** PFGE clusters from distinct sources. A dendrogram depicts the estimated relatedness of the PFGE patterns. The isolate identification number, isolate source, and *Campylobacter* species are depicted. RTM, retail poultry meat; poultry refers to isolates from poultry animal ceca.



**FIGURE 3.** Depiction of a high-virulence *Campylobacter* SA clone (DBRS16.0008). Isolate identification number, source of isolate, and species are shown. Some human patterns were included for comparison. RTM, retail poultry meat.

*Campylobacter* isolates with patterns indistinguishable from those from human patients based on source<sup>a</sup>

**TABLE 1.**

Source	No. of isolates ( <i>C. jejuni</i> / <i>C. coli</i> ) <sup>b</sup>	% isolates indistinguishable		
		<i>C. jejuni</i>	<i>C. coli</i>	Overall
Beef cattle	243/75	85.6	64.0	80.5
Dairy cattle	278/89	89.6	64.0	83.6
Pigs	6/357	66.7	12.6	13.7
Poultry animals and retail meat	194/115	74.2	47.0	64.1
Total	721/636	83.9	32.1	59.7

<sup>a</sup>Patterns indistinguishable are indistinguishable from those of human isolates in the CDC PulseNet database.

<sup>b</sup>Four *C. lari* were isolated from dairy cattle and one from pigs. One *C. fetus* was isolated from pigs. These six isolates were included in the overall data, with the five *C. lari* (but not the *C. fetus*) having patterns indistinguishable from those of human isolates.

**TABLE 2.** Number of PFGE patterns from different sources indistinguishable from patterns of human isolates<sup>a</sup>

Source	Total no. of patterns	Overall % patterns indistinguishable	No. of common patterns indistinguishable <sup>b</sup>
Beef cattle	129	56.6	23/23
Dairy cattle	157	65.0	34/34
Pigs	283	9.5	5/14
Poultry animals and retail meat	196	49.5	19/19

<sup>a</sup>Patterns indistinguishable are indistinguishable from those of human isolates in the CDC PulseNet database.

<sup>b</sup>Common patterns encompass three or more isolates from a particular source; this total is the denominator in each cell.