




# Prevalence of asymptomatic parasitaemia among household members of children under seasonal malaria chemoprevention coverage and comparison of the performance of standard rapid diagnostic tests *versus* ultrasensitive RDT for the detection of asymptomatic parasitaemia in Nanoro, Burkina Faso

Sié A. Elisée Kambou<sup>1</sup> · Kié Solange Millogo<sup>1</sup> · Paul Sondo<sup>1</sup> · Bérenger Kabore<sup>1</sup> · Amélé Fifi Chantal Kouevi<sup>1</sup> · Ismaila Bouda<sup>1</sup> · Toussaint Rouamba<sup>1</sup> · Karim Derra<sup>1</sup> · Mark Christian Tahita<sup>1</sup> · Hamidou Ilboudo<sup>1</sup> · Eli Rouamba<sup>1</sup> · Gauthier Tougri<sup>2</sup> · Sabina Dahlström Otienoburu<sup>3,4,5</sup> · Mehul Dhorda<sup>3,4,6</sup> · Sanata Bamba<sup>7</sup> · Philippe J Guerin<sup>3,4,6</sup>  · Halidou Tinto<sup>1</sup>

Received: 29 January 2024 / Accepted: 15 October 2024 / Published online: 15 November 2024  
© The Author(s) 2024

## Abstract

Asymptomatic carriers of *Plasmodium falciparum* represent important parasite reservoirs maintaining malaria transmission in the community. This study aimed on the one hand to screen the other household members living with children under seasonal malaria chemoprevention (SMC) coverage in order to determine the level of malaria infection in this population and on the other hand to determine the appropriate type of rapid diagnostic test (RDT) for this screening to detect these asymptomatic carriers in the community. During the 2022 SMC campaign (July to October), a cross-sectional survey was carried out in 745 participants who were screened by ultrasensitive rapid diagnostic test (usRDT), standard rapid diagnostic test (rRDT) and microscopy. Out of them, 395 had microscopy results available and were included in the data analysis. The prevalence of asymptomatic carriers of asexual forms of *Plasmodium falciparum* was 26.58% (105/395) while sexual forms were found in 5.32% (21/395) of the study population. Children from 5 to 15 years had the highest prevalence of *P. falciparum* asexual forms 35.76% (59/165) compared with older participants. Malaria positivity rate for rRDT and usRDT was 29.40% (219/745) and 40.49% (305/745) respectively. The usRDT had a higher sensitivity than the rRDT (72.38% (95% CI 62.8–80.66) vs. 60.95% (95% CI 50.94–70.33)). In terms of specificity, rRDT had a higher specificity 82.41% (95% CI 77.53–86.62) *versus* 69.66% (95% CI 64.01–74.89) for usRDT. This study reports a high prevalence of parasite carriers in household members of children under SMC coverage in Nanoro, Burkina Faso. In conclusion, usRDT seems more appropriate for strategies based on detection and treatment of parasite carriers within the community.

**Keywords** Malaria · Ultrasensitive RDT · Standard RDT · Asymptomatic carriers · Nanoro

Handling Editor: Julia Walochnik

✉ Sié A. Elisée Kambou  
eliseekambou96@gmail.com

<sup>1</sup> Institut de Recherche en Sciences de La Santé (IRSS),  
Clinical Research Unit of Nanoro (CRUN), Nanoro,  
Burkina Faso

<sup>2</sup> Ministry of Health of Burkina Faso: Permanent Secretary  
for Malaria Elimination (SP/Palu), Ouagadougou,  
Burkina Faso

<sup>3</sup> Infectious Diseases Data Observatory (IDDO), Oxford, UK

<sup>4</sup> Worldwide Antimalarial Resistance Network (WWARN),  
Oxford, UK

<sup>5</sup> College of STEM, Johnson C. Smith University, Charlotte,  
NC, USA

<sup>6</sup> Centre for Tropical Medicine and Global Health, Nuffield  
Department of Medicine, University of Oxford, Oxford, UK

<sup>7</sup> Institut Supérieur Des Sciences de La Santé (INSSA),  
Université Nazi Boni, Bobo Dioulasso, Burkina Faso

## Background

In Burkina Faso as in most countries in sub-Saharan Africa, malaria is rampant despite the various preventive and curative interventions implemented in recent years. In 2021, in Burkina Faso, an estimated 12,231,036 malaria cases resulting in 4355 deaths were reported (Ministry of Health BF 2021). Children under 5 years represented the most vulnerable population in the country. To reduce malaria-related mortality and morbidity in children, the World Health Organization (WHO) recommended in 2012 the seasonal malaria chemoprevention (SMC) as an innovative preventive strategy in areas with seasonal malaria transmission (World Health Organization 2012). SMC consists of monthly and intermittent administration of treatment with amodiaquine and sulfadoxine-pyrimethamine (AQSP) to children (aged 3–59 months) during the period of high malaria transmission. SMC has shown high level of protection against clinical malaria (Wilson and on behalf of the IPTc Taskforce 2011; Meremikwu et al. 2012; Cairns et al. 2021). Since 2018, SMC was implemented at national level in Burkina Faso. However, a high number of malaria cases continue to occur in children aged less than 5 years suggesting that the expected impact from SMC intervention is not achieved yet (Ministry of Health BF 2021). This may be attributable to various reasons including operational challenges resulting in suboptimal intervention coverage (Cairns et al. 2020), lack of observance of the second and/or third doses of AQ by parents or caregivers (Somé et al. 2022), inadequate for certain groups such as malnourished children who are at risk of subprotective drug concentrations (Oshikoya et al. 2010; De Kock et al. 2018), and increased selection of parasites highly resistant to SP or AQ (Roh et al. 2023). In addition, the parasite reservoirs present around children under SMC coverage may continually maintain the infection cycle. Indeed, the transmission of *Plasmodium* to healthy individuals requires the presence of a parasite reservoir (symptomatic and asymptomatic carriers). In sub-Saharan Africa, it was reported that the majority of malaria infections are asymptomatic due to transient immunity acquired during different exposures (Lindblade et al. 2013). In this context, targeting the parasite reservoirs, especially asymptomatic carriers surrounding children under SMC coverage, would be necessary for optimizing the impact of SMC intervention. However, the precise identification of these reservoirs in the community for adequate management remains a major challenge given the low sensitivity of standard RDT (rRDT) for the detection of asymptomatic infections, which consists mostly of low parasite density infections. The limit of detection (LOD) of rRDT is between 100 and 200 parasites/microliter of blood in field studies (Bousema et al. 2014; Adams

et al. 2015; Laban et al. 2015). Interestingly, ultrasensitive RDTs (usRDT) detecting the *hrp2* antigen were developed recently with the aim to improving the rapid detection of asymptomatic carriers of *Plasmodium falciparum* (Cunningham et al. 2019). usRDT are known to be able to detect *hrp2* antigen at concentrations as low as 10–40 pg/ml *hrp2* as opposed to 800–1000 pg/ml *hrp2* for rRDT (Kanwugu et al. 2019). Therefore, usRDT offer the possibility of detecting subpatent infections that will be missed by rRDT (Girma et al. 2019; Owalla et al. 2020). Nevertheless, there is limited data available on the effectiveness of usRDT in accurately identifying parasite reservoirs at the community level in endemic settings such as Burkina Faso. Therefore, this study aimed on the one hand to screen the other household members living with children under SMC coverage in order to determine the level of malaria infection in this population and on the other hand to determine the appropriate type of RDT for this screening to detect these asymptomatic carriers in the community.

## Materials and methods

### Study site

The present study was carried out in ten (10) villages of the rural commune of Soaw (Soaw, Kalwaka, Zoetgomde, Poesse, Rakalo, Kokolo, Kolokom, Mogdin, Bokin, Seguedin-Soaw) within the health district of Nanoro, Burkina Faso. The rural commune of Soaw is located in the center west of Burkina Faso, approximately 75 km from the capital Ouagadougou. According to data from the Nanoro health and demographic surveillance system (HDSS), the total population of the study area was estimated to be about 20,000 in 2018 (Derra et al. 2012). The climate there is of the Sudano-Sahelian type, with a rainy season from July to October followed by a dry season from November to June. Malaria is hyper-endemic with strong transmission during the rainy season, *P. falciparum* representing the dominant species (90%), followed by *Plasmodium malariae* (3–8%) and *Plasmodium ovale* (0.5–2%) (Sondo et al. 2015).

### Study design

This is an ancillary study nested to a large SMC-RST project (Sondo et al. 2022). In the SMC-RST study, 526 isolate households with a least one child under SMC coverage were included and assigned to one of the two study arms (control arm receiving SMC alone,  $n = 263$ ) or intervention arm (SMC + screening of household members and treatment if positive,  $n = 263$ ). The unit of randomization was the household, and the eligibility of a household was defined

by the presence of at least one child under SMC coverage. All roommates of children included in the intervention arm of the SMC-RST study were screened and treated. The inclusion and exclusion criteria for the SMC-RST study have been described elsewhere (Sondo et al. 2022).

This ancillary study was a cross-sectional study conducted from July to October 2022, i.e., during the 2022 SMC campaign, and focused on only households included in the intervention arm of the main SMC-RST study ( $n = 263$ ). The population of the present study was other household members referred to as roommates living with children under SMC coverage, in the intervention arm of the SMC-RST study. Other household members (roommates) were defined as any individual (not in SMC target population) sharing a household with at least one child under SMC coverage (aged 3–59 months). At each monthly visit, while all the roommates were screened with standard RDT and treated if positive as per the main SMC-RST intervention process, only one (01) roommate among them (other than those included in the previous months) was selected and benefited with additional testing with ultrasensitive RDT as per this ancillary study. The choice of roommate to be included at each visit was chosen from among the asymptomatic carriers presents at the time of the visit in the household. If more than one roommate was available and eligible, we drew lots to select the roommate to be included. In addition, in order to cover all eligible housemates in the study households by the end of the SMC campaign, it was necessary, during some visits, to select two roommates from certain households. All villages were systematically targeted at each SMC round. In the present study, asymptomatic carrier was defined as a person with no symptoms, with a body temperature  $< 37.5$  °C, and who reported no fever during the 2 days before blood collection. Participants were visited at home, and data on the sociodemographic characteristics were collected through a standardized numerical questionnaire. Capillary blood samples were taken from each participant for the detection of *P. falciparum* infection by rRDT, usRDT, and microscopy.

### Malaria diagnosis by RDT

The AdvDx™ Malaria Pf test kit (004ADFEF025KI-1, Advy Chemical Pvt. Ltd, India) served as the standard RDT, and the NxTek™ Eliminate Malaria Pf test kit (05FK140, Abbot, Korea) was used as the ultrasensitive RDT. Both tests are based on in vitro diagnostic immuno-chromatographic assay for the qualitative detection of the *hrp2* antigen specific of *Plasmodium falciparum* malaria in human whole blood. The tests were performed by trained field workers following the instructions of each manufacturer. Briefly, approximately 5  $\mu$ l ( $\mu$ l) of capillary blood was collected using a sample dropper and then transferred to the sample port, followed by the addition of four drops of the buffer solution to the buffer

port. The interpretation of the results was carried out after an incubation of 20 min. A sample was considered positive for *P. falciparum* malaria if the “Pf” test band and the “C” control band appeared in the result window. The presence of only the control band was considered a negative result. Results were declared invalid if the control band did not appear in the result window, thus warranting a new test.

### Malaria diagnosis by microscopy

Thick and thin blood smear slides were taken in the field and sent to the parasitology laboratory/Soaw station of the Clinical Research Unit of Nanoro (CRUN) for microscopic diagnosis. This diagnosis consisted of the detection of asexual and sexual forms of the parasites and their enumeration in the thick smear, followed by an identification of *Plasmodium* species on the thin smear. To do this, the slides were dried and then stained with Giemsa (diluted at 3%) for 45 min according to the standard operating procedures of the CRUN. They were then examined on a light microscope Olympus CX23 (Olympus corporation, Japan) at  $\times 100$  oil immersion objective. Thick smears were declared negative when no asexual parasites were encountered after running at least 100 microscopic fields. When the slide was positive, the parasite density (DP) was determined by counting the number of asexual parasites per 200 white blood cells and calculated per microliter of blood by assuming the number of white blood cells to be at 8000/ $\mu$ l. Gametocytes were counted against 500 white blood cells, and densities were estimated using a factor of 8000 leukocytes/microliter. Each blood slide was independently blinded and read by two expert microscopists of CRUN qualified by the National Institute for Communicable Diseases (NICD, South Africa) (Tinto et al. 2014). In case of significant discordance (negative vs. positive; difference in *Plasmodium* species; difference between the two readers greater than twice the DP value), a third reading was performed by another expert microscopist who had no knowledge of the previous results. The two closest readings were selected as the matching result.

### Sample size estimation

Details about the sample size calculation for the larger SMC-RST project were described previously (Sondo et al. 2022). Briefly, 526 isolate households with a least one child under SMC coverage were included, i.e., 263 households each of the two arms (control and intervention) in order to have 80% power to detect a 20% decrease in incidence of malaria after 1 year in comparison to a baseline incidence rate in the control arm (SMC) between 1.5 and 2.0 malaria cases per year, assuming a one-sided test with a significance level of 0.025 and large-sample  $z$ -test of

the Poisson event rate difference (PASS software) (Sondo et al. 2022). The sample size of this study was calculated considering that the prevalence of asymptomatic malaria parasite carriage is estimated at 30% in the study area. As the study is ancillary to the SMC-RST study, the sample size was calculated assuming a finite population of 2630 participants and also so as to be 95% certain that the sample size estimate will be within 10% of the actual proportion of asymptomatic carriers of *Plasmodium*. Taking these assumptions into account, the required sample size was estimated at 745 participants.

## Data management and statistical analysis

Sociodemographic data and RDT results were entered in a REDCap database. Sociodemographic characterization and age stratification of asymptomatic malaria were performed by analyzing data from the subpopulation of participants with available microscopy results. Descriptive statistics was conducted for the sociodemographic characteristics and presented as frequencies and percentages. The performance of each RDT (standard RDT and ultrasensitive RDT) was assessed regarding expert microscopy results (gold standard) by calculating parameters such as sensitivity (Se), specificity (Sp), and the positive and negative predictive values (PPV and NPV). The concordance between rRDT and expert microscopy and between usRDT and expert microscopy was calculated using Cohen's kappa coefficient of concordance ( $k$ ). The  $k$  values  $< 0$ , 0.01–0.20, 0.21–0.40, 0.41–0.60, 0.61–0.80, and 0.81–1 represent mediocre, slight, fair, moderate, substantial, and near-perfect degrees of agreement, respectively (Landis and Koch 1977; McHugh 2012).

## Results

### Baseline characteristics of study population

A total of 745 household members of children under SMC coverage from the SMC-RST study were included in this study. RDT data from 745 included participants were used to assess the prevalence of asymptomatic malaria by rRDT and usRDT. Of the 745 participants included, 395 had microscopy data available. The age of the participants in the study subpopulation ranged from 5 to 83 years with a median age of 18 years, with a majority above 22 years of age (180/395). The female household members represented 55.70% (220/395) of the study population and males represented 44.30% (175/395). The sociodemographic characteristics of study subpopulation are shown in Table 1.

**Table 1** Sociodemographic characteristics of household members with microscopy results

Characteristics	Frequency $n$ (%)	
Age (in years)	5–15	165 (41.77)
	16–26	74 (18.73)
	27–37	58 (14.68)
	$\geq 38$	98 (24.81)
Sex	Female	220 (55.70)
	Male	175 (44.30)

**Table 2** Prevalence of asymptomatic carriers of *P. falciparum* by microscopy, rRDT, and usRDT

Method of diagnosis	Positive $n$ (%)	Negative $n$ (%)	Total
usRDT	305 (40.94)	440 (59.06)	745
	164 (41.52)	231 (58.48)	395
rRDT	219 (29.40)	526 (70.60)	745
	115 (29.11)	280 (70.89)	395
Microscopy			
Asexual form	105 (26.58)	290 (73.42)	395


### Prevalence of asymptomatic carriers of malaria parasite

RDT positivity rate among household members of children under SMC coverage was 40.94% (305/745) determined by usRDT and 29.40% (219/745) using rRDT. The prevalence of asymptomatic carriers of asexual forms of *P. falciparum* detected by microscopy was 26.58% (105/395). Of the 105 asymptomatic carriers of asexual form of *P. falciparum*, 103 (98.09%) were *P. falciparum* mono-infection and 2 (1.09%) *P. falciparum* mixed infections (1 *P. falciparum* + *P. malariae* and 1 *P. falciparum* + *P. ovale*). Also, 3 *P. malariae* mono-infection and 1 *P. ovale* mono-infection were detected by microscopy.

The parasite density of *P. falciparum* infections ranged from 24 to 42,645 parasites/microliter with a geometric mean of 551.42 parasites/microliter. The prevalence of sexual form of *P. falciparum* among household members was 5.32% (21/395), and the geometric mean of gametocyte estimated at 43.98 parasites/microliter (Table 2).

### Prevalence of asymptomatic carriage of asexual forms and gametocyte of *P. falciparum* by age group

The prevalence of asymptomatic carriage of asexual form of *P. falciparum* was higher in children from 5 to 15 years old 35.76 (59/165). This was 32.43% (24/74) in the age group of 16 to 26 years old and 24.14% (14/58) and 8.16% (8/98) in the age groups 27 to 37 and  $\geq 38$  years old respectively



**Fig. 1** Prevalence of asymptomatic carriers of *P. falciparum* by age groups. The figure represents the prevalence of the carriers of parasite (detected by microscopy) in household members of children under

SMC coverage. **A** The carriers of the asexual form of *P. falciparum*. **B** The carriers of gametocyte

(Fig. 1A). The prevalence of *P. falciparum* gametocytes in children between 5 and 15 years was 7.88% (13/165) (Fig. 1B).


**Comparison of diagnostic performance of rRDT and usRDT versus microscopy**

Of the 105 participants with a positive *P. falciparum* microscopy result considered true positives, the usRDT detected 76 (72.38%) while the rRDT was detected only 64 (60.95%). The false-positive rate was 53.66% (88/164) and 44.35% (51/115) with usRDT and rRDT respectively. A total of 56 participants that tested negative by rRDT tested positive by usRDT. Out of these, 26.79% (15/56)

were confirmed to be *P. falciparum* positive by microscopy. In total, 25% (41/164) of the participants were diagnosed positive by usRDT and negative by microscopy and rRDT (Fig. 2).

The usRDT had a higher sensitivity than the rRDT (Se = 72.38% (95% confidence interval (CI) = 62.8 to 80.66) vs. 60.95% (95% CI = 50.94 to 70.33)). In terms of specificity, rRDT had a higher specificity (82.41% 95% CI = 77.53 to 86.62) compared to 69.66% (95% CI = 64.01 to 74.89) for usRDT). Both RDTs had a comparable negative predictive value (87.45% (95% IC = 82.47 to 91.43) for usRDT and 85.36 (95% IC = 80.66 to 89.28) for rRDT. usRDT and rRDT showed substantial agreement ( $k=0.70$  for the usRDT and  $k=0.76$  for the rRDT) with the microscopy results (Table 3).

**Fig. 2** Venn diagram showing the distribution of positive results according to the diagnostic test used with microscopy as reference test



**Table 3** Diagnostic performance of rRDT and usRDT using microscopy as a reference

Parameter	usRDT % [95% confidence interval]	rRDT (%) [95% confidence interval]
Sensitivity	72.38 [62.8–80.66]	60.95 [50.94–70.33]
Specificity	69.66[64.01–74.89]	82.41 [77.53–86.62]
Positive predictive value	46.34 [38.53–54.28]	55.65 [46.09–64.91]
Negative predictive value	87.45 [82.47–91.43]	85.36 [80.66–89.28]
Kappa coefficient ( <i>k</i> )	0.70	0.76

### Diagnostic performance of rRDT and usRDT according parasite density

The sensitivity of the different RDT decreased with the parasite density; however, the usRDT was more sensitive than rRDT in both high (PD  $\geq$  200) and low (PD < 200) parasite densities (Table 4).

### Discussion

This study reported on the prevalence of asymptomatic parasitaemia and on the performance of ultrasensitive *hrp2* RDT and standard *hrp2* RDT for detection of asymptomatic parasitaemia in household members of children under SMC coverage in Nanoro, Burkina Faso. High prevalence of asymptomatic carriers of asexual form of *P. falciparum* was observed in household members of children under SMC coverage. These asymptomatic carriers could compromise the effect of SMC intervention as these infections often go unnoticed, therefore untreated, and constitute major reservoirs of gametocytes for mosquito vectors, thus maintaining malaria transmission in the community (Alves et al. 2005; Lindblade et al. 2013). However, this finding is not surprising, especially in malaria-endemic areas where, despite the elimination of a large proportion of infected erythrocytes by immune-protective mechanisms, some may persist in the bloodstream and lead to asymptomatic parasitaemia (Dal-Bianco et al. 2007). Various studies reported higher prevalence of asymptomatic carriers in malaria endemic areas due to the transient immunity acquired over time (Laishram et al. 2012; Lindblade et al. 2013; Starzengruber et al. 2014). Malaria control strategies in endemic areas should therefore

consider asymptomatic parasitaemia as a major obstacle to control efforts. Therefore, simultaneous screening and treatment of other asymptomatic household members with an antimalarial drug to eliminate the parasite reservoir could maximize the expected impact of the SMC intervention and global malaria control efforts.

Regarding age, adults and older children are more likely to be asymptomatic carriers because of the development of partial immunity due to repeated exposure to malaria parasites. The high prevalence of asymptomatic carriage of *P. falciparum* in children from 5 to 15 years old compared to other ages suggests a vulnerability of this age group, which represents an important reservoir for maintaining malaria transmission in the community. This was pointed out in various previous studies, highlighting school-aged children as greater contribution to the infectious reservoir and thereby undermining malaria elimination efforts (Ouedraogo et al. 2016; Coalson et al. 2016; Gonçalves et al. 2017). Therefore, the implementation of intermittent preventive treatment of malaria in school-aged (IPTsc) children as recommended by the WHO in 2022 (World Health Organization 2022) could reduce the parasite reservoir and prevent significant morbidity in this age group. There is some evidence that administration of IPTsc can confer a community-level benefit to those not receiving intermittent preventive treatment. Indeed, the administration of IPTc has been associated with a significant reduction in gametocyte carriage, which has a positive effect on malaria indicators at the community level (Clarke et al. 2017; Staedke et al. 2018; Rehman et al. 2019).

Cissé et al. in a stepped-wedge trial in Senegal reported that the expansion of SMC to children under 10 was associated with a 27% reduction in malaria in persons who had not received SMC, an effect that was not seen when SMC was limited to children 3–59 months (Cissé et al. 2016).

In this study, the prevalence of sexual forms of *P. falciparum* in household members was low compared with findings from a previous study that showed a high prevalence of gametocyte carriage in settings with a high load of asymptomatic infections (Vantaux et al. 2018). Difference between the two studies results may be attributable to many factors. The higher sensitivity of the RT-PCR used in the previous study compared with the microscopy used in this study could be the most important factor, as gametocytes tend to occur at low densities (Churcher et al. 2013; Sturrock et al. 2013). Moreover, polyclonality, asexual

**Table 4** Diagnostic performance of rRDT and usRDT according parasite density

Diagnosis test	Parasite density	Sensitivity % [CI95%]	Specificity % [CI95%]
usRDT	< 200	51.52 [33.54–69.20]	59.39 [54.14–64.49]
	$\geq$ 200	40.61 [35.51–45.86]	48.48 [30.80–66.46]
rRDT	< 200	33.33 [17.96–51.83]	71.27 [66.31–75.88]
	$\geq$ 200	28.73 [24.12–33.69]	66.67 [48.17–82.04]

parasite density, multiplicity of infections, asexual genotype, acquired host immunity, immune responses, and seasonality of parasite transmission may also influence gametocytogenesis and therefore the prevalence of gametocyte (Nassir et al. 2005; Ouédraogo et al. 2008; Bousema and Drakeley 2011; Lamptey et al. 2018; Touray et al. 2021). Since gametocytemia is one of the indicators of intensity of transmission within an area, distinguishing gametocyte carriers in population would enable better characterization of the parasite reservoir (Lindblade et al. 2013). The use of more sensitive diagnostic tools for the detection of submicroscopic gametocytes and search for factors linked to gametocytogenesis would therefore be necessary to provide appropriate guidance for the future implementation of strategies aimed at reducing and interrupting transmission.

The findings showed that usRDT detected more asymptomatic *P. falciparum* carriers than the rRDT and microscopy confirming its superiority in detecting asymptomatic *P. falciparum* infections over the rRDT and microscopy. Similarly, Landier et al. are observed in a large field survey of asymptomatic carriers in Myanmar that usRDT were more sensitive than the rRDT and microscopy for the detection of asymptomatic parasitaemia (Landier et al. 2018). The superiority of the usRDT results indeed from its LOD which is relatively low compared to the LOD of the rRDT and the expert microscopy, thus offering it the capacity to detect very low parasite densities (Jimenez et al. 2017; Das et al. 2018; Mpina et al. 2022). Parasite density being controlled by acquired immunity in infected hosts (Bousema et al. 2014), populations in high-transmission areas are more likely to have submicroscopic infections (Okell et al. 2012; Mosha et al. 2013) which are not detectable by microscopy and rRDT. Low parasite density could therefore have affected the proportion of asymptomatic infections detected by microscopy and rRDT (Okell et al. 2009). Other studies have also documented similar findings (Das et al. 2017, 2018).

The high sensitivity of the usRDT compared to the rRDT in the detection of asymptomatic carriers as observed in this study corroborates with its ability to identify parasites below the detection threshold of the rRDT (Das et al. 2018). Similar results had been reported in different studies on different asymptomatic populations (Girma et al. 2019; Briand et al. 2020; Acquah et al. 2021; Manjurano et al. 2021; Yimam et al. 2022). However, the usRDT did not show increased sensitivity compared to rRDT as was the case in studies conducted in Indonesia (62% for rRDT vs. 84% for usRDT) (Unwin et al. 2020), Colombia (64.3% for rRDT vs. 71.4% usRDT) (Vásquez et al. 2018), and Myanmar (44% for usRDT vs. 0% for rRDT) (Das et al. 2017). In terms of usRDT, it had lower specificity than rRDT. This could be due to the influence of several parameters such as the persistence of the *hrp2* antigen in the blood (Mouatcho and Goldring 2013), the genetic variability of *hrp2* and its

homology with *hrp3* (Baker et al. 2005), and mutations or deletions in the *Pfhrp2* gene (Houzé et al. 2011; Koita et al. 2012; Maltha et al. 2012) to high malaria transmission in the study site (Hopkins et al. 2008; Baiden et al. 2012). Another source of variation in sensitivity could be the reference test used and its associated LOD (Ding et al. 2023). Factors such as transport and storage conditions and the inability of the parasite to express the *hrp2* target antigen can affect TDR performance. All this merits further studies. However, the high sensitivity of the usRDT compared to the rRDT makes it potentially more useful for community diagnosis of asymptomatic carriers especially in a perspective of malaria elimination.

### Limitations of the study

The lack of microscopy results in number of participants is the major limitation of this study. Indeed, at first, participants were including without slides (the two types of RDT only). The inclusion of microscopy resulted from a recommendation by steering committee, and this was caught up from month 2 leaving the firstly included participants without microscopy results. Another limitation was the use of microscopy as gold standard rather than molecular tools such as PCR that could have detected more asymptomatic infections.

### Conclusion

The present study reports higher prevalence of parasite carriers in other household members of children under SMC coverage in the health district of Nanoro, Burkina Faso. More specifically, school-aged children were mostly affected. This could lead to frequent reinfection of children under SMC coverage, compromising the effect of SMC intervention. Also, the usRDT seems more appropriate for strategies based on detection and treatment of parasite carriers within the community. Finally, this study supports the implementation of intermittent preventive treatment of malaria in school-aged (ITPsc) children in Burkina Faso.

**Acknowledgements** The authors would like to thank Expertise France L'Initiative for the financial supports of the project. Special thanks to Veronica Nosedá and Cyrielle Thomas for their kind support. We are grateful to the study participants and all the community of the department of Soaw for their acceptance in participating to the study. We thank the study staff i.e. field workers, fields supervisors, laboratory technicians, data managers and the personnel of the Clinical Research Unit of Nanoro (CRUN) in Burkina Faso who played an important role for the successful completion of the present study.

**Author contribution** SP, KB, RT, DK, MT, IH, RE, TG, SO, MD, GP, TH designed the study, KE, MS, SP, KB, KC, BI, RT, DK, MT, IH, RE, SB, TH implemented the study and supervised field work, KE, SP, IB, RT contributed in data management and statistical analysis, KE,

SP, RT,DK, SO,SB, TH contributed in drafting the manuscript and all authors read and approved the manuscript.

**Funding** This study is funded by Expertise France, L'Initiative, through the SMC-RST project (20SANIN204).

**Data availability** No datasets were generated or analysed during the current study.

## Declarations

**Ethics approval and consent to participate** This study was an ancillary to a larger study entitled "Boosting the impact of seasonal malaria chemoprevention (SMC) through simultaneous screening and treatment of household members of children receiving SMC in Burkina Faso" acronym SMC-RST which has been approved by the Ethics Committee for Health Research of Burkina Faso (Deliberation No.: 2021–03-059 of 10 March 2021). A signed informed consent was obtained from participants or their parents/guardians (if they are minors) before enrollment. An impartial, literate witness (not a member of the study staff) was presented in case the parents/guardians were illiterate. The parent(s)/guardian(s) and, if applicable, the witness signed the informed consent.

**Consent for publication** Not applicable.

**Competing interests** The authors declare no competing interests.

**Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

- Acquah FK, Donu D, Obboh EK et al (2021) Diagnostic performance of an ultrasensitive HRP2-based malaria rapid diagnostic test kit used in surveys of afebrile people living in Southern Ghana. *Malar J* 20:125. <https://doi.org/10.1186/s12936-021-03665-7>
- Adams M, Joshi SN, Mbambo G et al (2015) An ultrasensitive reverse transcription polymerase chain reaction assay to detect asymptomatic low-density *Plasmodium falciparum* and *Plasmodium vivax* infections in small volume blood samples. *Malar J* 14:520. <https://doi.org/10.1186/s12936-015-1038-z>
- Alves FP, Gil LHS, Marrelli MT et al (2005) Asymptomatic carriers of *Plasmodium* spp. as infection source for malaria vector mosquitoes in the Brazilian Amazon. *J Med Entomol* 42:777–779. <https://doi.org/10.1093/jmedent/42.5.777>
- Baiden F, Webster J, Tivura M et al (2012) Accuracy of rapid tests for malaria and treatment outcomes for malaria and non-malaria cases among under-five children in rural Ghana. *PLoS ONE* 7:e34073. <https://doi.org/10.1371/journal.pone.0034073>
- Baker J, McCarthy J, Gatton M et al (2005) Genetic diversity of *Plasmodium falciparum* histidine-rich protein 2 (PfHRP2) and its effect on the performance of PfHRP2-based rapid diagnostic tests. *J INFECT DIS* 192:870–877. <https://doi.org/10.1086/432010>
- Bousema T, Drakeley C (2011) Epidemiology and infectivity of *Plasmodium falciparum* and *Plasmodium vivax* gametocytes in relation to malaria control and elimination. *Clin Microbiol Rev* 24:377–410. <https://doi.org/10.1128/CMR.00051-10>
- Bousema T, Okell L, Felger I, Drakeley C (2014) Asymptomatic malaria infections: detectability, transmissibility and public health relevance. *Nat Rev Microbiol* 12:833–840. <https://doi.org/10.1038/nrmicro3364>
- Briand V, Cottrell G, Tuike Ndam N et al (2020) Prevalence and clinical impact of malaria infections detected with a highly sensitive HRP2 rapid diagnostic test in Beninese pregnant women. *Malar J* 19:188. <https://doi.org/10.1186/s12936-020-03261-1>
- Cairns M, Ceesay SJ, Sagara I et al (2021) Effectiveness of seasonal malaria chemoprevention (SMC) treatments when SMC is implemented at scale: case–control studies in 5 countries. *PLoS Med* 18:e1003727. <https://doi.org/10.1371/journal.pmed.1003727>
- Cairns ME, Sagara I, Zongo I et al (2020) Evaluation of seasonal malaria chemoprevention in two areas of intense seasonal malaria transmission: secondary analysis of a household-randomised, placebo-controlled trial in Houndé District, Burkina Faso and Bougouni District. *Mali Plos Med* 17:e1003214. <https://doi.org/10.1371/journal.pmed.1003214>
- Churcher TS, Bousema T, Walker M et al (2013) Predicting mosquito infection from *Plasmodium falciparum* gametocyte density and estimating the reservoir of infection. *eLife* 2:e00626. <https://doi.org/10.7554/eLife.00626>
- Cissé B, Ba EH, Sokhna C et al (2016) Effectiveness of seasonal malaria chemoprevention in children under ten years of age in Senegal: a stepped-wedge cluster-randomised trial. *PLoS Med* 13:e1002175. <https://doi.org/10.1371/journal.pmed.1002175>
- Clarke SE, Rouhani S, Diarra S et al (2017) Impact of a malaria intervention package in schools on *Plasmodium* infection, anaemia and cognitive function in schoolchildren in Mali: a pragmatic cluster-randomised trial. *BMJ Glob Health* 2:e000182. <https://doi.org/10.1136/bmjgh-2016-000182>
- Coalson JE, Walldorf JA, Cohee LM et al (2016) High prevalence of *Plasmodium falciparum* gametocyte infections in school-age children using molecular detection: patterns and predictors of risk from a cross-sectional study in southern Malawi. *Malar J* 15:527. <https://doi.org/10.1186/s12936-016-1587-9>
- Cunningham J, Jones S, Gatton ML et al (2019) A review of the WHO malaria rapid diagnostic test product testing programme (2008–2018): performance, procurement and policy. *Malar J* 18:387. <https://doi.org/10.1186/s12936-019-3028-z>
- Dal-Bianco MP, Köster KB, Kombila UD et al (2007) High prevalence of asymptomatic *Plasmodium falciparum* infection in Gabonese adults. *Am J Trop Med Hyg* 77:939–942
- Das S, Jang IK, Barney B et al (2017) Performance of a high-sensitivity rapid diagnostic test for *Plasmodium falciparum* malaria in asymptomatic individuals from Uganda and Myanmar and naive human challenge infections. *Am J Trop Med Hyg* 97:1540–1550. <https://doi.org/10.4269/ajtmh.17-0245>
- Das S, Peck RB, Barney R et al (2018) Performance of an ultrasensitive *Plasmodium falciparum* HRP2-based rapid diagnostic test with recombinant HRP2, culture parasites, and archived whole blood samples. *Malar J* 17:118. <https://doi.org/10.1186/s12936-018-2268-7>
- De Kock M, Tarning J, Workman L et al (2018) Population pharmacokinetic properties of sulfadoxine and pyrimethamine: a pooled analysis to inform optimal dosing in African children with uncomplicated malaria. *Antimicrob Agents Chemother* 62:e01370-e1417. <https://doi.org/10.1128/AAC.01370-17>



- Derra K, Rouamba E, Kazienga A et al (2012) Profile: Nanoro health and demographic surveillance system. *Int J Epidemiol* 41:1293–1301. <https://doi.org/10.1093/ije/dys159>
- Ding XC, Incardona S, Serra-Casas E et al (2023) Malaria in pregnancy (MiP) studies assessing the clinical performance of highly sensitive rapid diagnostic tests (HS-RDT) for *Plasmodium falciparum* detection. *Malar J* 22:60. <https://doi.org/10.1186/s12936-023-04445-1>
- Girma S, Cheaveau J, Mohon AN et al (2019) Prevalence and epidemiological characteristics of asymptomatic malaria based on ultrasensitive diagnostics: a cross-sectional study. *Clin Infect Dis* 69:1003–1010. <https://doi.org/10.1093/cid/ciy1005>
- Gonçalves BP, Kapulu MC, Sawa P et al (2017) Examining the human infectious reservoir for *Plasmodium falciparum* malaria in areas of differing transmission intensity. *Nat Commun* 8:1133. <https://doi.org/10.1038/s41467-017-01270-4>
- Hopkins H, Bebell L, Kambale W et al (2008) Rapid diagnostic tests for malaria at sites of varying transmission intensity in Uganda. *J INFECT DIS* 197:510–518. <https://doi.org/10.1086/526502>
- Houzé S, Hubert V, Le Pessec G et al (2011) Combined deletions of *pflhrp2* and *pflhrp3* genes result in *Plasmodium falciparum* malaria false-negative rapid diagnostic test. *J Clin Microbiol* 49:2694–2696. <https://doi.org/10.1128/JCM.00281-11>
- Jimenez A, Rees-Channer RR, Perera R et al (2017) Analytical sensitivity of current best-in-class malaria rapid diagnostic tests. *Malar J* 16:128. <https://doi.org/10.1186/s12936-017-1780-5>
- Kanwugu ON, Helegbe GK, Aryee PA et al (2019) Prevalence of asymptomatic malaria among children in the Tamale Metropolis: how does the PfHRP2 CareStart™ RDT perform against microscopy? *J Trop Med* 2019:1–7. <https://doi.org/10.1155/2019/6457628>
- Koita OA, Doumbo OK, Ouattara A et al (2012) False-negative rapid diagnostic tests for malaria and deletion of the histidine-rich repeat region of the *hrp2* gene. *Am J Trop Med Hyg* 86:194–198. <https://doi.org/10.4269/ajtmh.2012.10-0665>
- Laban NM, Kobayashi T, Hamapumbu H et al (2015) Comparison of a PfHRP2-based rapid diagnostic test and PCR for malaria in a low prevalence setting in rural southern Zambia: implications for elimination. *Malar J* 14:25. <https://doi.org/10.1186/s12936-015-0544-3>
- Laishram DD, Sutton PL, Nanda N et al (2012) The complexities of malaria disease manifestations with a focus on asymptomatic malaria. *Malar J* 11:29. <https://doi.org/10.1186/1475-2875-11-29>
- Lamptey H, Ofori MF, Kusi KA et al (2018) The prevalence of sub-microscopic *Plasmodium falciparum* gametocyte carriage and multiplicity of infection in children, pregnant women and adults in a low malaria transmission area in Southern Ghana. *Malar J* 17:331. <https://doi.org/10.1186/s12936-018-2479-y>
- Landier J, Haohankunnatham W, Das S et al (2018) Operational performance of a *Plasmodium falciparum* ultrasensitive rapid diagnostic test for detection of asymptomatic infections in Eastern Myanmar. *J Clin Microbiol* 56:e00565–e618. <https://doi.org/10.1128/JCM.00565-18>
- Landis JR, Koch GG (1977) The measurement of observer agreement for categorical data. *Biometrics* 33:159. <https://doi.org/10.2307/2529310>
- Lindblade KA, Steinhardt L, Samuels A et al (2013) The silent threat: asymptomatic parasitemia and malaria transmission. *Expert Rev Anti Infect Ther* 11:623–639. <https://doi.org/10.1586/eri.13.45>
- Maltha J, Gamboa D, Bendezu J et al (2012) Rapid diagnostic tests for malaria diagnosis in the Peruvian Amazon: impact of *pflhrp2* gene deletions and cross-reactions. *PLoS ONE* 7:e43094. <https://doi.org/10.1371/journal.pone.0043094>
- Manjurano A, Omolo JJ, Lyimo E et al (2021) Performance evaluation of the highly sensitive histidine-rich protein 2 rapid test for *Plasmodium falciparum* malaria in North-West Tanzania. *Malar J* 20:58. <https://doi.org/10.1186/s12936-020-03568-z>
- McHugh ML (2012) Interrater reliability: the kappa statistic. *Biochem Med (Zagreb)* 22:276–282
- Meremikwu MM, Donegan S, Sinclair D et al (2012) Intermittent preventive treatment for malaria in children living in areas with seasonal transmission. *Cochrane Database Syst Rev*. <https://doi.org/10.1002/14651858.CD003756.pub4>
- Ministry Of Health BF (2021) Statistical yearbook 2021. Burkina Faso. [https://www.sante.gov.bf/fileadmin/annuaire\\_2021\\_mshp.pdf](https://www.sante.gov.bf/fileadmin/annuaire_2021_mshp.pdf)
- Mosha JF, Sturrock HJ, Greenhouse B et al (2013) Epidemiology of subpatent *Plasmodium falciparum* infection: implications for detection of hotspots with imperfect diagnostics. *Malar J* 12:221. <https://doi.org/10.1186/1475-2875-12-221>
- Mouatcho JC, Goldring JPD (2013) Malaria rapid diagnostic tests: challenges and prospects. *J Med Microbiol* 62:1491–1505. <https://doi.org/10.1099/jmm.0.052506-0>
- Mpina M, Stabler TC, Schindler T et al (2022) Diagnostic performance and comparison of ultrasensitive and conventional rapid diagnostic test, thick blood smear and quantitative PCR for detection of low-density *Plasmodium falciparum* infections during a controlled human malaria infection study in Equatorial Guinea. *Malar J* 21:99. <https://doi.org/10.1186/s12936-022-04103-y>
- Nassir E, Abdel-Muhsin A-MA, Suliaman S et al (2005) Impact of genetic complexity on longevity and gametocytogenesis of *Plasmodium falciparum* during the dry and transmission-free season of eastern Sudan. *Int J Parasitol* 35:49–55. <https://doi.org/10.1016/j.ijpara.2004.10.014>
- Okell LC, Bousema T, Griffin JT et al (2012) Factors determining the occurrence of submicroscopic malaria infections and their relevance for control. *Nat Commun* 3:1237. <https://doi.org/10.1038/ncomms2241>
- Okell LC, Ghani AC, Lyons E, Drakeley CJ (2009) Submicroscopic infection in *Plasmodium falciparum* –endemic populations: a systematic review and meta-analysis. *J INFECT DIS* 200:1509–1517. <https://doi.org/10.1086/644781>
- Oshikoya KA, Sammons HM, Choonara I (2010) A systematic review of pharmacokinetics studies in children with protein-energy malnutrition. *Eur J Clin Pharmacol* 66:1025–1035. <https://doi.org/10.1007/s00228-010-0851-0>
- Ouédraogo AL, De Vlas SJ, Nébié I et al (2008) Seasonal patterns of *Plasmodium falciparum* gametocyte prevalence and density in a rural population of Burkina Faso. *Acta Trop* 105:28–34. <https://doi.org/10.1016/j.actatropica.2007.09.003>
- Ouédraogo AL, Gonçalves BP, Gnémé A et al (2016) Dynamics of the human infectious reservoir for malaria determined by mosquito feeding assays and ultrasensitive malaria diagnosis in Burkina Faso. *J Infect Dis* 213:90–99. <https://doi.org/10.1093/infdis/jiv370>
- Owalla TJ, Okurut E, Apungia G et al (2020) Using the ultrasensitive Alere *Plasmodium falciparum* malaria Ag HRP-2™ rapid diagnostic test in the field and clinic in Northeastern Uganda. *Am J Trop Med Hyg* 103:778–784. <https://doi.org/10.4269/ajtmh.19-0653>
- Rehman AM, Maiteki-Sebuguzi C, Gonahasa S et al (2019) Intermittent preventive treatment of malaria delivered to primary school-children provided effective individual protection in Jinja, Uganda: secondary outcomes of a cluster-randomized trial (START-IPT). *Malar J* 18:318. <https://doi.org/10.1186/s12936-019-2954-0>
- Roh ME, Zongo I, Haro A et al (2023) Seasonal malaria chemoprevention drug levels and drug resistance markers in children with or without malaria in Burkina Faso: a case-control study. *J Infect Dis* 228:926–935. <https://doi.org/10.1093/infdis/jiad172>
- Somé AF, Zongo I, Sagara I et al (2022) Factors influencing second and third dose observance during seasonal malaria chemoprevention (SMC): a quantitative study in Burkina Faso. *Mali and Niger Tropicalmed* 7:214. <https://doi.org/10.3390/tropicalmed7090214>

- Sondo P, Bihoun B, Valea I, et al (2015) La part du paludisme dans les maladies fébriles en saison sèche dans la région de Nanoro, Burkina Faso. *West African Journal of Research for Health*. <https://www.researchgate.net/publication/321058112>
- Sondo P, Tahita MC, Ilboudo H et al (2022) Boosting the impact of seasonal malaria chemoprevention (SMC) through simultaneous screening and treatment of household members of children receiving SMC in Burkina Faso: a protocol for a randomized open label trial. *Arch Public Health* 80:41. <https://doi.org/10.1186/s13690-022-00800-x>
- Staedke SG, Maiteki-Sebuguzi C, Rehman AM et al (2018) Assessment of community-level effects of intermittent preventive treatment for malaria in schoolchildren in Jinja, Uganda (START-IPT trial): a cluster-randomised trial. *Lancet Glob Health* 6:e668–e679. [https://doi.org/10.1016/S2214-109X\(18\)30126-8](https://doi.org/10.1016/S2214-109X(18)30126-8)
- Starzengruber P, Fuehrer H-P, Ley B et al (2014) High prevalence of asymptomatic malaria in south-eastern Bangladesh. *Malar J* 13:16. <https://doi.org/10.1186/1475-2875-13-16>
- Sturrock HJW, Hsiang MS, Cohen JM et al (2013) Targeting asymptomatic malaria infections: active surveillance in control and elimination. *PLoS Med* 10:e1001467. <https://doi.org/10.1371/journal.pmed.1001467>
- Tinto H, Valea I, Sorgho H et al (2014) The impact of clinical research activities on communities in rural Africa: the development of the Clinical Research Unit of Nanoro (CRUN) in Burkina Faso. *Malar J* 13:113. <https://doi.org/10.1186/1475-2875-13-113>
- Touray AO, Mobegi VA, Wamunyokoli F et al (2021) Prevalence of asymptomatic *P falciparum* gametocyte carriage among school children in Mbita, Western Kenya and assessment of the association between gametocyte density, multiplicity of infection and mosquito infection prevalence. *Wellcome Open Res* 5:259. <https://doi.org/10.12688/wellcomeopenres.16299.2>
- Unwin VT, Ahmed R, Noviyanti R et al (2020) Use of a highly-sensitive rapid diagnostic test to screen for malaria in pregnancy in Indonesia. *Malar J* 19:28. <https://doi.org/10.1186/s12936-020-3110-6>
- Vantaux A, Samreth R, Piv E et al (2018) Contribution to malaria transmission of symptomatic and asymptomatic parasite carriers in Cambodia. *J Infect Dis* 217:1561–1568. <https://doi.org/10.1093/infdis/jiy060>
- Vásquez AM, Medina AC, Tobón-Castaño A et al (2018) Performance of a highly sensitive rapid diagnostic test (HS-RDT) for detecting malaria in peripheral and placental blood samples from pregnant women in Colombia. *PLoS ONE* 13:e0201769. <https://doi.org/10.1371/journal.pone.0201769>
- Wilson AL, on behalf of the IPTc Taskforce (2011) A systematic review and meta-analysis of the efficacy and safety of intermittent preventive treatment of malaria in children (IPTc). *PLoS ONE* 6:e16976. <https://doi.org/10.1371/journal.pone.0016976>
- World Health Organization (2012) WHO policy recommendation: seasonal malaria chemoprevention (SMC) for *Plasmodium falciparum* malaria control in highly seasonal transmission areas of the Sahel sub-region in Africa. World Health Organization. <https://iris.who.int/handle/10665/337978>
- Yimam Y, Mohebali M, Abbaszadeh Afshar MJ (2022) Comparison of diagnostic performance between conventional and ultrasensitive rapid diagnostic tests for diagnosis of malaria: a systematic review and meta-analysis. *PLoS ONE* 17:e0263770. <https://doi.org/10.1371/journal.pone.0263770>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.