

# New particle formation from isoprene under upper-tropospheric conditions

<https://doi.org/10.1038/s41586-024-08196-0>

Received: 5 April 2024

Accepted: 10 October 2024

Published online: 4 December 2024

Open access

 Check for updates

Jiali Shen<sup>1,2,33</sup>, Douglas M. Russell<sup>3,33</sup>, Jenna DeVivo<sup>4</sup>, Felix Kunkler<sup>5</sup>, Rima Baalbaki<sup>1</sup>, Bernhard Mentler<sup>6</sup>, Wiebke Scholz<sup>6</sup>, Wenjuan Yu<sup>1</sup>, Lucía Caudillo-Plath<sup>3</sup>, Eva Sommer<sup>7,8</sup>, Emelda Ahongshangbam<sup>1,9</sup>, Dina Alfaouri<sup>1</sup>, João Almeida<sup>7,10</sup>, Antonio Amorim<sup>10</sup>, Lisa J. Beck<sup>3</sup>, Hannah Beckmann<sup>6,11</sup>, Moritz Berntheusel<sup>3</sup>, Nirvan Bhattacharyya<sup>4</sup>, Manjula R. Canagaratna<sup>12</sup>, Anouck Chassaing<sup>13</sup>, Romulo Cruz-Simbron<sup>14,15</sup>, Lubna Dada<sup>16</sup>, Jonathan Duplissy<sup>1,2</sup>, Hamish Gordon<sup>4</sup>, Manuel Granzin<sup>3</sup>, Lena Große Schulte<sup>3</sup>, Martin Heinritzi<sup>3</sup>, Siddharth Iyer<sup>17</sup>, Hannah Klebach<sup>3</sup>, Timm Krüger<sup>3</sup>, Andreas Kürten<sup>3</sup>, Markus Lampimäki<sup>1</sup>, Lu Liu<sup>16</sup>, Brandon Lopez<sup>18</sup>, Monica Martinez<sup>5</sup>, Aleksandra Morawiec<sup>8</sup>, Antti Onnela<sup>7</sup>, Maija Peltola<sup>1</sup>, Pedro Rato<sup>3,7</sup>, Mago Reza<sup>14,15</sup>, Sarah Richter<sup>3</sup>, Birte Rörup<sup>1</sup>, Milin Kaniyodical Sebastian<sup>19</sup>, Mario Simon<sup>3</sup>, Mihnea Surdu<sup>16</sup>, Kalju Tamme<sup>11</sup>, Roseline C. Thakur<sup>1</sup>, António Tomé<sup>20</sup>, Yandong Tong<sup>14,15</sup>, Jens Top<sup>16</sup>, Nsikanabasi Silas Umo<sup>19</sup>, Gabriela Unfer<sup>21</sup>, Lejish Vettikkat<sup>22</sup>, Jakob Weissbacher<sup>6</sup>, Christos Xenofontos<sup>23</sup>, Boxing Yang<sup>16</sup>, Marcel Zauner-Wieczorek<sup>3</sup>, Jiangyi Zhang<sup>1</sup>, Zhensen Zheng<sup>6,24</sup>, Urs Baltensperger<sup>16</sup>, Theodoros Christoudias<sup>23</sup>, Richard C. Flagan<sup>25</sup>, Imad El Haddad<sup>16</sup>, Heikki Junninen<sup>11</sup>, Ottmar Möhler<sup>19</sup>, Ilona Riipinen<sup>13</sup>, Urs Rohner<sup>26</sup>, Siegfried Schobesberger<sup>22</sup>, Rainer Volkamer<sup>14,15</sup>, Paul M. Winkler<sup>8</sup>, Armin Hansel<sup>6,24</sup>, Katrianne Lehtipalo<sup>1,27</sup>, Neil M. Donahue<sup>4,18,28,29</sup>, Jos Lelieveld<sup>5,23</sup>, Hartwig Harder<sup>5</sup>, Markku Kulmala<sup>1,2,30,31</sup>, Doug R. Worsnop<sup>1,12</sup>, Jasper Kirkby<sup>3,7,32</sup>, Joachim Curtius<sup>3,33</sup> & Xu-Cheng He<sup>1,27,32,33</sup>

Aircraft observations have revealed ubiquitous new particle formation in the tropical upper troposphere over the Amazon<sup>1,2</sup> and the Atlantic and Pacific oceans<sup>3,4</sup>. Although the vapours involved remain unknown, recent satellite observations have revealed surprisingly high night-time isoprene mixing ratios of up to 1 part per billion by volume (ppbv) in the tropical upper troposphere<sup>5</sup>. Here, in experiments performed with the CERN CLOUD (Cosmics Leaving Outdoor Droplets) chamber, we report new particle formation initiated by the reaction of hydroxyl radicals with isoprene at upper-tropospheric temperatures of  $-30\text{ °C}$  and  $-50\text{ °C}$ . We find that isoprene-oxygenated organic molecules (IP-OOM) nucleate at concentrations found in the upper troposphere, without requiring any more vapours. Moreover, the nucleation rates are enhanced 100-fold by extremely low concentrations of sulfuric acid or iodine oxoacids above  $10^5\text{ cm}^{-3}$ , reaching rates around  $30\text{ cm}^{-3}\text{ s}^{-1}$  at acid concentrations of  $10^6\text{ cm}^{-3}$ . Our measurements show that nucleation involves sequential addition of IP-OOM, together with zero or one acid molecule in the embryonic molecular clusters. IP-OOM also drive rapid particle growth at  $3\text{--}60\text{ nm h}^{-1}$ . We find that rapid nucleation and growth rates persist in the presence of  $\text{NO}_x$  at upper-tropospheric concentrations from lightning. Our laboratory measurements show that isoprene emitted by rainforests may drive rapid new particle formation in extensive regions of the tropical upper troposphere<sup>1,2</sup>, resulting in tens of thousands of particles per cubic centimetre.

Aerosol particles are important for climate because they scatter and absorb incoming solar radiation and seed cloud droplets by acting as cloud condensation nuclei (CCN). More CCN make clouds more reflective and may increase their extent and lifetime. Around half of CCN globally, and almost all in the upper troposphere<sup>6</sup>, arise from new particle formation, which involves the spontaneous condensation of low-volatility vapours in the atmosphere to form liquid or solid particles (particle nucleation). The initial stable molecular clusters form at diameters slightly above 1 nm. To become CCN, the new particles

should not be scavenged by pre-existing aerosol but grow by further vapour condensation to a size of around 50 nm and larger (particle growth). Although new particle formation has been extensively studied at ground-based sites<sup>7</sup>, little is known about the precursor vapours responsible for new particles in the remote upper troposphere and in marine regions. In particular, high concentrations of freshly formed particles are observed in the upper free troposphere over the Amazon<sup>1,2</sup> and the tropical Atlantic and Pacific oceans<sup>3,4</sup>. Chemical-transport models indicate that new particle formation persists across the tropical

upper troposphere over a latitude band covering about 40% of Earth's surface<sup>3</sup> and provides a global supply of CCN for low-altitude clouds in the sub-tropics and tropics<sup>3,8</sup>. However, the source of these particles has remained a puzzle for the past 20 years.

Early studies proposed that convective clouds could transport vapours from the boundary layer and form new particles in cold cloud outflows at high altitudes<sup>4,9</sup>. In the absence of an established mechanism, it was suggested that the oxidation products of isoprene ( $C_5H_8$ ) could contribute to the high mass concentrations of freshly formed particles observed in the upper troposphere over the Amazon<sup>10</sup>. Recent modelling studies have speculated that pure biogenic new particle formation from monoterpenes is the source of these particles<sup>11,12</sup>, but insufficient monoterpene concentrations have been found to account for the high particle-number concentrations<sup>13</sup>.

On the other hand, recent satellite observations have revealed high concentrations of isoprene in the upper troposphere over tropical South America, Central Africa and Southeast Asia, reaching up to around 1 ppbv during night-time<sup>5</sup> (1 ppbv is equivalent to around  $0.6 \times 10^{10}$  molecules  $cm^{-3}$  at 10 km altitude and  $2.5 \times 10^{10}$  molecules  $cm^{-3}$  at 0 km). The isoprene concentrations fall during daytime owing to high hydroxyl radical (OH) concentrations, which result in an isoprene lifetime of around 1–2 h. Furthermore, observations at 5,240 m altitude in the Bolivian Andes have found isoprene oxidation products in both gas and particle phases in air masses originating from the Amazon free troposphere<sup>14</sup>. Isoprene is emitted by vegetation, especially deciduous and broad-leaved evergreen trees, and is the most abundant hydrocarbon emitted into the atmosphere, after methane<sup>15</sup>. Median isoprene mixing ratios in the Amazon rainforest vary between 0.5 and 2.0 ppbv at night and 2.0–6.0 ppbv during the day<sup>16</sup>. Modelling studies show that isoprene is efficiently transported from the tropical boundary layer to the upper troposphere in deep convective clouds<sup>17</sup>.

Isoprene influences the oxidation capacity of the atmosphere<sup>18,19</sup> and contributes to the formation of secondary organic aerosol<sup>20–24</sup> particle mass, which—in turn—affects the climate<sup>25,26</sup>. The contribution of isoprene to secondary organic aerosol particle mass is at present believed to be dominated by the reactive uptake of isoprene dihydroxy epoxide (IEPOX)<sup>21,27</sup> and other IP-OOM<sup>28</sup>. Lower contributions are thought to arise from the condensation of low-volatility compounds formed during oxidation of the first-generation product, isoprene hydroxy hydroperoxide (ISOPOOH)<sup>29–31</sup>. IEPOX and ISOPOOH are isomers of  $C_5H_{10}O_3$  and cannot be separated by normal chemical-ionization mass spectrometry. The ability of isoprene to form new particles is considered negligible<sup>24</sup> and, moreover, isoprene inhibits new particle formation from monoterpenes under boundary-layer conditions<sup>32–34</sup>. However, it has so far remained unknown whether IP-OOM<sup>35</sup> can form new particles in the upper troposphere, where it is extremely cold and scavenging losses are small, and—if so—what are the associated nucleation and growth rates.

## CLOUD experiment

Here we report experiments performed in the CERN CLOUD chamber<sup>36</sup> to study new particle formation from the reaction of OH with isoprene at upper-tropospheric concentrations and temperatures of  $-30$  °C and  $-50$  °C. The experiments were performed during the CLOUD15 and CLOUD16 campaigns, September–November 2022 and 2023, respectively. Before injection into the chamber, the isoprene vapour was passed through a cryo-trap at  $-53$  °C to eliminate low-volatility contaminants (as confirmed by mass-spectrometer measurements). Further details of the CLOUD facility and its analysing instruments are provided in Methods.

The range of experimental parameters is summarized in Extended Data Table 1, together with the ambient upper-tropospheric conditions over the Amazon measured by Curtius et al.<sup>13</sup> during research flight (RF) 19. To maximize the IP-OOM detection efficiency, we combined the measurements of three mass spectrometers that use ammonium,

nitrate and bromide chemical-ionization (see Methods for further details), whereas the research flight uses nitrate ionization alone. In general, there is good overlap of the CLOUD experiments with the ambient conditions for this single flight (see Methods for further discussion). We note that, although the CLOUD chamber operates at atmospheric pressure, this does not affect the simulation of particle-nucleation dynamics at upper-tropospheric conditions. We performed experiments with nitrogen oxide (NO) concentrations varied between zero and around  $7 \times 10^9$   $cm^{-3}$ , characteristic of the outflow from electrified deep convective clouds. Trace amounts of sulfuric acid ( $H_2SO_4$ ), methanesulfonic acid ( $CH_3SO_3H$ ) and iodine oxoacids<sup>37</sup> ( $HIO_x$ ,  $x = 2, 3$ ) also exist in the upper troposphere from the oxidation of vapours such as sulfur dioxide ( $SO_2$ ), dimethyl sulfide (DMS;  $CH_3SCH_3$ ) and iodine ( $I_2$ ), respectively. For some experiments, therefore, we introduced upper-tropospheric concentrations of sulfuric acid or iodine oxoacids to explore their interactions with IP-OOM.

## IP-OOM


Hydroxyl radicals preferentially attack one of the terminal carbon atoms of isoprene, forming isoprene peroxy radicals (ISOPOO,  $C_5H_8(OH)(OO)$ ). In the absence of NO, ISOPOO reacts with hydroperoxy radicals ( $HO_2$ ) to form ISOPOOH ( $C_5H_8(OH)(OOH)$ )<sup>35,38</sup>. In the presence of NO, ISOPOO also reacts to form isoprene hydroxy nitrate (IHN,  $C_5H_8(OH)(ONO_2)$ ) and further products<sup>35</sup>. Further reaction of IHN with OH forms isoprene dihydroxy dinitrate ( $C_5H_8(OH)_2(ONO_2)_2$ ) and other second-generation products containing one or two nitrogen atoms. Further reactions of ISOPOOH with OH include the formation of its isomer, IEPOX<sup>21</sup>, as well as second-generation products containing zero or one nitrogen atom.

Oxidation by hydroxyl radicals of ISOPOOH<sup>29,30</sup>, IEPOX<sup>21,39,40</sup> and IHN are the main channels that feed the second-generation IP-OOM,  $IP_{0-2N}$ . Because the volatility of IP-OOM largely depends on their oxygen number<sup>41</sup>, we define  $IP_{0-2N}$  as  $C_iH_jO_kN_l$  with the requirements that  $i, j \geq 4$ ,  $l = 0-2$  and with a minimum oxygen number that takes the nitrogen content into account. As a nitrate group ( $-O-NO_2$ ) is considered to decrease the volatility of an organic compound by about the same factor as a hydroxyl group ( $-OH$ )<sup>41</sup>, we require  $(k-2l) \geq 4$ . We define  $IP_{0N}$ ,  $IP_{1N}$  and  $IP_{2N}$  as IP-OOM containing 0, 1 or 2 nitrogen atoms, respectively. In our study, therefore,  $IP_{0-2N}$  ( $= IP_{0N} + IP_{1N} + IP_{2N}$ ) excludes ISOPOOH and IEPOX but may include some first-generation IP-OOM from ISOPOO<sup>35</sup>. The  $IP_{0-2N}$  include IP-OOM monomers with up to five carbon atoms ( $C_5$ ) and dimers with up to ten carbons ( $C_{10}$ ), formed by reactions between two isoprene peroxy radicals ( $RO_2$ ), which produce covalently bound molecules.

As with other condensable vapours, both ISOPOOH and IEPOX deposit irreversibly on the CLOUD chamber walls at  $-50$  °C (ref. 42) and also at  $-30$  °C. The latter is confirmed by our measurements of a wall-loss lifetime for  $C_5H_{10}O_3$  at  $-30$  °C, which indicates irreversible loss on impact (Extended Data Fig. 1a). Although we cannot distinguish ISOPOOH from IEPOX, chemical model calculations<sup>35,43</sup> indicate that more than 80% of our  $C_5H_{10}O_3$  signal is ISOPOOH at  $5.5 \times 10^6$   $cm^{-3}$  OH (Methods and Extended Data Fig. 1b). Assuming this ISOPOOH fraction, we measure the total molar yield of  $IP_{0N}$  from the reaction of hydroxyl radicals with  $C_5H_{10}O_3$  in the absence of nitrogen oxides ( $NO_x$ ) to be 46% at  $-30$  °C and 55% at  $-50$  °C, with a systematic uncertainty of a factor of two (Extended Data Fig. 2). In the presence of  $NO_x$ , the  $IP_{0N}$  yield falls to around 38% owing to  $RO_2$  termination by NO or by further reactions that produce  $IP_{1N}$  and  $IP_{2N}$  (Extended Data Fig. 2). At upper-tropospheric NO concentrations of up to  $7 \times 10^9$   $cm^{-3}$ , we measure the nitrate IP-OOM fraction to be  $IP_{1-2N}/IP_{0-2N} = 23-88\%$ , in which  $IP_{1-2N} = IP_{1N} + IP_{2N}$ .

## Example experiments

A typical run sequence in the absence of  $NO_x$  is shown in Fig. 1. The experiment started at stage 1 by setting the internal mixing fans from 100% (high wall-loss rate) to 12% (standard operation) and switching on the



**Fig. 1 | Example new particle formation experiment from IP-OOM at  $-50\text{ }^{\circ}\text{C}$ , without  $\text{NO}_x$ .** **a–c**, Evolution of vapour concentrations (**a**), particle-nucleation rates at  $1.7\text{ nm}$ ,  $J_{1.7}$  (**b**), and naturally charged negative particle number size distribution ( $dN/d\log D_p$ ) and growth rates measured between  $3.2$  and  $8.0\text{ nm}$ ,  $\text{GR}_{3.2-8}$  ( $\text{nm h}^{-1}$ ), for total (naturally charged + neutral) particles (**c**). The black lines in **c** depict the linear fits of 50% appearance time of particles between  $3.2$  and  $8.0\text{ nm}$ . The vertical dashed lines and labels indicate the start of a new stage, at which the experimental conditions were adjusted. Trace sulfuric and iodic acid contaminants are present at the start of the run at concentrations of  $1\text{--}5 \times 10^4\text{ cm}^{-3}$ . Sulfur dioxide is injected and progressively increased during stages 5–8, which produces steps in  $\text{H}_2\text{SO}_4$ . The dotted black curve in **b** shows the expected  $\text{H}_2\text{SO}_4\text{--NH}_3$  nucleation rate, conservatively assuming that  $\text{NH}_3$  is present at the  $4\text{ pptv}$  limit of detection. Stages 1 and 3 are, respectively, under neutral (ion-free) and beam (ion-enhanced) conditions, whereas all of the other stages are under galactic cosmic ray (GCR; natural ion concentrations) conditions. The experimental conditions are: isoprene =  $0.20\text{--}0.41\text{ ppbv}$  ( $6.2\text{--}13.0 \times 10^9\text{ cm}^{-3}$ ),  $\text{O}_3 = 84\text{--}96\text{ ppbv}$  ( $2.6\text{--}3.0 \times 10^{12}\text{ cm}^{-3}$ ),  $\text{I}_2 = 3.7\text{--}23.0 \times 10^5\text{ cm}^{-3}$ ,  $\text{SO}_2 = 0\text{--}4 \times 10^9\text{ cm}^{-3}$ ,  $\text{OH} = 1\text{--}3 \times 10^6\text{ cm}^{-3}$ ,  $\text{HO}_2 = 4.7\text{--}6.0 \times 10^7\text{ cm}^{-3}$ ,  $\text{HO}_2/\text{OH}$  ratio =  $18\text{--}37$ ,  $\text{RH} = 62\%$ ,  $\text{NO}$  less than limit of detection ( $7\text{ pptv}$ ) and temperature =  $-49\text{ }^{\circ}\text{C}$ .

ultraviolet (UV) light, which photolysed ozone to produce  $2 \times 10^6\text{ cm}^{-3}$  OH and  $6 \times 10^7\text{ cm}^{-3}$  HO<sub>2</sub> radicals. Stages 1, 2 and 3 were, respectively, under different ionization conditions: neutral (all ions swept from the chamber by high-voltage electrodes); natural ionization from galactic cosmic rays (high voltage switched off); and pion beam (upper-tropospheric ion concentrations). The nucleation rate,  $J_{1.7}$ , is the measured flux of particles passing a  $1.7\text{-nm}$  threshold size. Because  $J_{1.7}$  varied little between these stages (Fig. 1b), it indicates relatively little sensitivity to ion concentrations. Before stage 5, the nucleation rate was  $0.48\text{--}0.73\text{ cm}^{-3}\text{ s}^{-1}$ , whereas IP<sub>ON</sub> were around  $1.6 \times 10^7\text{ cm}^{-3}\text{ s}^{-1}$  and sulfuric acid and iodic oxoacids were both below  $10^5\text{ cm}^{-3}$ . These nucleation rates far exceed those expected for HIO<sub>x</sub> and H<sub>2</sub>SO<sub>4</sub>, conservatively assuming  $1 \times 10^8\text{ cm}^{-3}$  contaminant

ammonia (NH<sub>3</sub>) (refs. 37,44,45). As no other condensable vapours were present, this experiment shows that IP-OOM form new particles under upper-tropospheric conditions at  $-50\text{ }^{\circ}\text{C}$ .

During stages 5–8, SO<sub>2</sub> was introduced into the chamber in steps, while keeping all other experimental conditions fixed (Fig. 1a). This increased the H<sub>2</sub>SO<sub>4</sub> concentration from  $1.5 \times 10^4\text{ cm}^{-3}$  to between  $2.8 \times 10^5$  and  $1.3 \times 10^6\text{ cm}^{-3}$ . Despite these extremely low acid concentrations, which are equivalent to between 0.05 and 0.2 parts per trillion by volume (pptv) at  $10\text{ km}$  altitude, the nucleation rate increased from  $6.2$  to  $37\text{ cm}^{-3}\text{ s}^{-1}$  between stages 5 and 8, respectively (Fig. 1b). Once again, these nucleation rates far exceed those expected for H<sub>2</sub>SO<sub>4</sub>–NH<sub>3</sub>. The particle growth rates between  $3.2$  and  $8.0\text{ nm}$  increased from  $2.6$  to  $4.1\text{ nm h}^{-1}$  between stages 5 and 8 (Fig. 1c). Because the expected particle growth rate for  $1.3 \times 10^6\text{ cm}^{-3}$  acid is around  $0.15\text{ nm h}^{-1}$  (ref. 46), these high growth rates must largely result from condensation of IP<sub>ON</sub>.


In other experiments, we introduced small concentrations of HIO<sub>x</sub> in the presence of negligible H<sub>2</sub>SO<sub>4</sub>. Extended Data Fig. 3 shows an example at  $-50\text{ }^{\circ}\text{C}$ . Here IP<sub>ON</sub> were held constant at  $1.6 \times 10^7\text{ cm}^{-3}\text{ s}^{-1}$  and contaminant H<sub>2</sub>SO<sub>4</sub> was less than  $5.0 \times 10^4\text{ cm}^{-3}$ . The HIO<sub>x</sub> concentration was increased in steps from  $1.2 \times 10^5$  to  $8.6 \times 10^5\text{ cm}^{-3}$  and the nucleation rate increased from  $3.3$  to  $18\text{ cm}^{-3}\text{ s}^{-1}$ . These nucleation rates closely match those measured for the IP<sub>ON</sub>–H<sub>2</sub>SO<sub>4</sub> system, suggesting that either inorganic acid plays a similar role in nucleation by stabilizing the embryonic molecular clusters.

In Extended Data Fig. 4, we show two further examples at  $-50\text{ }^{\circ}\text{C}$ , without (left panels) and with (right panels) NO<sub>x</sub>. Both experiments started with acid concentrations below the limit of detection ( $2 \times 10^4\text{ cm}^{-3}$ ). Extended Data Fig. 4c shows a threefold enhancement in  $J_{1.7}$  with the transition from neutral to galactic cosmic ray conditions, reaching  $0.7\text{ cm}^{-3}\text{ s}^{-1}$  at  $2.3 \times 10^7\text{ cm}^{-3}$  non-nitrate IP-OOM (IP<sub>ON</sub>) but showing very little further increase at higher (beam) ionization rates. This indicates that ions can enhance the stability of IP-OOM molecular clusters at especially low concentrations of acids or IP-OOM. When NO<sub>x</sub> is present and under galactic cosmic ray conditions (Extended Data Fig. 4d), the initial nucleation rate is  $1.7\text{ cm}^{-3}\text{ s}^{-1}$  at  $3.5 \times 10^7\text{ cm}^{-3}$  IP<sub>ON</sub> plus  $1.3 \times 10^8\text{ cm}^{-3}$  IP<sub>1-2N</sub>. This comparison suggests that nitrate IP-OOM are less effective for nucleation than non-nitrate IP-OOM. In Extended Data Fig. 4b, sulfuric acid was increased in steps from  $2 \times 10^4$  to  $3 \times 10^6\text{ cm}^{-3}$  and  $J_{1.7}$  increased from  $1.7$  to  $43\text{ cm}^{-3}\text{ s}^{-1}$ , demonstrating that the synergy between IP-OOM and sulfuric acid also occurs in the presence of NO<sub>x</sub>.

## New particle formation rates

We show in Extended Data Fig. 5 our measurements of  $J_{1.7}$  versus IP<sub>ON</sub>, IP<sub>1-2N</sub> and IP<sub>0-2N</sub> at  $-50\text{ }^{\circ}\text{C}$ , in the absence of acids (near or below the limit of detection). Extended Data Fig. 5a shows that the nucleation rates increase from around  $0.006\text{ cm}^{-3}\text{ s}^{-1}$  to  $48\text{ cm}^{-3}\text{ s}^{-1}$  by increasing IP<sub>ON</sub> from  $6 \times 10^6$  to  $9 \times 10^7\text{ cm}^{-3}$ . By contrast, there is a relatively weak dependence of the nucleation rate on IP<sub>1-2N</sub> (Extended Data Fig. 5b,c, which cover a larger IP<sub>1-2N</sub> range from  $2.4 \times 10^6$  to  $2.5 \times 10^8\text{ cm}^{-3}$ ). Nevertheless, careful inspection of the data in Extended Data Fig. 5b at the lowest IP<sub>ON</sub> concentrations does show some dependence of  $J_{1.7}$  on nitrate IP-OOM. The combined measurements suggest that IP<sub>ON</sub> are more effective for nucleation than IP<sub>1-2N</sub>, or—equivalently—that nitrate IP-OOM have higher volatilities than non-nitrate IP-OOM. However, at colder temperatures in the upper troposphere, the volatilities of all IP-OOM will decrease and nitrate IP-OOM will contribute more strongly to nucleation.

In Fig. 2, we present our measurements of  $J_{1.7}$  versus vapour concentrations at  $-30\text{ }^{\circ}\text{C}$  and  $-50\text{ }^{\circ}\text{C}$ . Here we consider only the IP<sub>ON</sub> component of IP<sub>0-2N</sub>, following the discussion above. In our experiments, the range of acid concentrations (mostly below  $3 \times 10^6\text{ cm}^{-3}$ ) is representative of the upper free troposphere<sup>47,48</sup> and the isoprene concentrations ( $0.14\text{--}4.2 \times 10^{10}\text{ cm}^{-3}$ ) correspond to those measured in the upper troposphere over tropical rainforests<sup>5</sup>, as do the oxidant concentrations<sup>18</sup> ( $\text{OH} = 0.11\text{--}6.9 \times 10^7\text{ cm}^{-3}$ ,  $\text{HO}_2 = 0.6\text{--}17 \times 10^8\text{ cm}^{-3}$  and  $\text{HO}_2/\text{OH}$



**Fig. 2 | Particle-nucleation rates from IP-OOM at -30 °C and -50 °C, with variable NO<sub>x</sub>.** **a–d**, Nucleation rates at 1.7 nm,  $J_{1,7}$ , versus  $IP_{ON}$  without NO<sub>x</sub> present (**a**),  $IP_{ON}$  with NO<sub>x</sub> (**b**),  $HIO_x + H_2SO_4$  with and without NO<sub>x</sub> (**c**) and the product  $(HIO_x + H_2SO_4) \times IP_{ON}$  with and without NO<sub>x</sub> (**d**). Measurements without NO<sub>x</sub> are indicated by diamonds (without acids) and circles (with acids). Measurements with NO<sub>x</sub> are indicated by triangles (without acids) and squares (with acids). Hollow symbols indicate -30 °C and solid symbols indicate -50 °C. In **c**, nucleation rates measured at contaminant acid concentrations are assigned the acid limit of detection (around  $2 \times 10^4$  cm<sup>-3</sup>). The solid lines in **c** show the nucleation rates expected for H<sub>2</sub>SO<sub>4</sub> with 4 pptv NH<sub>3</sub> at -30 °C (red) and -50 °C (blue), both at 60% RH (ref. 44). The dashed and solid lines in **d** represent fits to the equation  $10^{a + \log_{10}(x) + b}$ , in which, for the dashed line,  $a = 1.241$  and  $b = -15.065$ , and for the solid line,  $a = 1.505$  and  $b = -19.948$ . Panels **a–c** show that

ratio = 11–118). These vapour concentrations are sufficient to drive rapid nucleation rates between 0.006 and 48 cm<sup>-3</sup> s<sup>-1</sup>, which greatly exceed HIO<sub>x</sub> and H<sub>2</sub>SO<sub>4</sub>-NH<sub>3</sub> nucleation under these conditions<sup>37,44,45</sup>. Such fast nucleation rates can readily account for the high particle-number concentrations of up to 20,000 cm<sup>-3</sup> observed between 8 and 14 km over the Amazon<sup>2</sup>.


The nucleation rates depend on the concentrations of both  $IP_{ON}$  (Fig. 2a,b) and total acid ( $HIO_x + H_2SO_4$ ; Fig. 2c). Figure 2d suggests that the nucleation rates are linearly dependent on the product of  $IP_{ON}$  and total acid concentration, indicating that the critical step is dimer formation of an inorganic acid with a single  $IP_{ON}$ . We note that the  $J_{1,7}$  measurements in Fig. 2d at -50 °C with acids and NO<sub>x</sub> (filled square

both  $IP_{ON}$  and total acid ( $HIO_x + H_2SO_4$ ) contribute to the nucleation rate. Panel **d** indicates that it is the product of  $IP_{ON}$  and total acid (that is, the dimer formation rate) that best describes the nucleation rate,  $J_{1,7}$ , as the data points cluster into two groups primarily characterized by temperature alone.  $IP_{1,2N}$  also contribute to particle nucleation but they are less effective than  $IP_{ON}$  (Extended Data Fig. 5). The experimental conditions are: isoprene = 0.04–1.50 ppbv ( $0.1–4.2 \times 10^{10}$  cm<sup>-3</sup>), O<sub>3</sub> = 1–590 ppbv ( $3.7 \times 10^{10}$  to  $1.8 \times 10^{13}$  cm<sup>-3</sup>), I<sub>2</sub> =  $0–7.5 \times 10^7$  cm<sup>-3</sup>, SO<sub>2</sub> =  $0–4.6 \times 10^9$  cm<sup>-3</sup>, OH =  $0.11–6.90 \times 10^7$  cm<sup>-3</sup>, HO<sub>2</sub> =  $0.6–17.0 \times 10^8$  cm<sup>-3</sup>, HO<sub>2</sub>/OH ratio = 11–118, NO = 0–0.22 ppbv, NO<sub>2</sub> = 0–0.77 ppbv, RH = 29–70% and temperature = -30 °C and -50 °C. The error bars represent the standard deviation of the measurement at steady state. All measurements are made under galactic cosmic ray conditions (natural ionization amounts).

symbols) are systematically around a factor of 2–3 lower than the measurements without NO<sub>x</sub> (filled circles). This is an artefact resulting from a systematic uncertainty in those measurements (filled square symbols), as it is absent when  $J_{2,5}$  is measured for the same events but with a different particle counter (Extended Data Fig. 6). We measure a 20-fold increase of  $J_{1,7}$  between -30 °C and -50 °C (Fig. 2d) owing to decreasing IP-OOM volatilities.

### Molecular content of nucleating clusters

We have confirmed the nucleation mechanism inferred from Fig. 2d by direct molecular measurements with an Atmospheric Pressure



**Fig. 3 | Molecular composition of charged and neutral clusters during IP-OOM nucleation without acids at  $-50^{\circ}\text{C}$ .** **a–d**, Mass defect (difference from integer mass) versus  $m/z$  during IP-OOM nucleation events without acid and without added  $\text{NO}_x$  (**a, b**) or with  $\text{NO}_x$  (**c, d**). The data points are coloured by the number of carbon (**a, b**) or nitrogen (**c, d**) atoms. The symbol area in **a** and **c** is proportional to the normalized signal intensity by total signal, whereas that in **b** and **d** is proportional to IP-OOM concentrations. The charger ions ( $\text{Br}^-$ ,  $\text{NO}_3^-$  and  $\text{NH}_4^+$ ) are removed from the molecular formula in **b** and **d**. The data show that IP-OOM—which are around  $\text{C}_5$  for the monomer or  $\text{C}_{10}$  for the dimer—nucleate at  $-50^{\circ}\text{C}$ . The experimental conditions in **a–d** are: isoprene = 0.19,


0.07, 0.28 and 0.07 ppbv,  $\text{O}_3$  = 196.0, 129.0, 1.9 and 1.8 ppbv,  $\text{OH}$  = 0.3, 0.3, 1.6 and  $4.6 \times 10^7 \text{ cm}^{-3}$ ,  $\text{HO}_2$  = 1.8, 2.2, 3.4 and  $6.8 \times 10^8 \text{ cm}^{-3}$ ,  $\text{NO}$  = 0, 0, 0.07 and 0.18 ppbv,  $\text{NO}_2$  = 0, 0, 0.10 and 0.54 ppbv,  $\text{RH}$  = 31, 57, 66 and 29% and temperature =  $-49$ ,  $-49$ ,  $-48$  and  $-48^{\circ}\text{C}$ , respectively. The concentrations of  $\text{IP}_{0\text{N}}$  were unmeasured (instrument was not available),  $3.4 \times 10^7$ , unmeasured and  $2.5 \times 10^7 \text{ cm}^{-3}$  and the concentrations of  $\text{IP}_{1-2\text{N}}$  were unmeasured, 0, unmeasured and  $1.5 \times 10^8 \text{ cm}^{-3}$ , respectively. The bromide chemical-ionization mass spectrometer was converted to measure charged clusters for the experiments shown in **a** and **c**, under ground-level and upper-tropospheric ion concentrations, respectively.

Interface Time-of-Flight (APi-TOF) mass spectrometer during nucleation events without acids at  $-50^{\circ}\text{C}$  (Fig. 3a,c). Negatively charged (ion-induced) nucleation involves sequential accretion of IP-OOM monomers ( $\text{C}_5$  band) or dimers ( $\text{C}_{10}$  band) to an initial  $\text{C}_i\text{H}_j\text{O}_x^-$  or  $\text{NO}_3^-$  ion. When  $\text{NO}_x$  is present, a sharp reduction can be seen in the concentration of the  $\text{C}_{15}$  clusters compared with  $\text{C}_{10}$  (Fig. 3c). By contrast, the no- $\text{NO}_x$  data (Fig. 3a) show a smooth sequential growth of the clusters out to the detection limit. This indicates that the  $\text{C}_{15}$  clusters with  $\text{NO}_x$  are relatively unstable and have a high evaporation rate back to  $\text{C}_{10}$  clusters, hence the high concentration of the latter. We infer that the  $\text{IP}_{1-2\text{N}}$  have higher volatility than  $\text{IP}_{0\text{N}}$  and are less effective for nucleation. Nevertheless,  $\text{C}_{10}$  and  $\text{C}_{15}$  molecular clusters are seen in Fig. 3c with more than 3N (including  $\text{NO}_3^-$  core ion), which must include contributions from  $\text{IP}_{1-2\text{N}}$  as well as  $\text{IP}_{0\text{N}}$ . This confirms that  $\text{IP}_{1-2\text{N}}$  do indeed participate in ion-induced nucleation, but they are less effective than  $\text{IP}_{0\text{N}}$ , which leads to a rate-limiting step from  $\text{C}_{10}$  to  $\text{C}_{15}$ . This is consistent with the previous conclusions drawn from measurements of nucleation rate.

The composition of neutral clusters and molecules during IP-OOM nucleation without and with  $\text{NO}_x$  is shown in Fig. 3b,d, respectively. Here no signal is detected above the  $\text{C}_{10}$  band owing to the relatively

low charging efficiency of the chemical-ionization mass spectrometers, compared with unit efficiency for ions and ion-induced (charged) clusters in the APi-TOF mass spectrometer. Nevertheless, the neutral data reveal a clear shift towards (higher-mass)  $\text{IP}_{1\text{N}}$  and  $\text{IP}_{2\text{N}}$  compounds after the addition of  $\text{NO}_x$ . Furthermore, comparison with the corresponding charged clusters (Fig. 3a,c) shows that nucleation favours the more highly oxygenated compounds with lower volatility.

In Extended Data Fig. 7, we show the molecular composition of charged clusters in the presence of trace amounts of sulfuric acid (Extended Data Fig. 7a,c,e) and iodic acid (Extended Data Fig. 7b,d) during IP-OOM nucleation at  $-50^{\circ}\text{C}$ . The data are without  $\text{NO}_x$  (Extended Data Fig. 7a–e) and with  $\text{NO}_x$  (Extended Data Fig. 7f). These measurements confirm that the same nucleation mechanism occurs after addition of acids as seen previously without acids, that is, sequential addition of IP-OOM monomers ( $\text{C}_5$ ) or dimers ( $\text{C}_{10}$ ) to a core acid ion. Here the negatively charged core ions comprise the pure monomer, dimer or trimer of sulfuric acid (Extended Data Fig. 7a,c) with  $\text{HSO}_4^-$ , or iodic acid with  $\text{IO}_3^-$  (Extended Data Fig. 7b,d), whereas the positively charged core ions (Extended Data Fig. 7e,f) comprise an  $\text{IP-OOM}^+$ . The



**Fig. 4 | Particle growth rates and particle composition.** **a, b**, Particle growth rates from 3.2 to 8.0 nm,  $GR_{3.2-8}$ , versus  $IP_{ON}$  (**a**) and  $IP_{0-2N}$  (**b**). The solid lines show the predicted  $GR_{3.2-8}$  at the kinetic limit assuming monomer condensation and without considering any dipole enhancement. The prediction assumes that IP-OOM have a general formula of C<sub>5</sub>H<sub>12</sub>O<sub>6</sub> and a density of 1.34 g cm<sup>-3</sup> (ref. 50). The dashed lines are linear fits to the experimental data of the form  $GR_{3.2-8} = 2.04 \times 10^{-7} \times IP_{ON}$  (**a**) and  $GR_{3.2-8} = 1.23 \times 10^{-7} \times IP_{0-2N}$  (**b**). **c, d**, Mass-defect plots showing the molecular composition of particles measured with the FIGAERO at -30 °C (Br-FIGAERO) and -50 °C (I-FIGAERO), with particle geometric mean sizes ranging from 6 to 20 nm. The colour legends indicate the number of atoms of carbon (**c**) and nitrogen (**d**). The symbol area is proportional to the normalized signal by the sum of measured signals. The charger ions are removed from the molecular formula. The annotations show the molecular formula of the particle-phase compounds. Our measurements show that both  $IP_{ON}$  and  $IP_{1-2N}$  drive rapid early particle growth and that C<sub>5</sub>H<sub>12</sub>O<sub>5-6</sub> are the main


condensing vapours in  $IP_{ON}$ . The experimental conditions in **a** and **b** are: isoprene = 0.05–1.50 ppbv (0.15–4.40 × 10<sup>10</sup> cm<sup>-3</sup>), O<sub>3</sub> = 1–592 ppbv (3.6 × 10<sup>10</sup> to 1.8 × 10<sup>13</sup> cm<sup>-3</sup>), I<sub>2</sub> = 0–1.1 × 10<sup>8</sup> cm<sup>-3</sup>, SO<sub>2</sub> = 0–4.9 × 10<sup>9</sup> cm<sup>-3</sup>, OH = 0.09–7.40 × 10<sup>7</sup> cm<sup>-3</sup>, HO<sub>2</sub> = 0.6–19.0 × 10<sup>8</sup> cm<sup>-3</sup>, NO = 0–0.26 ppbv, NO<sub>2</sub> = 0–0.80 ppbv, RH = 29–72% and temperature = -30 °C and -50 °C. The experimental conditions for in **c** are: isoprene = 1.1 ppbv, O<sub>3</sub> = 184 ppbv, I<sub>2</sub> = 2.9 × 10<sup>7</sup> cm<sup>-3</sup>, SO<sub>2</sub> = 2.7 × 10<sup>7</sup> cm<sup>-3</sup>, OH = 2.3 × 10<sup>6</sup> cm<sup>-3</sup>, HO<sub>2</sub> = 0.8 × 10<sup>8</sup> cm<sup>-3</sup>, NO = 0 ppbv, NO<sub>2</sub> = 0 ppbv, RH = 72%, temperature = -30 °C and DMS = 0 ppbv and those for in **d** are: isoprene = 0.2 ppbv, O<sub>3</sub> = 1 ppbv, I<sub>2</sub> = 0, SO<sub>2</sub> = 1.2 × 10<sup>8</sup> cm<sup>-3</sup>, OH = 5.2 × 10<sup>7</sup> cm<sup>-3</sup>, HO<sub>2</sub> = 1.2 × 10<sup>9</sup> cm<sup>-3</sup>, NO = 0.11 ppbv, NO<sub>2</sub> = 0.65 ppbv, RH = 38%, temperature = -48 °C and DMS = 0.16 ppbv. The vertical error bars represent the statistical uncertainty in the appearance-time growth-rate measurements derived from the 95% confidence interval on the growth-rate fit. The horizontal error bars represent the standard deviation of measured  $IP_{ON}$  or  $IP_{0-2N}$  during the growth period.

core acid anion serves only to stabilize the initial molecular cluster with a single IP-OOM; there is no evidence for accretion of further acid molecules as the clusters grow, as expected at these very low acid concentrations (3–7 × 10<sup>5</sup> cm<sup>-3</sup>).

### Particle growth rates

Our measurements indicate that  $IP_{ON}$  (and, less effectively,  $IP_{1-2N}$ ) are rapidly nucleating together with H<sub>2</sub>SO<sub>4</sub> and HIO<sub>x</sub> at -30 °C and -50 °C (Fig. 2). We therefore expect that IP-OOM will also drive rapid particle growth at larger sizes as the Kelvin (curvature) barrier falls

and progressively higher-volatility compounds are able to condense onto the particles—as previously seen for new particle formation from  $\alpha$ -pinene oxidation products<sup>49</sup>. We show in Fig. 4 our measurements of particle growth rates between 3.2 and 8.0 nm versus  $IP_{ON}$  (Fig. 4a) and  $IP_{0-2N}$  (Fig. 4b) for experiments with and without NO<sub>x</sub> at -30 °C and -50 °C. Figure 4a shows that  $IP_{1-2N}$  are strongly contributing to early growth of particles; the measured growth rates cannot be explained by  $IP_{ON}$  alone. This is confirmed in Fig. 4b, in which all data (with or without NO<sub>x</sub> and at either temperature) are consistent with having the same dependency of growth rate on  $IP_{0-2N}$ .



**Fig. 5 | Schematic of new particle formation from isoprene in the upper troposphere.** Isoprene from forests is efficiently transported at night by deep convective clouds into the upper troposphere. During daylight, the isoprene accumulated overnight, together with daytime-convected isoprene, reacts with hydroxyl radicals and  $\text{NO}_x$  from lightning to produce IP-OOM. The IP-OOM combine with trace ambient acids to produce high particle-number

concentrations at cold temperatures below  $-30^\circ\text{C}$ . The newly formed particles grow rapidly over several hours to days while following the descending air masses. This mechanism may provide an extensive source of CCN for shallow continental and marine clouds, which strongly influence Earth's radiative balance.

We find that isoprene at upper-tropospheric concentrations will drive particle growth rates between 3 and  $56\text{ nm h}^{-1}$ . These rapid growth rates imply that new IP-OOM particles can reach several tens of nanometres in size within a few hours, which will help to prevent them from evaporation when descending to lower altitudes and warmer temperatures. Within measurement uncertainties, particle growth rates at all temperatures below  $-30^\circ\text{C}$  are linearly dependent on the  $\text{IP}_{0-2\text{N}}$  concentration, and they reach the kinetic limit. As expected from their low concentrations, the growth rates show no correlation with  $\text{H}_2\text{SO}_4$  and  $\text{HIO}_x$  (not shown).

We have verified these observations by direct particle-phase measurements made with the Filter Inlet for Gases and AEROsols (FIGAERO; Fig. 4c,d). The main  $\text{IP}_{0-2\text{N}}$  compounds in the particles are two second-generation oxidation products ( $\text{C}_5\text{H}_{12}\text{O}_5$  and  $\text{C}_5\text{H}_{12}\text{O}_6$ ) formed from reactions of ISOPOOH with OH, which together constitute 30% of the total signal. The same two compounds have been previously identified in the particle phase by isoprene experiments at room temperature<sup>30</sup>. The remaining particle-phase compounds are largely  $\text{C}_{10}$  IP-OOM dimers, in agreement with their measured low volatilities. The non-nitrate  $\text{C}_{10}$  dimers are suppressed by the presence of  $\text{NO}_x$ , as seen by comparing Fig. 4c with Fig. 4d. We note that the nitrate IP-OOM in Fig. 4d are probably under-represented because we measured that they have lower evaporation temperatures and so may escape thermal-desorption measurement in the FIGAERO.

## Upper-tropospheric particle formation

In summary, we find that IP-OOM rapidly form new particles at upper-tropospheric concentrations and temperatures below  $-30^\circ\text{C}$ . Moreover, the nucleation rates are up to 100 times faster in the presence of extremely low concentrations of sulfuric acid or iodine oxoacids, reaching rates around  $30\text{ cm}^{-3}\text{ s}^{-1}$  at  $-50^\circ\text{C}$  and acid concentrations of  $10^6\text{ cm}^{-3}$ . In the presence of  $\text{NO}_x$ , a large fraction of IP-OOM—around 23–88%—are found to contain either one or two nitrogen atoms. We find that nitrate IP-OOM contribute relatively weakly to particle nucleation compared with non-nitrate IP-OOM at the same concentration. However, cooler temperatures will favour nucleation from nitrate isoprene products<sup>13</sup>. Both non-nitrate and nitrate IP-OOM are equally effective at driving rapid particle growth at several tens of  $\text{nm h}^{-1}$ , at all temperatures below  $-30^\circ\text{C}$ . We present in Extended Data Fig. 8 four nucleation rate measurements that schematically encapsulate the effect of acids and  $\text{NO}_x$  on IP-OOM nucleation.

Our measured nucleation and growth rates provide the mechanistic missing link connecting the presence of abundant isoprene in the tropical upper troposphere<sup>5</sup> with the high particle-number concentrations found at high altitudes over the Amazon<sup>2</sup>. Our findings reveal a new mechanism (Fig. 5) that switches on rapid particle nucleation in extensive regions of the upper troposphere. Isoprene emitted by tropical rainforests is efficiently transported by deep

convective clouds and released at cloud outflows in the upper free troposphere<sup>10,17</sup>. During night-time, high isoprene concentrations build up in the upper troposphere<sup>5</sup> as a result of relatively slow oxidation by ozone and nitrate radicals<sup>35</sup>. During daytime, the isoprene is rapidly oxidized by hydroxyl radicals and mixed with NO<sub>x</sub> from lightning to produce IP-OOM. These mix with trace ambient acids to drive rapid particle nucleation and growth at cold temperatures below around -30 °C. Peak IP-OOM concentrations—and therefore the fastest new particle formation rates—will occur shortly after sunrise when the isoprene accumulated during the night is oxidized during the first 1–2 h of daylight<sup>5</sup>. However, later in the day, the increase of OH and HO<sub>2</sub> will accelerate reactions that form acids and favour production of non-nitrate IP-OOM from daytime-convected isoprene, which may lead to further particle nucleation. The newly formed particles grow over periods of several days by further condensation of low-volatility vapours, including acids. Model studies show that particles nucleated in the upper free troposphere over the Amazon are gradually transported downwards on horizontal scales much larger than 1,000 km (ref. 8).

Isoprene is the most abundant non-methane hydrocarbon emitted into the atmosphere, but its ability to nucleate particles in the boundary layer is considered negligible. Our findings show, however, that isoprene emitted by forests can drive rapid particle nucleation and growth in the upper troposphere. After further growth and descent to lower altitudes, these particles may represent a globally important source of CCN for shallow continental and marine clouds, and so influence Earth's radiative balance. Isoprene from forests may therefore provide a major source of biogenic particles in both the present-day and pre-industrial atmospheres that are at present unaccounted in atmospheric chemistry and climate models.

## Online content

Any methods, additional references, Nature Portfolio reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41586-024-08196-0>.

- Krejci, R. Evolution of aerosol properties over the rain forest in Surinam, South America, observed from aircraft during the LBA-CLAIRE 98 experiment. *J. Geophys. Res. Atmos.* **108**, 4561 (2003).
- Andreae, M. O. et al. Aerosol characteristics and particle production in the upper troposphere over the Amazon Basin. *Atmos. Chem. Phys.* **18**, 921–961 (2018).
- Williamson, C. J. et al. A large source of cloud condensation nuclei from new particle formation in the tropics. *Nature* **574**, 399–403 (2019).
- Clarke, A. D. et al. Particle production in the remote marine atmosphere: cloud outflow and subsidence during ACE 1. *J. Geophys. Res. Atmos.* **103**, 16397–16409 (1998).
- Palmer, P. I., Marvin, M. R., Siddans, R., Kerridge, B. J. & Moore, D. P. Nocturnal survival of isoprene linked to formation of upper tropospheric organic aerosol. *Science* **375**, 562–566 (2022).
- Gordon, H. et al. Causes and importance of new particle formation in the present-day and preindustrial atmospheres. *J. Geophys. Res. Atmos.* **122**, 8739–8760 (2017).
- Kerminen, V.-M. et al. Atmospheric new particle formation and growth: review of field observations. *Environ. Res. Lett.* **13**, 103003 (2018).
- Wang, X., Gordon, H., Grosvenor, D. P., Andreae, M. O. & Carslaw, K. S. Contribution of regional aerosol nucleation to low-level CCN in an Amazonian deep convective environment: results from a regionally nested global model. *Atmos. Chem. Phys.* **23**, 4431–4461 (2023).
- Kulmala, M. et al. Deep convective clouds as aerosol production engines: role of insoluble organics. *J. Geophys. Res. Atmos.* **111**, 2005JD006963 (2006).
- Ekman, A. M. L. et al. Do organics contribute to small particle formation in the Amazonian upper troposphere? *Geophys. Res. Lett.* **35**, 2008GL034970 (2008).
- Zhu, J. et al. Decrease in radiative forcing by organic aerosol nucleation, climate, and land use change. *Nat. Commun.* **10**, 423 (2019).
- Zhao, B. et al. Global variability in atmospheric new particle formation mechanisms. *Nature* **631**, 98–105 (2024).
- Curtius, J. et al. Isoprene nitrates drive new particle formation in Amazon's upper troposphere. *Nature* <https://doi.org/10.1038/s41586-024-08192-4> (2024).
- Zha, Q. et al. Oxidized organic molecules in the tropical free troposphere over Amazonia. *Natl Sci. Rev.* **11**, nwad138 (2023).
- Guenther, A. et al. A global model of natural volatile organic compound emissions. *J. Geophys. Res. Atmos.* **100**, 8873–8892 (1995).
- Andreae, M. O. et al. The Amazon Tall Tower Observatory (ATTO): overview of pilot measurements on ecosystem ecology, meteorology, trace gases, and aerosols. *Atmos. Chem. Phys.* **15**, 10723–10776 (2015).
- Bardakov, R., Thornton, J. A., Riipinen, I., Krejci, R. & Ekman, A. M. L. Transport and chemistry of isoprene and its oxidation products in deep convective clouds. *Tellus B Chem. Phys. Meteorol.* **73**, 1–21 (2021).
- Lelieveld, J. et al. Atmospheric oxidation capacity sustained by a tropical forest. *Nature* **452**, 737–740 (2008).
- Squire, O. J. et al. Influence of isoprene chemical mechanism on modelled changes in tropospheric ozone due to climate and land use over the 21st century. *Atmos. Chem. Phys.* **15**, 5123–5143 (2015).
- Clayes, M. Formation of secondary organic aerosols through photooxidation of isoprene. *Science* **303**, 1173–1176 (2004).
- Paulot, F. et al. Unexpected epoxide formation in the gas-phase photooxidation of isoprene. *Science* **325**, 730–733 (2009).
- Kroll, J. H., Ng, N. L., Murphy, S. M., Flagan, R. C. & Seinfeld, J. H. Secondary organic aerosol formation from isoprene photooxidation. *Environ. Sci. Technol.* **40**, 1869–1877 (2006).
- Massoli, P. et al. Ambient measurements of highly oxidized gas-phase molecules during the Southern Oxidant and Aerosol Study (SOAS) 2013. *ACS Earth Space Chem.* **2**, 653–672 (2018).
- Pandis, S. N., Paulson, S. E., Seinfeld, J. H. & Flagan, R. C. Aerosol formation in the photooxidation of isoprene and  $\beta$ -pinene. *Atmos. Environ. Part A Gen. Top.* **25**, 997–1008 (1991).
- Weber, J. et al. Chemistry-driven changes strongly influence climate forcing from vegetation emissions. *Nat. Commun.* **13**, 7202 (2022).
- Kanakidou, M. et al. Organic aerosol and global climate modelling: a review. *Atmos. Chem. Phys.* **5**, 1053–1123 (2005).
- Budisulistiorini, S. H. et al. Real-time continuous characterization of secondary organic aerosol derived from isoprene epoxydiols in downtown Atlanta, Georgia, using the Aerodyne Aerosol Chemical Speciation Monitor. *Environ. Sci. Technol.* **47**, 5686–5694 (2013).
- Lamkaddam, H. et al. Large contribution to secondary organic aerosol from isoprene cloud chemistry. *Sci. Adv.* **7**, eabe2952 (2021).
- St. Clair, J. M. et al. Kinetics and products of the reaction of the first-generation isoprene hydroxy hydroperoxide (ISOPPOOH) with OH. *J. Phys. Chem. A* **120**, 1441–1451 (2016).
- Krechmer, J. E. et al. Formation of low volatility organic compounds and secondary organic aerosol from isoprene hydroxyhydroperoxide low-NO oxidation. *Environ. Sci. Technol.* **49**, 10330–10339 (2015).
- Liu, J. et al. Efficient isoprene secondary organic aerosol formation from a non-IEPOX pathway. *Environ. Sci. Technol.* **50**, 9872–9880 (2016).
- Kiendler-Scharr, A. et al. New particle formation in forests inhibited by isoprene emissions. *Nature* **461**, 381–384 (2009).
- Heinritzi, M. et al. Molecular understanding of the suppression of new-particle formation by isoprene. *Atmos. Chem. Phys.* **20**, 11809–11821 (2020).
- McFiggans, G. et al. Secondary organic aerosol reduced by mixture of atmospheric vapours. *Nature* **565**, 587–593 (2019).
- Wennberg, P. O. et al. Gas-phase reactions of isoprene and its major oxidation products. *Chem. Rev.* **118**, 3337–3390 (2018).
- Kirkby, J. et al. Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation. *Nature* **476**, 429–433 (2011).
- He, X.-C. et al. Role of iodine oxoacids in atmospheric aerosol nucleation. *Science* **371**, 589–595 (2021).
- Crutzen, P. J. et al. High spatial and temporal resolution measurements of primary organics and their oxidation products over the tropical forests of Surinam. *Atmos. Environ.* **34**, 1161–1165 (2000).
- Bates, K. H. et al. Gas phase production and loss of isoprene epoxydiols. *J. Phys. Chem. A* **118**, 1237–1246 (2014).
- Jacobs, M. I., Darer, A. I. & Elrod, M. J. Rate constants and products of the OH reaction with isoprene-derived epoxides. *Environ. Sci. Technol.* **47**, 12868–12876 (2013).
- Kroll, J. H. & Seinfeld, J. H. Chemistry of secondary organic aerosol: formation and evolution of low-volatility organics in the atmosphere. *Atmos. Environ.* **42**, 3593–3624 (2008).
- Liu, Y. J., Herdinger-Blatt, I., McKinney, K. A. & Martin, S. T. Production of methyl vinyl ketone and methacrolein via the hydroperoxyl pathway of isoprene oxidation. *Atmos. Chem. Phys.* **13**, 5715–5730 (2013).
- Wolfe, G. M., Marvin, M. R., Roberts, S. J., Travis, K. R. & Liao, J. The Framework for O-D Atmospheric Modeling (FOAM) v3.1. *Geosci. Model Dev.* **9**, 3309–3319 (2016).
- Dunne, E. M. et al. Global atmospheric particle formation from CERN CLOUD measurements. *Science* **354**, 1119–1124 (2016).
- He, X.-C. et al. Iodine oxoacids enhance nucleation of sulfuric acid particles in the atmosphere. *Science* **382**, 1308–1314 (2023).
- Stolzenburg, D. et al. Enhanced growth rate of atmospheric particles from sulfuric acid. *Atmos. Chem. Phys.* **20**, 7359–7372 (2020).
- Lucas, D. D. & Prinn, R. G. Tropospheric distributions of sulfuric acid-water vapor aerosol nucleation rates from dimethylsulfide oxidation. *Geophys. Res. Lett.* **30**, 2136 (2003).
- Zauner-Wieczorek, M. et al. Mass spectrometric measurements of ambient ions and estimation of gaseous sulfuric acid in the free troposphere and lowermost stratosphere during the CAFE-EU/BLUESKY campaign. *Atmos. Chem. Phys.* **22**, 11781–11794 (2022).
- Stolzenburg, D. et al. Rapid growth of organic aerosol nanoparticles over a wide tropospheric temperature range. *Proc. Natl Acad. Sci.* **115**, 9122–9127 (2018).
- Clark, C. H. et al. Temperature effects on secondary organic aerosol (SOA) from the dark ozonolysis and photo-oxidation of isoprene. *Environ. Sci. Technol.* **50**, 5564–5571 (2016).



**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024

<sup>1</sup>Institute for Atmospheric and Earth System Research/Physics, Faculty of Science, University of Helsinki, Helsinki, Finland. <sup>2</sup>Helsinki Institute of Physics, University of Helsinki, Helsinki, Finland. <sup>3</sup>Institute for Atmospheric and Environmental Sciences, Goethe University Frankfurt, Frankfurt am Main, Germany. <sup>4</sup>Center for Atmospheric Particle Studies, Carnegie Mellon University, Pittsburgh, PA, USA. <sup>5</sup>Atmospheric Chemistry Department, Max Planck Institute for Chemistry, Mainz, Germany. <sup>6</sup>Institute for Ion Physics and Applied Physics, University of Innsbruck, Innsbruck, Austria. <sup>7</sup>CERN, the European Organization for Nuclear Research, Geneva, Switzerland. <sup>8</sup>Faculty of Physics, University of Vienna, Wien, Austria. <sup>9</sup>Department of Chemistry, University of Helsinki, Helsinki, Finland. <sup>10</sup>CENTRA and Faculdade de Ciências da

Universidade de Lisboa, Lisboa, Portugal. <sup>11</sup>Department of Environmental Physics, University of Tartu, Tartu, Estonia. <sup>12</sup>Aerodyne Research Inc., Billerica, MA, USA. <sup>13</sup>Department of Environmental Science, Stockholm University, Stockholm, Sweden. <sup>14</sup>Department of Chemistry, University of Colorado Boulder, Boulder, CO, USA. <sup>15</sup>Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, CO, USA. <sup>16</sup>Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, Villigen, Switzerland. <sup>17</sup>Aerosol Physics Laboratory, Faculty of Engineering and Natural Sciences, Tampere University, Tampere, Finland. <sup>18</sup>Department of Chemical Engineering, Carnegie Mellon University, Pittsburgh, PA, USA. <sup>19</sup>Institute of Meteorology and Climate Research, Atmospheric Aerosol Research, Karlsruhe Institute of Technology, Karlsruhe, Germany. <sup>20</sup>Instituto Dom Luiz (IDL), Universidade da Beira Interior, Covilhã, Portugal. <sup>21</sup>Atmospheric Microphysics Department, Leibniz Institute for Tropospheric Research (TROPOS), Leipzig, Germany. <sup>22</sup>Department of Technical Physics, University of Eastern Finland, Kuopio, Finland. <sup>23</sup>Climate and Atmosphere Research Centre (CARE-C), The Cyprus Institute, Nicosia, Cyprus. <sup>24</sup>IONICON Analytik GmbH, Innsbruck, Austria. <sup>25</sup>Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA, USA. <sup>26</sup>TOFWERK, Thun, Switzerland. <sup>27</sup>Finnish Meteorological Institute, Helsinki, Finland. <sup>28</sup>Department of Engineering and Public Policy, Carnegie Mellon University, Pittsburgh, PA, USA. <sup>29</sup>Department of Chemistry, Carnegie Mellon University, Pittsburgh, PA, USA. <sup>30</sup>Joint International Research Laboratory of Atmospheric and Earth System Sciences, School of Atmospheric Sciences, Nanjing University, Nanjing, China. <sup>31</sup>Aerosol and Haze Laboratory, Beijing Advanced Innovation Center for Soft Matter Science and Engineering, Beijing University of Chemical Technology, Beijing, China. <sup>32</sup>Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge, UK. <sup>33</sup>These authors contributed equally: Jiali Shen, Douglas M. Russell. <sup>✉</sup>e-mail: [Jasper.Kirkby@cern.ch](mailto:Jasper.Kirkby@cern.ch); [curtius@iauw.uni-frankfurt.de](mailto:curtius@iauw.uni-frankfurt.de); [xucheng.he@helsinki.fi](mailto:xucheng.he@helsinki.fi)

## Methods

### The CLOUD experiments

The CERN CLOUD chamber<sup>36</sup> was used to conduct the experiments presented in this study. CLOUD is an electropolished, stainless-steel, 26.1-m<sup>3</sup> chamber designed to study new particle formation under the full range of tropospheric and lower-stratospheric conditions. The thermal housing around the chamber is able to control the temperature from 208 to 373 K with high precision ( $\pm 0.1$  K)<sup>51</sup>. CLOUD was operated at a pressure of approximately  $965 \pm 5$  mbar in this study. To avoid cross-contamination between different experimental programmes and to achieve extremely low NH<sub>3</sub> concentrations, the chamber is cleaned by rinsing the chamber walls with ultrapure water and heating to 373 K for more than 24 h. To maintain cleanliness and ensure minimal contamination, ultrapure synthetic air—derived from mixing cryogenic liquids (21% oxygen and 79% nitrogen)—is continuously injected into the chamber. The chamber is characterized by a low loss rate, with condensation sink values comparable with those observed in pristine environments.

Various light sources are positioned in the CLOUD chamber to selectively drive photochemistry. OH production is initiated by illuminating O<sub>3</sub> with a UV fibre-optic system, a combination of four 200-W Hamamatsu Hg-Xe lamps with wavelengths spanning 250 and 450 nm, a krypton fluoride (KrF) excimer UV laser at 248 nm and a 52-W low-pressure mercury lamp centred at 254 nm. As well as O<sub>3</sub> photolysis, OH radicals are also produced by photochemical production from nitrous acid (HONO) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). Both the HONO and H<sub>2</sub>O<sub>2</sub> generators were designed specifically for CLOUD experiments. Following the same principle as an earlier study<sup>52</sup>, a gas–liquid mixture of HONO is synthesized from continuous mixing of H<sub>2</sub>SO<sub>4</sub> (Sigma Aldrich, 99%) with sodium nitrite (NaNO<sub>2</sub>, Sigma Aldrich, 99%) in a stainless-steel reactor<sup>53</sup>. HONO is transferred from liquid phase to gas phase by flowing nitrogen gas (1–2 l min<sup>-1</sup>) through the reactor. HONO is then introduced into the CLOUD chamber and photolysed by a UV light source centred at 385 nm to produce OH radicals and NO. The HONO reactor is continuously cooled to 5 °C and a cryo-trap is placed between the reactor and the chamber to remove excess water vapour and avoid ice blockage of the chamber input pipe. Gaseous H<sub>2</sub>O<sub>2</sub> is produced from bubbling N<sub>2</sub> gas through a H<sub>2</sub>O<sub>2</sub> solution. The H<sub>2</sub>O<sub>2</sub> solution is stored in a glass beaker contained in a stainless-steel container at a constant temperature of 5 °C. A different combination of UV sources is used to photolyse H<sub>2</sub>O<sub>2</sub> to produce different amounts of OH radicals.

A green light source centred at 528 nm is used to photolyse molecular iodine (I<sub>2</sub>). All light systems are continuously monitored by a spectrometer and an array of photodiodes at the bottom of the chamber. Dedicated actinometry experiments allow quantitative determination of actinic fluxes of the light system at different intensities.

Particle formation under different ionization regimes is simulated by combining a strong electric field ( $\pm 30$  kV) and the pion beam produced by the CERN Proton Synchrotron. The electric field eliminates natural ions in less than 1 s, thus creating ion-free conditions (neutral experiments). The pion beam produced by the CERN Proton Synchrotron enhances ion production on top of the galactic cosmic rays. Two magnetically coupled stainless-steel fans mounted at the top and bottom of the chamber enable uniform spatial mixing of particles and vapours within a few minutes. Iodine is injected into the chamber from a temperature-controlled evaporator containing crystalline iodine (I<sub>2</sub>, Sigma-Aldrich, 99.999% purity) at the bottom of the chamber. The SO<sub>2</sub> (Carbagas, 100 parts per million by volume (ppmv) in N<sub>2</sub>) and isoprene (PanGas, 1,000 ppmv in N<sub>2</sub>) are injected into the chamber from pressurized gas cylinders and the O<sub>3</sub> is introduced to the chamber by passing O<sub>2</sub> through an ozone generator.

The data presented in this study were collected in two consecutive CLOUD campaigns (CLOUD15 and CLOUD16). The CLOUD15 and CLOUD16 campaigns were carried out from September to November

in 2022 and 2023, respectively. Because the experiments reported in this study were carried out at extremely low temperatures ( $-30$  °C and  $-50$  °C), heat-insulation systems (CLOUD15) and active cooling systems (CLOUD16) were used to reduce measurement systematic error. The heat-insulation systems were primarily made with thermal insulation foam to isolate the instrument inlet system from ambient air. The active cooling systems involved circulating the air inside the chamber thermal housing, at the same temperature as the chamber, around the inlet systems of different instruments. The active cooling systems were also wrapped with thermal insulation foam to allow for more effective inlet cooling. These cooling systems were applied to all mass spectrometers and particle counters, except a butanol condensation particle chamber (CPC; TSI 3776), a nano-scanning mobility particle sizer (nano-SMPS, TSI 3938) and a long-SMPS (TSI 3082), which used a heat-insulation system in both campaigns to act as a standard to avoid systematic errors resulting from changing from the heat-insulation system to the active cooling system.

### Measurement of chemical composition

**Ozone (O<sub>3</sub>).** O<sub>3</sub> was monitored using a gas monitor (Thermo Environmental Instruments, TEI 49C).

**Hydroxyl radicals (OH).** The OH radical was measured by HORUS<sup>54</sup> (Hydroxyl Radical measurement Unit based on fluorescence Spectroscopy).

**Hydroperoxyl radical (HO<sub>2</sub>).** The HO<sub>2</sub> radical was primarily measured using the bromide chemical-ionization mass spectrometer coupled with a multi-scheme chemical-ionization inlet-2 (Br-MION2-CIMS)<sup>55</sup> and HORUS in both CLOUD15 and CLOUD16 campaigns. HORUS measures HO<sub>2</sub> by chemically converting it to OH by NO. However, the RO<sub>2</sub> radical (organic peroxy radicals) produced from isoprene oxidation may also contribute to the HO<sub>2</sub> signal measured by HORUS, as the reaction between RO<sub>2</sub> + NO can also produce OH radicals. By contrast, the HO<sub>2</sub> measurement by Br-MION2-CIMS is less ambiguous, as it is defined by the peak HO<sub>2</sub>Br<sup>-</sup> (ref. 55). However, the measurement of HO<sub>2</sub> by Br-MION2-CIMS is severely affected by air–water content<sup>55</sup>, making offline calibration difficult. Therefore, the HO<sub>2</sub> measurement by Br-MION2-CIMS was calibrated by HORUS under RO<sub>2</sub> radical-free conditions. The online calibration was carried out for every absolute humidity condition reported in this manuscript. During a small section in which the primary ions of Br-MION2-CIMS were saturated by either HONO or H<sub>2</sub>O<sub>2</sub>, either the low-pressure bromide chemical-ionization mass spectrometer or HORUS was used to complement the HO<sub>2</sub> measurement after intercomparing the data with Br-MION2-CIMS and HORUS during experiments without HONO and H<sub>2</sub>O<sub>2</sub>. The precision of OH and HO<sub>2</sub> data acquired by the HORUS instrument is quantified at 13% and 7%, respectively, with uncertainties calculated at  $1\sigma$  over a 10-min averaging period. Furthermore, the systematic error of the measurement is calculated to be 12% for OH and 30% for HO<sub>2</sub>.

**Nitrogen oxide (NO) and dioxide (NO<sub>2</sub>).** NO was measured by detecting the chemiluminescence of NO and O<sub>3</sub> using a chemiluminescence detector (ECO PHYSICS, CLD 780TR). This instrument was calibrated by a second NO monitor (ECO PHYSICS, CLD 780TR), which—in turn—was calibrated using the CMK5 Touch dilution system (Umwelttechnik MCZ GmbH) with a NO bottle (Praxair, 1.00 ppmv in N<sub>2</sub>) and synthetic air (Nippon Gases, hydrocarbon-free). The first detector, which provides data for this study, was found to contain background values that have been subtracted in this study. NO<sub>2</sub> was measured by a cavity-attenuated phase-shift nitrogen dioxide monitor (CAPS NO<sub>2</sub>, Aerodyne Research Inc.). Hourly, the instrument undergoes a 5-min background measurement of pure N<sub>2</sub> gas. During the 5-min background measurements, data have been interpolated to give a continuous time series. The NO<sub>2</sub> monitor was calibrated using a custom-made cavity-enhanced differential

optical absorption spectroscopy instrument<sup>56</sup>. After the subtraction of an average instrument background concentration, the final NO<sub>2</sub> concentration was obtained.

**Nitrous acid (HONO) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>).** Both HONO and H<sub>2</sub>O<sub>2</sub> were detected using bromide chemical-ionization mass spectrometry<sup>55</sup>, as they exhibit reasonable affinity with the bromide anion. Direct calibrations of these two species were not carried out on-site and the current estimation assumes that they share the same detection sensitivity as H<sub>2</sub>SO<sub>4</sub> (a low-limit estimation). Because these species serve as the precursors of OH and NO radicals, which were reliably traced, the concentrations of HONO and H<sub>2</sub>O<sub>2</sub> are not crucial to the reported results and are therefore omitted from this study.

Two bromide chemical-ionization systems were used to detect HONO and H<sub>2</sub>O<sub>2</sub>. The first system, Br-MION2-CIMS, offers sensitive detection of both species at concentrations below about 10<sup>10</sup> cm<sup>-3</sup>, with a detection limit of around 6 × 10<sup>6</sup> cm<sup>-3</sup> (H<sub>2</sub>O<sub>2</sub>) and 1.6 × 10<sup>5</sup> cm<sup>-3</sup> (HONO). However, in some experiments, the estimated HONO and H<sub>2</sub>O<sub>2</sub> concentrations exceeded 10<sup>10</sup> cm<sup>-3</sup>. The second system, Br-AIM-CIMS, uses bromide chemical-ionization at low pressure in combination with an active water feedback loop to control the Br-hydration in the ion molecule reactor and avoids saturation. Br-AIM-CIMS was used to measure concentrations from above the detection limit of 4.8 × 10<sup>7</sup> cm<sup>-3</sup> (HONO) and 3.3 × 10<sup>7</sup> cm<sup>-3</sup> (H<sub>2</sub>O<sub>2</sub>), based on a calibration factor of 3 × 10<sup>12</sup> for HONO and H<sub>2</sub>O<sub>2</sub>.

**Sulfur dioxide (SO<sub>2</sub>).** To measure the concentration of SO<sub>2</sub>, a gas monitor (Thermo Fisher Scientific Model 42i-TLE) was used. However, as the SO<sub>2</sub> concentrations in our experiments were usually below 5 × 10<sup>9</sup> cm<sup>-3</sup> (150 pptv), we also used the Br-MION2-CIMS to measure SO<sub>2</sub> (ref. 55) in both CLOUD15 and CLOUD16 campaigns. The measurement of SO<sub>2</sub> by Br-MION2-CIMS is substantially affected by air–water content, so we conducted online SO<sub>2</sub> calibration using the SO<sub>2</sub> monitor at both –30 °C and –50 °C. The derived calibration factors are 1.7 × 10<sup>13</sup> at –30 °C and 1.5 × 10<sup>11</sup> at –50 °C for CLOUD15 and 3.1 × 10<sup>11</sup> at –50 °C for CLOUD16. During the experiments, when the primary ions of Br-MION2-CIMS were saturated by either HONO or H<sub>2</sub>O<sub>2</sub>, the Br-AIM-CIMS was used to complement the SO<sub>2</sub> measurement. With an active water sensitivity control, Br-AIM-CIMS measures SO<sub>2</sub> concentrations from above the detection limit of 3 × 10<sup>7</sup> cm<sup>-3</sup> with a constant calibration factor of 20 × 10<sup>12</sup> at –30 °C and –50 °C.

**Sulfuric acid (H<sub>2</sub>SO<sub>4</sub>).** To ensure the quality of the reported data, we monitored H<sub>2</sub>SO<sub>4</sub> concentrations using two chemical-ionization mass spectrometers: the nitrate chemical-ionization mass spectrometer (NO<sub>3</sub>-CIMS) and the MION2-CIMS operating in bromide chemical-ionization mode (Br-MION2-CIMS<sup>55</sup>). Furthermore, isotopically labelled H<sup>15</sup>NO<sub>3</sub> was used during the CLOUD16 campaign to distinguish the nitrogen atom originating from the analyte with the reagent ion. The H<sub>2</sub>SO<sub>4</sub> calibration was carried out by two independent calibration systems. The first set-up used the original calibration box designed by Kürten et al.<sup>57</sup> along with their in-house calibration scripts. The second set-up is similar to the original version but with different physical dimensions. Also, the recently developed open-source MARFORCE model is used to simulate H<sub>2</sub>SO<sub>4</sub> production in both calibration set-ups<sup>55</sup>.

In total, we conducted seven calibration experiments at different stages of the CLOUD15 campaign, and each CIMS instrument was calibrated using both calibration set-ups. Two calibrations were performed for the Br-MION2-CIMS, resulting in equivalent H<sub>2</sub>SO<sub>4</sub> calibration factors of 157% and 149%. For the NO<sub>3</sub>-CIMS, five calibrations were carried out, resulting in equivalent calibration factors of 88%, 100%, 95%, 154% and 164%. Given that the NO<sub>3</sub>-CIMS provided most of the H<sub>2</sub>SO<sub>4</sub> concentration in this study, we use the calibration carried out immediately after the experiments for this study. This results in a

calibration factor of 6.2 × 10<sup>9</sup> cm<sup>-3</sup> for the NO<sub>3</sub>-CIMS and an equivalent calibration factor of 9.0 × 10<sup>9</sup> cm<sup>-3</sup> for the Br-MION2-CIMS. We use the minimum and maximum of the seven calibrations, ranging from 88% to 164%, as the systematic error of the H<sub>2</sub>SO<sub>4</sub> detection for CLOUD15. It is important to note that we had to change the optimal inlet flow rates of the Br-MION2-CIMS at –30 °C and –50 °C. The varying temperatures and flow rates result in different inlet loss rates, all of which have been accounted for in this dataset.

As well as the normal H<sub>2</sub>SO<sub>4</sub> calibration, we conducted a set of iodine oxoacid nucleation experiments at –10 °C, similar to those presented in ref. 37. The nucleation rates in these experiments are comparable with all of our earlier experiments, further enhancing our confidence in the reported acid concentrations.

In the CLOUD16 campaign, a total of seven calibration experiments were carried out. Two calibration experiments were conducted for the Br-MION2-CIMS, before and after the presented experiments. The results yield equivalent H<sub>2</sub>SO<sub>4</sub> calibration factors of 120% and 118%. For the labelled NO<sub>3</sub>-CIMS, six calibrations were performed in total, three before the isoprene experiments, resulting in equivalent calibration factors of 100%, 99% and 88%. It is important to note that, during the last few days of the isoprene experiments, the NO<sub>3</sub>-CIMS suffered from a pump failure that may have caused a shift (by up to 20%) in the calibration factor owing to a slight change in the sample flow. This potentially affects only two experiments in this study. To correct for this, we have assumed a linear correlation between the sample flow and calibration factor. The failing pumps were then replaced and the data from the rest of the experiments were calibrated after the presented experiments, with two calibrations that yielded equivalent calibration factors of 190% and 185%. This yields a calibration factor of 1 × 10<sup>10</sup> cm<sup>-3</sup> for the labelled NO<sub>3</sub>-CIMS and an equivalent calibration factor of 1.9 × 10<sup>10</sup> cm<sup>-3</sup> for the Br-MION2-CIMS. By considering all of the calibration experiments, the systematic error of H<sub>2</sub>SO<sub>4</sub> detection for CLOUD16 is estimated to range from 88% to 120%. Furthermore, using these two instruments, after applying their respective calibration factors, we compared the measured methanesulfonic acid concentrations from the CLOUD chamber at –50 °C. This comparison demonstrated a good agreement, confirming the accuracy of the calibrations.

**Iodine species.** We measured iodic acid (HIO<sub>3</sub>) and iodous acid (HIO<sub>2</sub>) using both the NO<sub>3</sub>-CIMS and Br-MION2-CIMS and we use the same calibration factor as H<sub>2</sub>SO<sub>4</sub> in the data analysis, similar to our earlier studies<sup>37,45,55,58,59</sup>. We used Br-MION2-CIMS to measure I<sub>2</sub>, which is detected at the collision limit, shown by our recent studies<sup>55,60</sup>.

**Isoprene.** Isoprene was measured by a proton transfer reaction mass spectrometer using the hydronium chemical-ionization method<sup>61</sup> (H<sub>3</sub>O-PTR-MS). This particular instrument used in this study is an adapted version, which is explained in greater detail previously<sup>62</sup>.

**ISOPOOH and IEPOX detection and separation.** Measuring and distinguishing between ISOPOOH and IEPOX can be experimentally challenging owing to their identical molecular formula (C<sub>8</sub>H<sub>10</sub>O<sub>3</sub>). As a result, mass-spectrometric methods often detect them together at the same exact mass in the same peak<sup>35</sup>. To address this issue, techniques such as tandem mass spectrometry have been used to separate ISOPOOH and IEPOX from each other<sup>29</sup>.

In this study, these two isomeric compounds were measured both by the Br-MION2-CIMS and the proton transfer reaction mass spectrometer 3 (ref. 63) operating in ammonium chemical-ionization mode (NH<sub>4</sub>-PTR3-CIMS<sup>64</sup>). NH<sub>4</sub>-PTR3-CIMS measured ISOPOOH and IEPOX primarily as clusters with ammonium cation, as the proton affinity (see the ‘Quantum-chemical calculations’ section) of NH<sub>3</sub> (204.25 kcal mol<sup>-1</sup>) is higher than that of 1,2-ISOPOOH (198.31 kcal mol<sup>-1</sup>), 4,3-ISOPOOH (195.51 kcal mol<sup>-1</sup>) and cis-β-IEPOX (204.11 kcal mol<sup>-1</sup>). In this study, we also aim to investigate the capability of the Br-MION2-CIMS in detecting

ISOPOOH and IEPOX. We calculate the formation free enthalpies of 1,2-ISOPOOH ( $-27.5 \text{ kcal mol}^{-1}$ ), 4,3-ISOPOOH ( $-26.9 \text{ kcal mol}^{-1}$ ) and cis- $\beta$ -IEPOX ( $-28.0 \text{ kcal mol}^{-1}$ ) with the bromide anion, respectively. We find that the formation free enthalpies are almost equal to the value of hypoiodous acid (HOI) clustered with the bromide anion ( $26.9 \text{ kcal mol}^{-1}$ ), as presented in ref. 55. Because the instrument used in ref. 55 and in this study is the same and the instrument tuning is identical, the fragmentation of these bromide anion cluster ions should be comparable. He et al.<sup>55</sup> calibrated both the  $\text{H}_2\text{SO}_4$  and the HOI, and the calibration factor of HOI was approximately two times larger than that of  $\text{H}_2\text{SO}_4$ . Therefore, the calibration factor used for  $\text{C}_5\text{H}_{10}\text{O}_3$  is two times the calibration factor for  $\text{H}_2\text{SO}_4$  in this study.

As neither the  $\text{NH}_4$ -PTR3-CIMS nor the Br-MION2-CIMS are able to distinguish between ISOPOOH and IEPOX, the reported  $\text{C}_5\text{H}_{10}\text{O}_3$  in this study is the sum of ISOPOOH and IEPOX. Earlier studies have shown that ISOPOOH is effectively lost to metal surfaces by converting it to methyl vinyl ketone (MVK) and methacrolein (MACR)<sup>42,65,66</sup>, whereas IEPOX is not affected by metal surfaces<sup>67</sup>. However, as the experiments in this study focus on extremely low temperatures ( $-30 \text{ }^\circ\text{C}$  and  $-50 \text{ }^\circ\text{C}$ ), the chamber wall itself may also serve as a cryo-trap<sup>68</sup> for both ISOPOOH and IEPOX. Therefore, it prevents us from using wall-loss-rate perturbation experiments to separate these two species at these temperatures.

To understand the distribution of ISOPOOH and IEPOX in  $\text{C}_5\text{H}_{10}\text{O}_3$ , we carry out a kinetic simulation using the reduced isoprene oxidation mechanism provided in ref. 35. The results are presented in Extended Data Fig. 1b. The simulation is carried out by the FOAM model<sup>43</sup>. The model requires input parameters such as isoprene, OH,  $\text{HO}_2$  and  $\text{O}_3$  concentrations measured by our instruments.

Another important parameter is the wall-loss rate of IP-OOM. We present an experiment in which we manipulate the loss rate of IP-OOM by turning off the light source and increasing the mixing fan spinning rate from 12% to 100% from the equilibrium conditions in Extended Data Fig. 1a. By turning off the light source, the production of IP-OOM stops. Furthermore, by increasing the fan speed, we increase the maximum wall-loss rate from approximately  $1.6 \times 10^{-3} \text{ s}^{-1}$  to  $8.5 \times 10^{-3} \text{ s}^{-1}$ . The decay rates of  $\text{C}_5\text{H}_{10}\text{O}_3$  and  $\text{C}_5\text{H}_{12}\text{O}_6$ , with lifetimes of 137 s and 112 s, respectively, are similar to the decay rate of  $\text{HIO}_3$  (129 s) and also, from previous measurements,  $\text{H}_2\text{SO}_4$ . Because  $\text{HIO}_3$  has an accommodation coefficient of unity to the chamber wall, we conclude that  $\text{C}_5\text{H}_{10}\text{O}_3$  and other species with lower volatilities have similar wall-loss rates. In this study, we apply a general wall-loss rate for these species of  $1.6 \times 10^{-3} \text{ s}^{-1}$ . This wall-loss rate is calculated from the measured  $\text{H}_2\text{SO}_4$  wall-loss rate by correcting the diffusivity of  $\text{C}_5\text{H}_{12}\text{O}_6$  at  $-30 \text{ }^\circ\text{C}$  using the method described by our earlier study<sup>58</sup>.

We further conduct simulations for all of our experiments using the same procedure, and the ratio of IEPOX in  $\text{C}_5\text{H}_{10}\text{O}_3$  versus OH concentration is presented in Extended Data Fig. 1b. As anticipated, the IEPOX ratio is positively correlated with OH concentrations. For further analysis, a fit with an expression of ratio of  $10^{(0.58 \times \log_{10}(\text{OH}) - 4.6)}$  is plotted.

**Gas-phase oxidized isoprene products.** The gas-phase measurement of IP-OOM was achieved by using a combination of  $\text{NO}_3$ -CIMS, Br-MION2-CIMS and  $\text{NH}_4$ -PTR3-CIMS. As defined in this study, only the species with carbon and oxygen numbers equal to or larger than 4 are considered in the  $\text{IP}_{0-2\text{N}}$ , which are primarily produced from OH oxidation of ISOPOOH and IEPOX with and without involving nitrogen oxides. Furthermore, the particle-phase  $\text{IP}_{0-2\text{N}}$  were monitored by a FIGAERO<sup>69</sup>, which operates with the bromide chemical-ionization method<sup>60</sup> in CLOUD15 (Br-FIGAERO-CIMS) and with the iodide chemical-ionization method in CLOUD16 (I-FIGAERO-CIMS). These chemical-ionization methods exhibit varying preferences for analytes. For example, the  $\text{NO}_3$ -CIMS is renowned for detecting highly oxygenated organic molecules<sup>70</sup> that contain more than 5 oxygen atoms. The  $\text{H}_3\text{O}$ -PTR-MS is the only one that can detect isoprene, whereas both the  $\text{NH}_4$ -PTR3-CIMS

and the Br-MION2-CIMS are capable of detecting semi-volatile organic compounds. Consequently, the combination of these CIMS methods enables the measurement of IP-OOM at different oxidation states.

It is worth mentioning our specialized approach to measuring IP-OOM using Br-MION2-CIMS during experiments involving excess HONO and/or  $\text{H}_2\text{O}_2$ , as described previously. In these experiments, the primary ions ( $\text{Br}^-$  and  $\text{H}_2\text{OBr}^-$ ) were substantially transformed into product ions such as  $\text{HONOBr}^-$ ,  $\text{H}_2\text{O}_2\text{Br}^-$  and  $(\text{H}_2\text{O}_2)_2\text{Br}^-$ . Consequently, the measurement of IP-OOM could be compromised if HONO and  $\text{H}_2\text{O}_2$  strongly bind with  $\text{Br}^-$ , thereby impeding the ligand exchange with IP-OOM. Therefore, we extensively compared the Br-MION2-CIMS measurements with those of  $\text{NO}_3$ -CIMS and  $\text{NH}_4$ -PTR3-CIMS during experiments with and without such primary ion saturation to ensure reliable measurements. We found that the Br-MION2-CIMS measurement remained uncompromised when we included  $\text{HONOBr}^-$ ,  $\text{H}_2\text{O}_2\text{Br}^-$  and  $(\text{H}_2\text{O}_2)_2\text{Br}^-$  as the primary ions. This is probably because of the relatively weak bonding of HONO and  $\text{H}_2\text{O}_2$  with  $\text{Br}^-$ , which enables effective charging of IP-OOM by allowing ligand exchange reaction. Quantum-chemical calculations further suggest that the formation free enthalpies of  $\text{HONOBr}^-$  and  $\text{H}_2\text{O}_2\text{Br}^-$  are  $-23.6$  and  $-21.2 \text{ kcal mol}^{-1}$ , respectively. These numbers are sufficiently lower than other molecules that are detected at the collision limit by Br-MION2-CIMS<sup>55</sup>.

To produce  $\text{IP}_{0-2\text{N}}$ , we conducted a set of experiments in which we varied the concentrations of isoprene (ranging from  $1.4 \times 10^9$  to  $4.2 \times 10^{10} \text{ cm}^{-3}$ ) and OH (ranging from 0.1 to  $6.9 \times 10^7 \text{ cm}^{-3}$ ) to alter the distribution of oxidation products<sup>35</sup>. To analyse the results of these experiments, we present a generic algorithm to calculate the total sum of gaseous  $\text{IP}_{0-2\text{N}}$  produced, with a focus on those with carbon and oxygen numbers greater than 3:

1. IP-OOM are independently identified by each of the CIMS instruments. Their responses to the isoprene oxidation in the chamber are observed to distinguish them from any background contaminations originating from either the chamber or the ion sources. If an individual  $\text{IP}_{0-2\text{N}}$  is affected by contaminants of the same molecular formula, its background, derived from the nearest cleaning stage, is subtracted from its concentrations.
2. If an  $\text{IP}_{0-2\text{N}}$  is detected by only one of the three CIMS, it is added to the total sum directly.
3. If several CIMS detect species with the same molecular formula, their measured signals are compared in pairs to derive a correlation coefficient. A pair is considered to measure identical molecules if the correlation coefficient is greater than 0.5. However, owing to the transfer of the  $\text{H}_2\text{SO}_4$  calibration factors to the measured  $\text{IP}_{0-2\text{N}}$  ( $\text{NO}_3$ -CIMS and Br-MION2-CIMS), the concentration of any molecule with a lower detection efficiency than  $\text{H}_2\text{SO}_4$  may be underestimated. The extent of this underestimation depends on the chemical-ionization method used, as the binding enthalpies of the analyte- $\text{Br}^-$ , analyte- $\text{NO}_3^-$  and analyte- $\text{NH}_4^+$  may differ. To address this, we add the highest measured concentration of the three CIMS to the  $\text{IP}_{0-2\text{N}}$  and discard the rest, as the highest concentration is probably the closest to the actual concentration.
4. If the correlation coefficient is less than 0.5, we consider that this pair represents two different molecular structures, that is, two isomers or conformers. In this case, both will be added to the  $\text{IP}_{0-2\text{N}}$ .

However, maintaining all three instruments to be operational throughout all experiments presents a challenge, for instance, the Br-MION2-CIMS operated in the API-TOF mode to measure charged clusters. Therefore, we excluded data collected during periods in which any one of the instruments was not available.

**Charged clusters.** Naturally charged clusters were measured with two API-TOF mass spectrometers (Aerodyne Research Inc.) operating at negative and positive ion mode<sup>71</sup>. The first instrument was equipped with a MION2 operating in the API-TOF mode (MION2-API-TOF)<sup>55,72</sup> by

deactivating the inlet voltages responsible for directing charged reagent ions into the sample flow. The second device was coupled with an ion-molecule reaction chamber (APi-TOF). Overall, the APi-TOF was less sensitive than the MION2-APi-TOF. The charged clusters reported in Fig. 3 were measured with the MION2-APi-TOF, which was validated by the APi-TOF. Because the MION2 inlet was operated in bromide chemical-ionization mode in some experiments, part of the data reported in Extended Data Fig. 7 was measured by the APi-TOF.

**Particle-phase measurements.** We measured the chemical composition of small particles using a FIGAERO coupled to a chemical-ionization mass spectrometer<sup>69</sup>. Particles were sampled from the CLOUD chamber onto a 5- $\mu\text{m}$ -pore polytetrafluoroethylene (PTFE) filter (Millipore-Sigma). Filter mass loading is dependent on particle distribution in the chamber, collection flow rate (typically 7–8 l min<sup>-1</sup>) and total collection time (1–2 h in this study). After particle collection, the filter was automatically moved to in front of the ion molecule reactor. The filter aligned with a sealed port that constantly flushes pure N<sub>2</sub>. In CLOUD15, the flow rate during chemical measurement was 3 l min<sup>-1</sup> and it was increased to 5 l min<sup>-1</sup> in CLOUD16 for more efficient heat transfer in a longer port. Pure N<sub>2</sub> was heated from room temperature up to 180 °C using programmed thermal desorption controlled by eyeon software v2.1.4.5. As the filter temperature increased, we detected lower-volatility molecules partitioning back into the gas phase. For the particle filter loadings in this study, we observed that all signals decreased back to the baseline by the end of the heating cycle, indicating no notable remaining mass.

Typically, FIGAERO-CIMS is operated using I<sup>-</sup> chemical-ionization in a reduced-pressure ion molecule reactor (about 120–150 mbar). Pure N<sub>2</sub> is flowed around a CH<sub>3</sub>I permeation tube (Vici) and through a 210Po ionizer (NRD LLC) to produce iodide ions. These polarizable ions effectively form adducts with oxygenated organic compounds, with a small fraction of interactions leading to charge transfer between the ion and neutral compound. In CLOUD15, we used Br<sup>-</sup> chemical-ionization to distinguish between our chemical-ionization reagent and iodine species inside the CLOUD chamber. The set-up is the same as iodide ionization mode except we exchange a CH<sub>2</sub>Br<sub>2</sub> permeation tube and heat it to 40 °C to increase permeation rates. These chemical-ionization techniques are both sensitive to oxygenated organic compounds, organics with nitrate and sulfate functional groups and inorganic acids<sup>60,69</sup>. Compounds chemically transformed through deprotonation or thermal decomposition have been excluded, as their parent molecule is unknown.

**Particle number size distribution.** The Neutral cluster and Air Ion Spectrometer<sup>73,74</sup> (NAIS) was used to measure the naturally charged particle number size distribution from 0.8 to 41 nm and the particle number size distribution (naturally charged + neutral) from 2 to 42 nm in both negative and positive polarities. The nano-condensation nucleus counter was used to measure the particle number size distribution between 1 and 3 nm. It consists of a particle size magnifier<sup>75</sup> (PSM, Airmodus Oy). The PSM, which is an aerosol pre-conditioner, uses diethylene glycol to grow aerosol particles as small as 1 nm to sizes that can be easily detected by a CPC<sup>75</sup>. Furthermore, a butanol CPC (TSI 3776) was used to measure the total number concentration of particles with diameters greater than 2.5 nm. A nano-scanning mobility particle sizer (TSI 3938)<sup>76</sup> coupled to a butanol CPC (TSI 3776), was used to measure the particle-size distribution within the range 6–65 nm, whereas particles larger than 65 nm were measured using a commercially available long-SMPS (TSI 3082) coupled to a water butanol CPC (TSI 3775).

#### Yield of IP<sub>ON</sub> from ISOPOOH

As shown in a previous section, the IP<sub>ON</sub> in this study is defined as species with C<sub>5</sub>O<sub>3</sub>. Therefore, ISOPOOH and IEPOX are not included in the IP<sub>ON</sub>. ISOPOOH and IEPOX are treated as the direct precursors of

IP<sub>ON</sub>, which in turn contribute to isoprene new particle formation. It is worth noting that both ISOPOOH and IEPOX undergo oxidation, producing compounds with C<sub>5</sub>O<sub>3</sub> ≥ 4. However, the reaction rate of ISOPOOH is approximately ten times larger than IEPOX<sup>35</sup>. To account for the difference in reaction-rate coefficients, we predict the ratio of IEPOX in C<sub>5</sub>H<sub>10</sub>O<sub>3</sub> using the data shown in Extended Data Fig. 1b based on the OH concentrations. Assuming that the concentration of IP<sub>ON</sub> is at equilibrium, the primary mechanism for IP<sub>ON</sub> loss is wall deposition, which is approximately equal to the production of IP<sub>ON</sub> from ISOPOOH and IEPOX. Therefore,

$$[\text{IP}_{\text{ON}}] \times k_{\text{wall}} = R \times (k_{\text{OH-ISOPOOH}} \times [\text{OH}] \times [\text{ISOPOOH}] + k_{\text{OH-IEPOX}} \times [\text{OH}] \times [\text{IEPOX}])$$

in which  $k_{\text{OH-ISOPOOH}}$  and  $k_{\text{OH-IEPOX}}$  are the reaction-rate coefficients of ISOPOOH (10<sup>-10</sup> cm<sup>3</sup> s<sup>-1</sup>) and IEPOX (10<sup>-11</sup> cm<sup>3</sup> s<sup>-1</sup>) with OH (ref. 35), respectively; [IP<sub>ON</sub>], [OH], [ISOPOOH] and [IEPOX] show concentrations and  $k_{\text{wall}}$  is the wall-loss rate of C<sub>5</sub>H<sub>10</sub>O<sub>3</sub>;  $R$  represents the yield of IP<sub>ON</sub> from C<sub>5</sub>H<sub>10</sub>O<sub>3</sub>.

We then define the reacted C<sub>5</sub>H<sub>10</sub>O<sub>3</sub> (cm<sup>-3</sup>) as:

$$\text{Reacted C}_5\text{H}_{10}\text{O}_3 = \frac{k_{\text{OH-ISOPOOH}} \times [\text{OH}] \times [\text{ISOPOOH}] + k_{\text{OH-IEPOX}} \times [\text{OH}] \times [\text{IEPOX}]}{k_{\text{wall}}}$$

The yield of IP<sub>ON</sub> from reacted C<sub>5</sub>H<sub>10</sub>O<sub>3</sub> is depicted in Extended Data Fig. 2. We find that the yields of IP<sub>ON</sub> are approximately 46% at -30 °C and 55% at -50 °C. However, it is essential to note that the detection of C<sub>5</sub>H<sub>10</sub>O<sub>3</sub>, IP<sub>ON</sub> and OH has various uncertainties. We estimate that the derived yield has an uncertainty of at least a factor of two, with the quantification of IP<sub>ON</sub> being the main source of uncertainty.

One further source of error in determining the yield is the contribution of highly oxygenated molecule production from the first-generation isoprene hydroxy peroxy radical (ISOPOO, C<sub>5</sub>H<sub>9</sub>O<sub>3</sub>) through auto-oxidation or dimer formation. For example, the reaction between two ISOPOO radicals can generate C<sub>10</sub>H<sub>18</sub>O<sub>4</sub>, and intramolecular H-shift followed by HO<sub>2</sub> termination of ISOPOO produces C<sub>5</sub>H<sub>10</sub>O<sub>5</sub>. Although these two molecules only contribute to a small fraction of IP<sub>ON</sub> in this study, other similar channels may contribute to a greater extent to IP<sub>ON</sub>, thereby reducing the yield of IP<sub>ON</sub> from C<sub>5</sub>H<sub>10</sub>O<sub>3</sub>. As disentangling first-generation and second-generation highly oxygenated molecules from isoprene oxidation is not the objective of this study, future research is necessary to investigate this direction.

#### Quantum-chemical calculations

Quantum-chemical methods are used to compute cluster formation free enthalpies and proton affinities. Initially, the Spartan'18 program is used for the conformational sampling with the MMFF method. Subsequently, density function theory (DFT) methods are used to optimize the molecules first at the B3LYP/6-31+G(d) level of theory, followed by optimization and frequency calculations at the  $\omega$ B97X-D/aug-cc-pVTZ-PP level of theory<sup>77,78</sup> on conformers within 2 kcal mol<sup>-1</sup> in relative electronic energies. Bromine pseudopotential definitions are obtained from the Environmental Molecular Sciences Laboratory (EMSL) basis set library<sup>79,80</sup>. The DFT calculations are carried out using the Gaussian 16 program<sup>81</sup>. To refine the DFT-calculated enthalpies, an extra coupled-cluster single-point energy correction is performed at the DLPNO-CCSD(T)/def2-QZVPP level of theory on the lowest-energy conformers. This coupled-cluster calculation is conducted using the ORCA program version 5.0.3 (ref. 82).

#### Calculation of the nucleation and growth rates

The nucleation rate,  $J_{1,7}$ , is calculated on the basis of PSM measurement of particles at a mobility diameter of 1.7 nm (1.4 nm in physical

diameter<sup>83</sup>), which are generally considered to be larger than their critical cluster sizes and thus stable.

To determine the nucleation rates, the time evolution of the particle concentration is analysed, taking into account various loss processes that also affect the concentration. However, because loss processes in a chamber setting differ from those in the atmosphere, the calculation method must be adjusted for chamber experiments<sup>84</sup>. Specifically, the nucleation rate ( $J_{1.7}$ ) is calculated by factoring in losses specific to the CLOUD chamber, such as dilution, wall and coagulation losses. We calculated  $J_{dp}$  as follows:

$$J_{1.7} = \frac{dN}{dt} + S_{dil} + S_{wall} + S_{coag}$$

in which  $dN/dt$  is the time derivative of the total particle concentration above a certain particle size (here  $>1.7$  nm for  $J_{1.7}$ ) and  $S_{dil}$ ,  $S_{wall}$  and  $S_{coag}$  are the particle losses owing to dilution, wall and coagulation. The details can be found in ref. 84. To calculate the coagulation sink, we used the combined particle-size distribution from three instruments (NAIS, nano-SMPS and long-SMPS).

Furthermore, the nucleation rate at  $2.5$  nm,  $J_{2.5}$ , derived from the butanol CPC and corrected by the same method described above, is calculated. The results are presented in Extended Data Fig. 6 in the same format as in Fig. 2. Because the CPC was not affected by the systematic upgrade in the cooling system between CLOUD15 and CLOUD16, it serves to distinguish subtle changes in our data. For example, the nucleation rates from experiments with  $\text{NO}_x$  (filled squares in Fig. 2) seem to be similar to the experiments without  $\text{NO}_x$  (filled circles in Fig. 2). This is probably a result of systematic errors introduced by changing either the cooling system or the instrument-calibration experiments. On the other hand, Extended Data Fig. 6 shows that experiments with  $\text{NO}_x$  have nucleation rates higher than the experiments without  $\text{NO}_x$ , therefore, isoprene nitrates ( $\text{IP}_{1-2\text{N}}$ ) do contribute, despite to a lesser extent compared with  $\text{IP}_{\text{ON}}$ , to particle nucleation.

To calculate particle growth rates, we use the 50% appearance-time method, as outlined in previous studies<sup>58,84,85</sup>. It is worth noting that the appearance-time method can overestimate growth rates when the impacts of coagulation (coagulation sink, coagulation source and particle coagulation growth) are non-negligible compared with the condensation growth, but the coagulation impact is rather small in the CLOUD experiments. For a deeper understanding of the molecular-level theory behind the method, we refer to the theoretical derivation presented in ref. 58. The particle number size distribution data used to calculate growth rates between  $3.2$  and  $8.0$  nm are measured by the NAIS. During previous experiments with  $\alpha$ -pinene and sulfuric acid, we have confirmed that the growth rates measured with the NAIS in total mode are similar to those measured with the DMA-train<sup>86</sup>.

### Comparison of experimental and ambient conditions

To compare the CLOUD experimental conditions with ambient conditions in the tropical upper troposphere, we summarize in Extended Data Table 1 the key chemical and physical parameters of the CLOUD experiments and the CAFE-Brazil (CB) flight campaign<sup>13</sup>. The CLOUD statistics are summarized from all experiments presented in this study, separated into two temperature conditions at  $-30$  °C and  $-50$  °C, respectively. Statistics of the CB flight campaign are derived from a single research flight, RF 19, samples T4 and T9, shown in Fig. 1 of ref. 13. All vapour concentrations are presented in units of molecules per  $\text{cm}^{-3}$ —the quantities as measured—for both CLOUD and CB. The values from the CB campaign are not corrected to their values at standard temperature and pressure, to allow for a direct comparison of the chemical and aerosol formation kinetics between the CLOUD experiments and the flight measurements. We

elaborate below on three aspects of this comparison: (1) isoprene non-nitrates ( $\text{IP}_{\text{ON}}$ ) and nitrates ( $\text{IP}_{1-2\text{N}}$ ); (2) atmospheric acids; and (3) impact of atmospheric pressure on particle nucleation.

**Distribution of  $\text{IP}_{\text{ON}}$  and  $\text{IP}_{1-2\text{N}}$ .** In general, the CLOUD experiments were designed to mimic ambient conditions as closely as possible. Key parameters such as temperature, relative humidity (RH), isoprene,  $\text{O}_3$ ,  $\text{NO}$  and  $\text{HO}_2/\text{OH}$  ratios are directly comparable between the CLOUD experiments and the CB measurements. The largest differences between CLOUD and CB are higher atmospheric pressure in CLOUD and higher  $\text{OH}/\text{HO}_2$  concentrations, resulting in a higher  $\text{HO}_2/\text{NO}$  ratio in CLOUD. The higher  $\text{OH}$  concentration in the CLOUD chamber is required to reproduce ambient  $\text{IP}$ -OOM concentrations at a chamber-wall-loss rate of approximately  $2 \times 10^{-3} \text{ s}^{-1}$ . In the upper troposphere, the condensation sink for low-volatility gaseous species could be several times or even up to one order of magnitude lower than the chamber-wall-loss rate.


Because this study aims to investigate the contribution of  $\text{IP}_{\text{ON}}$  and  $\text{IP}_{1-2\text{N}}$  to particle nucleation and growth, these parameters are critical for CLOUD to reproduce at atmospheric concentrations. A consequence of the relatively higher  $\text{HO}_2/\text{NO}$  ratio in CLOUD is that the  $\text{IP}_{\text{ON}}$  to  $\text{IP}_{1-2\text{N}}$  ratio is elevated compared with CB. However, because the higher operating pressure in CLOUD favours the formation of  $\text{IP}_{1-2\text{N}}$  by accelerating the reaction of organic peroxy radical ( $\text{RO}_2$ ) with  $\text{NO}$  as well as enhancing the organic nitrate formation branching ratio<sup>35</sup> (see Extended Data Table 1), this effect largely compensates for the higher  $\text{HO}_2/\text{NO}$  ratio in CLOUD. Nevertheless, regardless of the chemical details, which will be presented by follow-up studies, CLOUD successfully reproduces isoprene oxidation products  $\text{IP}_{\text{ON}}$  and  $\text{IP}_{1-2\text{N}}$ , in terms of both absolute values and relative ratios (Extended Data Table 1). The wide range of  $\text{IP}_{\text{ON}}$  and  $\text{IP}_{1-2\text{N}}$  concentrations and  $\text{IP}_{\text{ON}}/\text{IP}_{1-2\text{N}}$  ratios covered by CLOUD experiments enables CLOUD to reasonably simulate particle nucleation and growth dynamics, consistent with CB measurements during dawn hours (T4 period in Extended Data Table 1) and morning (T9 period in Extended Data Table 1). As the daylight hours proceed beyond the period measured by CB, both  $\text{OH}$  and  $\text{HO}_2$  concentrations will increase, whereas  $\text{NO}_x$  concentrations decrease, favouring the formation of  $\text{IP}_{\text{ON}}$  over  $\text{IP}_{1-2\text{N}}$ . Thus, the importance of  $\text{IP}_{\text{ON}}$  may be further enhanced after noon. It is also noteworthy that chemical distribution and nucleation dynamics might differ in other seasons and locations from those covered by CB measurements, such as the cases presented in Fig. S11 of ref. 5. Therefore, we believe that the wide range of conditions explored by CLOUD provides valuable data to enable global models to evaluate the impact of isoprene on new particle formation in other upper-tropospheric environments in which  $\text{NO}_x$  concentrations may differ from those measured by CB.

**Impact of sulfuric acid and iodine oxoacids.** In this study, we observe a large enhancement of  $\text{IP}$ -OOM nucleation rates from trace amounts of atmospheric acids (specifically,  $\text{H}_2\text{SO}_4$  and  $\text{HIO}_x$ ). The enhancement starts at acid concentrations of  $10^5 \text{ cm}^{-3}$  and, at an acid concentration of  $2 \times 10^6 \text{ cm}^{-3}$ , the particle-nucleation rate is approximately 100-fold faster than without added acids.

The CB measurements are unable to comment on this synergistic role of acids for  $\text{IP}$ -OOM particle nucleation owing to their  $\text{H}_2\text{SO}_4$  detection limit of several times  $10^6 \text{ cm}^{-3}$ . Nevertheless, acid enhancement of  $\text{IP}$ -OOM nucleation can be expected to occur in the atmosphere. Aircraft measurements indicate that approximately 10 pptv  $\text{SO}_2$  is ubiquitous in the global atmosphere between the marine boundary layer and the upper troposphere<sup>87,88</sup>. Global simulations also suggest that  $\text{H}_2\text{SO}_4$  concentrations greater than  $10^5 \text{ cm}^{-3}$  are widespread throughout the troposphere<sup>44</sup>. The global distribution of  $\text{HIO}_x$  is less well known and global simulations and aircraft measurements are needed to quantify its concentrations in

the upper troposphere. However, recent measurements of iodine oxide and particle-phase iodine in the upper troposphere<sup>89</sup> suggest that HIO<sub>x</sub> may also play a role in enhancing IP-OOM particle nucleation.

**Impact of atmospheric pressure on particle nucleation.** Cluster-forming interactions (such as nucleation) are of the form:



in which A and B are two molecules (or small clusters) forming the cluster C and C\* is a vibrationally excited state of the cluster containing the cluster energy  $E_{AB}$  as part of its internal vibrational energy. The excited cluster C\* will lose energy to the bath gas, M, with a concentration given by pressure through the ideal gas law. Cluster-forming interactions can therefore depend on ambient pressure<sup>90</sup>. When pressure is relatively 'low', reaction R3 will be the rate-limiting step for cluster formation and the overall rate will be of the third order (in A, B and M). However, when pressure is relatively high, reaction R1 will be the rate-limiting step and the rate will be of the second order (in A and B) and be independent of pressure, that is, at the so-called 'high-pressure limit'. The critical pressure occurs when the rates of R2 and R3 are equal and therefore depends on the lifetime (evaporation rate) of C\* through reaction R2, as well as the ambient pressure through reaction R3.

In practice, only systems with very few heavy atoms (four or fewer heavy atoms, in which 'heavy' excludes hydrogen) would show any meaningful pressure dependence in the atmosphere<sup>91</sup>, as quantified below. Because IP-OOM nucleation typically involves more than 15 heavy atoms, measurements in the CLOUD chamber at near 1 bar are directly applicable to the upper troposphere near 0.2 bar, provided the results are interpreted in terms of number concentrations (thus accounting for the dilution effect of reduced pressure) and not mixing ratios.

The (microcanonical) decomposition rates (inverse lifetimes) of the cluster are given by Rice–Ramsperger–Kassel–Marcus theory<sup>90</sup>. It depends strongly on the number of internal vibrational modes in C\*. As a rule, it can be estimated (by Rice, Ramsperger and Kassel theory)<sup>90</sup> roughly as the 'fractional excess free energy',  $\nu \left( \frac{e - e_0}{e} \right)^s$ , in which  $\nu$  is a typical frequency, 100 THz or so,  $e$  is the cluster energy above the ground vibrational state and  $e_0$  is the critical energy for decomposition (the cluster energy  $E_{AB}$ ) and  $s$  is an effective number of vibrational modes in the cluster. Approximately,  $e - e_0$  will be on the order  $kT$  (200 cm<sup>-1</sup>), whereas for the systems nucleating under atmospheric conditions,  $e_0$  will be on the order 800 cm<sup>-1</sup> or more. Thus,  $e$  will be on the order 1,000 cm<sup>-1</sup> and  $(e - e_0)/e$  will be on the order 0.2. Again, approximately and conservatively,  $s$  is  $3N - 7$ , in which  $N$  is the number of heavy (non-H) atoms in C\*. The '7' excludes external modes as well as the reaction coordinate. A system with five heavy atoms would have a decay coefficient of roughly  $2.6 \times 10^8$  s<sup>-1</sup>, whereas the collision frequency at 1 atm is near  $10^{10}$  s<sup>-1</sup>. Such a system would (barely) show some pressure dependence. By contrast, for isoprene oxidation products and H<sub>2</sub>SO<sub>4</sub>,  $N$  is probably 15, so  $s = 3N - 7 = 38$ . Given this, the microcanonical decay coefficients for these clusters will be on the order  $3 \times 10^{-13}$  s<sup>-1</sup>. This is extremely slow. In practice, it means that the energy distribution in cluster C will be entirely thermal, that is, given by a Boltzmann term at the ambient temperature, and the rates of cluster formation (and decomposition or evaporation) will be unaffected by pressure.

## Data availability

The full dataset shown in the figures and tables is available open access at <https://doi.org/10.5281/zenodo.13736557> (ref.92).

## Code availability

Codes conducting the analysis presented here can be obtained on reasonable request to the corresponding authors.

- Dias, A. et al. Temperature uniformity in the CERN CLOUD chamber. *Atmos. Meas. Tech.* **10**, 5075–5088 (2017).
- Taira, M. & Kanda, Y. Continuous generation system for low-concentration gaseous nitrous acid. *Anal. Chem.* **62**, 630–633 (1990).
- Xiao, M. et al. The driving factors of new particle formation and growth in the polluted boundary layer. *Atmos. Chem. Phys.* **21**, 14275–14291 (2021).
- Martinez, M. et al. Hydroxyl radicals in the tropical troposphere over the Suriname rainforest: airborne measurements. *Atmos. Chem. Phys.* **10**, 3759–3773 (2010).
- He, X.-C. et al. Characterisation of gaseous iodine species detection using the multi-scheme chemical ionisation inlet 2 with bromide and nitrate chemical ionisation methods. *Atmos. Meas. Tech.* **16**, 4461–4487 (2023).
- Thalman, R. & Volkamer, R. Inherent calibration of a blue LED-CE-DOAS instrument to measure iodine oxide, glyoxal, methyl glyoxal, nitrogen dioxide, water vapour and aerosol extinction in open cavity mode. *Atmos. Meas. Tech.* **3**, 1797–1814 (2010).
- Kürten, A., Rondo, L., Ehrhart, S. & Curtius, J. Calibration of a chemical ionization mass spectrometer for the measurement of gaseous sulfuric acid. *J. Phys. Chem. A* **116**, 6375–6386 (2012).
- He, X.-C. et al. Determination of the collision rate coefficient between charged iodine clusters and iodic acid using the appearance time method. *Aerosol Sci. Technol.* **55**, 231–242 (2021).
- Finkenzeller, H. et al. The gas-phase formation mechanism of iodic acid as an atmospheric aerosol source. *Nat. Chem.* **15**, 129–135 (2023).
- Wang, M. et al. Measurement of iodine species and sulfuric acid using bromide chemical ionization mass spectrometers. *Atmos. Meas. Tech.* **14**, 4187–4202 (2021).
- Graus, M., Müller, M. & Hansel, A. High resolution PTR-TOF: quantification and formula confirmation of VOC in real time. *J. Am. Soc. Mass. Spectrom.* **21**, 1037–1044 (2010).
- Canaval, E., Hyytiäinen, N., Schmidbauer, B., Fischer, L. & Hansel, A. NH<sub>4</sub><sup>+</sup> association and proton transfer reactions with a series of organic molecules. *Front. Chem.* **7**, 191 (2019).
- Breitenlechner, M. et al. PTR3: an instrument for studying the lifecycle of reactive organic carbon in the atmosphere. *Anal. Chem.* **89**, 5824–5831 (2017).
- Hansel, A., Scholz, W., Mentler, B., Fischer, L. & Berndt, T. Detection of RO<sub>2</sub> radicals and other products from cyclohexene ozonolysis with NH<sub>4</sub><sup>+</sup> and acetate chemical ionization mass spectrometry. *Atmos. Environ.* **186**, 248–255 (2018).
- Bernhammer, A.-K., Breitenlechner, M., Keutsch, F. N. & Hansel, A. Technical note: Conversion of isoprene hydroxy hydroperoxides (ISOPROOHs) on metal environmental simulation chamber walls. *Atmos. Chem. Phys.* **17**, 4053–4062 (2017).
- Rivera-Rios, J. C. et al. Conversion of hydroperoxides to carbonyls in field and laboratory instrumentation: observational bias in diagnosing pristine versus anthropogenically controlled atmospheric chemistry. *Geophys. Res. Lett.* **41**, 8645–8651 (2014).
- Mentler, B. *Measuring Isoprene Hydroxy Hydroperoxides and Isoprene Epoxidiols Using a Novel Funnel SRI-TOF-MS*. Thesis, Univ. Innsbruck (2017).
- Liu, Y. et al. Isoprene photochemistry over the Amazon rainforest. *Proc. Natl Acad. Sci.* **113**, 6125–6130 (2016).
- Lopez-Hilfiker, F. D. et al. A novel method for online analysis of gas and particle composition: description and evaluation of a Filter Inlet for Gases and AEROSols (FIGAERO). *Atmos. Meas. Tech.* **7**, 983–1001 (2014).
- Ehn, M. et al. A large source of low-volatility secondary organic aerosol. *Nature* **506**, 476–479 (2014).
- Junninen, H. et al. A high-resolution mass spectrometer to measure atmospheric ion composition. *Atmos. Meas. Tech.* **3**, 1039–1053 (2010).
- Rissanen, M. P., Mikkilä, J., Iyer, S. & Hakala, J. Multi-scheme chemical ionization inlet (MION) for fast switching of reagent ion chemistry in atmospheric pressure chemical ionization mass spectrometry (CIMS) applications. *Atmos. Meas. Tech.* **12**, 6635–6646 (2019).
- Manninen, H. E., Mirme, S., Mirme, A., Petäjä, T. & Kulmala, M. How to reliably detect molecular clusters and nucleation mode particles with Neutral cluster and Air Ion Spectrometer (NAIS). *Atmos. Meas. Tech.* **9**, 3577–3605 (2016).
- Mirme, S. & Mirme, A. The mathematical principles and design of the NAIS – a spectrometer for the measurement of cluster ion and nanometer aerosol size distributions. *Atmos. Meas. Tech.* **6**, 1061–1071 (2013).
- Vanhanen, J. et al. Particle size magnifier for nano-CN detection. *Aerosol Sci. Technol.* **45**, 533–542 (2011).
- Tröstl, J. et al. Fast and precise measurement in the sub-20 nm size range using a Scanning Mobility Particle Sizer. *J. Aerosol Sci.* **87**, 75–87 (2015).
- Chai, J.-D. & Head-Gordon, M. Long-range corrected hybrid density functionals with damped atom–atom dispersion corrections. *Phys. Chem. Chem. Phys.* **10**, 6615 (2008).
- Kendall, R. A., Dunning, T. H. & Harrison, R. J. Electron affinities of the first-row atoms revisited. Systematic basis sets and wave functions. *J. Chem. Phys.* **96**, 6796–6806 (1992).
- Feller, D. The role of databases in support of computational chemistry calculations. *J. Comput. Chem.* **17**, 1571–1586 (1996).
- Peterson, K. A., Figgen, D., Goll, E., Stoll, H. & Dolg, M. Systematically convergent basis sets with relativistic pseudopotentials. II. Small-core pseudopotentials and correlation consistent basis sets for the post-d group 16–18 elements. *J. Chem. Phys.* **119**, 11113–11123 (2003).
- Frisch, M. J. et al. Gaussian 16, Revision C.01 (Gaussian, Inc., 2016).
- Neese, F. The ORCA program system: the ORCA program system. *Wiley Interdiscip. Rev. Comput. Mol. Sci.* **2**, 73–78 (2012).
- Ku, B. K. & de la Mora, J. F. Relation between electrical mobility, mass, and size for nanodrops 1–6.5 nm in diameter in air. *Aerosol Sci. Technol.* **43**, 241–249 (2009).

84. Dada, L. et al. Formation and growth of sub-3-nm aerosol particles in experimental chambers. *Nat. Protoc.* **15**, 1013–1040 (2020).
85. Lehtipalo, K. et al. Methods for determining particle size distribution and growth rates between 1 and 3 nm using the Particle Size Magnifier. *Boreal Environ. Res.* **19**, 215–236 (2014).
86. Stolzenburg, D., Steiner, G. & Winkler, P. M. A DMA-train for precision measurement of sub-10 nm aerosol dynamics. *Atmos. Meas. Tech.* **10**, 1639–1651 (2017).
87. Thompson, C. R. et al. The NASA Atmospheric Tomography (ATom) mission: imaging the chemistry of the global atmosphere. *Bull. Am. Meteorol. Soc.* **103**, E761–E790 (2022).
88. Ranjithkumar, A. et al. Constraints on global aerosol number concentration, SO<sub>2</sub> and condensation sink in UKESM1 using ATom measurements. *Atmos. Chem. Phys.* **21**, 4979–5014 (2021).
89. Koenig, T. K. et al. Quantitative detection of iodine in the stratosphere. *Proc. Natl Acad. Sci.* **117**, 1860–1866 (2020).
90. Marcus, R. A. Unimolecular dissociations and free radical recombination reactions. *J. Chem. Phys.* **20**, 359–364 (1952).
91. Atkinson, R. et al. Evaluated kinetic and photochemical data for atmospheric chemistry: supplement III. IUPAC Subcommittee on Gas Kinetic Data Evaluation for Atmospheric Chemistry. *J. Phys. Chem. Ref. Data* **18**, 881–1097 (1989).
92. Shen, J. et al. New particle formation from isoprene under upper-tropospheric conditions: data sources. *Zenodo* <https://doi.org/10.5281/zenodo.13736557> (2024).
93. Pfeifer, J. et al. Measurement of ammonia, amines and iodine compounds using protonated water cluster chemical ionization mass spectrometry. *Atmos. Meas. Tech.* **13**, 2501–2522 (2020).

**Acknowledgements** We thank CERN for supporting CLOUD with important technical and financial resources and for providing a particle beam from the CERN Proton Synchrotron. Funding: this work was supported by Research Council of Finland ACCC Flagship projects 337550, 337549 and 337552; Research Council of Finland professorship (302958); Research Council of Finland's Centre of Excellence 346371; Research Council of Finland projects 349659, 346371, 345982, 1325656, 311932, 334792, 316114, 325647, 325681, 339489, 347782 and 355966; the Strategic Research Council (SRC) at the Research Council of Finland (352431); European Research Council project ATM-GTP (742206); European Union through Non-CO<sub>2</sub> Forcers and their Climate, Weather, Air Quality and Health Impacts (FOCI); European Research Council Consolidator Grant INTEGRATE grant 865799; EU MSCA Doctoral Network CLOUD-DOC 101073026; German Federal Ministry of Education and Research project

CLOUD-22 01LK2201A; US National Science Foundation grants AGS-2132089, AGS-2215489, CHE-2336463, AGS-1602086, AGS-1801329, AGS-1951514, AGS-1801280, AGS-2027252 and AGS-2215489; NASA ROSES programme grant 80NSSC19K0949; Swiss National Science Foundation grants 200021\_213071 and 216181; Knut and Alice Wallenberg Foundation projects 2021.0169 and 2021.0298; Federal Ministry of Education and Research (BMBF) financed project CLOUD-22 (01LK2201B); Untersuchung von Aerosolnukleation, Aerosolwachstum und Wolkenaktivierung an der CLOUD-Kammer am CERN zur Erforschung des Einflusses auf das Klima; Estonian Research Council project RVTT3 – 'CERN Science Consortium of Estonia' and project PRG714; Horizon 2020 research and innovation programme under grant 856612. We thank L. Yang for her contribution to Fig. 5. We thank the Jane and Aatos Erkkö Foundation and the 'Gigacity' project financed by Wihuri foundation. A.M. and P.M.W. acknowledge financial support by the Vienna Doctoral School in Physics (VDSF).

**Author contributions** X.-C.H., J.S., D.M.R., J.K., J.C. and M.H. planned the experiments. J.S., D.M.R., J.De., F.K., R.B., B.M., W.S., W.Y., L.C.-P., E.S., E.A., D.A., J.A., A.A., L.J.B., H.B., M.B., N.B., M.R.C., A.C., L.D., J.D., M.G., L.G.S., M.H., H.K., T.K., A.K., L.L., M.M., A.M., A.O., M.P., P.R., M.R., B.R., M.K.S., M.Si., M.Su., K.T., R.C.T., A.T., Y.T., J.T., N.S.U., J.W., B.Y., M.Z.-W., J.Z., Z.Z., T.C., I.E.H., R.C.F., H.J., U.R., S.S., R.V., P.M.W., A.H., K.L., N.M.D., J.L., H.H., M.K., D.R.W., J.K., J.C. and X.-C.H. prepared the CLOUD facility or measuring instruments. J.S., D.M.R., J.De., F.K., R.B., B.M., W.S., W.Y., L.C.-P., E.S., E.A., J.A., A.A., L.J.B., H.B., N.B., A.C., R.C.-S., J.D., M.G., L.G.S., M.H., H.K., T.K., M.L., L.L., B.L., M.M., A.M., M.P., P.R., M.R., S.R., B.R., M.Si., M.Su., K.T., R.C.T., A.T., Y.T., J.T., G.U., L.V., J.W., C.X., B.Y., M.Z.-W., J.Z., Z.Z., I.E.H., O.M., R.V., P.M.W., A.H., K.L., H.H., J.K., J.C. and X.-C.H. collected the data. S.I. carried out quantum-chemical calculations. J.S., X.-C.H., D.M.R., J.De., F.K., R.B., B.M., W.S., W.Y., L.C.-P., E.S., H.K., T.K., L.L., P.R., M.Si., M.Su., Y.T., J.W., B.Y., M.Z.-W., J.Z., Z.Z., H.H. and J.K. analysed the data. J.K. and X.-C.H. wrote the manuscript, with contributions from J.S., D.M.R., J.De. and N.M.D. Finally, J.S., D.M.R., J.De., F.K., R.B., B.M., W.S., E.S., R.C.-S., L.D., H.G., B.R., M.Si., Y.T., L.V., C.X., B.Y., U.B., I.E.H., R.C.F., I.R., S.S., R.V., A.H., N.M.D., J.L., M.K., D.R.W., J.K., J.C. and X.-C.H. commented on and edited the manuscript.

**Competing interests** The authors declare no competing interests.


#### Additional information

**Correspondence and requests for materials** should be addressed to Jasper Kirkby, Joachim Curtius or Xu-Cheng He.

**Peer review information** *Nature* thanks the anonymous reviewers for their contribution to the peer review of this work.


**Reprints and permissions information** is available at <http://www.nature.com/reprints>.






**Extended Data Fig. 1 | Wall-loss rates of C<sub>5</sub>H<sub>10</sub>O<sub>3</sub>, C<sub>5</sub>H<sub>12</sub>O<sub>6</sub> and HIO<sub>3</sub> at -30 °C and the IEPOX fraction of C<sub>5</sub>H<sub>10</sub>O<sub>3</sub> at -30 °C and -50 °C. a.** Wall-loss lifetimes (1/(loss rate)) of C<sub>5</sub>H<sub>10</sub>O<sub>3</sub>, C<sub>5</sub>H<sub>12</sub>O<sub>6</sub> and HIO<sub>3</sub> at 100% mixing fan speed (increased loss rate). At stage 1, the mixing fan speed is set to 12% (standard operation). At stage 2, the mixing fan speed is set to 100% to increase the wall-loss rate. All three vapours have wall-loss rates consistent with that measured for H<sub>2</sub>SO<sub>4</sub> (not shown), indicating irreversible loss on wall impact. **b.** Simulation of the fraction of IEPOX in C<sub>5</sub>H<sub>10</sub>O<sub>3</sub> using the reduced Wennberg et al. mechanism<sup>35</sup>. The dashed grey line is a fit to the data of the form IEPOX fraction = 10<sup>(0.58 × log<sub>10</sub>([OH]) - 4.6)</sup>. Our simulation predicts that roughly 30% of the measured C<sub>5</sub>H<sub>10</sub>O<sub>3</sub> signal is IEPOX

under the conditions of panel a and that both the ISOPOOH and IEPOX (summed as C<sub>5</sub>H<sub>10</sub>O<sub>3</sub>) are lost to the chamber wall on impact, at -30 °C and at -50 °C. The experimental conditions for panel a are: isoprene = 0.13 ppbv (3.7 × 10<sup>9</sup> cm<sup>-3</sup>), O<sub>3</sub> = 30 ppbv (8.6 × 10<sup>11</sup> cm<sup>-3</sup>), I<sub>2</sub> = 3.1 × 10<sup>7</sup> cm<sup>-3</sup>, SO<sub>2</sub> = 3 × 10<sup>8</sup> cm<sup>-3</sup>, OH = 8.9 × 10<sup>6</sup> cm<sup>-3</sup>, HO<sub>2</sub> = 1.8 × 10<sup>8</sup> cm<sup>-3</sup>, RH = 57% and temperature = -30 °C. The experimental conditions for panel b are: isoprene = 0.04–1.50 ppbv (0.1–4.2 × 10<sup>10</sup> cm<sup>-3</sup>), O<sub>3</sub> = 1–590 ppbv (3.7 × 10<sup>10</sup> to 1.8 × 10<sup>13</sup> cm<sup>-3</sup>), I<sub>2</sub> = 0–7.5 × 10<sup>7</sup> cm<sup>-3</sup>, SO<sub>2</sub> = 0–4.6 × 10<sup>9</sup> cm<sup>-3</sup>, OH = 0.11–6.90 × 10<sup>7</sup> cm<sup>-3</sup>, HO<sub>2</sub> = 0.6–17.0 × 10<sup>8</sup> cm<sup>-3</sup>, HO<sub>2</sub>/OH ratio = 11–118, NO = 0–0.22 ppbv, NO<sub>2</sub> = 0–0.77 ppbv, RH = 29–70% and temperature = -30 °C and -50 °C.




**Extended Data Fig. 2 |  $IP_{ON}$  yield from OH reaction with  $C_5H_{10}O_3$ .** Measured  $IP_{ON}$  concentration versus reacted  $C_5H_{10}O_3$  (see Methods for details). The dashed lines show the predictions for several  $IP_{ON}$  molar yields; a yield of 100% implies that each OH reaction with  $C_5H_{10}O_3$  produces one  $IP_{ON}$  molecule. The data indicate that the yield is between 20% and 50%, with an overall systematic uncertainty of a factor of two. When  $NO_x$  is added to the system, the yield of  $IP_{ON}$  is reduced, owing to NO terminating isoprene peroxy radicals. The ratios of IEPOX and ISOPOOH in  $C_5H_{10}O_3$  are determined from the measured OH

concentrations and the fit to the simulation shown in Extended Data Fig. 1b. For simplification, we apply a general reaction-rate coefficient of  $10^{-10}$  and  $10^{-11} \text{ cm}^3 \text{ s}^{-1}$  for the ISOPOOH + OH and IEPOX + OH reactions, respectively<sup>35</sup>. The experimental conditions are: isoprene = 0.04–1.50 ppbv ( $0.1\text{--}4.2 \times 10^{10} \text{ cm}^{-3}$ ),  $O_3 = 1\text{--}590 \text{ ppbv}$  ( $3.7 \times 10^{10}$  to  $1.8 \times 10^{13} \text{ cm}^{-3}$ ),  $I_2 = 0\text{--}7.5 \times 10^7 \text{ cm}^{-3}$ ,  $SO_2 = 0\text{--}4.6 \times 10^9 \text{ cm}^{-3}$ ,  $OH = 0.11\text{--}6.90 \times 10^7 \text{ cm}^{-3}$ ,  $HO_2 = 0.6\text{--}18.0 \times 10^8 \text{ cm}^{-3}$ ,  $HO_2/OH$  ratio = 11–117,  $NO = 0\text{--}0.22 \text{ ppbv}$ ,  $NO_2 = 0\text{--}0.77 \text{ ppbv}$ ,  $RH = 29\text{--}70\%$  and temperature =  $-30 \text{ }^\circ\text{C}$  and  $-50 \text{ }^\circ\text{C}$ .




**Extended Data Fig. 3 | Example new particle formation experiment at -50 °C with added HIO<sub>x</sub>.** Evolution of vapour concentrations (a), particle-nucleation rates at 1.7 nm,  $J_{1.7}$  (b), and naturally charged particle number size distribution ( $dN/d\log D_p$ ) and growth rate measured between 3.2 and 8.0 nm, GR<sub>3.2-8</sub> (nm h<sup>-1</sup>) (c). The black line in panel c depicts the linear fit of 50% appearance time of particles between 3.2 and 8.0 nm. The expected nucleation rates for HIO<sub>x</sub> at the

kinetic limit are less than  $2 \times 10^{-4} \text{ cm}^{-3} \text{ s}^{-1}$ . The results show that HIO<sub>x</sub> enhances IP<sub>0N</sub> nucleation in a similar way as H<sub>2</sub>SO<sub>4</sub>. Discontinuous lines show missing data. The experimental conditions are: isoprene = 0.07–0.16 ppbv ( $2.1\text{--}4.9 \times 10^9 \text{ cm}^{-3}$ ), O<sub>3</sub> = 95 ppbv ( $3 \times 10^{12} \text{ cm}^{-3}$ ), I<sub>2</sub> = 0.4–6.7 × 10<sup>7</sup> cm<sup>-3</sup>, OH = 3.2 × 10<sup>6</sup> cm<sup>-3</sup>, HO<sub>2</sub> = 7.4 × 10<sup>7</sup> cm<sup>-3</sup>, RH = 63% and temperature = -49 °C.




**Extended Data Fig. 4 | Aerosol formation from pure IP<sub>ON</sub> products and mixture of IP<sub>ON</sub>, IP<sub>1-2N</sub> and H<sub>2</sub>SO<sub>4</sub>.** Evolution of vapour concentrations (a,b), particle-nucleation rates at 1.7 nm,  $J_{1.7}$  (c,d), and naturally charged particle number size distribution ( $dN/d\log D_p$ ) and growth rates measured between 3.2 and 8.0 nm,  $GR_{3.2-8}$  (nm h<sup>-1</sup>) (e,f). The black lines in panel f depict the linear fits of 50% appearance time of particles between 3.2 and 8.0 nm. The results show that pure IP<sub>ON</sub> aerosol formation is feasible and atmospheric ions can enhance the nucleation rate by a few times. The addition of low concentrations of H<sub>2</sub>SO<sub>4</sub> further enhances IP<sub>0-2N</sub> nucleation by up to two orders of magnitude. The vertical dashed lines and labels indicate the start of a new stage in which the experimental conditions were adjusted. The dotted black curve in panel d shows the expected H<sub>2</sub>SO<sub>4</sub>-NH<sub>3</sub> nucleation rate, conservatively assuming

that NH<sub>3</sub> is present at the 4 pptv limit of detection<sup>44,93</sup>. Stages 1 and 3 are, respectively, under neutral (ion-free) and beam (ion-enhanced) conditions, whereas all of the other stages are under galactic cosmic ray (GCR; natural ion concentrations) conditions. The SO<sub>2</sub> concentration is increased in steps during stages 5–12. The experimental conditions for the right panels are: isoprene = 0.08–0.63 ppbv ( $2.4\text{--}20.0 \times 10^9 \text{ cm}^{-3}$ ), O<sub>3</sub> = 189 ppbv ( $6 \times 10^{12} \text{ cm}^{-3}$ ), OH =  $2.4 \times 10^6 \text{ cm}^{-3}$ , HO<sub>2</sub> =  $1.8 \times 10^8 \text{ cm}^{-3}$ , RH = 32% and temperature = -49 °C. The experimental conditions for the left panels are: isoprene = 0.05–0.50 ppbv ( $1.6\text{--}16.0 \times 10^9 \text{ cm}^{-3}$ ), O<sub>3</sub> = 1.2 ppbv ( $3.8 \times 10^{10} \text{ cm}^{-3}$ ), SO<sub>2</sub> = 0– $1.9 \times 10^8 \text{ cm}^{-3}$ , OH =  $6 \times 10^7 \text{ cm}^{-3}$ , HO<sub>2</sub> =  $1 \times 10^9 \text{ cm}^{-3}$ , NO = 0.12–0.81 ppbv, NO<sub>2</sub> = 0.57–0.84 ppbv, RH = 52% and temperature = -48 °C.




**Extended Data Fig. 5 | Particle-nucleation rates from IP-OOM at  $-50\text{ }^{\circ}\text{C}$ , in the absence of acids.** Nucleation rates at  $1.7\text{ nm}$ ,  $J_{1.7}$ , versus IP-OOM without nitrogen,  $\text{IP}_{0\text{N}}$  (a), with nitrogen,  $\text{IP}_{1\text{N}} + \text{IP}_{2\text{N}}$  (b) and the sum,  $\text{IP}_{0-2\text{N}}$  (c). Panel a indicates a mild enhancement of the nucleation rates with increased  $\text{IP}_{1\text{N}} + \text{IP}_{2\text{N}}$ . Panels b and c confirm that the nucleation rates are only weakly dependent on  $\text{IP}_{1\text{N}} + \text{IP}_{2\text{N}}$ . These data show that non-nitrate IP-OOM are primarily responsible for particle nucleation, with a smaller contribution from nitrate-containing

isoprene products. The experimental conditions are: isoprene =  $0.04\text{--}0.50\text{ ppbv}$  ( $0.14\text{--}1.7 \times 10^{10}\text{ cm}^{-3}$ ),  $\text{O}_3 = 1.2\text{--}590\text{ ppbv}$  ( $4 \times 10^{10}$  to  $1.8 \times 10^{13}\text{ cm}^{-3}$ ),  $\text{OH} = 0.14\text{--}6.40 \times 10^7\text{ cm}^{-3}$ ,  $\text{HO}_2 = 1.6\text{--}17.0 \times 10^8\text{ cm}^{-3}$ ,  $\text{HO}_2/\text{OH}$  ratio =  $11\text{--}118$ ,  $\text{NO} = 0\text{--}0.18\text{ ppbv}$ ,  $\text{NO}_2 = 0\text{--}0.74\text{ ppbv}$ ,  $\text{RH} = 29\text{--}61\%$  and temperature =  $-50\text{ }^{\circ}\text{C}$ . All experiments are carried out under galactic cosmic ray ionization (ambient-boundary-layer conditions). The error bars represent the standard deviation of the measurement at steady state.




**Extended Data Fig. 6 | Particle-nucleation rates at 2.5 nm versus vapour concentrations.** Nucleation rates at 2.5 nm,  $J_{2.5}$ , versus  $IP_{ON}$  (a,b),  $HIO_x + H_2SO_4$  (c) and the product  $(HIO_x + H_2SO_4) \times IP_{ON}$  (d). Panel a and b show experiments without and with added  $NO_x$ . Panel c shows all experiments and panel d shows experiment with acids, both with and without  $NO_x$ .  $IP_{ON}$  excludes nitrogen-containing isoprene-oxygenated molecules. The solid lines in panel c show the nucleation rates at 1.7 nm expected for  $H_2SO_4$  with 4 pptv  $NH_3$  at 60% RH (ref. 44). The dashed and solid lines in panel d represent fits to the equation  $10^{a \times \log_{10}(x)+b}$ , in which, for the dashed line,  $a = 1.167$  and  $b = -14.743$ , and for the solid line,

$a = 1.669$  and  $b = -22.707$ . Panels a and b show that both total acid ( $HIO_x + H_2SO_4$ ) and  $IP_{ON}$  contribute to the nucleation rate. The experimental conditions are: isoprene = 0.04–1.50 ppbv ( $0.1$ – $4.2 \times 10^{10} \text{ cm}^{-3}$ ),  $O_3 = 1$ –590 ppbv ( $3.7 \times 10^{10}$  to  $1.8 \times 10^{13} \text{ cm}^{-3}$ ),  $I_2 = 0$ – $7.5 \times 10^7 \text{ cm}^{-3}$ ,  $SO_2 = 0$ – $4.6 \times 10^9 \text{ cm}^{-3}$ ,  $OH = 0.11$ – $6.90 \times 10^7 \text{ cm}^{-3}$ ,  $HO_2 = 0.6$ – $18.0 \times 10^8 \text{ cm}^{-3}$ ,  $HO_2/OH$  ratio = 11–118,  $NO = 0$ – $0.22$  ppbv,  $NO_2 = 0$ – $0.77$  ppbv,  $RH = 29$ – $70\%$  and temperature =  $-30$  °C and  $-50$  °C. All experiments are carried out under galactic cosmic ray ionization (ambient conditions). The error bars represent the standard deviation of the measurement at steady state.



**Extended Data Fig. 7 | Molecular composition of charged clusters with sulfuric acid, iodic acid and no acid during IP-OOM nucleation at  $-50\text{ }^{\circ}\text{C}$ .** Mass defect (difference from integer mass) versus  $m/z$  (mass-to-charge ratio) during nucleation events for ions and negatively charged (**a-d**) and positively charged (**e,f**) molecular clusters measured with two API-TOF instruments (see Methods), coloured by the number of carbon (**a,b,e**), sulfur (**c**), iodine (**d**) and nitrogen (**f**) atoms. The data are without  $\text{NO}_x$  (**a-e**) and with  $\text{NO}_x$  (**f**). In panel **d**, the (purple) clusters with no iodine contain a  $\text{NO}_3^-$  ion. The symbol area is proportional to the ion signal rate (cps, counts  $\text{s}^{-1}$ ). It should be noted that the API-TOF used in this experiment is less sensitive than the MION2-API-TOF and thus the absolute ion signals and cluster population should not be compared between the instruments. Together with Fig. 3, our results show that ion-induced nucleation occurs both with pure IP-OOM and with further  $\text{H}_2\text{SO}_4$

and  $\text{HIO}_3$ . The experimental conditions for **a,c,e** are: isoprene =  $0.19\text{ ppbv}$  ( $5.9 \times 10^9\text{ cm}^{-3}$ ),  $\text{O}_3 = 93\text{ ppbv}$  ( $2.9 \times 10^{12}\text{ cm}^{-3}$ ),  $\text{I}_2 = \text{unmeasured}$ ,  $\text{SO}_2 = \text{unmeasured}$ ,  $\text{OH} = 3.0 \times 10^6\text{ cm}^{-3}$ ,  $\text{HO}_2 = \text{unmeasured}$ ,  $\text{RH} = 63\%$  and temperature =  $-49\text{ }^{\circ}\text{C}$ . The concentrations of  $\text{IP}_{\text{ON}} = \text{unmeasured}$ ,  $\text{H}_2\text{SO}_4 = 3.0 \times 10^5\text{ cm}^{-3}$  and  $\text{HIO}_x = 2.8 \times 10^4\text{ cm}^{-3}$ . The experimental conditions for **b,d** are: isoprene =  $0.09\text{ ppbv}$  ( $2.8 \times 10^9\text{ cm}^{-3}$ ),  $\text{O}_3 = 97\text{ ppbv}$  ( $3 \times 10^{12}\text{ cm}^{-3}$ ),  $\text{I}_2 = 5.1 \times 10^7\text{ cm}^{-3}$ ,  $\text{SO}_2 = 0$ ,  $\text{OH} = 3.4 \times 10^6\text{ cm}^{-3}$ ,  $\text{HO}_2 = 7.6 \times 10^7\text{ cm}^{-3}$ ,  $\text{RH} = 63\%$  and temperature =  $-49\text{ }^{\circ}\text{C}$ . The concentrations of low-volatility vapours are  $\text{IP}_{\text{ON}} = 1.3 \times 10^7\text{ cm}^{-3}$ ,  $\text{H}_2\text{SO}_4 = 8.3 \times 10^3\text{ cm}^{-3}$  and  $\text{HIO}_x = 6.7 \times 10^5\text{ cm}^{-3}$ . The experimental conditions for **f** are: isoprene =  $0.12\text{ ppbv}$  ( $3.7 \times 10^9\text{ cm}^{-3}$ ),  $\text{O}_3 = 1.2\text{ ppbv}$  ( $3.7 \times 10^{10}\text{ cm}^{-3}$ ),  $\text{SO}_2 = 1.2 \times 10^8\text{ cm}^{-3}$ ,  $\text{OH} = 6.2 \times 10^7\text{ cm}^{-3}$ ,  $\text{HO}_2 = 1 \times 10^9\text{ cm}^{-3}$ ,  $\text{RH} = 42\%$ ,  $\text{NO} = 0.25\text{ ppbv}$ ,  $\text{NO}_2 = 0.78\text{ ppbv}$  and temperature =  $-48\text{ }^{\circ}\text{C}$ . The concentrations of low-volatility vapours are  $\text{IP}_{\text{ON}} = 1.2 \times 10^7\text{ cm}^{-3}$ ,  $\text{H}_2\text{SO}_4 = 2.4 \times 10^5\text{ cm}^{-3}$  and  $\text{HIO}_x = 0\text{ cm}^{-3}$ .



**Extended Data Fig. 8 | Four experiments that demonstrate the influence of acids and  $\text{NO}_x$  on IP-OOM nucleation at  $-50^\circ\text{C}$ .** Four experiments showing nucleation rates at  $1.7\text{ nm}$ ,  $J_{1.7}$ , measured at similar  $\text{IP}_{0\text{N}}$  concentrations,  $1.2$ – $3.3 \times 10^7\text{ cm}^{-3}$ , but with added acids,  $\text{NO}_x$ , and acids +  $\text{NO}_x$ , respectively. The marker area is proportional to the  $\text{IP}_{0\text{N}}$  concentration. The experimental conditions are shown next to the data points. Circles correspond to absence of

acids and squares include acids. The data show that extremely low  $\text{H}_2\text{SO}_4$  concentrations,  $1.3$ – $2.8 \times 10^6\text{ cm}^{-3}$ , strongly enhance the nucleation rates regardless of the absence or presence of  $\text{NO}_x$ . On the other hand, additions of high nitrate IP-OOM concentrations around  $10^8\text{ cm}^{-3}$  only produce relatively small increases in nucleation rates.



**Extended Data Table 1 | Comparison of CLOUD experimental conditions with the ambient conditions during CAFE-Brazil research flight RF19, periods T4 and T9 (Curtius et al.<sup>13</sup>)**

Parameter (units)	CLOUD		CAFE-Brazil (CB)	
	Range: min-max (Median)		Flight RF19-T4 Median	Flight RF19-T9 Median
Local time	2022-Sep to 2023-Nov	2022-Sep to 2023-Nov	2023-Jan-23 08:10-08:11	2023-Jan-23 10:06-10:10
Temperature (°C)	-50 to -46 (-48)	-30 to -29 (-30)	-58	-58
Pressure (hPa)	965	965	187	187
Relative humidity (RH) (%)	29-63 (57)	47-70 (67)	45	43
Isoprene (cm <sup>-3</sup> )	1.4×10 <sup>9</sup> -1.7×10 <sup>10</sup> (6.2×10 <sup>9</sup> )	2.4×10 <sup>9</sup> -4.2×10 <sup>10</sup> (2.1×10 <sup>9</sup> )	1.7×10 <sup>9</sup>	3.4×10 <sup>8</sup>
IP <sub>0N</sub> <sup>*,†</sup> (cm <sup>-3</sup> )	1.4×10 <sup>7</sup> -6.2×10 <sup>7</sup> (2.9×10 <sup>7</sup> )	5.7×10 <sup>7</sup> -3.3×10 <sup>8</sup> (2.1×10 <sup>8</sup> )	7.6×10 <sup>7</sup>	2.3×10 <sup>8</sup>
IP <sub>1-2N</sub> <sup>*,†</sup> (cm <sup>-3</sup> )	1.4×10 <sup>7</sup> -2.5×10 <sup>8</sup> (7.0×10 <sup>7</sup> )	5.3×10 <sup>6</sup> -3.7×10 <sup>7</sup> (6.1×10 <sup>6</sup> )	8.9×10 <sup>7</sup>	3.4×10 <sup>8</sup>
IP <sub>0N</sub> /IP <sub>1-2N</sub> ratio <sup>*</sup>	0.12-0.77 (0.3)	0.6-0.98 (0.96)	0.46	0.40
H <sub>2</sub> SO <sub>4</sub> <sup>‡,§,  </sup> (cm <sup>-3</sup> )	<2×10 <sup>4</sup> -2.8×10 <sup>6</sup> (2.4×10 <sup>5</sup> )	1.2×10 <sup>4</sup> -7.6×10 <sup>6</sup> (2.9×10 <sup>4</sup> )	<2×10 <sup>6</sup> (LOD)	<2×10 <sup>6</sup> (LOD)
HIO <sub>3</sub> <sup>¶</sup> (cm <sup>-3</sup> )	1.8×10 <sup>4</sup> -1.1×10 <sup>6</sup> (3.5×10 <sup>4</sup> )	5.2×10 <sup>4</sup> -1.4×10 <sup>6</sup> (3.5×10 <sup>5</sup> )	n/a	n/a
O <sub>3</sub> (cm <sup>-3</sup> )	3.7×10 <sup>10</sup> -1.8×10 <sup>13</sup> (2.8×10 <sup>12</sup> )	4.3×10 <sup>11</sup> -6.7×10 <sup>12</sup> (1.0×10 <sup>12</sup> )	1.8×10 <sup>11</sup>	2×10 <sup>11</sup>
OH (cm <sup>-3</sup> )	1.4×10 <sup>6</sup> -6.7×10 <sup>7</sup> (8.6×10 <sup>6</sup> )	1.2×10 <sup>6</sup> -6.9×10 <sup>7</sup> (1.2×10 <sup>7</sup> )	4.7×10 <sup>5</sup>	1.6×10 <sup>6</sup>
HO <sub>2</sub> (cm <sup>-3</sup> )	5.7×10 <sup>7</sup> -1.7×10 <sup>9</sup> (2.3×10 <sup>8</sup> )	8.9×10 <sup>7</sup> -1.7×10 <sup>9</sup> (3.3×10 <sup>8</sup> )	2.1×10 <sup>7</sup>	2.3×10 <sup>7</sup>
NO <sup>*</sup> (cm <sup>-3</sup> )	LOD <sup>#</sup> -6.8×10 <sup>9</sup> (1.8×10 <sup>9</sup> )	LOD <sup>#</sup> -3.6×10 <sup>8</sup> (LOD <sup>#</sup> )	1.1×10 <sup>9</sup>	8.0×10 <sup>8</sup>
NO <sub>2</sub> <sup>*</sup> (cm <sup>-3</sup> )	2.7×10 <sup>9</sup> -2.4×10 <sup>10</sup> (1.1×10 <sup>10</sup> )	1.1×10 <sup>9</sup> -3.9×10 <sup>9</sup> (1.5×10 <sup>9</sup> )	7.2×10 <sup>8</sup>	9.0×10 <sup>8</sup>
HO <sub>2</sub> /OH ratio	11-118 (24)	14-80 (29)	45.2	14.4
HO <sub>2</sub> /NO ratio <sup>*</sup>	0.06-19 (0.63)	0.92-83 (33)	0.02	0.03
(k(RO <sub>2</sub> ×HO <sub>2</sub> ) <sup>★</sup> ×[HO <sub>2</sub> ])/ (k(RO <sub>2</sub> ×NO)× α(RO <sub>2</sub> ×NO) <sup>**</sup> ×[NO]) ratio <sup>*</sup>	0.7-236 (7.6)	19-1759 (698)	0.66	0.95

The times of the CB flights are in local time. <sup>\*</sup>Only CLOUD experiments with added NO<sub>x</sub>. <sup>†</sup>The definitions of IP<sub>0N</sub> and IP<sub>1-2N</sub> in the CB study refer to the measurement of the NO<sub>3</sub><sup>-</sup> chemical ionization alone, whereas the CLOUD results refer to measurements using the combination of NO<sub>3</sub><sup>-</sup>, Br<sup>-</sup> and NH<sub>4</sub><sup>+</sup> chemical-ionization methods. <sup>‡</sup>Only CLOUD experiments with H<sub>2</sub>SO<sub>4</sub>. <sup>§</sup>The CLOUD limit-of-detection (LOD) for H<sub>2</sub>SO<sub>4</sub> is 2×10<sup>4</sup> cm<sup>-3</sup>. <sup>||</sup>The CB LOD for H<sub>2</sub>SO<sub>4</sub> is 2×10<sup>6</sup> cm<sup>-3</sup> at altitudes above 8 km. <sup>¶</sup>Only experiments with HIO<sub>3</sub>. <sup>★</sup>The CLOUD LOD for NO is 2×10<sup>9</sup> cm<sup>-3</sup>. <sup>\*\*</sup>Using equation 7 in ref. 35. <sup>††</sup>Using equations 1-4 in ref. 35.