



## CORRECTIONS

The reversible immobilization of proteins on nylon activated through the formation of a substituted imidoester, and its unusual properties

By P. V. SUNDARAM

Volume 183 (1979)

p. 446, Scheme 1:



Scheme 1. Alkylation of nylon with dimethyl sulphate (DMS) or triethyloxonium tetrafluoroborate (TTFB) and subsequent amidination by an amine  
X=CH<sub>3</sub>(DMS) or C<sub>2</sub>H<sub>5</sub> (TTFB).

Scheme 1. Alkylation of nylon with dimethyl sulphate (DMS) or triethyloxonium tetrafluoroborate (TTFB) and subsequent amidination by an amine  
X=CH<sub>3</sub>(DMS) or C<sub>2</sub>H<sub>5</sub> (TTFB).

Effects of manganese ions and magnesium ions on the activity of soya-bean ribulose biphosphate carboxylase/oxygenase

J. T. CHRISTELLER and W. A. LAING

Volume 183 (1979)

p. 749, Table 1:

Delete  $V_{\text{max.}(O_2)}$  from the heading of the fifth column

p. 749, Title of Figure 3:

*for* oxygenase *read* carboxylase

p. 749, Legend of Figure 3:

The symbols ○ and ● should be transposed

Identification of *O*-acetyl-5-methoxytryptophol in the pineal gland by gas chromatography–mass spectrometry

By I. SMITH, P. FRANCIS, R. M. LEONE and P. E. MULLEN

Volume 185 (1980)

p. 537, Title, line 1:

*for* methoxytryptophenol *read* methoxytryptophol

p. 537, first column, lines 15 and 16:

*for* *N*-[2-(5-methoxytryptophol-3-yl)ethyl]acetamide  
*read* *N*-[2-(5-methoxyindol-3-yl)ethyl]acetamide

p. 539, second column, lines 12 and 13:

*for* *O*-acetyl-5-hydroxymethoxytryptophol *read* melatonin