



Comparison of External Circular Fixation and Plate and Screw Fixation Methods in Tibial Pilon Fractures

Comparação entre métodos de fixação externa circular e de fixação com placa e parafuso nas fraturas de pilão tibial

Marco Antonio Schueda^{1,2} Leonardo Maranhão Gubert^{2,3} Yan Celuppi Dal Vesco²
Fernanda Fossa Dal Piva² Felipe Guglielmi Niada^{1,2} Ryad Fayez Mehanna²

¹ Orthopedics and Traumatology Service, Hospital do Coração Balneário Camboriú, Balneário Camboriú, SC, Brazil

² Orthopedics and Traumatology Service, Hospital e Maternidade Marieta Konder Bornhausen, Itajaí, SC, Brazil

³ Orthopedics Service, Hospital Unimed Litoral, Balneário Camboriú, SC, Brazil,

Address for correspondence Ryad Fayez Mehanna, Serviço de Ortopedia e Traumatologia, Hospital e Maternidade Marieta Konder Bornhausen, Itajaí, SC, Brazil (e-mail: ryad.mehanna@hotmail.com).

Rev Bras Ortop 2024;59(5):e663–e671.

Abstract

Objective To compare the fixation methods for tibial pilon fractures among patients treated in a hospital.

Methods We analyzed the medical records of 28 patients who underwent a surgical procedure for tibial pilon fracture, among whom 15 subjects received a circular external fixator, and 13 underwent internal fixation using a plate and screws. We assessed age, sex, aggravating factors, trauma energy (high or low), presence of soft tissue injuries, associated fractures, and clinical outcomes.

Results Most patients were male, aging between 40 and 60 years. The most common trauma mechanism was car accident, and the associated injury was a fracture of the distal third of the fibula. The fracture patterns in patients treated with a circular external fixator were AO 43C3 and 43C2. As for the prevalent fracture patterns in the internal fixation group, we identified AO 43C1, 43C2, and 43C3.

Conclusion An individualized therapeutic choice is critical for a better functional outcome. Additionally, it is essential to highlight that the profile of fractures and patients in the circular external fixator and internal fixation groups tends to be quite heterogeneous, because the treatment of fractures with the worst classification and most frequently associated with soft tissue injuries often uses circular external fixation;

Keywords

- ▶ external fixators
- ▶ Ilizarov technique
- ▶ tibial fractures

Work carried out at the Orthopedics and Traumatology Service, Hospital e Maternidade Marieta Konder Bornhausen, Itajaí, SC, Brazil.

received
November 6, 2023
accepted
June 23, 2024

DOI <https://doi.org/10.1055/s-0044-1790213>.
ISSN 0102-3616.

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by/4.0/>).

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

meanwhile, those with less severe fractures and a lower incidence of soft tissue injuries are usually managed with open reduction and internal fixation. We noted that the clinical and radiographic outcomes tended to be similar between both groups despite the particularities of each method.

Resumo

Objetivo Comparar os métodos de fixação de fraturas do pilão tibial de pacientes atendidos em um hospital.

Métodos Foram analisados os prontuários de 28 pacientes que realizaram procedimento cirúrgico para fratura do pilão tibial, sendo 15 deles tratados com fixador externo circular e 13 com fixação interna por placa e parafusos. Foram analisados idade, sexo, fatores agravantes, energia do trauma (alta ou baixa), presença de lesão de tecidos moles, ocorrência de fraturas associadas e desfechos clínicos.

Resultados O sexo mais acometido foi o masculino, na faixa etária entre 40 e 60 anos. O mecanismo de trauma em sua maioria foi acidentes automobilísticos e a lesão associada em 100% dos casos foi a fratura do terço distal da fíbula. O padrão de fratura observado nos pacientes tratados com fixador externo circular foram AO 43C3 e 43C2. Já os padrões de fratura prevalentes observado no grupo de fixação interna foram AO 43C1, 43C2 e 43C3.

Conclusão A escolha terapêutica individualizada é de extrema importância para um melhor desfecho funcional. Também é imprescindível ressaltar que o perfil da fratura e dos pacientes que do grupo fixador externo circular e de grupo fixação interna por placa e parafusos tende a ser bastante heterogêneo, pois as fraturas de pior classificação e mais frequentemente associadas a lesões de partes moles costumam ser manejadas com fixação externa circular, enquanto aqueles com fraturas de menor gravidade e menor incidência de lesão de partes moles tendem a ser manejados por redução aberta e fixação interna. Observa-se que os desfechos clínicos e radiográficos tendem a ser similares entre ambos os grupos apesar das particularidades de cada método.

Palavras-chave

- ▶ fixadores externos
- ▶ fraturas da tibia
- ▶ técnica de Ilizarov

Introduction

The term tibial pilon was introduced in 1911 by the French radiologist Destot, who described fractures involving the weight-bearing joint surface of the distal third of the tibia but not necessarily affecting it.¹ Radiographic evaluation requires three views: anteroposterior (AP), lateral (L), and AP with 20° of internal rotation (mortise). In cases of doubt, one may request an oblique or stress radiograph. Additionally, one may ask for a radiograph of the contralateral ankle for comparison. In cases of joint involvement, especially in fractures caused by high-energy trauma, computed tomography (CT) helps to evaluate the fracture pattern and joint comminution degree, as well as preoperative planning.²

Tibial pilon fractures predominantly affect young men between 30 and 40-years-old, and high-energy trauma accounts for 10 to 30% of open fractures.³ Currently these fractures are common because of high-energy traumas, such as traffic accidents or falls from height. Other fractures, including in the talus, calcaneus, tibial plateau, pelvis, acetabulum, and spine, may accompany tibial pilon fractures.⁴

The trauma mechanism is longitudinal compression of the talus on the distal surface of the tibia, potentially associated with rotational forces. The fracture pattern results from the direction and speed of the damaging energy, along with the foot's positioning during load application.⁵

Furthermore, it is important to assess soft tissue involvement and consider whether the fracture is open or closed. We know many factors deserve consideration when choosing the definitive treatment⁶ since the soft tissue involvement may suffer exacerbation by major surgical procedures involving large incisions, bone deperiostization, and separation of its soft components, which can lead to necrosis of the overlying tissue and increase the risk of pseudarthrosis, infection, or both.⁷

Materials and Methods

Our institution approved this study under CAEE number 71275623.0.0000.0120.

This cross-sectional and observational study analyzed the medical records of patients hospitalized with a tibial pilon fracture who underwent surgical treatment from

January 1st, 2021, to January 1st, 2023. Using the medical records, we assessed age, sex, aggravating factors, trauma energy, presence of soft tissue injury, and occurrence of associated fractures. We analyzed imaging tests (AP, L, and oblique radiographs of the ankle and preoperative CTs), transoperative, immediate postoperative, and postoperative follow-up radiographies in ambulatory. For the functional analysis, the American Association for Foot and Ankle Surgery's (AFOS) functional scores were used.

Polytrauma patients underwent damage control treatment with mono or biplanar fixation and a review for internal or external circular fixation within 7 days of their emergency treatment.

Results

We analyzed the medical records from 30 patients operated on for tibial pilon fracture, including 15 treated with a circular external fixator and 15 treated with internal fixation using a plate and screws. We lost 2 patients from the internal fixation group at follow-up and excluded them from the analyses. Thus, this subgroup had 13 patients, and the total research sample comprised 28 subjects.

The patients included 21 males and 7 females (►Fig. 1). The external fixation subgroup treated using the method proposed by Ilizarov had 15 patients, including 10 males and 5 females. The internal fixation subgroup treated using a nonlocking trefoil plate and screws had 13 patients, with 11 males and 2 females.

We observed a high prevalence of patients in the age group from 40 to 60-years-old, with an average of 44 (►Fig. 2). The average age was 43 in males and 46 in females.

Regarding the mechanism of trauma in the circular external fixator group, we observed a higher tendency for high-energy trauma (11 patients). Car accidents accounted for 7 of the 11 cases, while the remaining 4 resulted from falls from height. From that last group, all 4 patients suffered lower-energy trauma resulting from falls from their height. Among patients from the circular external fixator group, there were 60 open and 40% closed fractures.




Fig. 1 Sex of the participants.

Regarding the trauma mechanism in the internal fixation group, a high tendency towards high-energy trauma was noted (10 patients). Car accidents accounted for 5 of the 10 cases, while another 4 cases resulted from falls from height, and 1 involved a gunshot wound. Among the 3 patients suffering low-energy trauma, 2 had fractures from physical aggression, and 1 case was due to a fall from their level (►Fig. 3). Therefore, the internal fixation group had approximately 77% closed and 23% open fractures.

According to the AO group classification for fracture patterns, we noted that 60% of patients treated with a circular external fixator had 43C3 fractures with frank joint comminution, and another 40% had 43C2. No subject treated with the circular external fixation method had 43C1 fractures.

Using the same classification, we observed that patients treated with internal fixation included 5 subjects with 43C1 fractures, 5 with 43C2 fractures, and 3 with 43C3 fractures.

In both groups, the associated injury in 100% of cases was a fracture of the distal third of the fibula. Other injuries in the circular external fixation group included tibial plateau (4 cases), calcaneus (2 cases), tibial diaphysis, contralateral ankle, and hand bone fractures (1 case each). A clavicle fracture was the single other injury observed in the internal fixation group.

Regarding the outcomes of the circular external fixator group, 5 patients presented complete fracture consolidation, 1 had pseudarthrosis, and 2 malunions. ►Figs. 4–7 show examples of the circular external (I) and internal fixation (II) groups.

A total of 6 patients still remain in follow-up, but their progress is satisfactory, with radiographic consolidation signs. A single patient was lost at follow-up (►Fig. 8).

As for the outcomes of the open reduction and internal fixation group, 10 patients presented complete fracture consolidation, 1 had pseudarthrosis, 1 presented with malunion. Finally, one of the cases had a frank infection in the topography requiring synthetic material removal and multiple surgical interventions (►Fig. 9).

To facilitate the functional assessment of patients according to the American Orthopedic Foot & Ankle Society (AOFAS) score for the ankle and hindfoot (►Fig. 10), we subdivided the patients as follows:

- Group 1: 0 to 20 points
- Group 2: 20 to 40 points
- Group 3: 40 to 60 points
- Group 4: 60 to 80 points
- Group 5: 80 to 100 points

As such, we noted that the distribution of the groups was as follows: group 1–2 patients; group 2–6 patients; group 3–7 patients; group 4–8 patients (this group presented the highest prevalence); and group 5–5 patients.

The group treated with the Ilizarov external fixator contained 15 patients, distributed as follows: group 1–1 patient;




Fig. 2 Age of the patients.




Fig. 3 Mechanism of trauma.

group 2–3 patients; group 3–4 patients; group 4–4 patients; and group 5–3 patients.

In contrast, the group treated with internal fixation (plate and screws) comprised 13 patients with the following distribution: group 1–1 patient; group 2–3 patients; group 3–3 patients; group 4–4 patients; and group 5–2 patients.

Discussion

Our study revealed a prevalence of males, which is consistent with the literature. However, the prevalent age group was older than the one reported by Marsh and Saltzman.⁵ Pimenta et al.⁸ observed an average age of 42, similar to our study.



Fig. 4 Tibial pilon fracture (AO 43C2) before internal fixation with plate and screw.




Fig. 5 Tibial pilon fracture (AO 43C2) after internal fixation with plate and screw.




Fig. 6 Tibial pilon fracture (AO 43C2) before circular external fixation.




Fig. 7 Tibial pilon fracture (AO 43C2) after circular external fixation.




Fig. 8 Circular external fixation outcome.




Fig. 9 Internal fixation (plate and screw) outcome.

In the circular external fixator group, the rate of open fractures was above the percentiles found in the literature.⁹ In contrast, the internal fixation group had a rate of soft tissue injuries with fracture exposure well below the literature, in which open fracture rates are at around 50%.

In our service, the main mechanism of trauma was car accidents, followed by falls from a height, which is consistent with the literature. Furthermore, it was observed that the

highest rate of fracture exposure among the 28 patients occurred in the group classified as AO 43C2, representing 60% of subjects, compared to 38% in the AO 43C3 group. These rates differ from the literature, which reports a higher percentile of open fractures in the group with the most complex fractures, classified as AO 43C3.²

We detected no statistically significant difference when evaluating the presence of a fibula fracture and correlating it

AOFAS score for the ankle and hindfoot (total: 100 points)	
Pain (40 points)	
• None	0
• Mild, occasional	30
• Moderate, daily	20
• Severe, virtually always present	0
Function (50 points)	
Activity limitations, requiring support	
• No limitations, no support	10
• No daily activity limitation, recreational activity limitation, no support	7
• Limited daily and recreational activities, cane	4
• Severe limitation of daily and recreational activities, walker, crutches, wheelchair, orthoses (ankle bracer/immobilizer)	0
Maximum walking distance, blocks	
• Over 6	5
• 4 to 6	4
• 1 to 3	2
• Less than 1	0
Walking surfaces	
• No difficulty on any surface	5
• Some difficulty on uneven terrain, stairs, inclines, ladders	3
• Severe difficulty on uneven terrain, stairs, inclines, ladders	0
Gait abnormality	
• None, slight	8
• Obvious	4
• Marked	0
Sagittal motion (flexion + extension)	
• Normal or mild restriction (30° or more)	8
• Moderate restriction (15–29°)	4
• Severe restriction (lower than 15°)	0
Hindfoot motion (inversion + eversion)	
• Normal or mild restriction (75–100% from normal)	6
• Moderate restriction (25–74% from normal)	3
• Marked restriction (lower than 25% from normal)	0
Ankle-hindfoot stability (anteroposterior, varusvalgus)	
• Stable	8
• Unstable	0
Alignment (10 points)	
• Good, plantigrade foot, ankle-hindfoot well aligned	10
• Fair, plantigrade foot, some degree of ankle-hindfoot malalignment, no symptoms	5
• Poor, nonplantigrade foot, severe malalignment, with symptoms	0
TOTAL SCORE : _____	

Fig. 10 American Orthopedic Foot & Ankle Society's (AOFAS) score for the ankle and hindfoot (total: 100 points).

with high- and low-energy traumas. This lack of difference occurred because, in our study, all patients presented an associated distal fibula fracture, even those suffering low-energy traumas. Furthermore, the literature reports that ipsilateral talus and calcaneal fractures accompanying tibial pilon fractures are extremely rare. However, we had 2 patients with concomitant calcaneal fractures, which may negatively influence long-term outcomes.^{9,10}

Regarding radiographic outcomes, it appears that studies evaluated them differently. Here, we concluded that among patients receiving a circular external fixator and with a definitive radiographic outcome, 62.5% presented fracture consolidation, 25 had malunion, and 12.5 presented nonunion. Among those undergoing internal fixation, 84% presented union, 8% had malunion, and 8% presented pseudarthrosis.

The full assessment of the remaining subjects was not feasible, leading to their exclusion.

As for this series' general median, considering all 28 patients, the average AFAOS score was 55.7 points (circular external fixator group: 56.6; internal fixation: 54.6). This score is lower than the average of 65 points reported by Moura Junior et al.²

Conclusion

The present study retrospectively assessed 28 cases of tibial pilon fracture, including 15 patients undergoing surgical treatment with a circular external fixator and 13 with internal plate fixation (simple, cloverleaf type, with no locking mechanism). There was a wide range of variants directly influencing the characteristics of each fracture and its clinical-radiographic outcome.

Although there are many therapeutic alternatives for tibial pilon fractures, the internal fixation methods using unlocked plates and screws, as well as the circular external fixation method, remain the most available, mainly in hospitals within the public health system. Therefore, discussing these methods is critical and inevitable.

It is essential to highlight that the profile of fractures and patients in the circular external fixator and internal fixation groups tended to be quite heterogeneous. This occurs because the treatment of fractures with worst AO classifications and more frequent association with soft tissue injuries (resulting from open fractures or presenting significant edema or blistering) often employs circular external fixation. Meanwhile, management of less-severe fractures, with a lower incidence of soft tissue injuries, usually relies on open reduction and internal fixation. Clinical and radiographic outcomes tended to be similar between both groups, despite the particularities of each method.

We conclude that it is critical to perform an attentive, individualized, and detailed assessment. This will enable a more assertive therapeutic choice, leading to better functional outcomes for the particularities of each case and lower rates of socioeconomic impact for those with a tibial pilon fracture.

Financial Support

This authors declare that they did not receive financial support from agencies in the public, private, or non-profit sectors to conduct the present study.

Conflict of Interests

The authors have no conflict of interests to declare.

References

- 1 Netter FH. Atlas de Anatomia Humana. 7ª. ed. Rio de Janeiro: Elsevier; 2019

- 2 Moura Júnior AF, Machado Filho AA, Campos ET, Martins JS. Tibial pilon fractures: functional results and evaluation of quality of life. *Sci J Foot Ankle* 2018;12(01):42–48
- 3 Koval KJ, Zucherman JD. *Fraturas: Manual para consulta rápida*. 3ª ed. Rio de Janeiro: Di Livros Editora Ltda; 2008
- 4 Pozzi I, Reginaldo S, de Almeida MV, Cristante AF. *Manual de Trauma Ortopédico*. São Paulo: Sociedade Brasileira de Ortopedia e Traumatologia; 2011
- 5 Marsh JL, Saltzaman CL. Fraturas do Tornozelo. In: Bucholz RW, Heckman JD, editors. *Fraturas em adultos de Rockwood e Green*. 5ª ed. Barueri: Manole; 2006:2051–2078
- 6 Catagni MA, Carvalho Filho G. Tratamento da fratura tibial em pilão com fixador externo circular de Ilizarov. *Rev Bras Ortop* 1996;31(08):643–648
- 7 Matias VEG, Brito GV, Spósito AL, Zamboni C, Mercadante M, Christian RW. “Preliminary evaluation of a prospective randomized case series comparing plate and screws versus Ilizarov circular external fixator for distal tibial fractures” [Internet]. *SciELO Preprints*. 2023 [cited 2023 Nov. 9]. Available from: <https://preprints.scielo.org/index.php/scielo/preprint/view/6785>
- 8 Pimenta LSM, Costa ARF, Baptista MV, Daher SS. Fraturas do pilão tibial: avaliação do tratamento em 25 pacientes. *Rev Bras Ortop* 1997;32(07):497–502
- 9 Borges VQ, Moraes LVM, Ferraz GF, Stéfani KC. Fraturas de pilão tibial tipo AO 43C – o que influencia o resultado funcional? *Sci J Foot Ankle* 2018;12(03):233–239
- 10 Gobezie R, Ponce B, Vrahas M. Pilon fractures: Use of the posterolateral approach for ORIF. *Oper Tech Orthop* 2003;13(02):113–119