

# SYSTEMATIC REVIEW

# **REVISED** Toward person-centred measures of contraceptive

# demand: a systematic review of the relationship between

# intentions to use and actual use of contraception.

[version 3; peer review: 2 approved, 1 approved with reservations]

Victoria Boydell<sup>1</sup>, Kelsey Quinn Wright<sup>2</sup>, Shatha Elnakib<sup>3</sup>, Christine Galavotti<sup>4</sup>

<sup>1</sup>University College London, London, England, UK
 <sup>2</sup>University of Helsinki, Helsinki, Uusimaa, Finland
 <sup>3</sup>Johns Hopkins University, Baltimore, Maryland, USA
 <sup>4</sup>Bill & Melinda Gates Foundation, Seattle, Washington, USA

 V3 First published: 03 Jan 2024, 8:1 https://doi.org/10.12688/gatesopenres.15078.1
 Second version: 25 Jun 2024, 8:1 https://doi.org/10.12688/gatesopenres.15078.2
 Latest published: 28 Feb 2025, 8:1 https://doi.org/10.12688/gatesopenres.15078.3

### Abstract

#### Background

Understanding people's interest in using modern contraception is critical to ensuring programs align with people's preferences and needs. Current measures of demand for contraception are misinterpreted. More direct measures of intention to use (ITU) contraception do exist but remain underexplored. This systematic review examines the relationship between intention to use and actual use of contraception.

### Methods

We searched PubMed, PsycInfo, Web of Science, and the Cochrane Collaboration to identify studies published from 1975-2020 that: (1) examined contraceptive behaviour, (2) included measures of ITU and future contraceptive use, and (3) included at least one quantitative measure of association between ITU and actual use. The inclusion criteria were: 1) examined contraceptive behaviour (excluding condom use only), (2) included disaggregated integral measures of ITU contraceptives and later contraceptive use, (3) included at least one quantitative measure of the association between ITU contraceptives and actual contraceptive use, (4) study population was women of

#### **Open Peer Review**

# Approval Status 🗹 🗸 ?

	1	2	3
version 3			
(revision)	viow		
28 Feb 2025			
version 2	Ĩ ?	~	?
(revision)	view	view	view
25 Jun 2024	ſ	t	
version 1	?	?	
03 Jan 2024	view	view	

- Emily R Boniface D, Oregon Health & Science University, Portland, USA OHSU-PSU School of Public Health, Portland, USA
- 2. Anastasia J Gage (D), Tulane University, New Orleans, USA
- 3. **Jean François Régis Sindayihebura** (D), University of Burundi, Bujumbura, Burundi

Any reports and responses or comments on the article can be found at the end of the article.

reproductive age, (5) were peer-reviewed, and (6) written in English.

#### Results

10 prospective cohort studies met the inclusion criteria; these provided 28,749 person-years of data (N=10,925). Although we could pool the data for unadjusted odds ratios, a metanalysis was not possible. We calculated that 6 of the 10 studies indicated significant, increased, unadjusted odds of subsequent contraceptive use after reporting ITU. Of those, 3 study analyses reported significant, positive adjusted odds ratios for the relationship between intention to use and later contraceptive use across varying covariates. The range of confounding factors, particularly around sub-populations, points to the need for more research so that a meta-analysis can be done in the future.

#### Conclusions

People's self-reported ITU contraception has the potential to be a strong predictor of subsequent contraceptive use. Few studies directly examined the relationship between ITU and contraceptive uptake and recruitment was primarily pregnant or postpartum samples.

#### **Keywords**

Systematic review, contraception, intention, preferences

#### This article is included in the International

**Conference** on Family Planning gateway.

#### Corresponding author: Victoria Boydell (vickyboydell@gmail.com)

Author roles: Boydell V: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Wright KQ**: Data Curation, Formal Analysis, Investigation, Methodology, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Elnakib S**: Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Galavotti C**: Conceptualization, Methodology, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

**Grant information:** This work was supported by the Bill and Melinda Gates Foundation [INV-020683]. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.* 

**Copyright:** © 2025 Boydell V *et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Boydell V, Wright KQ, Elnakib S and Galavotti C. Toward person-centred measures of contraceptive demand: a systematic review of the relationship between intentions to use and actual use of contraception. [version 3; peer review: 2 approved, 1 approved with reservations] Gates Open Research 2025, 8:1 https://doi.org/10.12688/gatesopenres.15078.3

First published: 03 Jan 2024, 8:1 https://doi.org/10.12688/gatesopenres.15078.1

#### **REVISED** Amendments from Version 2

We are grateful for the feedback from the two reviewers. We have made the clarifications around what we mean by "strength of association", we have simplified the argument being made to be less ambitious in scope and added further detail to Table 1. Several of the methodological clarifications requested were made in the last set of revisions. All editorial and grammatical issues were also addressed.

Any further responses from the reviewers can be found at the end of the article

#### Introduction

Understanding people's desire to use modern contraception is critical to ensuring programs support people to achieve their reproductive needs and preferences. Since the 1970s 'unmet need for contraception' has been the main measure of demand for contraception, with some revisions along the way<sup>1-3</sup>. Unmet need is defined as the number or percentage of women currently married or in a union who are fecund and desire to either terminate, limit, or postpone childbearing but who are not currently using a contraceptive method<sup>4</sup>. Unmet need has been misinterpreted as a desire to use contraception when it actually measures a person's fertility intentions and then assumes because they are not using contraception that they have a "need" or want to use it5,6. However, people's fertility desires may or may not lead them to desire contraception, and thus "unmet need" may not necessarily align with people's desires to use contraception<sup>7-10</sup>. In addition to this misinterpretation, recent research has shown further limitations of unmet need: the calculations used for global estimates differ<sup>4,8,11,12</sup> and the focus on women in unions miscategorises and excludes many women in other arrangements<sup>7,11,13–18</sup>.

Ilene Spiezer et al., in considering how better to apply a human rights and reproductive rights lens, suggest we need to advance person-centred measures that better reflect people's needs and preferences<sup>6</sup>. As such, if we want to understand the relationship between the desire or intention to use contraception and contraceptive use, we need measures that ask women whether they desire or intend to use. Intention-to-use (ITU) contraception captures a person's interest in using contraception in the future by directly asking respondents what their stated preferences are about using contraception. This may better predict future contraceptive use and could potentially be a way to estimate programmatic gaps more accurately for those who face barriers<sup>12</sup>. Though ITU data has been collected since the 1970s, it has yet to receive the same attention as other key family planning metrics (e.g., unmet need, additional/new users, discontinuation)<sup>14,19,20</sup>.

To test the potential scope of ITU as a more person-centred measure to support more responsive contraceptive programme, we first conducted a scoping review and found that scholars working on ITU suggest that contraceptive intentions as a proximate predictor of future contraceptive use merits further research<sup>5,12,15,16,19–24</sup>. The earlier scoping review included a wider range of evidence and identified 112 papers and their operationalizations of ITU; here we build off of that work to examine a subset of the studies where the data collection design and reporting was sufficient to be able to assess whether ongoing and continued measurement of ITU has the potential to accurately predict subsequent contraceptive use for those who desire it. The research protocol is registered in PROSPERO<sup>25</sup>.

#### Methods

#### Search strategy

The search strategy was informed by the earlier scoping review that examined the extent, range, and nature of the evidence on measuring ITU<sup>5</sup>. This scoping review indicated that further analysis was needed to better understand whether ITU has significant effects on subsequent contraceptive uptake, so we performed a systematic review to examine this relationship. For this systematic review, we followed the PRISMA guidelines for reporting systematic reviews and meta-analyses<sup>26</sup>. Please see Figure 1. We searched PubMed, PsycInfo, Web of Science, and the Cochrane Collaboration for studies published between 1975 and August 2020 using search terms relevant to intent-to-use and contraceptive use. The search terms and strategy are shown in the protocol<sup>25</sup>.

#### Inclusion and exclusion criteria

The study design included in the review were experimental, quasi-experimental, or observational studies with either a pre/post or treatment/control comparison. Studies were eligible for inclusion if they: (1) examined contraceptive behaviour (excluding condom use only), (2) included disaggregated integral measures of ITU contraceptives and later contraceptive use, (3) included at least one quantitative measure of the association between ITU contraceptives and actual contraceptive use, (4) the study population was women of reproductive age, (5) were peer-reviewed, and (6) were written in the English language. There were no limits to study inclusion based on the study setting. Studies were excluded if the full text was not accessible, not published in a journal (e.g., dissertations), or not written in English.

#### Study selection and data extraction

We exported the search results into Endnote21 to remove duplicates and then imported the de-duplicated results into Excel 2021. Two authors (VB and SE) independently screened 1,464 titles and abstracts<sup>27</sup>. Where discrepancies arose, the authors resolved disagreements through discussion between the reviewers. Subsequently, SE and VB independently reviewed 39 full-text articles to ascertain their eligibility for inclusion and resolved disagreements through discussion. Data extracted included the year of publication, study purpose, location, study design, sample size, participant characteristics, follow-up period in months, type of contraceptive used, measurement of ITU, measurement of contraceptive use, attrition, number of participants who reported ITU contraception who subsequently did and did not use contraception, the number of participants who reported no ITU contraception



Figure 1. PRISMA.

who then did and did not use contraception, and effect measure and size (See Table 1). Data were then independently extracted from the 10 included articles by one author (SE) using a predesigned data extraction form<sup>27</sup>. One author (KW) reviewed the full papers and checked the data extraction. We calculated unadjusted odds ratios for the included studies, as several did not report adjusted odds ratios for the relationship between ITU and contraceptive use. We report both our calculations of the unadjusted odds ratios and author's adjusted odds ratios with the variables adjusted for in our presented results.

#### Assessment of risk of bias

One author (SE) assessed the risk of bias using the Joanna Briggs Institute Critical Appraisal Checklist for Cohort Studies<sup>28</sup>, which assesses the trustworthiness, relevance and results of cohort studies. A scoring system assigns a score of 1 or 0 against each risk of bias domain. The scores were assigned and then summed across each domain, and studies were given a score ranging from 1 to 11. Subsequently, studies were classified into low (score below 5), medium (score of 6 to 8) and high quality (score above 8). Table 2 outlines the results of the assessment for each study.

#### Data synthesis

Although some of the included papers did report relationships between intention to use and contraceptive use adjusted for a variety of covariates, these covariates are not the same across different studies. This means that either different studies included completely different covariates in their adjusted models or the way similar covariates were measured was not comparable across studies. Therefore, we calculated unadjusted odds ratios for the relationship between ITU and contraceptive use and reported on the adjusted ratios reported by authors. Despite the small sample size, we attempted to run a meta-analysis that combined the results of the studies for which we were able to calculate unadjusted odds ratios, as this would have generated a more robust source of evidence. However, meta-analysis diagnostics indicated that the high degree of variation across studies in follow up times, predictor and outcome measures, and sample populations (See Table 2) precluded pooling the data for a meta-analysis. This is the first attempt to systematically synthesise this information, and more studies that assess the longer-term relationship between reported intent to use and contraceptive use are needed for any future meta-analyses.

Table 1. Description of included papers.

Measure of contraceptive use	Not described	Self-reported use	Use of modem contraception	Self-reported use	Self-reported use		
Measure of intention	All ever-married respondents who contraceptive method were asked: "Do you imend to use a method to delay or avoid pregnancy at any time in the fuure/in the next 12 months?"	Not described	Non- contracepting asked whether asked whether they would use contraceptives in the future	Contraceptive methods she was planning to use in the first year after delivery	Not described		
What Significance Test is Testing For	Whether contraceptive use significantly increased anong those reporting ITU compared to those not reporting	Same as Curtis and Westhoff, 1996	Same as Curtis and Westhoff, 1996	Same as Curtis and Westhoff, 1996	Testing whether those who those who intention to use had different actual use actual use compared to those who were use a method use a method		
Calculated Unadjusted Odds Ratio (CI)	7.40 (5.51,9.33)	2.17(1.11,4.25)	3 years 4.48 (3.13, 6.42) 30 months 3.75 (2.62, 5.38) 24 months 3.22 (2.24, 4.62) 18 months 2.59 (1.79, 3.75) 12 months 2.27 (1.55, 3.33)	1.05 (.67, 1.64)	4.55 (3.00, 6.92)		
Results	OR: 6.78**** aOR: 2.6**** aOR (with interactions): 2.40	aOR (any method); 1.549 aOR (any modern method);1.085	HR: 1.65* aHR: 1.45*	HR (implant use only): 1.88** aHR (implant use only): 1.95*	OR (4 months): 7.89*** OR (12 months): 3.32*** aOR (4 months): 4.60*** aOR (12 months): 2.38		
Effects Measure Reported in Study	Odds Ratio	Odds Ratio	Hazard Ratio	Hazard Ratio	Odds Ratio		
Quality Rating	High (10)	High (10)	(9) High (9)	(9) High (9)	Medium (7)		
Follow up Period	3 years	1 year	6,12,18, 24, and 36 months	3, 6, and 12 months after delivery	months months		
Study Design	Longitudinal (cohort)	Longitudinal (cohort)	Longitudinal (cohort)	Longitudinal (cohort)	Longitudinal (cohort)		
Study Location	Morocco	Ghana	Uganda	Malawi	Cambodia		
Contraceptive type	Any moder n contraceptive	Any modern contraceptive	Any modern contraceptive	Long-Acting Reversible Methods	Oral contraceptive PIII		
Participant Sample Size at Baseline and Follow Up(s)	908 women married to same partner at both surveys, non-users at initial survey	240 pregnant women at Antenatal care settings at baseline and 164 at endline	747 sexuality active, non- contracepting, fecund, women	539 postpartum women (3 months), 480 (6 months), and 331 (12 months)	500 postabortion patients		
Aim	To examine the relationship between stated ITU between stated ITU contraceptives and subsequent use during a three-year period	To examine the uptake and continuation of family planning following enrolment in group versus individual ANC	To assess the dynamic influence of unmet need on thre to contraceptive uptake, as compared with that of contraceptive intentions and their concordance	To (1) calculate the incidence of LARC use among postpartum Malawan women, and (2) assess if LARC involvedeg and ITU LARC were associated with LARC uptake.	To evaluate which characteristics collected at the point of abortion are associated with contraceptive postabortion period for women.		
Study	Kurtis & Westoff 1996	Lori <i>et al.</i> 2018	Sarnak et al. 2020	2016 2016	Adelman et al. 2019		

Measure of contraceptive use	Self-reported use	Self-reported use and for those who reported more than one method, the most efficient was used.	Self-reported use	Self-reported use				
Measure of intention	7-point scales responses to the statement 'ff do have intercourse in the next year, I no (very unkely) to ever use (method X) for birth control."	Women were asked while pregnant what type of contraceptive they intended to use after childbirth	Pregnant and nonpregnant nonpregnant younger han 50 were asked: "Do were asked: "Do up ou think you will we a method to delay or avoid delay or avoid any time in the future?" and were asked which method they intended to use	7-point Likert scale measuring from likely to unlikely response unlikely response to the statement: "I intend to use contraception within the next 2 years"				
What Significance Test is Testing For	Testing correlation of intention to use method with frequency of use in the following year	Only assess significance by demographic or pregnancy planning group, not overall between ITU and contraceptive use	Only assess demographic demographic planning group, not overall significance contraceblve use and Westhoff, 1996					
Calculated Unadjusted Odds Ratio (CI)	ź	1.48 (.54, 4.04)	7.25 (5.50, 9.50)	YZ YZ				
Results	Pill (female): 0.42*** Pill (male): 0.10 Diaphragm (female): 0.27**** Diaphragm (male): 0.27* Withdrawal (female): 0.20** Withdrawal (male): 0.46***	28.9% concordance between contraceptive preference and subsequent contraceptive use.	OR (women with unmet need): 8.29* OR (women with no unmet need): 7.17*	Correlation (for contraceptive use): 0.68**				
Effects Measure Reported in Study	Correlation coefficient	Concordance	Odds Ratio	Behavioural Intention B correlation				
Quality Rating	Medium (7)	Medium (6)	Medium (8)	Medium (6)				
Follow up Period	1 year	6 months after birth	3 years	2 years				
Study Design	Longitudinal (cohort)	Longitudinal (cohort)	Longitudinal (cohort)	Longitudinal (cohort)				
Study Location	O SA	Brazi	Bangladesh	∀S ∪				
Contraceptive type	Any modern contraceptive	Any modern contraceptive	Any modern contraceptive	Oral Contraceptive Pill				
Participant Sample Size at Baseline and Follow Up(s)	325 postpartum, low- income, breastfeeding contraceptive initiators	474 AnteNatal Care (ANC) patients	3.933 married women at baseline and 3,687 at endine	279 married women at baseline and 244 at endine				
Aim	To understand adolescent beliefs about contraception and their intention to use	To examine the effect of pregnancy planming staus on the relationship between TU and current use of contraceptives among postpartum women	To link women's contraceptive uptake and experience of unwanted pregnancy between 2006 and 2009 to their unwet need status and their stated TTU contraceptives in 2006	To examine whether within versus across- subject procedures are more accurate for predicting behaviour from attitudes				
Study	Adler et al. 1990	Borges et al. 2018	Callahan & Becker 2014	Davidson & Jaccard 1979				

What Measure of Measure of Significance intention contraceptive Test is Testing use For	Tests whether Respondents Self-reported difference intending to use between a birth control within and method during across subject the next year Phi-square were asked what coefficients is method they intended to use.								Tests whether Not described Not described LARC uptake at Site A (public PP services)	were different wera different (guaranteed implant and IUD services)		Testing whether Not described Self-reported those who intended to use contraceptives were significantly more likely to use compared to those who had not planned on using a method	Correlation Not described For the analysis, between prenatal contraceptive		
Unadjusted Odds Ratio (CI)	¥ Z	*								n 1.23 (.48, 3.21)			2.53 (1.53, 3.60)	0.75 (.47, 1.22)	
Results	Within and across subjects Condoms (within subjects): 0.86**	Condoms (across subjects): 0.63*	Pill (within subjects): 0.83**	Pill (across subjects): 0.77**	IUD: (within subjects): 0.94***	IUD: (across subjects): 0.85**	Diaphragm (within subjects):	0.92** Diaphragm (across subjects):	0.78**	53% pregnant women reported a intention to use a LARC or to be sterilised after delivery	72% of women who had intended to start using a LARC actually did so at a site offering LARCs compared to only 4% of women a	public FP sites***	Of the 421 women who were asked the NFHS question on contraceptive intentions, 127 stated that they would use a method in the future. More than hits (51%) of the women stating they would use a more do intersurvey prend compared to 29% of respond not practice family planming actually did so**	Prenatal contraceptive intention and postpartum in-hospital correlation: 0.41***	
Effects Measure Reported in Study	Phi coefficients	Phi coefficients								Percentages			Proportions	Correlation coefficient	
Quality Rating	Medium (6)						Medium (6)			Medium (7)	Low (5)				
Follow up Period	1 year						9 months after birth			6 years	Not specified				
Study Design	(cohort) (cohort)						Longitudinal (cohort)			(cohort) (cohort)	Longitudinal (cohort)				
Study Location	A SU								Rwanda			India	NSA		
Contraceptive type	Any modern contraceptive								Any modern contraceptive			Any modern contraceptive	Long-Acting Reversible Methods		
Participant Sample Size at Baseline and Follow Up(s)	aged 18-38 years								219 HIV-positive pregnant women at ANC settings at baseline and 205 at endline			421 female participants in the 1922-92 National Family Health Survey	223 postpartum women at baseline; 214 women postpartum in the	hospital and 119 women	
Aim	To understand factors that moderate the attitude-behaviour relation								To investigate unmet need for LARCs and sterilization among HIV-positive pregnant	women, and the impact of increased access to LARCs in the postpartum period on their contraceptive	uptake	To investigate women's TU a method as a measure of contraceptive demand	To understand how women's prenatal infant feeding and	contraception intentions	
Study	8 Morrison 1983								Dhont et al. 2009			Roy et al. 2003	Johnson et al. 2019		

Page 7 of 28

Study	Quality Rating	Calculated Unadjusted Odds Ratio (CI)	Author Reported Adjusted Odds Ratios (CI) for ITU coefficient on contraceptive use, and factors adjusted for				
Curtis & Westoff 1996	High (10)	7.40*** (5.51-9.93)	2.64*** (CI not given)	Categorical: fecundity, wanted last birth, fertility preference, prior contraceptive use, discussed family size with partner, attitudes about family planning messages in media, listened to radio weekly, education, residence, age, births, child deaths Continuous: number of living children <i>Note</i> : do not include results for interacted model			
Roy <i>et al.</i> 2003	Medium (7)	2.53*** (1.53-3.60)	Contraceptive use reported as regression outcome, intention to use not distinctive predictor variable but as a stratifier variable				
Dhont <i>et al.</i> 2009	Medium (6)	1.23 (0.48-3.21)	Contraceptive use not reported as regression outcome				
Callahan & Becker 2014	Medium (8)	7.25*** (5.50-9.56)	Contraceptive use not reported as regression outcome				
Tang <i>et al.</i> 2016	High (9)	1.05 (0.67-1.64)	HR: 1.95** (1.28-2.98)	Age, parity, education, having a friend using the implant, HIV status, having trouble obtaining food, clothing, or medications			
Borges <i>et al.</i> 2018	Medium (6)	1.48 (0.54-4.04)	Contraceptive use reported as regression outcome, intention to use not distinctive predictor variable				
Lori <i>et al.</i> 2018	High (10)	2.17* (1.11-4.25)	<i>Note</i> : postpartum, modern method only 1.085 (0.444-2.655)	Age, gravida, religion, highest level of education			
Adelman <i>et al.</i> 2019	Medium (7)	4.55*** (3.00-6.92)	Note: ITU not presented in final adjusted models Outcome is 80% "continued contraception use" over 4 month: 7.98*** (2.99-20.83) Note: outcome is 80% "continued contraception use" over 12 months: 3.32** (1.35-8.20)	Categorical: age, SES, residence, education, marital status, occupation, number of living children, number of previous abortions, abortion method, disclosure of abortion, previous contraception use, postabortion contraceptive intention, fertility intention, contraceptive decision making			
Johnson <i>et al.</i> 2019	Low (5)	0.75 (0.47-1.22)	Contraceptive use not reported as regression outcome				
Sarnak <i>et al.</i> 2020	High (9)	36 months 4.48*** (3.13-6.42)	36 months: 1.45*** (1.22-1.73)	Categorical variables: age, parity, education, residence, wealth quintile			

#### Table 2. Summary of the findings from the included papers.

\*p<.05 \*\*p<.01 \*\*\*p<.001

#### Results

This is the first attempt to systematically synthesise this information, and more studies that assess the longer-term relationship between reported intent to use and contraceptive use are needed for any future meta-analyses (see Table 1). The limited number of studies and the heterogeneity of the data in available studies made it impossible to conclusively demonstrate the effects of intention to use contraception on actual use across different contexts; instead, we outline what were the range of variables underpinning the heterogeneity.

#### Study characteristics

The search yielded 1,464 articles. Many papers were excluded because they did not have a clear definition of intention to use (732), did not state an association between intention to use and contraceptive use (235), did not meet the study design requirements (238), did not contain sufficient information in the text to be assessed against the inclusion criteria (30), focused on condoms (161), did not include a measure of contraceptive use (61) or focused on only on the drivers of intention to use and did not test the association with actual use (17).

After the initial abstract screening and full paper review, a total of 10 articles were included<sup>27</sup>. One of the 10 studies was conducted in the USA. The remaining studies were undertaken in low- and middle-income country (LMIC) settings: Bangladesh (n=1), Brazil (n=1), Cambodia (n=1), Ghana (n=1), India (n=1), Malawi (n=1), Morocco (n=1), Rwanda (n=1), and Uganda (n=1). All 10 studies were longitudinal cohort studies with pre-and post-tests or treatment and control groups. The characteristics of the studies, such as study aim, population, location, study design, follow up period, quality rating, effects measures, measure of ITU and measure of contraceptive use, are summarized in Table 1.

#### Number and characteristics of participants

The number of participants varied between studies from 219 to 3,933, while six papers had sample sizes of approximately 200 to 300 participants. The papers looked at a variety of different participants – either women as broad category (e.g., sexually active or married) or at different points in their reproductive career (e.g., pre and post-partum). Two papers sampled married women<sup>16,17</sup>; two papers sampled postpartum women<sup>29,30</sup>; two papers sampled pregnant women<sup>31,32</sup> and another two sampled sexually activity women<sup>7,33</sup>. Only one paper looked at women post-abortion<sup>34</sup>. These papers provide 28,749 person-years of data (N=10,925).

#### Definition of measures and outcomes

Half of the 10 included studies did not describe how exactly intention-to-use contraception was measured, and no details are provided on the exact wording of the items used to solicit information on the intention to use contraception<sup>29,31–34</sup>. Of the remaining studies, three used items that asked about the intention to use contraception in the future with no exact time frame specified<sup>7,16,33</sup>. Only one study used items that asked about intention to use contraception within a specific time; the time frame used was within the year<sup>30</sup>.

In contrast, the majority of included studies did outline how they captured the outcome measure, contraceptive use. All of the studies used self-reported contraceptive use as the outcome measure (n=10). However, Johnson *et al.* used clinical records and two studies did not specify how they captured contraceptive use<sup>17,29,32</sup>.

There was extensive heterogeneity in the measures used to report associations or effects in the included studies. Four papers used odds ratios to examine the relation between intention-to-use and use of contraception<sup>7,16,31,35</sup>. Across the studies that used odds ratios, researchers compared women who intended to use contraception to women who did not intend to use any method. These four studies found higher odds of women using contraception if they had planned to use it previously; this finding was statistically significant at p<.001 for three of the four studies. One paper used correlation coefficients<sup>29</sup>, and two papers used hazard ratios<sup>7,30</sup>. The remaining papers reported on their findings using "concordance"<sup>33</sup>, and simple percentages or proportions<sup>32,34</sup>.

#### Associations

Of the 10 studies for which we calculated unadjusted odds ratios of contraceptive use by intention to use status, six had significant, increased odds of subsequent contraceptive use after reporting an intention to do so at an earlier point, see Table 2. The unadjusted associations range from 0.75-7.40 based on odds ratios. Of the 10 included studies, five reported on an adjusted relationship between intent to use as a predictor variable and contraceptive use as an outcome variable. Of these, four found significantly increased odds or hazards of contraceptive use given stated intent to use at the initial measurement. These studies adjusted for a variety of covariates, with the most common being age, measures of the number of pregnancies, and education. As would be expected, the magnitude of significant unadjusted odds ratios generally decreases with adjustment for covariates, however the strength of the association, e.g, the precision with which the confidence intervals are estimated to be different than 1, does not. In one case, Tang et al. (2016), our unadjusted odds ratio was non-significant, while the author's calculation of an adjusted hazard ratio was. In the study conducted by Lori et al. (2018), our unadjusted calculation was significant at the p<.05 level while the authors' adjusted calculation is non-significant.

#### Specific contraceptive methods

Two of the included papers examined only long acting reversible method (LARC) use at follow up<sup>30,32</sup>. Three studies included only what would be considered modern contraceptive methods, including LARCS such as IUDs and implants, and shorter term methods like pills, injectables, vaginal rings, and condoms, alongside sterilization<sup>29,33,35</sup>. The remaining studies grouped contraceptive methods into various groupings, such as 'modern', 'modern and reversible', 'modern and permanent', and 'traditional'<sup>7,16,17,31,34</sup>.

#### Time frame

There were also significant differences in the intervals between baseline and follow-up within the included studies. Most of the studies examined the relationship between intention to use and contraceptive use over long-term (longer than one-year) periods, ranging from one-year follow up measurements to six years in between measurements<sup>7,16,17,31,34,35</sup>. Some of these studies of longer duration included intervening measurements at specified month-intervals<sup>7,30,35</sup>. The differences in odds ratios of contraceptive use at these intervals especially highlights the need for subsequent work to focus on specific intervals to better understand the duration range of intention to use reports. The remaining papers examined contraceptive use for less than one year, or the duration of follow up was unspecified<sup>29,32,33</sup>.

#### Population

Of the 10 studies included, six focused in and around pregnancy; this refers to the antenatal, postabortion, and postpartum period. Two of the 10 studies examined intention to use contraception among women in the postpartum period and followed up on whether women's intention had transformed into use over the following 12 months<sup>4,29,30</sup>. A further three studies examined women's choice to use contraception in the antenatal period and followed up six months to one year after to see if they were using a method<sup>31–33</sup>.

Only one study looked at the intention to use among women following an abortion<sup>35</sup>. In Cambodia, Adelman *et al.*, examined what characteristics collected at the point of abortion are associated with oral contraceptive use at four and 12 months after the abortion. Intention to use contraception was found to be positively associated with increased contraceptive use over the year<sup>35</sup>.

The remaining four studies looked at the intention to use contraception among women with partners, including married women<sup>7,16,17,34</sup>. Using longitudinal data from rural Bangladeshi women (n=2,500), Callahan and Becker found that intention to use a method was predictive of subsequent contraceptive use for women with and without an unmet need. Only two of these studies specified whether the women were non-users<sup>7,16,17</sup>. In Uganda, Sarnak et al., compared unmet need and contraceptive adoption to contraceptive intentions and use7. They found that women who intended to use contraception in the future used contraceptives significantly earlier (aHR = 1.45, 95% CI = 1.22-1.73) than those who did not intend to use contraception7. Interestingly, women with an intention to use but not classed as having no unmet need had the highest rate of adoption compared to those with no unmet need and no intention to use (aHR = 2.78, 95% CI = 1.48-5.2586. The follow-up period to see if married women's intentions had turned into actual contraceptive use was a one-to-three-year period in this set of studies<sup>7,16,17,34</sup>.

#### Quality of evidence in included studies

We used the Joanna Briggs Institute Critical Appraisal Checklist for Cohort Studies<sup>28</sup>, which assesses the trustworthiness, relevance and results of cohort studies, to rate the quality of each study using the following domains: the sample, exposure measures, confounding factors, outcome measures, follow-up time reported, and type of analysis used. Four studies were graded as high quality, and five were of medium quality. One study was classed as low quality.

#### Discussion

In this review, we found that there are significant positive associations between intention to use a contraceptive method and actual use in six medium- to high-quality studies. Yet the heterogeneity across the papers poses an analytical challenge for us to be able to really interrogate the potential of this person-centred measure; this in itself is a finding and speaks to the need for (1) refining the outcomes to measure intention to use, and (2) identifying a) which relevant variables need to be included in adjusted models and b) how these variables can be measured in ways so that they are comparably reported across studies.

#### Refining the outcomes

Reading across the papers, there is inconsistency in how ITU is currently operationalized and applied. This analysis found that five (n=5) papers did not provide details on the wording of the items used to measure ITU<sup>29,32-35</sup>. Based on what information is available from the included papers, five (n=5) papers captured goal intentions<sup>7,16,17,24,35</sup> whereas four (n=4) captured implementation intention<sup>20,30,31,34</sup>. This finding is significant because established behavioural theory suggests that distinguishing the type of intention may be helpful as implementation intentions are more likely to translate into the behaviour than goal intentions<sup>36</sup>. Gollwitzer and Sheeran helpfully distinguish between goal intention and what people plan to do some time in the future<sup>37</sup>. In contrast, implementation intentions are more specific regarding when, where, and how one's achievement of an intention will occur. Implementation intentions tend to be oriented towards a particular action, whereas goal intentions tend to be outcomes achieved by performing several actions37. Gollwitzer and Sheeran argue that goal intentions do not prepare people for dealing with the problems they face in initiating, maintaining, disengaging from, or overextending themselves in realizing their intentions<sup>37</sup>. In contrast, an implementation intention sets out the when, where, and how in advance and is a form of planning that bridges the intention-behaviour gap, increasing the likelihood of intentions being realized<sup>37</sup>. Unfortunately, none of the papers included distinguished between goal and implementation intentions. Additional research on how ITU is conceptualized and operationalized is needed to understand how different types of intentions (e.g., goal vs implementation) predict contraceptive use and continuation. To address this, further research in needed using standardized ITU and outcome measures and similar follow-up durations amongst similar populations to assess the magnitude and direction of associations between ITU and contraceptive use.

#### Adjusting for confounders

Given the heterogeneity, several potential confounding variables could affect whether an intention to use contraception leads to future contraceptive use. These possible confounding variables make it difficult to establish a causal link between ITU and contraceptive use. This review points to several potential confounding variables to consider in future work.

Several studies in this review focused on populations during and around pregnancy. This could be an artefact of research study design as recruiting women attending pregnancy-related services may be easier. It could be an artefact of programme design in that women are more likely to engage in healthcare during pregnancy. Similarly, parity and relationship status may also affect whether an intention to use contraception translates into actual use. Future research should examine how pregnancy status may affect intentions to use contraception compared to women seeking to prevent pregnancy who are not pregnant.

Another variable that may affect the relationship between intention to use and actual use is the type of contraception method being considered. For example, long-acting reversible contraceptive methods may require more commitment and planning, whereas short-acting methods may be easier to access and use. Hence, the specific type of method may differentially affect the ease or difficulty of a person transforming their intentions into action. Work on developing a psychometric scale on contraceptive intent highlighted that contraceptives are a form of medication, and the woman's desire and adherence to them are influenced by beliefs about the medicine<sup>10</sup>. Another variable we noted is how long it may take to move from intention to action and when to measure if this execution has taken place. Several studies reported different follow-up durations7,30,35. Our findings are too inconsistent in reporting the timeframe to make any generalizations about the appropriate time to move intention to action; the literature on behaviour implementation suggests that this is an important avenue for future study.

The range of potential interceding factors that emerged in the review point to the fact that contraceptive behaviour is a complex psychosocial process shaped by the confluence of individual and contextual factors<sup>10</sup>. Such factors may help explain how pregnancy and relationship status are related to intentions or use of specific methods, whether goal or implementation intentions result in actual use, and over what timeframe intentions to use contraception are likely to transform into action. In turn, this can contribute to better understand people's needs and preferences and how we can align programs to support them to achieve their reproductive goals and contraceptive goals.

There are several limitations to this review. There were relatively few studies that met the inclusion criteria. The relationship between ITU and contraceptive uptake was not the primary outcome of interest for those included papers. Thus, we had to calculate an odds ratio to estimate that relationship. Therefore, we treat our results as indicative. Another limitation is that the samples recruited for the included studies were primarily pregnant or postpartum samples-the desire to start sexual activity and contraception may be different for these populations compared to others. Geographic settings, particularly the difference in health systems and contraceptive access, may also explain the differences we found. In addition, other factors (e.g., cultural and social norms, knowledge about contraceptive methods, personal beliefs) may all contribute to reproductive and contraceptive intentions, decision-making, and subsequent use, and require further consideration.

#### Conclusion

Six studies indicated significant, increased odds of subsequent contraceptive use after reporting ITU and show a significant positive association between desire to use contraception and actual use. This suggests that self-reported ITU contraception may be a strong predictor of subsequent contraceptive use and a promising alternative measure of demand for contraception. As a person-centred measure, we need further high-quality research that measures the relationship between intent-to-use and contraceptive use using standardized measures and more fully considering the range of additional factors that may influence both ITU and subsequent use.

#### **Data availability** Underlying data

OSF: Toward person-centred measures of contraceptive demand: a systematic review of the intentions to use contraception and actual use. https://doi.org/10.17605/OSF.IO/6FXQT<sup>27</sup>.

The project contains the following underlying data:

- ITU Sys Review underlaying data citations (data citations for the systematic review).
- ITU Sys Review underlaying data citations screening too (screening tool).
- ITU Sys Review underlaying full papers (list of full papers for the systematic review).
- ITU Sys Review underlaying full paper screening tool (screening tool for full papers for the systematic review).

#### Extended data

OSF: Toward person-centred measures of contraceptive demand: a systematic review of the intentions to use contraception and actual use. https://doi.org/10.17605/OSF.IO/6FXQT<sup>27</sup>.

This project contains the following extended data:

- Supplementary Table 1. (Description of included studies)
- Supplementary Figure 1. (PRISMA flowchart)
- Data collection tool. (raw data used in analysis)

#### Reporting guidelines

OSF: PRISMA and PRISMA for abstracts checklists for 'Toward person-centred measures of contraceptive demand: a systematic review of the intentions to use contraception and actual use'. https://doi.org/10.17605/OSF.IO/6FXQT<sup>27</sup>.

Data are available under the terms of the Creative Commons Zero "No rights reserved" data waiver (CC0 1.0 Public domain dedication).

#### References

Westoff CF: The unmet need for birth control in five Asian countries. Fam 1 Plann Perspect. 1978; 10(3): 173-81. PubMed Abstract | Publisher Full Text

- Westoff CF, Ochoa LH: Unmet need and the demand for family planning. 2. Institute for Resource Development; Columbia, MD: DHS Comparative Studies No. 5, 1991
- Bradley SEK, Croft TN, Fishel JD, et al.: **Revising unmet need for family** planning. ICF International; Calverton, MD. DHS Analytical Studies No. 2. 2012. 3. Reference Source
- Measure Evaluation: Family planning and reproductive health indicators database. 2020; March 31st, 2021. 4. **Reference Source**
- Boydell V, Galavotti C: Getting intentional about intention to use: a scoping 5. review of person-centered measures of demand. Stud Fam Plann. 2022; 53(1): 61-132

PubMed Abstract | Publisher Full Text | Free Full Text

- Speizer IS, Bremner J, Farid S: Language and measurement of contraceptive 6. need and making these indicators more meaningful for measuring fertility intentions of women and girls. *Glob Health Sci Pract.* 2022; **10**(1): e2100450. PubMed Abstract | Publisher Full Text | Free Full Text
- Sarnak D, Tsui A, Makumbi F, et al.: The predictive utility of unmet need 7. on time to contraceptive adoption: a panel study of non-contracepting Ugandan women. Contracept X. 2020; 2: 100022. PubMed Abstract | Publisher Full Text | Free Full Text
- Singh S, Darroch JE: Adding it up: costs and benefits of contraceptive 8. services-estimates for 2012. New York: Guttmacher Institute and United Nations Population Fund (UNFPA), 2012. **Reference Source**
- Moreau C, Shankar M, Helleringer S, et al.: Measuring unmet need for 9 contraception as a point prevalence. BMJ Glob Health. 2019; 4(4): e001581. PubMed Abstract | Publisher Full Text | Free Full Text
- Raine-Bennett TR, Rocca CH: Development of a brief questionnaire to assess 10. contraceptive intent. Patient Educ Couns. 2015; **98**(11): 1425-30. PubMed Abstract | Publisher Full Text | Free Full Text
- Peterson SA: Marriage structure and contraception in Niger. J Biosoc Sci. 11. 1999; 31(1): 93-104.

PubMed Abstract | Publisher Full Text

- Moreau C, Hall K, Trussell J, et al.: Effect of prospectively measured 12. pregnancy intentions on the consistency of contraceptive use among young women in Michigan. *Hum Reprod.* 2013; **28**(3): 642–650. PubMed Abstract | Publisher Full Text | Free Full Text
- Alkema L, Kantorova V, Menozzi C, et al.: National, regional, and global 13. rates and trends in contraceptive prevalence and unmet need for family planning between 1990 and 2015: a systematic and comprehensive analysis. Lancet. 2013; 381(9878): 1642–52. PubMed Abstract | Publisher Full Text
- Ross JA, Winfrey WL: Contraceptive use, ITU and unmet need during the 14. extended postpartum period. Int Fam Plann Perspect. 2001; 27(1): 20-27. **Reference Source**
- Bradley SEK, Casterline JB: Understanding unmet need: history, theory, and 15. measurement. Stud Fam Plann. 2014; 45(2): 123-150. PubMed Abstract | Publisher Full Text | Free Full Text
- 16. Callahan R, Becker S: Unmet need, intention to use contraceptives and unwanted pregnancy in rural Bangladesh. Int Perspect Sex Reprod Health. 2014: 40(1): 4–10. PubMed Abstract | Publisher Full Text
- Curtis SL, Westoff CF: Intention to use contraceptives and subsequent 17. contraceptive behavior in Morocco. Stud Fam Plann. 1996; 27(5): 239-50. PubMed Abstract | Publisher Full Text
- Cavallaro FL, Benova L, Macleod D, et al.: Examining trends in family planning 18. among harder-to-reach women in Senegal 1992-2014. Sci Rep. 2017 41006 PubMed Abstract | Publisher Full Text | Free Full Text
- Khan MS, Hashmani FN, Ahmed O, et al.: Quantitatively evaluating the effect of social barriers: a case-control study of family members' opposition 19. and women's intention to use contraception in Pakistan. Emerg Themes

Epidemiol. 2015; 12(1): 2.

PubMed Abstract | Publisher Full Text | Free Full Text Babalola S, John N, Ajao B, et al.: Ideation and intention to use

- 20 contraceptives in Kenya and Nigeria. Demogr Res. 2015; 33(1): 211-238. PubMed Abstract | Publisher Full Text | Free Full Text
- Ross J, Heaton L: **Intended contraceptive use among women without an unmet need**. *Int Fam Plann Perspect*. 1997; **23**(4): 148–154. 21. **Publisher Full Text**
- Ross J, Heaton L: Intended contraceptive use among women without an 22. unmet need. Int Perspect Sex Reprod Health. 1997; 23(4): 148-154 **Reference Source**
- Ross J, Winfrey W: Contraceptive use, intention to use and unmet need 23. during the extended postpartum period. Int Perspect Sex Reprod Health. 2001; 27(1): 20-27. **Reference Source**
- Hanson JD, Nothwehr F, Yang JG, et al.: Indirect and direct perceived 24. behavioral control and the role of intention in the context of birth control behavior. *Matern Child Health J.* 2015; **19**(7): 1535–42. PubMed Abstract | Publisher Full Text | Free Full Text
- Boydell V, Galvotti C: The relationship between intent to use and modern 25 contraceptive use in countries globally: a systematic review and meta-analysis. PROSPERO 2020 CRD42020199730. **Reference Source**
- Page MJ, McKenzie JE, Bossuyt PM, et al.: The PRISMA 2020 statement: an 26. updated guideline for reporting systematic reviews. BMJ. 2021; 372: n71. PubMed Abstract | Publisher Full Text | Free Full Text
- Wright KQ, Boydell V: Toward person-centred measures of contraceptive 27. demand: a systematic review of the intentions to use contraception and actual use. Dataset. 2023. http://www.doi.org/10.17605/OSF.IO/6FXQT
- Joanna Briggs Institute: Checklist for cohort studies. 2017; March 31st, 2021. 28 **Reference Source**
- 29 Johnson NA, Fuell Wysong E, Tossone K, et al.: Associations between prenatal intention and postpartum choice: infant feeding and contraception decisions among inner-city women. *Breastfeed Med.* 2019; **14**(7): 456–464. PubMed Abstract | Publisher Full Text
- Tang JH, Kopp DM, Stuart GS, et al.: Association between contraceptive 30. implant knowledge and intent with implant uptake among postpartum Malawian women: a prospective cohort study. Contracept Reprod Med. 2016; 1(1): 13 PubMed Abstract | Publisher Full Text | Free Full Text
- 31. Lori IR, Chuev M, Munro-Kramer ML, et al.: Increasing postpartum family planning uptake through group antenatal care: a longitudinal prospective cohort design. *Reprod Health*. 2018; **15**(1): 208. PubMed Abstract | Publisher Full Text | Free Full Text
- Dhont N, Ndayisaba GF, Peltier CA, et al.: Improved access increases 32 postpartum uptake of contraceptive implants among HIV-positive women in Rwanda. Eur J Contracept Reprod Health Care. 2009; 14(6): 420-5. PubMed Abstract | Publisher Full Text
- Borges ALV, Dos Santos OA, Fujimori E: Concordance between intention to use and current use of contraceptives among six-month postpartum women in Brazil: the role of unplanned pregnancy. Midwifery. 2018; 56: 94-101

PubMed Abstract | Publisher Full Text

- Roy TK, Ram F, Nangia P, et al.: Can women's childbearing and contraceptive 34. intentions predict contraceptive demand? Findings from a longitudinal study in Central India. Int Fam Plan Perspect. 2003; 29(1): 25-31. PubMed Abstract
- Adelman S, Free C, Smith C: Predictors of postabortion contraception use in 35. Cambodia. Contraception. 2019; 99(3): 155–159. PubMed Abstract | Publisher Full Text
- 36. Cohen J: A power primer. Psychol Bull. 1992; 112(1): 155-9. PubMed Abstract | Publisher Full Text
- Gollwitzer PM, Sheeran P: Implementation intentions and goal 37. achievement: a meta-analysis of effects and processes. Adv Exp Soc Psychol. 2006: 38: 69-119 Publisher Full Text

# **Open Peer Review**

# Current Peer Review Status: 💙 🔹

Version 3

Reviewer Report 03 March 2025

https://doi.org/10.21956/gatesopenres.17707.r39461

© **2025 Boniface E.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



# Emily R Boniface 问

<sup>1</sup> Oregon Health & Science University, Portland, Oregon, USA <sup>2</sup> Health Systems & Policy, OHSU-PSU School of Public Health, Portland, Oregon, USA

The authors have successfully addressed all my previous comments.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Biostatistics, contraception use, person-centered contraceptive care

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

# Version 2

Reviewer Report 11 December 2024

https://doi.org/10.21956/gatesopenres.17363.r38605

© **2024 Sindayihebura J.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Jean François Régis Sindayihebura University of Burundi, Bujumbura, Burundi

# ABSTRACT

The authors have found that the measure of the effect of the demand for family planning methods

on the actual use of contraception has always been operationalized by Unmet Need. However, the latter can also characterize women who have no intention of using it, and therefore don't really participate in the expression of demand for FP methods (unmet needs include in the analysis people who didn't want the recent birth/pregnancy in progress, or who would not like to get pregnant in the future, without having the desire to use contraception). Thus, they would like to focus the analysis on the actual needs expressed by the intention to predict contraceptive use.

By carrying out a systematic review of previous studies, they selected, according to rigorous inclusion and selection criteria, 10 studies from which they show that intention to use contraception defines its actual use in the future. The conclusion is drawn from the results of calculating unadjusted odds ratios, as the control variables were not common to the all selected studies.

# **ANSWERS TO QUESTIONS**

# Are the rationale and objectives of the systematic review clearly stated?

The researchers justify their study of the impact of intention to use contraception on subsequent actual use by contrasting it with studies of unmet need for FP. According to these authors, unmet need is not sufficient as a measure of the desire to use contraception (unmet need includes people who did not want the recent birth/pregnancy or who would not like to become pregnant, without having the desire to use contraception). In this way, they intend to focus the measure on people who actually express the need for contraception, in order to assess its effect on actual use. However, it seems that a systematic review of the literature is not enough to achieve this objective. The reviewed works are the ones that best meet this objective. They simply justify why these analyzed studies were carried out, without being able to justify their own (this systematic review).

The authors reviewed a multitude of papers, from which they eventually retained only 10. They have the advantage of formulating a good problematic than that of the study focused on people in need of contraceptives (i.e. those who express the intention to use contraception).

# Are the methods and analysis detailed enough to be replicated by others?

The authors describe at length the objective process of inclusion and selection of the studies reviewed. They also cite the calculation of unadjusted odds ratios because the variables used to control for the effect of intention to use contraception on actual contraceptive use were not the same in the reviewed studies. However, no methodological approach to calculating these unadjusted odds ratios is mentioned to allow reproducibility. Mathematical details of the methodology are needed.

The reader will also wonder how this calculation was possible when no manipulation of the databases is mentioned. What was the target population, or what were the statistical units?

# Are the statistical analysis and its interpretation appropriate?

The statistical analysis and its interpretation are appropriate, but the lack of detail in the methodology expressed above prevents us from drawing any objective conclusions.

### Are the conclusions drawn adequately supported by the results presented in the review?

The results presented appear to be a continuation of the systematic review, the selection of studies to be included and the methodology. The reader will find that the results present in unclear and immense content the effect of intention to use contraception on its actual use. The authors would do well to be brief and concise.

The conclusion presents the essential results, but lacks the necessary elements. A reminder of the context, objective and methodological approach. It should also present the strengths and limitations.

The conclusion is therefore relatively short.

Are the rationale for, and objectives of, the Systematic Review clearly stated? Partly

Are sufficient details of the methods and analysis provided to allow replication by others?  $\ensuremath{\mathbb{No}}$ 

Is the statistical analysis and its interpretation appropriate?

Partly

Are the conclusions drawn adequately supported by the results presented in the review?  $\ensuremath{\mathsf{Yes}}$ 

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Demography ; Human geography ; Social Sciences

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 13 Feb 2025

#### victoria boydell

Thank you so much for your feedback on version 1 of the paper. We have made substantial revisions to the paper based on the input of the two other reviewers and many of these changes address the issues you raised. Your concerns about the framing of the paper have been addressed. We appreciate the input and think it will make for a much-improved paper.

Competing Interests: No competing interests were disclosed.

Reviewer Report 17 July 2024

https://doi.org/10.21956/gatesopenres.17363.r37271

© **2024 Gage A.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



# Anastasia J Gage 匝

Department of International Health and Sustainable Development, School of Public Health and Tropical Medicine, Tulane University, New Orleans, Louisiana, USA

The authors have adequately addressed the reviewers' comments. Table 1 now specifies the study designs, study populations, sample size and other study characteristics, which is helpful. The authors have provided a justification for the inclusion of a "low-quality" study in the systematic review and now have a more thorough discussion of the study's limitations.

Are the rationale for, and objectives of, the Systematic Review clearly stated? Yes

Are sufficient details of the methods and analysis provided to allow replication by others?  $\ensuremath{\mathsf{Yes}}$ 

# Is the statistical analysis and its interpretation appropriate?

Yes

Are the conclusions drawn adequately supported by the results presented in the review?  $\ensuremath{\mathsf{Yes}}$ 

Competing Interests: No competing interests were disclosed.

*Reviewer Expertise:* Family planning, reproductive and maternal health

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 10 July 2024

https://doi.org/10.21956/gatesopenres.17363.r37272

© **2024 Boniface E.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

# ? Emily R Boniface 匝

<sup>1</sup> Oregon Health & Science University, Portland, Oregon, USA <sup>2</sup> Health Systems & Policy, OHSU-PSU School of Public Health, Portland, Oregon, USA

Thank you for your work on this important topic and the revisions, particularly the inclusion of

Table 1, which have improved the clarity of the manuscript. I have a few additional suggestions to improve the interpretability of the study results.

Introduction:

- 2nd paragraph: the second sentence appears to be missing some words.
- 3rd paragraph, 1st sentence: "programme" should be "programmes".

## <u>Table 1:</u>

• ANC abbreviation should be defined.

### Results:

- Associations section: I appreciate the clarification regarding magnitude or odds ratios and strength of association in the response from authors. However, without any edits to the manuscript, the text as written remains unclear from a statistical perspective. I strongly suggest specifying that "strength of association" refers to the precision of the estimates, as that is not a typical use of the term and could easily be misinterpreted.
- Specific contraceptive methods: It would be very helpful to the reader to include these details in Table 1.
- There is a lot of redundancy between the "Number and characteristics of participants" and "Population" sections. Clarity would be significantly improved by combining them or removing duplicate information.

# Are the rationale for, and objectives of, the Systematic Review clearly stated? Yes

Are sufficient details of the methods and analysis provided to allow replication by others?  $\ensuremath{\mathsf{Yes}}$ 

# Is the statistical analysis and its interpretation appropriate?

Yes

# Are the conclusions drawn adequately supported by the results presented in the review? $\ensuremath{\mathsf{Yes}}$

*Competing Interests:* No competing interests were disclosed.

Reviewer Expertise: Biostatistics, contraception use, person-centered contraceptive care

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 13 Feb 2025

### victoria boydell

Thank you so much for your feedback on version 2 of the paper. We have reviewed all your comments and have incorporated all the suggested changes. We appreciate the input and think it will make for a much-improved paper.

*Competing Interests:* No competing interests were disclosed.

# Version 1

Reviewer Report 16 April 2024

# https://doi.org/10.21956/gatesopenres.16413.r36249

© **2024 Gage A.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

# 了 🔹 Anastasia J Gage 匝

Department of International Health and Sustainable Development, School of Public Health and Tropical Medicine, Tulane University, New Orleans, Louisiana, USA

The extent to which contraceptive intention translates into actual contraceptive use has long been a subject of debate, even though studies generally show a positive correlation between intention and behavior. The research question in the present systematic review was straightforward. The authors examined whether contraceptive intentions predict subsequent contraceptive use. The inclusion and exclusion criteria were clearly specified. While all studies included in the review were longitudinal, not all employed an experimental or quasi-experimental design. Some included studies were observational. My comments on the present study are outlined below.

- The literature search strategy was comprehensive. Search terms were provided in PROSPERO and included the following: ((intent\* OR intend\*) AND ("to use")) OR (intent\* OR intend\* OR willingness) AND (contracept\* OR "birth control" OR "family planning"). The literature was independently screened by two of the authors to determine the eligibility of studies for inclusion in the systematic review, and full text articles were independently reviewed by the same authors.
- While data were independently extracted from the 10 included articles by one author using a predesigned data extraction form, a second author reviewed the full papers and checked the data extraction.
- Of 39 articles that were retrieved, 25 articles were excluded after full text screen and 4 articles during data extraction, Unfortunately, the review authors did not fully account for the excluded articles. It would be instructive to know the likely impact of their exclusion on the conclusions of the systematic review.
- One consideration was the extent to which the authors described the studies in adequate detail. Although Table 1 does not provide details about research designs, study populations,

interventions (if applicable), and study settings, this information is described in the text and gives insights into variations in the study populations and study settings.

- As all 10 studies included in the present systematic review were longitudinal cohort studies (with pre-and post-tests or treatment and control/comparison groups), the authors used the Joanna Briggs Institute Critical Appraisal Checklist for Cohort Studies to assess risk of bias. One of the included studies had a "low quality" rating and was retained in the systematic review. Could the authors kindly justify its retention? If this study were to be omitted, what would be the implications for the interpretation of the results of the review?
- The authors provided a satisfactory discussion of observed heterogeneity in the results of the review. They reported heterogeneity in (a) measures used to report associations or effects in the included studies (odds ratios, hazard ratios, correlation coefficients, simple percentages/proportions); (b) study design (which included non-randomization); (c) analysis (non-adjustment or adjustment for covariates).
- In the discussion section, the authors highlighted the importance of adjusting for possible confounding variables, such as parity, relationship status, type of contraceptive method. Contraceptive decision making is also shaped by factors that were not mentioned, including cultural and social norms, knowledge about contraceptive methods, personal beliefs, and access to and supply of contraceptive methods. It is important to mention these factors when discussing the limitations of the study.
- The preceding comment (i.e., the importance of adjusting for confounding variables) begs the question as to whether the five studies that did not report an adjusted relationship between intent to use (predictor variable) and contraceptive use (outcome variable) should be included in the systematic review. I believe that these studies should not be included as they detract from the robustness of the results.

Overall, the present systematic review highlights (a) research gaps, (b) the need for standardized measures of intention to use contraception, and (c) the importance of distinguishing between goal intentions and implementation intentions when predicting subsequent contraceptive use, after adjusting for confounding variables.

# Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?  $\ensuremath{\mathsf{Yes}}$ 

# Is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

# Are the conclusions drawn adequately supported by the results presented in the review? $\ensuremath{\mathsf{Yes}}$

*Competing Interests:* No competing interests were disclosed.

*Reviewer Expertise:* Family planning, reproductive and maternal health

# I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have

## significant reservations, as outlined above.

# Author Response 10 Jun 2024 victoria boydell

The extent to which contraceptive intention translates into actual contraceptive use has long been a subject of debate, even though studies generally show a positive correlation between intention and behavior. The research question in the present systematic review was straightforward. The authors examined whether contraceptive intentions predict subsequent contraceptive use. The inclusion and exclusion criteria were clearly specified. While all studies included in the review were longitudinal, not all employed an experimental or quasi-experimental design. Some included studies were observational. My comments on the present study are outlined below.

The literature search strategy was comprehensive. Search terms were provided in PROSPERO and included the following: ((intent\* OR intend\*) AND ("to use")) OR (intent\* OR intend\* OR willingness) AND (contracept\* OR "birth control" OR "family planning"). The literature was independently screened by two of the authors to determine the eligibility of studies for inclusion in the systematic review, and full text articles were independently reviewed by the same authors.

While data were independently extracted from the 10 included articles by one author using a predesigned data extraction form, a second author reviewed the full papers and checked the data extraction.

Of 39 articles that were retrieved, 25 articles were excluded after full text screen and 4 articles during data extraction, Unfortunately, the review authors did not fully account for the excluded articles. It would be instructive to know the likely impact of their exclusion on the conclusions of the systematic review.

### **Response:** We have now added in this information.

One consideration was the extent to which the authors described the studies in adequate detail. Although Table 1 does not provide details about research designs, study populations, interventions (if applicable), and study settings, this information is described in the text and gives insights into variations in the study populations and study settings.

## **Response:** Table 1, which was mistakenly excluded, does provide this information.

As all 10 studies included in the present systematic review were longitudinal cohort studies (with pre-and post-tests or treatment and control/comparison groups), the authors used the Joanna Briggs Institute Critical Appraisal Checklist for Cohort Studies to assess risk of bias. One of the included studies had a "low quality" rating and was retained in the systematic review. Could the authors kindly justify its retention? If this study were to be omitted, what would be the implications for the interpretation of the results of the review?

**Response:** We review a low-quality study in the systematic review because it meets all of the a priori inclusion requirements, which is part of the process of systematic reviews. It was post-hoc given a quality rating as part of a typical quality review for a study like this. We are happy to add

this text regarding systematic review processes if it would aid in clarifying this for readers. If this study was removed, the existing findings would remain the same, we would just be removing one study with non-significant findings. It doesn't change the overall interpretation of the review, which is that there is not sufficient evidence to do a meta-analysis of ITU and further research should be conducted that would allow researchers to identify whether this is a successful and potentially more person-centered measure of contraceptive use.

The authors provided a satisfactory discussion of observed heterogeneity in the results of the review. They reported heterogeneity in (a) measures used to report associations or effects in the included studies (odds ratios, hazard ratios, correlation coefficients, simple percentages/proportions); (b) study design (which included non-randomization); (c) analysis (non-adjustment or adjustment for covariates).

In the discussion section, the authors highlighted the importance of adjusting for possible confounding variables, such as parity, relationship status, type of contraceptive method. Contraceptive decision making is also shaped by factors that were not mentioned, including cultural and social norms, knowledge about contraceptive methods, personal beliefs, and access to and supply of contraceptive methods. It is important to mention these factors when discussing the limitations of the study.

**Response:** Thank you for the observation and we have made the change: "In addition, other factors (e.g., cultural and social norms, knowledge about contraceptive methods, personal beliefs) may all contribute to reproductive and contraceptive intentions, decision-making, and subsequent use, and require further consideration."

The preceding comment (i.e., the importance of adjusting for confounding variables) begs the question as to whether the five studies that did not report an adjusted relationship between intent to use (predictor variable) and contraceptive use (outcome variable) should be included in the systematic review. I believe that these studies should not be included as they detract from the robustness of the results.

**Response:** We have included papers that did not report the adjusted relationship between the predictor and the outcome variable to ensure thoroughness in our analysis and avoid introducing bias.

Overall, the present systematic review highlights (a) research gaps, (b) the need for standardized measures of intention to use contraception, and (c) the importance of distinguishing between goal intentions and implementation intentions when predicting subsequent contraceptive use, after adjusting for confounding variables.

Competing Interests: No competing interests were disclosed.

Reviewer Report 16 April 2024

https://doi.org/10.21956/gatesopenres.16413.r36251

© **2024 Boniface E.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



# Emily R Boniface 问

<sup>1</sup> Oregon Health & Science University, Portland, Oregon, USA

<sup>2</sup> Health Systems & Policy, OHSU-PSU School of Public Health, Portland, Oregon, USA

**Overall Comments:** This systematic review attempts to examine how well intention to use (ITU) contraception predicts future contraceptive use. The goal of identifying a more person-centered measure of desire for contraception is an important one. Unfortunately, the small number of studies identified in the analysis and the broad range of associations don't support a claim that ITU is a better measure, and the authors rightly point out the clear need for more research.

# Abstract

- Background: suggest removing the last portion of the final sentence. The study never actually compares ITU's predictive ability to that of unmet need and I'd argue that the results don't allow for a definitive conclusion about how well ITU successfully predicts future use.
- It's a bit confusing to address study populations in the conclusion when they are not mentioned anywhere else in the abstract. Suggest including some mention of them in the results if they are an important part of the conclusion.

# Introduction

- 1<sup>st</sup> paragraph: the connection between understanding desire for contraception and access to a contraceptive program is unclear, as is the last sentence. Is the argument that people shouldn't be classified as having unmet need given that they state they intend to use contraception in the future but are not currently using a method? I would think that would be a more reasonable assumption than categorizing someone as having unmet need who states that they do *not* intend to use a method in the future.
- 3<sup>rd</sup> paragraph: "programme" should be "programmes". Clarify that the scoping review was conducted previously; as currently written, the statement about the scoping review could be interpreted as referring to the current study.
- It would be helpful to clarify why the scoping review included so many more studies than the systematic review
- Suggest considering PMID 36841972 is part of the background literature.

### Methods

- Please include the search terms used to identify papers to facilitate reproducibility
- Curious about the choice to exclude studies that looked at condoms given the fact that they
  were included as a method choice in several of the included studies, and in fact, some of the
  included studies even included "traditional" methods. I don't necessarily have an issue with
  it, but some justification would be appreciated.

### Results

• Data synthesis: the last sentence should be in the discussion rather than methods

- It would be helpful to include details on the reasons/n's for the 1425, 25, and 4 studies that were excluded after applying the inclusion/exclusion criteria at various steps. I'm a little surprised by how few studies were ultimately included and it would be useful to see why others were excluded.
- The text states that study aim, population, location, study design, and follow-up period are included in Table 1, however, this is not the case. Please include columns for each of these variables, as well as sample sizes and titles. It would also be nice to be able to see the definition of methods used in each study. Perhaps dividing the information into 2 tables would be useful.
- Suggest including full list of papers as supplemental material.
- Without being able to see the sample sizes and follow-up periods for each paper, the statement about person-years of data is unclear. I'm assuming the statement implies just a few months of follow-up for almost 5,000 study participants. Is that correct? If not, more clarification is needed.
- Associations section: the distinction between magnitude and strength of association doesn't have a statistical meaning, so it's unclear what is being communicated at the end of this paragraph. Why doesn't the strength of association change after adjustment if there are examples of significant unadjusted OR and non-significant aOR *and* vice versa? Please clarify.

## Discussion

- Appreciate the nuance about goal and implementation intentions. Is that a distinction that was recognized by any of the excluded studies that did not assess future contraception use following report of ITU?
- Another limitation is the variability of geographic settings for the included studies. The analysis seems to assume that the relationship between ITU and subsequent method use is generalizable across settings, which is a fairly strong assumption given the possible differences in health systems and contraceptive access. Could the wide range of ORs be explained by some of these differences?
- It would be helpful to address/compare the results to unmet need. If the motivation for this study is that unmet need is an inadequate measure for predicting contraception use, how does this study compare and what do the results add? It doesn't appear that there is currently enough evidence to support the claim that ITU is a better predictor of future use, so the conclusion seems to overstate it's strength as a predictive measure.

## References

1. Senderowicz L, Bullington BW, Sawadogo N, Tumlinson K, et al.: Assessing the Suitability of Unmet Need as a Proxy for Access to Contraception and Desire to Use It.*Stud Fam Plann*. 2023; **54** (1): 231-250 PubMed Abstract | Publisher Full Text

### Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

# Are sufficient details of the methods and analysis provided to allow replication by others? Partly

### Is the statistical analysis and its interpretation appropriate?

#### Yes

Are the conclusions drawn adequately supported by the results presented in the review? Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Biostatistics, contraception use, person-centered contraceptive care

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

#### Author Response 10 Jun 2024

#### victoria boydell

**Overall Comments:** This systematic review attempts to examine how well intention to use (ITU) contraception predicts future contraceptive use. The goal of identifying a more person-centered measure of desire for contraception is an important one. Unfortunately, the small number of studies identified in the analysis and the broad range of associations don't support a claim that ITU is a better measure, and the authors rightly point out the clear need for more research.

### Abstract.

Background: suggest removing the last portion of the final sentence. The study never actually compares ITU's predictive ability to that of unmet need and I'd argue that the results don't allow for a definitive conclusion about how well ITU successfully predicts future use.

**Response:** We have changed the wording to be more appropriate to the content of the paper and it now reads: "This systematic review examines the relationship between relationship between intentions to use and actual use of contraception and could potentially in developing responsive programs."

It's a bit confusing to address study populations in the conclusion when they are not mentioned anywhere else in the abstract. Suggest including some mention of them in the results if they are an important part of the conclusion.

**Response:** Noted and we have now included a mention of population in the conclusion of the abstract, which reads "The range of possible confounding factors, particularly around the different populations, points to the need for more research so that a meta-analysis can be done in the future."

## Introduction

1<sup>st</sup> paragraph: the connection between understanding desire for contraception and access to a contraceptive program is unclear, as is the last sentence. Is the argument that people shouldn't be classified as having unmet need given that they state they intend to use

contraception in the future but are not currently using a method? I would think that would be a more reasonable assumption than categorizing someone as having unmet need who states that they do *not* intend to use a method in the future.

**Response:** Thank you for the observation and we have made the change and we have removed this statement to prevent confusion.

3<sup>rd</sup> paragraph: "programme" should be "programmes".

Response: Change has been made.

Clarify that the scoping review was conducted previously; as currently written, the statement about the scoping review could be interpreted as referring to the current study.

**Response:** Thank you for the observation and we have made the change to distinguish this and the earlier review: "The scoping review included a wider range of evidence and identified 112 papers and their operationalizations of ITU; here we build off of that work to examine a subset of the studies where the data collection design and reporting was sufficient to be able to assess whether ongoing and continued measurement of ITU has the potential to accurately predict subsequent contraceptive use for those who desire it."

It would be helpful to clarify why the scoping review included so many more studies than the systematic review.

**Response:** Thank you for the observation and we have made the change: "The scoping review included a wider range of evidence and identified 112 papers and their operationalizations of ITU; here we build off of that work to examine a subset of the studies where the data collection design and reporting was sufficient to be able to assess whether ongoing and continued measurement of ITU has the potential to accurately predict subsequent contraceptive use for those who desire it."

Suggest considering PMID 36841972 is part of the background literature.

**Response:** Thank you for the observation and we have made the change. This has been included in the references and citations.

### Methods

Please include the search terms used to identify papers to facilitate reproducibility

**Response:** Thank you, we already direct the readers to the protocol – explicitly stating this is where they can find the search terms.

Curious about the choice to exclude studies that looked at condoms given the fact that they were included as a method choice in several of the included studies, and in fact, some of the included studies even included "traditional" methods. I don't necessarily have an issue with

it, but some justification would be appreciated.

**Response:** We have removed this phrase to avoid confusion. We excluded condoms because they do not require the same type of premeditation and planning to use them as a contraception.

## Results

Data synthesis: the last sentence should be in the discussion rather than methods

**Response:** Change has been made.

It would be helpful to include details on the reasons/n's for the 1425, 25, and 4 studies that were excluded after applying the inclusion/exclusion criteria at various steps. I'm a little surprised by how few studies were ultimately included and it would be useful to see why others were excluded.

**Response:** This is now included and it states: Many papers were excluded because they did not have a clear definition of intention to use (732), did not state an association between intention to use and contraceptive use (235), did not meet the study design requirements (238), did not contain sufficient information in the text to be assessed against the inclusion criteria (30), focused on condoms (161), did not include a measure of contraceptive use (61) or focused on only on the drivers of intention to use and did not test the association with actual use (17).

The text states that study aim, population, location, study design, and follow-up period are included in Table 1, however, this is not the case. Please include columns for each of these variables, as well as sample sizes and titles. It would also be nice to be able to see the definition of methods used in each study. Perhaps dividing the information into 2 tables would be useful.

**Response:** We have this Table, but we see that it was not included in the paper, only Table 2 was included. Apologies and we will rectify this with the production team.

Suggest including full list of papers as supplemental material.

**Response:** This information is included in Table 1.

Without being able to see the sample sizes and follow-up periods for each paper, the statement about person-years of data is unclear. I'm assuming the statement implies just a few months of follow-up for almost 5,000 study participants. Is that correct? If not, more clarification is needed.

**Response:** We have this information in Table, but I see that it was not included in the paper, only Table 2 was included. Apologies and we will rectify this. Yes, for most data included the follow-up periods were short.

Associations section: the distinction between magnitude and strength of association doesn't

have a statistical meaning, so it's unclear what is being communicated at the end of this paragraph. Why doesn't the strength of association change after adjustment if there are examples of significant unadjusted OR and non-significant aOR *and* vice versa? Please clarify.

**Response:** Thank you for this comment but we have not changed the text because the magnitude of a coefficient, here odds ratios, indicates the direction and degree of the relationship between the predictor and outcome variable, while the strength of association indicates how precisely the coefficient is measured. What is being said here is that, as expected, when one adds more variables to the model (adjusted ORs), the magnitude of the relationship between the predictor and outcome, or independent and dependent, variable is attenuated towards zero. This is expected, as the addition of new variables typically explains additional portions of the variance in the outcome. A reduction in the statistical significance, or strength of association, indicates that adding the new variable(s) has diluted the [magnitude] of the original association between the predictor and outcome variable, so that the estimate has become less precise. So, to answer the first part of the question, what is being communicated is that there is some relationship between the added variables and intention to use, thus reducing the magnitude of the coefficients, however these coefficients continue to be very precisely estimated and are significant in the presence of effect modifiers. For the second part of the question 'Why doesn't the strength of the association change...', this is a statistical question, and the answer is that the coefficient continues to be as or close to precisely estimated in adjusted models as it is in unadjusted models.

# Discussion

Appreciate the nuance about goal and implementation intentions. Is hat a distinction that was recognized by any of the excluded studies that did not assess future contraception use following report of ITU?

**Response:** Thank you for the observation and we have made the change: "None of the papers included distinguished between goal and implementation intentions."

Another limitation is the variability of geographic settings for the included studies. The analysis seems to assume that the relationship between ITU and subsequent method use is generalizable across settings, which is a fairly strong assumption given the possible differences in health systems and contraceptive access. Could the wide range of ORs be explained by some of these differences?

**Response:** Thank you for the observation and we have made the change: "Geographic settings, particularly the difference in health systems and contraceptive access, may also explain the differences we found."

It would be helpful to address/compare the results to unmet need. If the motivation for this study is that unmet need is an inadequate measure for predicting contraception use, how does this study compare and what do the results add? It doesn't appear that there is currently enough evidence to support the claim that ITU is a better predictor of future use, so the conclusion seems to overstate it's strength as a predictive measure.

**Response:** Thank you for pointing this out. We have removed the word 'alternative' from the conclusion as we do not draw a comparison.

*Competing Interests:* No competing interests were disclosed.

Author Response 13 Feb 2025

## victoria boydell

Thank you so much for your feedback on version 2 of the paper. We have reviewed all your comments and have incorporated all the suggested changes. We appreciate the input and think it will make for a much-improved paper.

*Competing Interests:* No competing interests were disclosed.