

Allopachria nigrocatta sp. nov. from Guangxi, with a key and checklist of Chinese species and additional records of *Allopachria* Zimmermann, 1924 from China (Coleoptera, Dytiscidae, Hydroporinae, Hyphydrini)

Zhuo-yin Jiang^{1,2} , Ya-ru Chen¹, Li-bo Han¹, Feng-long Jia² , Bei-xin Wang¹ 

1 Department of Entomology, College of Plant Protection, Nanjing Agricultural University, Nanjing 210095, China

2 Department of Entomology, College of Life Sciences, Sun Yat-sen University, Guangzhou 510275, China

Corresponding authors: Feng-long Jia (lssjfl@mail.sysu.edu.cn, fenglongjia@aliyun.com); Bei-xin Wang (wangbeixin@njau.edu.cn)

Abstract

A new species *Allopachria nigrocatta* sp. nov. from Guangxi, China is described. New records are provided for the following *Allopachria* species: *A. dudgeoni* Wewalka, 2000 and *A. weinbergeri* Wewalka, 2000 from Zhejiang, and *A. yiae* Bian, Guo & Ji, 2013 from Guizhou. Additional records are also given for some other Chinese species. The habitus and male genitalia of the new species are illustrated. An updated key and a checklist of all Chinese species of *Allopachria* are provided.



Academic editor: Mariano Michat

Received: 16 September 2024

Accepted: 30 October 2024

Published: 4 December 2024

ZooBank: <https://zoobank.org/B970895F-E1CE-4F42-81FD-1F03B7DE5680>

Citation: Jiang Z-yin, Chen Y-ru, Han L-bo, Jia F-long, Wang B-xin (2024) *Allopachria nigrocatta* sp. nov. from Guangxi, with a key and checklist of Chinese species and additional records of *Allopachria* Zimmermann, 1924 from China (Coleoptera, Dytiscidae, Hydroporinae, Hyphydrini). ZooKeys 1219: 287–302. <https://doi.org/10.3897/zookeys.1219.137179>

Key words: *Allopachria*, checklist, China, Dytiscidae, fauna, Hydroporinae, identification key, new record, new species, Oriental region, taxonomy

Introduction

Allopachria Zimmermann, 1924, are small to extremely small beetles, most of which bear maculae on their elytra. Male specimens of some species have antennomeres, protibiae or protarsomeres modified in various ways (Wewalka 2000). Species of the genus occur mainly in the Oriental region, from Indonesia in the south to southern China and Japan in the north (Wewalka 2010). *Allopachria* typically live by the margin of flowing waters, usually with a sand or stone bottom. The knowledge of *Allopachria* was very poor until Wewalka (2000) made a comprehensive revision of the genus. Then, the studies increased gradually, and the fauna of Chinese *Allopachria* was also supplemented (Bian and Ji 2010; Wewalka 2010; Bian and Ji 2012; Bian et al. 2013; Ji et al. 2014). Up to now, the genus contains 47 species worldwide (Nilsson and Hájek 2024a), of which 29 species occur in China (Nilsson and Hájek 2024b).

In the present study, we describe a new species from Guangxi, China and provide additional records for some Chinese *Allopachria* species. For the first time, an updated key to all known species from China is provided.

Copyright: © Zhuo-yin Jiang et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Materials and methods

Specimens were examined and measured under a Nikon SMZ800N stereomicroscope. Some of the specimens were dissected, and the genitalia were put into 10% KOH at room temperature for 8–10 h, then placed in a drop of glycerol on glass slides for photographing. Photographs of habitus and characters were taken with a Nikon DS-Ri2 mounted on a Nikon SMZ25; layers were captured and aligned in the NIS-Elements software. Photographs of genitalia were taken with a Zeiss AxioCam HRc mounted on a Zeiss AX10 microscope with the Axio Vision SE64 software, then stacked in Helicon Focus (ver. 7.0.2). After being photographed, the genitalia were transferred to a transparent plastic plate in a drop of glycerol and attached to the respective specimen. The images were edited and assembled with Adobe Photoshop CS6.

The following abbreviations were used in the descriptions: TL, total length, measurement of length from clypeal margin to apex of elytra; TL-h, total length minus head length, measurement of length from anterior margin of pronotum to apex of elytra; MW, maximum width of body measured at right angle to TL. The terminology follows Wewalka (2000) and Wewalka (2010), and the style of description of the new species follows Jiang et al. (2022). The terminology to denote the orientation of the genitalia follows Miller and Nilsson (2003). Exact label data are cited for the type material and given in quotation marks. Authors' additional remarks are provided in square brackets; [p]—preceding data are printed. Separate label lines are indicated by a slash (/), and separate labels by a double slash (//).

The specimens included in this study are deposited in the following collections:

- SYSU Biological Museum, Sun Yat-sen University, Guangzhou, China.
ZJCQ Zhuo-yin Jiang collection, Quzhou, China.

Taxonomy

Allopachria nigrocatta Jiang & Jia, sp. nov.

<https://zoobank.org/0D13517E-1D9A-4171-AFB3-CD19D5880446>

Figs 1A–E, 2A–C, 4A, 6, 7C

Type locality. China, Guangxi Zhuang Autonomous Region, Guilin, Ziyuan County, Maoershan Mt., Hui long Temple, c. 25.9125°N, 110.4656°E; 1557.2 m.

Type material. Holotype • male (SYSU), labelled: “广西桂林市资源县 / 猫儿山迴龙寺 / 25.9125N, 110.4656E / 1557.2m, 30.viii.2020 / 姜卓寅 [p] // CHINA: Guangxi, Guilin / Ziyuan County, Maoershan Mt. / Hui long Temple / 25.9125N, 110.4656E / 1557.2m, 30.viii.2020 / Zhuoyin Jiang leg. [p] // HOLOTYPE / ALLOPACHRIA / nigrocatta sp. nov. / Jiang & Jia det. 2024 [red label, p]”. **Paratype** • 1 female (SYSU), same label data as holotype, with a paratype label, labelled: “PARATYPE / ALLOPACHRIA / nigrocatta sp. nov. / Jiang & Jia det. 2024 [red label, p]”.

Description of male holotype. Habitus (Fig. 1A) regularly oval, with continuous outline, broadest in 1/3 of elytral length, moderately convex. **Colouration.** Head dark brown to black, somewhat paler at clypeal margin, with small yellowish-brown area behind eyes; pronotum black with lateral margins reddish-brown;




Figure 1. *Allopachria nigrocatta* sp. nov. (Maoershan, Guangxi) **A** habitus in dorsal view (holotype, male) **B** habitus in ventral view (holotype, male) **C** habitus in dorsal view (paratype, female) **D** male protarsus in dorsal view **E** male mesotarsus in dorsal view. Scale bars: 1.0 mm (**A-C**); 0.5 mm (**D, E**).

elytra black, apex with an irregularly shaped yellowish-brown spot not reaching suture; appendages yellowish-brown to reddish-brown; ventral side reddish-brown to black. **Head.** Moderately broad, c. 0.67 × width of pronotum, trapezoidal. Anterior margin of clypeus regularly rounded, without bead (Fig. 7C). Antenna with antennomeres long and slender. Shiny, microreticulation presents on anterior half of head and along eyes, consisting of well-impressed polygonal isodiametric meshes. Punctures spread sparsely and more distinct on vertex;




Figure 2. Male genitalia of *Allopachria nigrocatta* sp. nov. (holotype, male; Maoershan, Guangxi) **A** median lobe in ventral view **B** median lobe in lateral view **C** right paramere in lateral view. Scale bar: 0.1 mm.

setigerous punctures present along inner margin of eyes and anterolaterally to eyes in fronto-clypeal depressions. **Pronotum.** Strongly transverse (width/length ratio = 2.71), broadest between posterior angles. Lateral margins moderately curved, distinctly beaded. Shiny, microreticulation absent. Punctures and micropunctures spread sparsely and evenly; rows of setigerous punctures present along anterior margin; navel-like punctures present in posterior half. Posterior half also covered with some irregular longitudinal wrinkles. **Elytra.** Base as broad as pronotal base; lateral margins moderately curved. Shiny, microreticulation absent. Punctures and micropunctures spread sparsely and evenly; longitudinal rows of setigerous punctures incomplete. **Legs.** Protarsomere 1 minimally enlarged (Fig. 1D), mesotarsomere 1 distinctly enlarged (Fig. 1E), with adhesive setae on their ventral side; claws simple, metatarsal claws unequal. **Ventral side** (Fig. 1B). Prosternum sinuate anteriorly. Prosternal process heart-shaped, with distinct wide lateral beads in basal two thirds, apex obtuse; surface distinctly punctured. Metaventrite distinctly beaded on anterior margin; lateral parts of metaventrite (“metasternal wings”) narrow. Metacoxal lines prominent,

divergent anteriorly. Metacoxal plates with some indistinct wrinkles. Metacoxal processes with a small triangular lobe, obscuring part of metatrochanter. Abdomen with five ventrites (III–IV fused). Microreticulation presents on abdominal ventrites, consisting of well-impressed polygonal isodiametric meshes. Punctures and micropunctures spread sparsely and evenly on metaventrete, metacoxae and abdominal ventrites; setigerous punctures present along anterior margin of metaventrete and medially on ventrites II–IV. **Male genitalia.** Median lobe of aedeagus gradually narrowing from base to apex in ventral view, apex truncate (Fig. 2A); ‘L-shaped in lateral view, moderately curved, apex obtuse (Fig. 2B). Parameres moderately broad, distal portion with a large triangular process on dorsal side, with a tuft of setae apically (Fig. 2C).

Female (Fig. 1C). Identical to male in habitus. Dorsal surface submatt, microreticulation presents on head, pronotum and elytra, more densely and longitudinally stretched on elytra. Pro- and mesotarsomeres not enlarged, without adhesive setae.

Variability. Minor variability can be seen in dorsal colouration; there is an indistinct irregular reddish-brown spot on each elytral base in the paratype.

Measurements. TL: 2.5–2.7 mm (mean value: 2.6 ± 0.1 mm); holotype: 2.7 mm. TL-h: 2.2–2.4 mm (mean value: 2.3 ± 0.1 mm); holotype: 2.4 mm. MW: 1.7–1.8 mm (mean value: 1.75 ± 0.05 mm); holotype: 1.8 mm.

Differential diagnosis. *Allopachria nigrocatta* sp. nov. can be distinguished from other *Allopachria* species by its black colour pattern with a yellowish-brown spot on elytral apex. The shape of the median lobe and paramere is also unique. Female specimens can be easily identified by the submatt appearance.

Collection details. On Maoershan Mt., *Allopachria nigrocatta* sp. nov. was collected syntopically with *Platambus punctatipennis* Brancucci, 1984 from a pool with decaying leaves under a rock wall covered with flowing water (Fig. 4A).

Etymology. The species name is a combination of “*nigro-*” from Latin “*nigriculus*” (= blackish) and “-catta” (= cat, in Chinese “mao” means cat), referring to its black appearance and the fact that it was collected from Maoershan Mt. The gender of the name is feminine.

Distribution. So far, known only from the type locality on Maoershan Mt., northern Guangxi, China (Fig. 6).

Additional records of Chinese *Allopachria*

Allopachria biana Wewalka, 2010

Figs 3A, 4B, 6

Allopachria biana Wewalka, 2010: 31 (orig. descr.).

Material studied. CHINA: Guangdong • 11 spec. (SYSU, ZJCQ), Zhaoqing, Dinghu Distr., Dinghushan Mt., Qingyunsi management station, 23.1737°N, 112.5361°E, 241.2 m, 12.vii.2021, Z.Y. Jiang, Z.Q. Mai & Z.L. Liang leg. • 5 spec. (SYSU, ZJCQ), Zhaoqing, Dinghu Distr., Dinghushan Mt., Qingyunsi management station, 23.1732°N, 112.5354°E, 256.9 m, 17.x.2021, Z.Y. Jiang, Z.Q. Mai & W.C. Xie leg. • 5 spec. (SYSU, ZJCQ), Zhaoqing, Dinghu Distr., Dinghushan Mt., Dizhi management station, 23.1609°N, 112.5323°E, 88.7 m, 14.ix.2022, Z.Y. Jiang & W.C. Xie leg. • 18 spec. (SYSU, ZJCQ), Zhaoqing, Dinghu Distr., Ding-




Figure 3. Habitus of *Allopachria* species **A** *Allopachria bianae* (male; Dinghushan, Guangdong) **B** *Allopachria dudgeoni* (male; Tianmushan, Zhejiang) **C** *Allopachria jendeki* (male; Baihualing, Yunnan) **D** *Allopachria miaowangi* (male; Lingwei, Zhejiang) **E** *Allopachria schoenmanni* (male; Tianmushan, Zhejiang) **F** *Allopachria weinbergeri* (male; Lingwei, Zhejiang) **G** *Allopachria yiae* (male; Machang Town, Guizhou). Scale bar: 1.0 mm.




Figure 4. Habitat of *Allopachria* species **A** pool with decaying leaves under rock wall covered with flowing water, Maoershian, Guangxi (*A. nigrocatta* sp. nov.) **B** stream pool with decaying leaves, Dinghushan, Guangdong (*A. biana*) **C** small stream with decaying leaves, Baihualing, Yunnan (*A. jendeiki*) **D** small pool with flowing water surrounded by large stones, Tianmushan, Zhejiang (*A. schoenmanni*).

hushan Mt., Qingyunsi management station, 23.1732°N, 112.5354°E, 256.9 m, 15.ix.2022, Z.Y. Jiang & W.C. Xie leg.

Distribution. So far known only from its type locality on Dinghushan Mt. (Guangdong) (Fig. 6).

Collection details. On Dinghushan Mt., *Allopachria bianae* was collected syntopically with *Neptosternus coomani* Peschet, 1923 and *Neptosternus punctatus* Zhao, Hájek, Jia & Pang, 2012 among decaying leaves in a stream pool (Fig. 4B).

***Allopachria dudgeoni* Wewalka, 2000**

Figs 3B, 6, 7E

Allopachria dudgeoni Wewalka, 2000: 117 (orig. descr.); Wewalka 2010: 36 (fauna), Bian and Ji 2010: 64 (fauna), Bian and Ji 2012: 31 (fauna).

Material studied. CHINA: Zhejiang • 1 male (ZJCQ), Hangzhou, Lin'an Distr., Tianmushan Mt., Xiguan Stream S02, 30.3579°N, 119.4487°E, 609 m, x.2023, H.M. Zang & L.B. Han leg. • 1 female (ZJCQ), Hangzhou, Lin'an Distr., Tianmushan Mt., Waijunling, 30.3750°N, 119.4854°E, 537 m, xii.2023, H.M. Zang & L.B. Han leg.

Distribution. Southeastern China (Guangdong, Guangxi, Hong Kong, Jiangxi, Zhejiang). First record from Zhejiang Province (Fig. 6).

***Allopachria jendeki* Wewalka, 2000**

Figs 3C, 4C, 6, 7A

Allopachria jendeki Wewalka, 2000: 116 (orig. descr.); Wewalka 2010: 36 (fauna).

Material studied. CHINA: Yunnan • 1 male, 2 females (SYSU, ZJCQ), Baoshan, Baihualing Mt., Zaotang Stream, 25.3087°N, 98.7936°E, 1481.4 m, 11.v.2021, Z.Y. Jiang, Z.M. Yang, Z.Q. Mai & B.P. Huang leg.

Distribution. Southwestern China (Yunnan) (Fig. 6).

Collection details. The specimens were collected among decaying leaves and sand in a small stream on Baihualing Mt. (Fig. 4C).

***Allopachria miaowangi* Wewalka, 2010**

Figs 3D, 5A, B, 5Ca, 6, 7F

Allopachria miaowangi Wewalka, 2010: 29 (orig. descr.); Bian and Ji 2012: 34 (fauna).

Allopachria dieterlei Wewalka, 2000: Bian and Ji 2010: 64 (misidentification).

Material studied. CHINA: Zhejiang • 4 males, 1 female (SYSU, ZJCQ), Quzhou, Kecheng Distr., Lingwei, 28.8507°N, 118.9372°E, 177.4 m, 2.v.2024, Z.Y. Jiang & Z.X. Mao leg. • 1 female (ZJCQ), Quzhou, Kecheng Distr., Lingwei, 28.8507°N, 118.9372°E, 177.4 m, 13.iv.2020, Z.Y. Jiang & Z.X. Mao leg.



Figure 5. Habitat and live specimens of *Allopachria miaowangi* and *Allopachria weinbergeri* **A** small stream under the forest (Lingwei, Zhejiang) **B** sand and stones by the margin of the stream (Lingwei, Zhejiang) **C** live specimens of *Allopachria* (**a** *A. miaowangi* **b** *A. weinbergeri*).

Distribution. Eastern China (Hunan, Jiangxi, Zhejiang) (Fig. 6).

Collection details. *Allopachria miaowangi* was collected syntopically with *Allopachria weinbergeri* under sand and stones by the margin of a small stream in Lingwei (Zhejiang) (Fig. 5A, B).




Figure 6. Map of distribution of *Allopachria nigrocatta* sp. nov. and additional records of other *Allopachria* species in China.

***Allopachria schoenmanni* Wewalka, 2000**

Figs 3E, 4D, 6

Allopachria schoenmanni Wewalka, 2000: 113 (orig. descr.).

Material studied. CHINA: Zhejiang • 6 males, 2 females (SYSU, ZJCQ), Hangzhou, Lin'an Distr., Tianmushan Mt., Y437 roadside (Xiguan), 30.3483°N, 119.4509°E, 484.5 m, 20.xi.2023, Z.Y. Jiang leg. • 1 female (ZJCQ), Hangzhou, Lin'an Distr., Tianmushan Mt., Shuiduitang, 30.3682°N, 119.4255°E, 867 m, viii.2023, H.M. Zang & L.B. Han leg.

Distribution. Eastern China (Anhui, Zhejiang) (Fig. 6).

Collection details. Most specimens from Tianmushan Mt. were collected in a small pool with flowing water surrounded by large stones (Fig. 4D).

***Allopachria weinbergeri* Wewalka, 2000**

Figs 3F, 5A, B, 5Cb, 6, 7G

Allopachria weinbergeri Wewalka, 2000: 112 (orig. descr.); Wewalka 2010: 36 (fauna). *Allopachria weinbergerorum* Nilsson, 2007: 50 (as unjustified emendation of *weinbergeri*).

Material studied. CHINA: Zhejiang • 4 males, 2 females (SYSU, ZJCQ), Quzhou, Kecheng Distr., Lingwei, 28.8507°N, 118.9372°E, 177.4 m, 2.v.2024, Z.Y. Jiang & Z.X. Mao leg. • 1 female (ZJCQ), Quzhou, Kecheng Distr., Lingwei, 28.8507°N, 118.9372°E, 177.4 m, 13.iv.2020, Z.Y. Jiang & Z.X. Mao leg.




Figure 7. Morphological characters of *Allopachria* spp. A–C anterior aspect of head showing clypeus D–G male protarsus in dorsal view. A *Allopachria jendeki* B, D *Allopachria yiae* C *Allopachria nigrocatta* sp. nov. E *Allopachria dudgeoni* F *Allopachria miaowangi* G *Allopachria weinbergeri*. Scale bars: 0.5 mm.

Distribution. Eastern China (Fujian, Guangdong, Zhejiang). First record from Zhejiang Province (Fig. 6).

Collection details. *Allopachria weinbergeri* was collected syntopically with *Allopachria miaowangi* under sand and stones by the margin of a small stream in Lingwei (Zhejiang) (Fig. 5A, B).

***Allopachria yiae* Bian, Guo & Ji, 2013**

Figs 3G, 6, 7B, 7D

Allopachria yiae Bian, Guo & Ji, 2013: 85 (orig. descr.).

Material studied. CHINA: Guizhou • 3 males, 2 females (SYSU), Pingba, Machang Town, 13.viii.1982, Z.H. Huang leg.

Distribution. Southwestern China (Chongqing, Guizhou). First record from Guizhou Province (Fig. 6).

Key to Chinese *Allopachria* species (appropriate to males only)

- | | | |
|----|---|---------------------------------------|
| 1 | Anterior margin of clypeus with bead (Fig. 7A, B)..... | 2 |
| - | Anterior margin of clypeus without bead (Fig. 7C) | 10 |
| 2 | Anterior margin of clypeus with distinct bead (Fig. 7A) | 3 |
| - | Anterior margin of clypeus with very fine bead (Fig. 7B) | 7 |
| 3 | Elytra completely black, without spots (Fig. 3C)..... | |
| | <i>A. jendeiki</i> Wewalka, 2000 | |
| - | Elytra with two or three yellow to reddish-brown spots | 4 |
| 4 | Lateral margin of elytron with a longitudinal furrow | |
| | <i>A. manfredi</i> Wewalka, 2010 | |
| - | Lateral margin of elytron without a longitudinal furrow..... | 5 |
| 5 | Male antennae modified, antennomeres 4 and 5 distinctly enlarged | |
| | <i>A. scholzi</i> Wewalka, 2000 | |
| - | Male antennae not modified, with antennomeres long and slender..... | 6 |
| 6 | Size larger (TL: 2.1–2.2 mm); head yellow; elytra with three spots (see Wewalka 2000, fig. 25)..... | <i>A. ernsti</i> Wewalka, 2000 |
| - | Size smaller (TL: 1.7–1.8 mm); head black; elytra with two spots (see Wewalka 2000, fig. 8)..... | <i>A. taiwana</i> (Satô, 1990) |
| 7 | Size larger (TL: 2.7–2.9 mm); male protarsomere 1 distinctly enlarged with indentation on inner margin at basal half (Fig. 7D)..... | |
| | <i>A. yiae</i> Bian, Guo & Ji, 2013 | |
| - | Size smaller (TL: 1.5–1.8 mm); male protarsomere 1 minimally enlarged... | 8 |
| 8 | Male antennomere 5 distinctly enlarged, 'L'-shaped (see Wewalka 2000, fig. 107) | <i>A. froehlichi</i> Wewalka, 2000 |
| - | Male antennomere 5 moderately enlarged, with an obtuse tooth (see Wewalka 2000, fig. 108)..... | 9 |
| 9 | Median lobe of aedeagus in lateral view shorter and broader (see Wewalka 2000, fig. 44b) | <i>A. hautmanni</i> Wewalka, 2000 |
| - | Median lobe of aedeagus in lateral view longer and slender (see Wewalka 2000, fig. 43b)..... | <i>A. schillhammeri</i> Wewalka, 2000 |
| 10 | Size smaller (TL: 1.6 mm)..... | <i>A. jilanzhui</i> Wewalka, 2000 |
| - | Size larger (TL: ≥ 2.0 mm)..... | 11 |
| 11 | Apex of median lobe of aedeagus trifid..... | 12 |
| - | Apex of median lobe of aedeagus simple, not trifid | 16 |
| 12 | Elytra entirely microreticulate..... | 13 |
| - | Elytra without microreticulation | 14 |

- 13 Punctuation of pronotum regular in size, not navel-like; interstice of the tridentate apex of median lobe narrower (see Wewalka 2000, fig. 63a).....
..... ***A. wangii* Wewalka & Nilsson, 1994**
- Punctuation of pronotum irregular in size, partly coarse and navel-like; interstice of the tridentate apex of median lobe wider (see Wewalka 2000, fig. 64a)
..... ***A. jaechi* Wewalka, 2000**
- 14 Male protarsomere 1 not modified, minimally enlarged.....
..... ***A. flavomaculata* (Kamiya, 1838)**
- Male protarsomere 1 modified, distinctly enlarged
..... **15**
- 15 Elytra with two or three spots (see Wewalka 2010, fig. 7a, b); middle part of the tridentate apex of median lobe longer than lateral parts (see Wewalka 2010, fig. 15a).....
..... ***A. komareki* Wewalka, 2010**
- Elytra with two spots (see Bian and Ji 2010, fig. 1); middle part and lateral parts of the tridentate apex of median lobe equal in length (see Bian and Ji 2010, fig. 3)
..... ***A. wuzhifengensis* Bian & Ji, 2010**
- 16 Male antennomeres 4 and 5 distinctly enlarged (see Wewalka 2000, fig. 113).....
..... ***A. friedrichi* Wewalka, 2000**
- Male antenna not or minimally enlarged
..... **17**
- 17 Male protarsomere 1 modified, distinctly enlarged (Fig. 7E, F).....
..... **18**
- Male protarsomere 1 not modified, minimally enlarged (Fig. 7G).....
..... **22**
- 18 Male protarsomere 1 elongate (Fig. 7E)
..... **19**
- Male protarsomere 1 short (Fig. 7F)
..... **20**
- 19 Male protarsomere 1 about 1.5 times as long as protarsomere 2, insertion of protarsomere 2 at the apical part of protarsomere 1 (Fig. 7E).....
..... ***A. dudgeoni* Wewalka, 2000**
- Male protarsomere 1 about 2 times as long as protarsomere 2, insertion of protarsomere 2 at the middle of protarsomere 1 (see Bian and Ji 2012, figs 15, 17)
..... ***A. elongata* Bian & Ji, 2012**
- 20 Apex of median lobe of aedeagus pointed in ventral view (see Wewalka 2010, fig. 11a).....
..... ***A. miaowangi* Wewalka, 2010**
- Apex of median lobe of aedeagus truncate in ventral view (see Wewalka 2010, fig. 12a)
..... **21**
- 21 Punctuation of elytra stronger; median lobe of aedeagus in lateral view moderately curved (see Wewalka 2000, fig. 58b)
..... ***A. dieterlei* Wewalka, 2000**
- Punctuation of elytra finer; median lobe of aedeagus in lateral view almost straight at middle part (see Wewalka 2010, fig. 12b)
..... ***A. guangdongensis* Wewalka, 2010**
- 22 Elytra often completely black, sometimes with two reddish-brown spots (see Wewalka 2010, fig. 6a, b).....
..... ***A. hajeki* Wewalka, 2010**
- Elytra with one or two yellowish-brown to reddish-brown spots
..... **23**
- 23 Size larger (TL: 2.5–2.7 mm); male mesotarsomere 1 distinctly enlarged (Fig. 1E); elytra often with one spot at apex (Fig. 1A)
..... ***A. nigrocatta* sp. nov.**
- Size smaller (TL: 2.0–2.4 mm); male mesotarsomere 1 not or minimally enlarged; elytra with two spots: one at base and one at apex
..... **24**
- 24 Apex of median lobe of aedeagus rounded or truncate in ventral view ..
..... **25**
- Apex of median lobe of aedeagus pointed in ventral view
..... **27**

- 25 Apex of median lobe of aedeagus rounded in ventral view (see Wewalka 2000, fig. 51a).....*A. liselotteae* Wewalka, 2000
– Apex of median lobe of aedeagus truncate in ventral view.....26
26 Median lobe of aedeagus in ventral view distinctly narrowed behind apical fourth (see Wewalka 2010, fig. 13a)*A. bianaee* Wewalka, 2010
– Median lobe of aedeagus in ventral view with sides almost parallel over entire length (see Ji et al. 2014, fig. 3)
.....*A. longyanensis* Ji, Guo & Bian, 2014
27 Median lobe of aedeagus in ventral view broadened in middle, apex slender (see Bian and Ji 2012, fig. 10).....*A. yanfengi* Bian & Ji, 2012
– Median lobe of aedeagus in ventral view narrowed in middle, apex slightly broad28
28 Median lobe of aedeagus in ventral view slightly narrowed in middle (see Wewalka 2000, fig. 50a).....*A. schoenmanni* Wewalka, 2000
– Median lobe of aedeagus in ventral view distinctly narrowed in middle.29
29 Median lobe of aedeagus in ventral view broader at apex (see Bian and Ji 2010, fig. 6)*A. grandis* Bian & Ji, 2010
– Median lobe of aedeagus in ventral view narrower at apex (see Wewalka 2000, fig. 49a).....*A. weinbergeri* Wewalka, 2000

List of Chinese *Allopachria* Zimmermann species, including their distribution

- Allopachria bianaee* Wewalka, 2010 Guangdong
Allopachria dieterlei Wewalka, 2000 Hunan
Allopachria dudgeoni Wewalka, 2000 Guangdong, Guangxi, Hong Kong, Jiangxi, Zhejiang
Allopachria elongata Bian & Ji, 2012 Guangxi
Allopachria ernsti Wewalka, 2000 Guangxi; North Vietnam
Allopachria flavomaculata (Kamiya, 1838) Guangxi; Japan
Allopachria friedrichi Wewalka, 2000 Hunan
Allopachria froehlichi Wewalka, 2000 Hong Kong
Allopachria grandis Bian & Ji, 2010 Jiangxi
Allopachria guangdongensis Wewalka, 2010 Guangdong, Guangxi
Allopachria hajeki Wewalka, 2010 Yunnan
Allopachria hautmanni Wewalka, 2000 Anhui
Allopachria jaechi Wewalka, 2000 Hainan
Allopachria jendeki Wewalka, 2000 Yunnan
Allopachria jilanzhui Wewalka, 2000 Guangxi, Hunan
Allopachria komareki Wewalka, 2010 Guangdong
Allopachria liselotteae Wewalka, 2000 Guangxi
Allopachria longyanensis Ji, Guo & Bian, 2014 Fujian
Allopachria manfredi Wewalka, 2010 Guangdong
Allopachria miaowangi Wewalka, 2010 Hunan, Jiangxi, Zhejiang
Allopachria nigrocatta sp. nov. Guangxi
Allopachria schillhammeri Wewalka, 2000 Hunan
Allopachria schoenmanni Wewalka, 2000 Anhui, Zhejiang
Allopachria scholzi Wewalka, 2000 Yunnan
Allopachria taiwana (Satô, 1990) Taiwan

Allopachria wangii Wewalka & Nilsson, 1994 Taiwan
Allopachria weinbergeri Wewalka, 2000 Fujian, Guangdong, Zhejiang
Allopachria wuzhifengensis Bian & Ji, 2010 Hunan, Jiangxi
Allopachria yanfengi Bian & Ji, 2012 Guangxi
Allopachria yiae Bian, Guo & Ji, 2013 Chongqing, Guizhou

Acknowledgements

We thank Dr Bing-lan Zhang (SYSU) for guiding us in the use of photography equipment.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was supported by GDAS Special Project of Science and Technology Development (2020GDSYL-20200102021, 2020GDASYL-20200301003).

Author contributions

Writing manuscript: Zhuo-yin Jiang, Feng-long Jia. Taking photos and making plates: Zhuo-yin Jiang, Ya-ru Chen, Li-bo Han. Reviewing manuscript: Zhuo-yin Jiang, Bei-xin Wang.

Author ORCIDs

Zhuo-yin Jiang  <https://orcid.org/0000-0001-6825-4283>

Feng-long Jia  <https://orcid.org/0000-0003-2391-5038>

Bei-xin Wang  <https://orcid.org/0000-0002-5253-8799>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Bian DJ, Ji LZ (2010) *Allopachria* Zimmermann, 1924 from Jiangxi, China, with descriptions of two new species (Coleoptera: Dytiscidae). Zootaxa 2350: 59–65. <https://doi.org/10.11646/zootaxa.2350.1.6>
- Bian DJ, Ji LZ (2012) Notes on Chinese *Allopachria* Zimmermann, with descriptions of two new species from Guangxi (Coleoptera: Dytiscidae). Zootaxa 3151: 28–34. <https://doi.org/10.11646/zootaxa.3151.1.2>
- Bian DJ, Guo C, Ji LZ (2013) *Allopachria yiae* sp. n. from Chongqing, China (Coleoptera: Dytiscidae: Hydroporinae). Zootaxa 3681(1): 85–88. <https://doi.org/10.11646/zootaxa.3681.1.6>
- Brancucci M (1984) A new *Platambus* from China (Coleoptera, Dytiscidae). Mitteilungen der Schweizerischen Entomologischen Gesellschaft 57: 153–154.

- Ji LZ, Guo C, Bian DJ (2014) *Allopachria longyanensis* sp. n. from China (Coleoptera: Dytiscidae). Zootaxa 3755(2): 194–196. <https://doi.org/10.11646/zootaxa.3755.2.5>
- Jiang ZY, Zhao S, Yang ZY, Jia FL, Hájek J (2022) A review of *Copelatus* Erichson, 1832 of Mainland China, with description of ten new species from the *japonicus* complex (Coleoptera: Dytiscidae: Copelatinae). Zootaxa 5124(3): 251–295. <https://doi.org/10.11646/zootaxa.5124.3.1>
- Miller KB, Nilsson AN (2003) Homology and terminology: Communicating information about rotated structures in water beetles. Latissimus 17: 1–4.
- Nilsson AN (2007) Some necessary corrections of the spelling of species-group names within the family Dytiscidae (Coleoptera). Zootaxa 1615: 49–54. <https://doi.org/10.11646/zootaxa.1615.1.5>
- Nilsson AN, Hájek J (2024a) A World Catalogue of the family Dytiscidae, or the Diving Beetles (Coleoptera, Adephaga). Version 1.I.2024. Distributed as a PDF file via Internet. <http://www.waterbeetles.eu>
- Nilsson AN, Hájek J (2024b) Catalogue of Palearctic Dytiscidae (Coleoptera). Version 1.I.2024. Distributed as a PDF file via Internet. <http://www.waterbeetles.eu>
- Peschet R (1923) Dytiscidae et Haliplidae nouveaux (Col.). Bulletin de la Société Entomologique de France 1923: 175–181. <https://doi.org/10.3406/bsef.1923.27196>
- Wewalka G (2000) Taxonomic revision of *Allopachria* (Coleoptera: Dytiscidae). Entomological Problems 31(2): 97–128.
- Wewalka G (2010) New species and new records of *Allopachria* Zimmermann (Coleoptera: Dytiscidae). Koleopterologische Rundschau 80: 25–42.
- Zhao S, Hájek J, Jia FL, Pang H (2012) A taxonomic review of the genus *Neptosternus* Sharp of China with the description of a new species (Coleoptera: Dytiscidae: Laccophilinae). Zootaxa 3478: 205–212. <https://doi.org/10.11646/zootaxa.3478.1.22>