

A new species in *Begonia* sect. *Diploclinium* (Begoniaceae) from Guangxi, South China

Ping-Ting Hu¹, Yu-Ni Huang², Meng-Ni Zhang¹, Bi-Xuan Chen¹, Shu-Shu Huang³, Ren-Kun Li⁴, Wen-Guang Wang⁵, Xin-Xin Feng²

¹ School of Urban Construction and Intelligent Manufacturing, Dongguan City University, Dongguan 523419, China

² Dongguan Botanical Garden, Dongguan 523086, China

³ Department of Horticulture and Landscape Architecture, Zhongkai University of Agriculture and Engineering, Guangzhou 510225, China

⁴ Enshi Gesneriad and Begonia Nursery Co., Ltd., Enshi 445000, China

⁵ Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, Yunnan, 666303, China

Corresponding authors: Wen-Guang Wang (wangwenguang@xtbg.ac.cn); Xin-Xin Feng (xinxin8715@163.com)

Abstract

Although Guangxi represents one of the distribution centres of begonias in China, the sect. *Diploclinium* (Wright) A. DC is not well documented herein. In this article, we illustrate a new species belonging to this section, *Begonia fangchengensis* Y.N.Huang, X.X. Feng & R.K.Li, which has been discovered in southern Guangxi. *Begonia fangchengensis* mostly resembles *B. rotundilimba* in elliptic leaf blade, palmate veins, dichasial cyme, three styles, axile placentation, and unequally 3-winged capsule. On the other hand, *B. fangchengensis* is characterized by creeping rhizome, pilose stipule, red and densely pilose petiole, pubescent leaf blade and pilose inflorescence bract, sparsely pilose ovary and the zygomorphic androecium, differing from the erect stem at anthesis, the green and densely villous petiole, the glabrous stipule, leaf blade, inflorescence bract, ovary and actinomorphic androecium in *B. rotundilimba*. Considering its small population size and narrow distribution, its conservation status is categorized as 'Endangered (D)' according to the IUCN Red List Categories and Criteria.

Key words: Limestone karst, morphology, new taxon, south China, taxonomy



Academic editor: Hanno Schaefer

Received: 3 August 2024

Accepted: 27 October 2024

Published: 5 December 2024

Citation: Hu P-T, Huang Y-N, Zhang M-N, Chen B-X, Huang S-S, Li R-K, Wang W-G, Feng X-X (2024) A new species in *Begonia* sect. *Diploclinium* (Begoniaceae) from Guangxi, South China. *PhytoKeys* 249: 277–285. <https://doi.org/10.3897/phytokeys.249.133724>

Copyright: © Ping-Ting Hu et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

As one of the fastest-evolving genera in vascular plants around the world, *Begonia* L. species are now 2,151 in number (Hughes et al. 2015) and these are mainly distributed in the humid tropical and subtropical regions of Asia, America, and Africa. Over 270 species of *Begonia* have been reported in China and the taxa are expected to exceed 300 species in the future (Tian et al. 2018). The begonias are primarily distributed in southern and central China, with Yunnan and Guangxi representing the centers of natural distribution.

Approximately 90 species of *Begonia* have been documented in Guangxi and new species have been reported frequently in recent years (Liu et al. 2020; Feng et al. 2021; 2022; 2023; Tian et al. 2021; Zhou et al. 2024). The begonias in Guangxi are dominated by *B.* sect. *Coelocentrum* taxa from karst

landforms. Only nine species belonging to the sect. *Diploclinium* are distributed in Guangxi, including: *Begonia bambusetorum* H. Q. Nguyen, Y. M. Shui & W. H. Chen, *Begonia fimbristipula* Hance, *Begonia gigabracteata* H. Z. Li & H. Ma, *Begonia glechomifolia* C. M. Hu ex C. Y. Wu & T. C. Ku., *Begonia grandis* Dryander, *Begonia grandis* subsp. *sinensis* (A. Candolle) Irmscher, *Begonia hymenocarpa* C. Y. Wu, *Begonia obsolescens* Irmscher and *Begonia sinovietnamica* C. Y. Wu.

In October 2021, an unknown begonia was discovered growing on the slope by a stream during fieldwork in the Shiwanshan Mountain, southern Guangxi. This taxon is distinct from the commonly karst distributed sect. *Coelocentrum* begonias in consideration of its ovary locule and placentae type in the field. Some individuals have been well cultivated in the nursery of Dongguan Botanical Garden and flower characters were observed in January 2022. Based on detailed morphological comparisons with similar species, this *Begonia* species is identified as a new species of the section *Diploclinium*.

Materials and methods

Morphological characters were observed and measured from fresh samples in the field. Morphological comparisons with similar taxa were undertaken by consulting the literature, examining herbarium (**IBK** and **IBSC**) specimens and observing living collections cultivated in the nursery of Dongguan Botanical Garden. The specimens were deposited at the Herbarium of South China Botanical Garden (**IBSC**), CAS.

Taxonomy

***Begonia fangchengensis* Y.N.Huang, X.X.Feng & R.K.Li, sp. nov.**

urn:lsid:ipni.org:names:77346002-1

Figs 1, 2

Chinese name: 防城秋海棠

Type. CHINA • Guangxi Zhuang Autonomous Region (广西壮族自治区), Fangchenggang City (防城港市), Fangcheng District (防城区), Shiwanshan Yao Autonomous Town (十万山瑶族乡), Bajiao Mountain, 21°56'54"N, 108°14'20"E, (Fig. 3), 198 m alt., on the rock or slope beside the stream, 18 October 2021, Yu Ni Huang, Xin Xin Feng & Ren Kun Li (holotype: 0925761, IBSC!).

Diagnosis. *Begonia fangchengensis* mostly resembles *B. rotundilimba* in its elliptic leaf blade, palmate veins, dichasial cyme, three styles, axile placentation, and unequally 3-winged capsule. However, there are creeping rhizome, pilose stipule, red and densely pilose petiole, pubescent leaf blade and pilose inflorescence bract, sparsely pilose ovary and the zygomorphic androecium in *B. fangchengensis*. These characters differ from the erect stem at anthesis, the green and densely villous petiole, the glabrous stipule, leaf blade, inflorescence bract, ovary and actinomorphic androecium in *B. rotundilimba*.

Description. Perennial evergreen herb, monoecious, 15–30 cm height.

Rhizome creeping, ca. 6–8 cm long and 8–10 mm in diameter, internode short. **Stipules** membranous, reddish-green, translucent, triangular, 4.5–8 × 4–5 mm, pilose. **Leaves** all basal, petiole 10–15 cm long, red and densely




Figure 1. *Begonia fangchengensis* drawn by Yunxiao Liu **A** habitat **B** stipule **C, D** bracts of inflorescences **E, F** front and back view of staminate flower **G** androecium **H** stamens **I, J** pistillate flower with 3 or 5 tepals **K** styles **L** styles and ovary **M–Q** ovary section from upper to lower positions.




Figure 2. Habitat and morphology of *Begonia fangchengensis* **A** habitat **B** flowering plant **C** creeping rhizome **D** stipules **E** leaf blade (adaxial) **F, G** leaf blade (abaxial) **H** juvenile leaf **I** inflorescence **J** bracts **K, L** front and side views of staminate flower **M, N** front and side views of androecium **O** stamens **P** pistillate flower with 3 tepals **Q, R, S** pistillate flower with 5 tepals **T** stigmas and ovary **U** ovary sections from upper to lower positions **V** capsule (Photos by Z.X. Liu).




Figure 3. Distribution of *B. fangchengensis*, *B. rotundilimba* and *B. sinovietnamica*.

white pilose; leaf blade asymmetric, widely ovate to elliptic, 12–17 × 8–10 cm; fleshy; adaxially bright green and smooth; abaxially pale-green, densely covered with white pilose along the veins; base oblique-cordate; margin irregularly denticulate; apex caudate acuminate; venation palmate, primary veins 8, adaxially slightly concave, abaxially convex. **Inflorescences** arising directly from rhizome, dichasial cymes, peduncle 6.5–8 cm long, pilose; flowers unisexual, 2–5 flowers per inflorescence; bracts membranous, pale green, triangular to subcircular, 2–3 lobed apically, margin serrate with cilia, 7–8 × 4–6 mm. **Staminate flower** tepals 4, white; outer 2 larger, widely ovate, 17–21 × 15–20 mm, abaxially sparsely pubescent; inner 2, lanceolate, margin entire, 10–12 × 4–6 mm, glabrous; androecium zygomorphic, sub-globose, ca. 6 mm in diameter; stamens numerous, 3–3.5 mm long, filaments fused at base; anthers yellow, ca. 1.5 mm long, cuneiform, apex slightly bent and retuse. **Pistillate flower** tepals 3–5, white, glabrous; outer 2, widely ovate to subcircular, margin entire, 11–12 × 9–10 mm; inner 1–3, widely ovate, 8–11 × 4–7 mm; ovary yellowish-green, trigonous-ellipsoid, 8.0–8.5 × 5.2–5.5 mm (wings excluded), sparsely pilose, 3-locular, upper 1 (abaxial wing side) degenerated occasionally; placentation axile, bilamellate. **Styles** 3, fused at base, yellow, ca. 3 mm in diameter, stigma spirally U-shaped twisted. **Capsules** nodding, trigonous-ellipsoid, 9–10 × 6.0–6.5 mm (wings excluded), yellowish-green, sparsely pilose, wings 3, unequal, 2-lateral wings smaller, crescent-shaped, abaxial wing semilune-shaped.

Distribution and habitat. Currently only known from one locality in Fangcheng District, Fangchenggang City, Guangxi, China (Fig. 3). It grows in

shaded environments along the stream or near a waterfall under the broad-leaved forest.

Phenology. November-December (Flowering), January- February (Fruiting).

Etymology. The specific epithet "*fangchengensis*" refers to the Fangcheng District, Guangxi where the species is distributed.

Conservation. Only two populations with approximately 200 individuals have been found in the type locality. According to the IUCN Red List Categories and Criteria (IUCN 2024), *B. fangchengensis* should be assessed as 'Endangered (D)' due to its small geographic distribution and low population size.

Discussion

The putative new *Begonia* species is morphologically characterized by having three styles, 3-locular ovary with axile and bifid placentation, representing the typical circumstance of the *Begonia* section *Diploclinium* (Lindl. 1846: 319) de Candolle (1859: 129). A total of 53 begonias of this section have been reported in China and most of them occur in Yunnan and Tibet, southeast China.

The putative new *Begonia* species mostly resembles *B. rotundilimba* from Pingbian, Yunnan in its elliptic leaf blade, palmate veins, dichasial cyme, three styles, unequally 3-winged capsule (Table 1, Fig. 4; Huang and Shui (1994)). In *B. rotundilimba*, there are erect stem at anthesis, green and villous petiole, and the stipule, leaf blade and inflorescence bract are all glabrous, being different from the creeping rhizome, pilose stipule and petiole, pubescent leaf blade and pilose inflorescence bract of *B. fangchengensis*. Furthermore, morphological dissimilarities also occur in reproductive organs, like the glabrous ovary and actinomorphic androecium of *B. rotundilimba*, compared with sparsely pilose ovary and the zygomorphic androecium in *B. fangchengensis*.

Besides the similarity to *B. rotundilimba* in morphology, the putative new *Begonia* species is closest to *B. sinovietnamica* in geographic distribution and altitude (200–400 m) (Fig. 3). In *B. sinovietnamica*, leaf blade is adaxially hirsute, tepals

Table 1. Morphological comparisons of *B. fangchengensis* and relevant taxa.

Character	<i>B. rotundilimba</i>	<i>B. sinovietnamica</i>	<i>B. fangchengensis</i>
Rhizome	creeping, erect stem at anthesis; internode 2-3 cm long	elongate; internode 0.3-0.5 cm long	creeping; internode 0.3-1 cm long
Stipule	ovato-triangular, glabrous	oblong, pilose	triangular, pilose
Petiole	green, densely villous	reddish brown, densely villous	red, densely pilose
Leaves	blade adaxially green, glabrous	blade adaxially green, hirsute	blade adaxially bright green, smooth
Inflorescence bract	bracts glabrous, apex acuminate	bracts glabrous, apex acuminate	bracts apex 2-3 lobed, both pilose
Staminate flower	androecium actinomorphic, anthers oblong	anthers oblong, apex obtuse	androecium zygomorphic, anthers cuneiform, apex slightly bent and retuse
Pistillate flower	tepals 5, styles 3, ovary pilose, unequally 3-winged	tepals 4, styles 3, ovary sparsely pilose, subequally 3-winged	tepals 3-5, styles 3, ovary sparsely pilose, unequally 3-winged
Phenology (flowering; fruiting)	Apr; Jul	Jul; Aug	Nov; Jan
Habitat	alt. 1600-1800 m, Yunnan	alt. 230-400 m, Guangxi	alt. 198 m, Guangxi




Figure 4. Habitat and morphology of *Begonia rotundilimba* **A** habitat **B** flowering plant with erect stem at anthesis **C** leaf blade (adaxial) **D** leaf blade (abaxial) **E** petiole **F, G** front and back views of staminate flower **H** androecium **I, J** front and lateral view of pistillate flower **K** ovary section (Photos by W. G. WANG).

of pistillate flower are 4 and the 3-winged ovary is subequal, in contrast with the adaxially smooth blade, 3–5 tepals of pistillate flower and unequally 3-winged ovary of the new species. Furthermore, the two species also differ from each other in phenology, including flowering and fruiting time (Table 1; Wu (1997)).

Acknowledgements

We thank Zhixian Liu for providing the photographs. We are also grateful to the editor and the reviewers for improving our manuscript.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was supported by the Dongguan Municipal Project for Improvement of Social Science and Technology (20231800940632); Guangdong Provincial Education Department Young Innovative Talent Project for Colleges and Universities (2022WQNCX124); the Science and Technology Innovation Project in Department of Housing and Urban Rural Development of Guangdong Province (2023-K13-384726) and the Guangdong Rural Revitalization Strategic Foundation (20211800400042).

Author contributions

Conceptualization: XXF. Data curation: MNZ, BXC. Formal analysis: SSH. Investigation: PTH, YNH, GWG. Resources: XXF, RKL.

Author ORCIDs

Ping-Ting Hu  <https://orcid.org/0009-0001-2656-167X>

Yu-Ni Huang  <https://orcid.org/0000-0001-5421-4192>

Bi-Xuan Chen  <https://orcid.org/0009-0006-5022-6229>

Xin-Xin Feng  <https://orcid.org/0000-0002-7421-2773>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- De Candolle A (1859) Mémoire sur la famille des Bégoniacées. In: Brongniart AD, Decaisne J (Eds) Annales des Sciences Naturelles Botanique, serie 4, 11: 93–149. [Fortin, Masson, Paris]
- Feng XX, Xiao Y, Liu ZX, Li RK, Wei D, Tian DK (2021) *Begonia pseudoedulis*, a new species in *Begonia* sect. *Platycentrum* (Begoniaceae) from southern Guangxi of China. *PhytoKeys* 182: 113–124. <https://doi.org/10.3897/phytokeys.182.69074>
- Feng XX, Huang XF, Huang YN, Liu ZX, Li RK, Zhou JY, Guo W, Chen XY, Tian DK (2022) *Begonia parvibracteata*, a new species in *Begonia* sect. *Platycentrum* (Begoniaceae) from Guangxi of China, based on morphological and molecular evidence. *PhytoKeys* 214: 27–38. <https://doi.org/10.3897/phytokeys.214.90004>
- Feng XX, Chen YM, Liu JX, Huang YN, Liu ZX, Li RK, Zhao T, Guo W (2023) *Begonia barosma*, a new species in *Begonia* sect. *Coelocentrum* (Begoniaceae) from limestone

- karsts across Sino-Vietnamese Border. *Taiwania* 68(3): 349–354. <https://doi.org/10.6165/tai.2023.68.349>
- Huang SH, Shui YM (1994) New taxa of *Begonia* from Yunnan. *Acta Botanica Yunnanica* 16(4): 333–342.
- Hughes M, Moonlight PW, Jara-Muñoz A, Tebbitt MC, Wilson HP, Zhong K, Mills T, Pullan M (2015) *Begonia* Resource Centre. [Online database] <http://padme.rbge.org.uk/begonia/> [Updated: 7th July 2024]
- IUCN (2024) Guidelines for Using the IUCN Red List Categories and Criteria. Version 16, prepared by the Standards and Petitions Committee of the IUCN Species Survival Commission. <https://www.iucnredlist.org/resources/redlistguidelines> [accessed 18th October 2024]
- Lindley J (1846) *The vegetable kingdom; or, the structure, classification, and uses of plants, illustrated upon the natural system*. Bradbury and Evans, London, 911 pp. <https://doi.org/10.5962/bhl.title.7756>
- Liu Y, Tseng YH, Yang HA, Hu AQ, Xu WB, Lin CW, Kono Y, Chang CC, Peng CI, Chung KF (2020) Six new species of *Begonia* from Guangxi, China. *Botanical Studies* 61(1): 21. <https://doi.org/10.1186/s40529-020-00298-y>
- Tian DK, Xiao Y, Tong Y, Fu NF, Liu QQ, Li C (2018) Diversity and conservation of Chinese wild begonias. *Plant Diversity* 40(3): 75–90. <https://doi.org/10.1016/j.pld.2018.06.002>
- Tian DK, Ge BJ, Xiao Y, Tian Q, Li C (2021) *Begonia scorpiuroloba*, a new species in *Begonia* sect. *Platycentrum* (Begoniaceae) from southern Guangxi of China. *Phytotaxa* 479(2): 191–197. <https://doi.org/10.11646/phytotaxa.479.2.5>
- Wu CY, Ku TC (1997) New Taxa of the *Begonia* L. (Begoniaceae) from China (Cont.). *Acta Phytotaxonomica Sinica* 35(1): 43–56.
- Zhou JY, Song Q, Zhao FC, Hoang TS, Tao D, Guan S, Yan H, Tian DK (2024) *Begonia floriprolifera*, a new species of *Begonia* (Begoniaceae) from Southwestern Guangxi of China and Northern Vietnam *Taiwania* 69(1): 83–88. <https://doi.org/10.6165/tai.2024.69.83>