

Innate Antimicrobial Activity of Nasal Secretions

ALEXANDER M. COLE, PUNEET DEWAN, AND TOMAS GANZ*

*Departments of Medicine and Pathology and the Will Rogers Institute for Pulmonary Research,
UCLA School of Medicine, Los Angeles, California 90095*

Received 9 February 1999/Returned for modification 19 March 1999/Accepted 2 April 1999

Minimally manipulated nasal secretions, an accessible form of airway surface fluid, were tested against indigenous and added bacteria by using CFU assays. Antimicrobial activity was found to vary between donors and with different target bacteria and was markedly diminished by dilution of the airway secretions. Donor-to-donor differences in electrophoresis patterns of nasal secretions in sodium dodecyl sulfate-polyacrylamide gel electrophoresis (PAGE) and acid urea-PAGE analyses were readily observed, suggesting that polymorphic genes encode the secreted proteins. Three donors (of twenty-four total), whose nasal fluid yielded similar protein band patterns and did not kill indigenous bacteria, were determined to be heavy nasal carriers of *Staphylococcus aureus*. Their fluid was deficient in microbicidal activity toward a colonizing strain of *S. aureus* but the defect was corrected in vitro by a 1:1 addition of nasal fluid from noncarriers. The microbicidal activity of normal fluid was inactivated by heating it for 10 min to 100°C and could not be restored solely by the addition of two major nasal antimicrobial proteins, lysozyme and lactoferrin. Several other known antimicrobial proteins and peptides, including statherin, secretory phospholipase A₂, and defensins, were identified in nasal secretions and likely contribute to their total antimicrobial properties. Nasal fluid may serve as a useful model for the analysis of lower-airway secretions and their role in host defense against airway colonization and pulmonary infections.

Host defense is a prominent function of nasal secretions. Among the components of the secretions, mediators of adaptive immunity include immunoglobulin A (IgA) and IgG, which are produced by plasma cells juxtaposed to submucosal glands (29). The immunoglobulins are thought to act in the lumen of the airway and on the mucosa to prevent the attachment and invasion of pathogenic organisms. Mediators of innate mucosal host defense are also found in nasal secretions and include substances that selectively disrupt bacterial cell walls and membranes, sequester microbial nutrients, or act as decoys for microbial attachment. Alexander Fleming discovered the intrinsic antimicrobial properties of human nasal secretions in 1922 and attributed them to lysozyme (8). Since then, additional components of nasal secretions have been identified, including lactoferrin, mucus glycoproteins, secretory leukoprotease inhibitor (SLPI), uric acid, peroxidase, aminopeptidase, immunoglobulins, and neutral endopeptidase (reviewed in references 15 and 16). Systematic analysis of the individual and combined antimicrobial effects of these and other components has not been reported. Pathological alterations in antimicrobial properties of airway secretions may contribute to epithelial colonization by bacteria in such disorders as cystic fibrosis, chronic bronchitis, and chronic sinusitis.

The epithelial lining fluid covering the airways derives its proteins from plasma transudate, mucous and serous cells in submucosal glands, goblet cells, Clara cells, epithelial cells, and other cells within the mucosa (plasma cells, mast cells, phagocytes, and fibroblasts). The epithelial lining fluid is a bilaminar mucous layer consisting of an outer viscous gel (mucus) and a periciliary serous layer (42). Nasal epithelial cilia continually transport mucus with ensnared particulate matter and microbes toward the oropharynx, where they are swallowed. Consequently, in the normal individual, the nasal mucus blanket

turns over every 10 to 20 min and must be replenished. However, fluid is relatively stagnant in the anterior portion of the nasal mucosa (15), and there the elimination of impacted microbes must be largely dependent on its intrinsic antimicrobial properties. This stagnant anterior microenvironment might serve as a useful model of torpid viscous secretions in the airways of patients with cystic fibrosis and other forms of bronchiectasis.

The limited available evidence suggests that nasal and bronchial secretions have many similarities in their composition (23). However, unlike bronchial secretions, nasal secretions can be easily collected and could provide a useful foundation for the study of other airway secretions. We reexamined the antimicrobial properties of human nasal secretions by using a CFU microassay, wherein bacteria are directly incubated with minimally manipulated nasal secretions. In contrast to recent studies that have focused on the antibacterial action of individual components of lower airway fluids (12, 27, 35, 44), the analysis of minimally manipulated fluid affords the opportunity to assess the natural antimicrobial activity within the airways.

MATERIALS AND METHODS

Collection of nasal secretions. Nasal secretions were collected from 24 healthy volunteer donors according to a protocol approved by the University of California at Los Angeles (UCLA) Institutional Review Board. From most donors, nasal secretions were collected by vacuum-aided suction, without chemical stimulation, to avoid introducing foreign substances into the nasal fluids. Gentle manipulation of a narrow rubber-tipped vacuum device inside the nasal passages mildly stimulated nasal secretions. The range of secretion volumes collected varied between 150 μ l and 14 ml, depending on the donor. For selected collections, the secretion of nasal fluids was induced by 25 to 50 mg of methacholine chloride (Spectrum Pharmaceuticals, Irvine, Calif.) administered into one nostril by an atomizing dosimeter. The range of secretion volume varied between 0.5 and 15 ml per donor. Secretions were stored at 4°C for prompt usage or at -20°C for longer-term storage. The microbicidal activity and protein profile by acid urea-polyacrylamide gel electrophoresis (AU-PAGE) of secretions stored for 2 weeks at 4°C or >1 month at -20°C was unchanged compared to fresh samples.

Minimally manipulated fluid. Vacuum-aspirated nasal fluid collected without methacholine was sonicated for 2 h at 4 to 8°C in a Branson 1200 sonicating water bath (Branson Cleaning Equipment Co., Shelton, Conn.) to disrupt the mucoprotein aggregates and to facilitate accurate handling. Unless otherwise stated, fluid samples with a high number (>10⁵/ml) of indigenous bacteria were

* Corresponding author. Mailing address: Department of Medicine, Division of Pulmonary and Critical Care, UCLA School of Medicine, CHS 37-055, 10833 Le Conte Ave., Los Angeles, CA 90095-1690. Phone: (310) 825-7499. Fax: (310) 206-8766. E-mail: tganzz@ucla.edu.




FIG. 1. The fate of indigenous bacteria in minimally manipulated nasal secretions. (A) The majority of nasal fluid samples were inhibitory to indigenous microbes. (B) Nasal fluid samples from a minority of donors were permissive for indigenous microbes. Asterisk indicates $n = 1$, due to insufficient amounts of nasal secretions from the indicated donors.

were employed to ascertain the effect of bacterial species and donor-to-donor variability on the antimicrobial capacity of nasal fluid. The differential effects of a single donor's secretions (donor 4) against multiple species and strains of bacteria is presented in Fig. 2 and 3, while the various effects of multiple donors' secretions against an individual bacterial strain is shown in Fig. 4. An inoculum of 2×10^4 to 4×10^4 CFU/ml was used to model the host-defense challenge by moderate numbers of inhaled microbes (Fig. 1A). To avoid selection bias, we included samples from donors from whom only scant amounts of nasal secretions could be collected, but their secretions were tested in a single assay (indicated by an asterisk).

Nasal fluid samples obtained on different days from the same donor showed similar antimicrobial activity against *P. aeruginosa* CF at 3 h of incubation (Fig. 2A). By 24 h, minor regrowth of bacteria was observed in two of the three samples (b and c); however, the differences in CFU counts compared to input were not significant. Thus, variance in the composition of nasal secretions from the same donor may contribute to small fluctuations in bacterial regrowth.

The antimicrobial effect of nasal secretions from a single collection was seen to depend on the properties of the bacterial target (Fig. 2B). An LPS mutant strain, H234, and a mucoid CF strain, H246, were effectively cleared at 1 h with no regrowth at 24 h ($P < 0.003$). The PAO1 strain, H103, was not cleared but did not significantly grow throughout the observed time period. The nonmucoid revertant strain of H246, H247, slowly grew throughout the entire period of observation with a doubling time of approximately 8 h. Similar differences were observed in the effect of nasal fluid on non-*Pseudomonas* strains (Fig. 2C). *S. aureus* (clinical isolate) was effectively eliminated at 24 h compared with time zero ($P < 0.003$). In contrast, *E. coli* and *S. typhimurium* CFU counts slowly increased ($P < 0.004$ and $P < 0.00007$, respectively), with a doubling time of 10 to 16 h.

To assess the effect of dilution in a CFU microassay, donor 4's nasal secretions were diluted 1:1 and 1:2 with $1 \times$ NB. Dilution had little effect on antimicrobial activity against *S. aureus* (clinical isolate), *E. coli*, and *S. typhimurium* (Fig. 3A to C), as well as on *P. aeruginosa* H234, *P. aeruginosa* H246, and *P. aeruginosa* H247 (data not shown). However, *S. aureus* (nasal carrier isolate), *P. aeruginosa* CF, and *P. aeruginosa* H103 proliferated in diluted nasal fluid (Fig. 3D to F). In this latter

group, the increased dilution resulted in significantly increased bacterial growth either at all of the timepoints tested (Fig. 3D, $P < 0.02$), at 1 and 3 h (Fig. 3E, $P < 0.02$), or at 3 h (Fig. 3F, $P < 0.003$). The observation that thinning of nasal fluids can increase bacterial proliferation indicates that the factors that determine microbial growth in nasal fluid are primarily suppressive and concentration limited.

To detect the variation of the antimicrobial effect between donors, minimally manipulated nasal secretions from seven different donors were tested in a CFU microassay against several strains of *P. aeruginosa*. Four strains were employed for this study: strains CF (CF isolate), H246 (CF isolate), H234 (LPS mutant), and H103 (wild type) (Fig. 4). Several patterns of antimicrobial effect emerged. All samples tested against strain CF decreased CFU counts by >2 log of the inoculum by 1 h ($P < 0.0002$) and nearly cleared the nasal secretions of exogenous bacteria by 3 h (Fig. 4A, $P < 0.0004$). Bacterial regrowth at 24 h appeared to be a donor-specific phenomenon. The wild type, H103, was at least transiently inhibited by nasal secretions from all donors tested, but regrowth at 24 h ($P < 0.01$) was again observed in secretions from several donors (Fig. 4B). The mucoid (alginate-producing) *P. aeruginosa* strain, H246, and the LPS-deficient mutant strain, H234, were intermediate in their susceptibilities to nasal secretions (data not shown). Thus, alginate production and LPS structure affect the sensitivity of bacteria to normal nasal fluid.

Identification of antimicrobial (poly)peptides in nasal secretions. We screened the nasal secretions by Western blotting with specific antibodies against known antimicrobial (poly) peptides (Table 1). Among the proteins so identified, lysozyme, lactoferrin, SLPI, and secretory phospholipase A_2 (sPLA $_2$) are antimicrobial components previously noted in nasal secretions. Prominent protein bands that could not have been identified by Western blotting were separated by AU-SDS/2-dimensional PAGE and transferred to PVDF membranes for protein sequencing. Several abundant (poly)peptides were identified as fragments of albumin and the light chain of IgG. Newly described cationic antimicrobial components of nasal secretions included two β -defensins (HBD-1 and HBD-2), an α -defensin (human neutrophil peptide 1 [HNP-1]), statherin, and β -microseminoprotein. The latter three peptides were identified from larger-scale (>5 -ml) preparations of acid-extracted nasal se-






FIG. 2. Susceptibility of various strains of bacteria to a single donor's nasal fluid. Nasal secretions from donor 4 were subjected to a modified CFU microassay of nasal fluid targeting four strains of *P. aeruginosa*, *E. coli* ML-35p, *S. typhimurium* wild type, and *S. aureus* clinical isolate. (A) Secretions tested at three different times showed similar activities against *P. aeruginosa* CF between 0 and 3 h but varied in the amount of regrowth at 24 h. (B) Fluid incubated with four different strains of *P. aeruginosa* showed strain-specific antimicrobial activity. (C) Nasal secretions permitted the slow growth of *E. coli* and *S. typhimurium*; however, *S. aureus* was effectively cleared. Init, inoculum (input) at time zero for each sample tested, without the addition of nasal fluid.

cretions and were present at much lower concentrations than lysozyme, lactoferrin, or SLPI.

Variation in antimicrobial (poly)peptide content. Differences in the antimicrobial activities of nasal secretions could result from donor-to-donor variations in the protein composition of the secretions. Equal volumes of eight donors' nasal secretions were analyzed by AU-PAGE (Fig. 5A) and stained with Coomassie brilliant blue (Sigma) or analyzed by Tricine-SDS-PAGE and stained with silver stain (Fig. 5B). The variability of the protein patterns among the donors is readily seen. Although a unique protein fingerprint is seen in each donor's fluid, the electrophoretic patterns can be classified into three groups. Group "a" (donors 8, 20, and 24) had the highest protein concentration, group "b" (donors 9 and 11) had a moderate protein concentration, and group "c" (donors 2, 4, and 19) had the lowest overall protein concentration. Group a also had a number of low-molecular-weight (poly)peptides, several of which are possible antimicrobials. In agreement with previous reports (17), the total protein concentration in our nasal fluid samples ($n = 3$), as performed by the UCLA Clinical Laboratories by the pyrogallol red molybdate method (Synchro CX Microprotein Calibrator; Beckman Instruments, Inc., Brea, Calif.), was found to be 1 to 3 mg/ml. Concentrations of lysozyme, lactoferrin, and SLPI in crude nasal fluid ($n = 5$) as tested by semiquantitative Western analysis are given in Table 1. A β -defensin, HBD-2, was also detected by Western blot in three of five donors. Donor-to-donor differences in electro-

phoretic patterns of nasal secretions in SDS-PAGE and AU-PAGE may indicate that polymorphic genes encode many of the proteins of nasal secretions.

Nasal *S. aureus* carriage. Since the nasal fluid of donors 20 and 24 was permissive for the growth of indigenous bacteria (Fig. 1B) and the two protein band patterns were similar (Fig. 5), the indigenous bacteria were typed to identify microbiological similarities between these two donors. With a Bacto-Staph Latex rapid slide agglutination kit (Difco), both donors were found to have numerous colonies that tested positive for *S. aureus*. Twenty-two other donors' secretions were also subjected to the Bacto-Staph Latex agglutination test; only one of these donors had high numbers of colonies positive for *S. aureus* (donor 8), and two had a few colonies positive for *S. aureus* (donors 3 and 23). Samples of secretions from donors 8, 20, and 24, suspected of having high *S. aureus* colonization, were submitted to standard clinical microbiological typing, which confirmed this finding. Heavy *S. aureus* nasal carriage in 3 of 24 total donors (12.5%) is within the expected frequency based on previous estimates of carriage rates of 8 to 18% (18).

In principle, carriage could result either if the colonizing strains of *S. aureus* are extraordinarily resistant to secretions or if the carriers' nasal fluid is favorable to *S. aureus* proliferation. *S. aureus*, isolated from a carrier's nose (donor 24), was tested in CFU microassays of carrier and noncarrier nasal secretions (Fig. 6A). The nasal fluids from four donors, two noncarrier (donors 4 and 11) and two carrier (donors 20 and 24, secretions gamma-irradiated), were tested. While the noncarriers' secretions were bacteriostatic (donor 4) or bactericidal (donor 11), both carriers' secretions permitted a 2- to 3-log increase in CFU counts of the *S. aureus* isolate compared to time zero ($P < 0.05$). This suggests that the *S. aureus* strain is not extraordinarily resistant to nasal secretions but that the carriers' secretions may be deficient in antimicrobial activity.

Noncarrier and carrier nasal secretions, with their indigenous flora, were mixed to determine whether the antimicrobial




FIG. 3. Effect of fluid dilution on microbicidal activity. Six strains of bacteria were subjected to a CFU microassay of undiluted, 1:1-diluted, and 1:2-diluted minimally manipulated nasal fluid from donor 4. The CFU counts of *S. aureus* (clinical isolate) (A), *E. coli* (B), and *S. typhimurium* (C) were little affected. The antimicrobial effects of nasal secretions on *S. aureus* (isolate from nasal carrier donor 24) (D), *P. aeruginosa* CF (E), and *P. aeruginosa* H103 (F) were decreased with increasing dilution. The effects of undiluted, 1:1-diluted, and 1:2-diluted secretions are represented by closed circles, open triangles, and closed squares, respectively. The input inoculum (closed triangles) is shown for each graph.

activity of carrier fluid could be restored by the addition of noncarrier fluid. Figure 6B reveals that the indigenous flora of noncarrier secretions (donor 4) was killed throughout incubation ($P < 0.01$). When carrier fluid (donor 24) was incubated alone, its bacteria proliferated to 1.5 log units above the initial CFU count. Indigenous bacteria in carrier secretions incubated with heat-inactivated (boiling for 10 min) noncarrier fluid showed growth kinetics similar to that of carrier fluid incubated alone. However, a 1:1 (vol/vol) mixture of minimally manipulated carrier and noncarrier fluids was bactericidal for indigenous bacteria in the carrier's secretions ($P < 0.02$ compared to carrier fluid with or without the addition of heat-in-

activated normal secretions). A similar pattern of activity was seen with the same noncarrier fluid (donor 4) and different carrier fluid (donor 20) (data not shown). A 1:1 (vol/vol) mixture of carrier (donor 20) and noncarrier fluids was bactericidal ($P < 0.04$ compared to either carrier fluid alone or carrier fluid mixed with heat-inactivated noncarrier fluid). Thus, the non-carrier nasal secretions contain antimicrobial substances that are either deficient or inactive in the carrier's nasal secretions.

Lysozyme and lactoferrin did not restore the microbicidal activity of heat-inactivated nasal secretions. Physiological concentrations of human milk lysozyme and human milk lactoferrin (Sigma), reported as the major antimicrobial agents in




FIG. 4. Susceptibility of *P. aeruginosa* to multiple donors' nasal fluids. Activities of multiple donors' nasal secretions were tested individually against two strains of *P. aeruginosa*. Different donors' fluids were tested in a CFU microassay against a cystic fibrosis clinical isolate (A) and the wild-type strain H103 (B). Init, inoculum (input) at time zero for each donor tested, without the addition of nasal fluid. Asterisks indicate $n = 1$, due to scant fluid from the indicated donors.

TABLE 1. Eleven (poly)peptides identified in nasal secretions by Western analysis, microsequencing, and/or mass spectrometry

Protein	Molecular size (kDa)	Concn ($\mu\text{g/ml}$) ^a	Method of detection ^b
Lysozyme	14.7	250–500	a, b, c
Lactoferrin	78	80–200	a
SLPI	11.7	10–80	a, b, c
β -Microseminoprotein	10.8	ND	b, c
sPLA ₂	14.1	ND	d
Statherin	5.2	ND	b, c
Light-chain IgG	~28	ND	b
Albumin	~6 (fragment)	ND	b
Albumin	~30 (fragment)	ND	b
Albumin	~31 (fragment)	ND	b
HBD-1 ^c	3.9	ND	a
HBD-2 ^c	4.3	0.3–4	a
HNP-1	3.4	ND	b

^a Concentration was determined from five samples by using semi quantitative Western analysis, with each sample containing lysozyme, lactoferrin, and SLPI. Note that three of five donor samples contained measurable amounts ($>0.3 \mu\text{g/ml}$) of HBD-2. ND, not determined.

^b a, Western analysis; b, microsequencing by N-terminal Edman degradation; c, mass spectrometry; d, PLA₂ activity determined by using radiolabeled *E. coli* ML-35p substrate as described previously (31).

^c Some β -defensin comigrated with higher-molecular-weight proteins.

nasal secretions (32), were added to heat-inactivated fluid in an attempt to reconstitute the antimicrobial activity of nasal secretions (Fig. 7). While unaltered nasal fluid was bactericidal, *P. aeruginosa* H246 was insensitive to heat-inactivated secretions. Lysozyme activity in heat-inactivated nasal fluid, as measured by the lysoplate assay, was found to be >25 -fold less than in nonboiled control ($n = 4$; data not shown). The addition of 200 μg of lactoferrin per ml alone, 500 μg of lysozyme per ml alone, or the combination of 200 μg of lactoferrin and 500 μg of lysozyme per ml to heat-inactivated nasal fluid did not restore its antimicrobial activity ($P < 0.03$). The addition of lysozyme restored the enzymatic activity of nasal fluid towards *Micrococcus lysodeikticus* in the lysoplate assay (data not shown), but lysozyme added alone increased the growth of *P. aeruginosa* H246 at 24 h compared to growth in boiled nasal fluid ($P < 0.04$). The difference in the effect of lactoferrin

alone and the lactoferrin-lysozyme combination was not statistically significant at 24 h compared to the boiled nasal fluid alone. Thus, other (poly)peptides besides lysozyme and lactoferrin contribute to the total microbicidal activity of nasal fluid.

DISCUSSION

Since the pioneering studies of Fleming (8), scant attention has been paid to the antimicrobial activity of nasal secretions and their role in the prevention of microbial colonization. We were motivated to reexamine the antimicrobial activity of nasal fluid because of recent evidence that the bacterial colonization of the airways in cystic fibrosis is due to a local defect in epithelial host defense (26). Our study confirms Fleming's observation that nasal secretions have intrinsic antimicrobial activity, but it also shows that this activity is precarious. Activity varies depending on the donor and the target bacteria; is often time limited, with regrowth observed by 24 h of incubation; and is ablated by dilution. Under normal conditions in vivo, the contributions of mechanical factors such as mucociliary clearance, phagocytic clearance, or sneezing may shorten the time available to microbes to adapt to the deleterious effects of nasal secretions. It is easy to conceive how the failure of one or more of these mechanisms could permit the colonization of airways by microbes.

Nasal *S. aureus* carriage has been identified as a risk factor for the pathogenesis of surgical and other hospital-associated infection. Persistent nasal carriers of *S. aureus* account for approximately 20% of the population, with 8 to 18% heavily colonized ($>10^5$ CFU/ml; reviewed in reference 18). Although various epithelial and mucus host factors, such as surface glycoproteins and proteoglycans, have been shown to mediate binding, the precise adhesive molecules on host and bacteria have not been identified. *S. aureus* appears to attach to cell-associated and cell-free secretions (34) and to involve receptor sites of secretory IgA (3), glycolipids (20), and surfactant protein A (28). Various bacteria, including *Staphylococcus epidermidis*, are capable of reducing nasal ciliary activity in vitro (6). Enhanced adhesion and diminished mucociliary clearance could explain the retention of *S. aureus* within the nasal passageway but have no direct bearing on bacterial proliferation. Our observations suggest that the nasal fluid of at least some




FIG. 5. Donor-to-donor variability in the protein pattern of nasal secretions. (A) AU-PAGE of eight donor's nasal secretions shows variations in lactoferrin, albumin, β -microseminoprotein (beta-MSP), SLPI, lysozyme, and statherin concentrations. (B) Reducing Tricine-SDS-PAGE of eight donors' nasal secretions shows the variability in low-molecular-weight proteins. Superscripts indicate three donor groupings based on protein band similarities. The positions of identified components of nasal secretions are shown at the gel margins.




FIG. 6. Nasal carriers of *S. aureus* are deficient in antimicrobial components that are restored by the addition of noncarrier secretions. (A) *S. aureus*, isolated from donor 24, was used as the target bacterium in a CFU microassay of nasal fluid from two noncarriers (donors 4 and 11) and two carriers (donors 20 and 24, gamma-irradiated fluid). While the noncarriers' fluids were bacteriostatic or bactericidal, the carriers' fluids did not inhibit bacterial growth. (B) The 1:1 mix of nasal secretions from an *S. aureus* carrier (donor 24) with fluid from a noncarrier (donor 4) is bactericidal to indigenous flora, while the mix of nasal secretions from an *S. aureus* carrier with heat-inactivated (boiling for 10 min) secretions from a noncarrier promoted the growth of indigenous bacteria.

carriers is deficient in antimicrobial activity. Because of the natural donor-to-donor variation in the composition of nasal fluid, larger studies will be necessary to determine whether this mechanism is the sole or common determinant of nasal colonization by *S. aureus* and to identify the specific (poly)peptides or other substances that may be deficient in carriers.

Early colonization of cystic fibrosis airways predominantly involves *S. aureus*, *Haemophilus influenzae*, or *P. aeruginosa*. The initially infecting *P. aeruginosa* organisms are typically nonmucoid; however, they become highly mucoid after continued exposure to the environment of the cystic fibrosis airways. Colonization with *P. aeruginosa* may induce further airway injury, perhaps due to destructive virulence factors, such as exotoxin A and proteases, and the recruitment of the host's neutrophils with their proteases, leading to the progressively destructive bronchitis and bronchiolitis. Some studies have attributed *Pseudomonas* colonization in cystic fibrosis to abnormally glycosylated cell surface receptors with enhanced affinity for pseudomonal adhesive structures (13, 43). Other work suggests that abnormal mucin and the lack of hydration of respiratory secretions entraps bacteria, permitting colonization (22, 26, 36). It has also been proposed that defects in intrinsic antimicrobial activity of airway secretions may be induced by alterations in the salt concentration due to defective chloride transport in cystic fibrosis (37). Our study of nasal secretions suggests that their antimicrobial activity is naturally variable from donor to donor and is often time limited. Factors, such as thick mucus or ciliary dysmotility, that delay the mechanical clearance of particulates from the airway may permit microbes to adapt to the damaging effects of secretions and to recover and resume growth. The ablation of the antimicrobial effects of nasal fluid by dilution *ex vivo* points to the possibility that reduced concentrations of antimicrobial substances due to abnormalities of fluid movement across epithelia could impair the antimicrobial activity of the airway secretions. The well-recognized phenotypic variability of cystic fibrosis among patients with identical mutations may in part be due to interactions with genes that control the (poly)peptide composition of nasal (and presumably other airway) fluid. Our observations on the variability of the protein patterns in nasal secretions are reminiscent of similar studies with human saliva, whose (poly)

peptides are remarkably polymorphic (2). The effects of nasal (poly)peptide polymorphisms on host defense and their possible role as phenotypic modulators in cystic fibrosis should be a productive area for further investigation.

Nasal secretions contain antimicrobial proteins and peptides as a first-line host defense against microbial invaders. Lysozyme and lactoferrin, stored in and secreted from serous cells in nasal submucosal glands, are the most abundant antimicrobial proteins of nasal fluid. They are bactericidal alone for gram-positive bacteria and in combination for some gram-negative bacteria (5, 32). Given their strong combined antibacterial properties and since few other major microbicidal proteins have been previously described in nasal fluid, we expected they would reconstitute the antibacterial activity in heat-inactivated secretions. Our experiments suggest that other factors,




FIG. 7. Physiological concentrations of lysozyme and lactoferrin added to heat-inactivated nasal fluid (donor 4) did not restore microbicidal activity. While nonboiled fluid exhibited the expected bactericidal activity, heat-inactivated (boiling for 10 min) did not inhibit CFU growth. Physiological concentrations of lactoferrin and lysozyme, added separately and concurrently to heat-inactivated secretions, did not restore the antimicrobial activity.

or cofactors acting with lysozyme and lactoferrin, contribute to the antimicrobial properties of nasal fluid.

sPLA₂ has been previously identified from methacholine-induced nasal lavage fluid and mucosa as a far less abundant (poly)peptide by both immunostain and enzymatic activity (1, 38). Ca²⁺-dependent antimicrobial activity of sPLA₂ has been demonstrated against both gram-positive and gram-negative bacteria (reviewed in reference 10). In the present study we identified sPLA₂ in unstimulated nasal fluid as a potential contributor to the total microbicidal activity of nasal fluid.

The current study is the first to identify various small cationic antimicrobial peptides in nasal secretions. HBD-1 has recently been characterized in urogenital tissues (40) and bronchoalveolar lavage (35), while HBD-2 is expressed at sites of inflammation (24, 35). HNP-1 (9) was likely a degranulation product of transudated neutrophils. Statherin, a small antimicrobial phosphoprotein previously identified in saliva (14), was also found to be present in nasal secretions. Another low-molecular-weight protein, β -microseminoprotein, has been localized to several human and rat mucosal tissues (7, 41) and was identified here as a component of nasal fluid (Table 1). Its detection in nasal fluid suggests that β -microseminoprotein is a ubiquitous antimicrobial protein secreted as a component of mucosal defense. We also acknowledge the possibility that anionic antimicrobial (poly)peptides contribute to the antimicrobial activity of nasal fluid, as observed in ovine bronchoalveolar lavage fluid (4); however, our fractionation methodology was not designed to detect such peptides. The abundance of antimicrobial peptides and proteins present in nasal secretions, coupled with the inability of lysozyme and lactoferrin to restore the microbicidal activity of heat-inactivated nasal fluid toward the test strain of *P. aeruginosa*, suggest that the antimicrobial activity of nasal fluid results from complex effects of its many components.

ACKNOWLEDGMENTS

This work was supported by grants from the Cystic Fibrosis Foundation and the National Institutes of Health (HL46809 and AI40268).

We are grateful to Erika Valore and Christina Park for recombinant HBD-1 and -2 and polyclonal antisera and also for their guidance in protein analysis. We thank Jenny Lee for the purification of human milk lysozyme, Jishu Shi for his assistance in sequencing several peptides, and the UCLA Protein Microsequencing Facility, Mass Spectrometry Facility, and Clinical and Microbiological Laboratories for their invaluable help with this project. We also thank Edith M. Porter for continual discussions on microbiology and the properties of lysozyme.

REFERENCES

- Aho, H. J., R. Grenman, J. Sipila, H. Peuravuori, J. Hartikainen, and T. J. Nevalainen. 1997. Group II phospholipase A2 in nasal fluid, mucosa and paranasal sinuses. *Acta Otolaryngol.* **117**:860–863.
- Azen, E. A. 1993. Genetics of salivary protein polymorphisms. *Crit. Rev. Oral Biol. Med.* **4**:479–485.
- Biesbrock, A. R., M. S. Reddy, and M. J. Levine. 1991. Interaction of a salivary mucin-secretory immunoglobulin A complex with mucosal pathogens. *Infect. Immun.* **59**:3492–3497.
- Brögden, K. A., M. Ackermann, and K. M. Huttner. 1998. Detection of anionic antimicrobial peptides in ovine bronchoalveolar lavage fluid and respiratory epithelium. *Infect. Immun.* **66**:5948–5954.
- Ellison, R. T., III, and T. J. Giehl. 1991. Killing of gram-negative bacteria by lactoferrin and lysozyme. *J. Clin. Invest.* **88**:1080–1091.
- Ferguson, J. L., T. V. McCaffrey, E. B. Kern, and W. J. Martin. 1988. The effects of sinus bacteria on human ciliated nasal epithelium *in vitro*. *Otolaryngol. Head Neck Surg.* **98**:299–304.
- Fernlund, P., L. B. Granberg, and I. Larsson. 1996. Cloning of β -microseminoprotein of the rat: a rapidly evolving mucosal surface protein. *Arch. Biochem. Biophys.* **334**:73–82.
- Fleming, A. 1922. On a remarkable bacteriolytic element found in tissues and secretions. *Proc. R. Soc. Lond. (Biol.)* **93**:306–317.
- Ganz, T., M. E. Selsted, D. Szklarek, S. S. Harwig, K. Daher, D. F. Bainton, and R. I. Lehrer. 1985. Defensins: natural peptide antibiotics of human neutrophils. *J. Clin. Invest.* **76**:1427–1435.
- Ganz, T., and J. Weiss. 1997. Antimicrobial peptides of phagocytes and epithelia. *Semin. Hematol.* **34**:343–354.
- Hancock, R. E., L. M. Mutharia, L. Chan, R. P. Darveau, D. P. Speert, and G. B. Pier. 1983. *Pseudomonas aeruginosa* isolates from patients with cystic fibrosis: a class of serum-sensitive, nontypable strains deficient in lipopolysaccharide O side chains. *Infect. Immun.* **42**:170–177.
- Harder, J., J. Bartels, E. Christophers, and J.-M. Schroeder. 1997. A peptide antibiotic from human skin. *Nature* **387**:861–862.
- Imundo, L., J. Barasch, A. Prince, and Q. Al-Awqati. 1995. Cystic fibrosis epithelial cells have a receptor for pathogenic bacteria on their apical surface. *Proc. Natl. Acad. Sci. USA* **92**:3019–3023.
- Jensen, J. L., T. Xu, M. S. Lamkin, P. Brodin, H. Aars, T. Berg, and F. G. Oppenheim. 1994. Physiological regulation of the secretion of histatins and statherins in human parotid saliva. *J. Dent. Res.* **73**:1811–1817.
- Kaliner, M. A. 1991. Human nasal respiratory secretions and host defense. *Am. Rev. Respir. Dis.* **144**:S52–S56.
- Kaliner, M. A. 1992. Human nasal host defense and sinusitis. *J. Allergy Clin. Immunol.* **90**:424–430.
- Kaulbach, H. C., M. V. White, Y. Igarashi, B. K. Hahn, and M. A. Kaliner. 1993. Estimation of nasal epithelial lining fluid using urea as a marker. *J. Allergy Clin. Immunol.* **92**:457–465.
- Kluytmans, J., A. van Belkum, and H. Verbrugh. 1997. Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risks. *Clin. Microbiol. Rev.* **10**:505–520.
- Knowles, M. R., J. M. Robinson, R. E. Wood, C. A. Pue, W. M. Mentz, G. C. Wager, J. T. Gatzky, and R. C. Boucher. 1997. Ion composition of airway surface liquid of patients with cystic fibrosis as compared with normal and disease-control subjects. *J. Clin. Invest.* **100**:2588–2595.
- Krivan, H. C., D. D. Roberts, and V. Ginsburg. 1988. Many pulmonary pathogenic bacteria bind specifically to the carbohydrate sequence GalNAc beta 1-4Gal found in some glycolipids. *Proc. Natl. Acad. Sci. USA* **85**:6157–6161.
- Lee, I. H., Y. Cho, and R. I. Lehrer. 1997. Effects of pH and salinity on the antimicrobial properties of clavanins. *Infect. Immun.* **65**:2898–2903.
- Li, J. D., A. F. Dohrman, M. Gallup, S. Miyata, J. R. Gum, Y. S. Kim, J. A. Nadel, A. Prince, and C. B. Basbaum. 1997. Transcriptional activation of mucin by *Pseudomonas aeruginosa* lipopolysaccharide in the pathogenesis of cystic fibrosis lung disease. *Proc. Natl. Acad. Sci. USA* **94**:967–972.
- Lindahl, M., B. Stahlbom, and C. Tagesson. 1995. Two-dimensional gel electrophoresis of nasal and bronchoalveolar lavage fluids after occupational exposure. *Electrophoresis* **16**:1199–1204.
- Liu, L., L. Wang, H. P. Jia, C. Zhao, H. H. Q. Heng, B. C. Schutte, P. B. McCray, and T. Ganz. 1998. Structure and mapping of the human β -defensin HBD-2 gene and its expression at sites of inflammation. *Gene* **222**:237–244.
- Lorin, M. I., P. F. Gaerlan, and I. D. Mandel. 1972. Quantitative composition of nasal secretions in normal subjects. *J. Lab. Clin. Med.* **80**:275–281.
- Matsui, H., B. R. Grubb, R. Tarran, S. H. Randell, J. T. Gatzky, C. W. Davis, and R. C. Boucher. 1998. Evidence for periciliary liquid layer depletion, not abnormal ion composition, in the pathogenesis of cystic fibrosis airways disease. *Cell* **95**:1005–1015.
- McCray, P. B., J., and L. Bentley. 1997. Human airway epithelia express a β -defensin. *Am. J. Respir. Cell. Mol. Biol.* **16**:343–349.
- McNeely, T. B., and J. D. Coonrod. 1993. Comparison of the opsonic activity of human surfactant protein A for *Staphylococcus aureus* and *Streptococcus pneumoniae* with rabbit and human macrophages. *J. Infect. Dis.* **167**:91–97.
- Meredith, S. D., G. D. Raphael, J. N. Baraniuk, S. M. Banks, and M. A. Kaliner. 1989. The pathophysiology of rhinitis. III. The control of IgG secretion. *J. Allergy Clin. Immunol.* **84**:920–930.
- Osserman, E. F., and D. P. Lawlor. 1966. Serum and urinary lysozyme (muramidase) in monocytic and monomyelocytic leukemia. *J. Exp. Med.* **124**:921–952.
- Qu, X. D., K. C. Lloyd, J. H. Walsh, and R. I. Lehrer. 1996. Secretion of type II phospholipase A2 and cryptidin by rat small intestinal Paneth cells. *Infect. Immun.* **64**:5161–5165.
- Raphael, G. D., E. V. Jeney, J. N. Baraniuk, I. Kim, S. D. Meredith, and M. A. Kaliner. 1989. Pathophysiology of rhinitis. Lactoferrin and lysozyme in nasal secretions. *J. Clin. Invest.* **84**:1528–1535.
- Schagger, H., and G. von Jagow. 1987. Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for the separation of proteins in the range from 1 to 100 kDa. *Anal. Biochem.* **166**:368–379.
- Shuter, J., V. B. Hatcher, and F. D. Lowy. 1996. *Staphylococcus aureus* binding to human nasal mucin. *Infect. Immun.* **64**:310–318.
- Singh, P. K., H. P. Jia, K. Wiles, J. Hesselberth, L. Liu, B. D. Conway, E. P. Greenberg, E. V. Valore, M. J. Welsh, T. Ganz, B. F. Tack, and P. B. J. McCray. 1998. Production of β -defensins by human airway epithelia. *Proc. Natl. Acad. Sci. USA* **95**:14961–14966.
- Smith, A. 1997. Pathogenesis of bacterial bronchitis in cystic fibrosis. *Pediatr. Infect. Dis. J.* **16**:91–95.
- Smith, J. J., S. M. Travis, E. P. Greenberg, and M. J. Welsh. 1996. Cystic

- fibrosis airway epithelia fail to kill bacteria because of abnormal airway surface fluid. *Cell* **85**:229–236.
38. **Stadel, J. M., K. Hoyle, R. M. Naclerio, A. Roshak, and F. H. Chilton.** 1994. Characterization of phospholipase A2 from human nasal lavage. *Am. J. Respir. Cell. Mol. Biol.* **11**:108–113.
 39. **Steinberg, D. A., and R. I. Lehrer.** 1997. Designer assays for antimicrobial peptides. Disputing the “one-size-fits-all” theory. *Methods Mol. Biol.* **78**: 169–186.
 40. **Valore, E. V., C. H. Park, A. J. Quayle, K. R. Wiles, P. B. McCray, and T. Ganz.** 1998. Human β -defensin-1: an antimicrobial peptide of urogenital tissues. *J. Clin. Invest.* **101**:1633–1642.
 41. **Weiber, H., C. Lindstrom, H. Lilja, A. Bjartell, and P. Fernlund.** 1997. Immunohistochemical and *in situ* hybridization studies of β -microseminoprotein in the human gastric mucosa. *Histochem. J.* **29**:839–845.
 42. **Widdicombe, J. H., and J. G. Widdicombe.** 1995. Regulation of human airway surface liquid. *Respir. Physiol.* **99**:3–12.
 43. **Zar, H., L. Saiman, L. Quittell, and A. Prince.** 1995. Binding of *Pseudomonas aeruginosa* to respiratory epithelial cells from patients with various mutations in the cystic fibrosis transmembrane regulator. *J. Pediatr.* **126**:230–233.
 44. **Zhao, C. Q., I. Wang, and R. I. Lehrer.** 1996. Widespread expression of β -defensin HBD-1 in human secretory glands and epithelial cells. *FEBS Lett.* **396**:319–322.

Editor: J. R. McGhee