

REVIEW PAPER

From fibers to flowering to metabolites: unlocking hemp (*Cannabis sativa*) potential with the guidance of novel discoveries and tools

Francesco Pancaldi , Elma M. J. Salentijn, and Luisa M. Trindade*

Plant Breeding, Wageningen University & Research, Droevendaalsesteeg 1, 6708PB, Wageningen, The Netherlands

* Correspondence: luisa.trindade@wur.nl

Received 4 March 2024; Editorial decision 23 September 2024; Accepted 24 September 2024

Editor: Ricarda Jost, La Trobe University, Australia

Abstract

***Cannabis sativa* L. is an ancient crop, but its agricultural adoption has been interrupted to prevent the use of marijuana as a psychoactive drug. Nevertheless, hemp—the *C. sativa* type with low concentrations of intoxicating Δ^9 -tetrahydrocannabinoid—is experiencing a resurgence in interest due to loosened cultivation restrictions and its potential as a multipurpose bio-based crop. Hemp has valuable applications, including production of medicines from its non-intoxicating cannabinoids, food, medical, and industrial uses of its seed oil rich in polyunsaturated fatty acids, and production of fibers for textiles and industry from its stems. Recently, several hemp genomic and genetic resources have been developed, allowing significant expansion of our knowledge of major hemp traits, such as synthesis of cannabinoids, oil, and fibers, and regulation of flowering and sex determination. Still, hemp is an under-improved crop, and its development will depend on the ability to expand and collectively use the novel resources arising from fast advancements in bioinformatics and plant phenotyping. This review discusses current genetic and genomic knowledge of the most important hemp traits, and provides a perspective on how to further expand such knowledge and tackle hemp improvement with the most up-to-date tools for plant and hemp research.**

Keywords: Breeding tools, cannabinoids, fiber genetics, flowering genetics, hemp breeding, hemp genetics, hemp genomics, terpenes.

Introduction

Cannabis sativa L. is the only plant species within the *Cannabis* genus from the *Cannabaceae* family (Ren *et al.*, 2021). This species is an annual, herbaceous, diploid plant, with a genome of ~830 Mb made up of 10 chromosome pairs ($2n=20$) (Van Bakel *et al.*, 2011; Divashuk *et al.*, 2014). Of these, nine pairs are homomorphic autosomes, while the 10th pair groups semi-heteromorphic XX/XY sex chromosomes (Moliterni *et al.*, 2004; Petit *et al.*, 2020a; Prentout *et al.*, 2020). *Cannabis sativa* is naturally dioecious, with male (XY) and female (XX)

plants producing male and female flowers, respectively, which are fertilized through wind pollination (Salentijn *et al.*, 2019; Smart *et al.*, 2022). However, monoecious plants also occur, which are generally genetic females (XX) producing a variable number of both male and female flowers (Faux *et al.*, 2014; Salentijn *et al.*, 2019). *Cannabis sativa* has its center of origin in central Asia (Devkota, 2022), where it has been domesticated by humans since at least 6000 years ago (Hillig, 2005; Ren *et al.*, 2021). The initial use of this plant was for medicinal

purposes (Hillig, 2005; Liu *et al.*, 2017), but since ~3000 years ago it has also been used for fiber production (Liu *et al.*, 2017; Ren *et al.*, 2021). Nowadays, *C. sativa* is abundant worldwide, mostly in temperate regions, and is a promising but underimproved multipurpose crop that can be used to produce diverse products such as fibers, oil, cannabinoids, and terpenes (Andre *et al.*, 2016; Kovalchuk *et al.*, 2020; Devkota, 2022). Specifically, multipurpose hemp is the *C. sativa* type that can be cultivated and used for the aforementioned applications, due to the low amount of the intoxicating Δ^9 -tetrahydrocannabinoid (THC; <0.3% of inflorescence dry weight) (Small, 2015; Petit, 2020). Still, hemp plants can synthesize >100 types of cannabinoids, among which cannabidiolic acid (CBDA) is prevalent (Johnson, 2019; Lange and Zager, 2022). Hemp cannabinoids—especially CBD—have valuable medical applications in formulations to treat epilepsy, chronic pain, disorders from drugs abuse, anxiety, inflammation, and cancer (Blessing *et al.*, 2015; Überall, 2020; Morel *et al.*, 2021; O'Brien, 2022). Thus, these molecules constitute the basis of the modern use of hemp as a medicinal plant. Apart from cannabinoids, hemp seeds contain ~28–35% oil with high amounts of α -linolenic acid, γ -linolenic acid, tocopherols, and antioxidant polyphenols, which are all compounds with high value for use in food and cosmetic products (Cerino *et al.*, 2021; Devkota, 2022). Moreover, hemp oil can also be used to produce biofuels (Cerino *et al.*, 2021; Devkota, 2022). Finally, hemp biomass is also an excellent source of natural fibers, which mostly consist of bast fibers from hemp stems, and can be used to replace glass fibers, producing textiles, and synthesizing bioplastics and building materials, such as 'hempcrete' (Ranalli and Venturi, 2004; Andre *et al.*, 2016; Novakova and Sal, 2019).

Despite the potential of hemp as a multipurpose crop to produce several bio-based products, this species is currently underimproved and far from expressing its genetic potential for diverse applications (Salentijn *et al.*, 2015; Schluttenhofer and Yuan, 2017). This is mainly due to the restrictions to *C. sativa* cultivation to prevent illegal production of marijuana (i.e. *C. sativa* types with >0.3% THC) that for more than 80 years have also extensively blocked hemp research and breeding (Smart *et al.*, 2022). As such, the major available industrial hemp cultivars have been bred through conventional basic breeding methods, such as mass selection, without the use of any marker or genomic technology (Salentijn *et al.*, 2015; Schluttenhofer and Yuan, 2017). Still, large genetic variation for the content and quality of the major products extractable from hemp—fibers, cannabinoids, terpenes, and oil—has been recorded in multiple hemp populations (Kriese *et al.*, 2004; Galasso *et al.*, 2016; Petit *et al.*, 2020c; Johnson and Wallace, 2021; Stack *et al.*, 2021). Moreover, the resurgence of hemp research that has taken place over the last decades has produced a wealth of genomic and genetic resources—including multiple hemp genome assemblies (Van Bakel *et al.*, 2011; Braich *et al.*, 2020; Grassa *et al.*, 2021; Wei *et al.*, 2024), transcriptomic resources (Guerrero *et al.*, 2017; Braich *et al.*, 2019; Adal *et al.*, 2021;

Tang *et al.*, 2023), and molecular markers (Petit *et al.*, 2020a, b; Borin *et al.*, 2021)—that are extremely valuable to perform technology-driven breeding research on this crop. Finally, the increasing availability of genetic, genomic, and functional data on specific traits across all plant species, as well as of bioinformatic tools for large-scale comparative analyses, opens up novel scenarios for crop improvement through the translation of genetic information between species.

In the context just discussed, this paper reviews the advancements achieved in understanding the genetic basis of major target traits for the utilization of hemp in a bio-based economy. Moreover, a perspective is offered on how the further development of multipurpose hemp varieties can flourish by integrating those research advancements with new genomic and bioinformatic technologies. Major target hemp traits for a multipurpose use of hemp in a bio-based economy include the content and quality of hemp cannabinoids, terpenes, oil, and fibers. Moreover, regulation of sex determination and flowering behavior are also important traits, as they directly influence the expression of major industrial hemp characters, while in parallel allowing for specific breeding schemes (i.e. hybrid breeding) (Salentijn *et al.*, 2019). In the future, hemp improvement will integrate conventional breeding methods with novel genomic technologies, bioinformatic tools, and biotechnological approaches based on the genetics of target traits from both hemp and other crops.

The genetics of relevant hemp traits from the perspective of crop improvement

Cannabinoids

Maximizing the yield of non-intoxicating cannabinoids is a important objective in hemp research (Smart *et al.*, 2022), as these secondary metabolites—particularly CBDA—have several pharmaceutical applications with high economic value. *Cannabis sativa* synthesizes cannabinoids as phytoprotectants (Stack *et al.*, 2023), and the biosynthetic pathways of both THCA and CBDA largely overlap (Fig. 1) (Tahir *et al.*, 2021). Specifically, the common part of the THCA and CBDA biosynthetic pathway starts with the conversion of hexanoic acid into hexanoyl-CoA by an acyl-activating enzyme (AAE) (Lavery *et al.*, 2019; Kovalchuk *et al.*, 2020). Hexanoyl-CoA is then converted into cannabigerolic acid (CBGA) through a series of condensations, cyclations, and aromatizations catalyzed by the enzymes tetraketide synthase [TKS; also known as olivetol synthase (OLS)], olivetolic acid cyclase (OAC), and cannabigerolic acid synthase (CBGAS, an aromatic prenyltransferase) [reviewed by Tahir *et al.*, 2021 and Innes and Vergara, 2023]. CBGA is the precursor of both CBDA and THCA (Innes and Vergara, 2023), whose final synthesis is carried out by the oxidocyclase enzymes CBDA and THCA synthases (CBDAS and THCAS), respectively (Sirikantaramas *et al.*, 2004; Taura *et al.*, 2007).




Fig. 1. Genes taking part in biosynthesis of cannabidiolic acid and establishment of glandular trichomes in the context of hemp female flowers and glandular trichome development. (A) Drawing of a typical hemp female flower. (B) Microscopic structure of a typical hemp glandular trichome [adapted from Andre *et al.* (2016)]. (C) The biosynthetic pathway of cannabidiolic acid, with the main molecules (blue boxes) and genes/enzymes (arrows) involved, which takes place in the disk cells of glandular trichomes. AAE, acyl-activating enzyme; TKS, tetraketide synthase; OAC, olivetolic acid cyclase; CBGAS, cannabigerolic acid synthase; CBDAS, cannabidiolic acid synthase. (D) The major genes involved in the promotion (left) and repression (right) of glandular trichome development based on research on *Arabidopsis* and tomato, as described in the text.

Over the years, several genes encoding the enzymes of the cannabinoid pathway have been identified in hemp [see the works of Van Bakel *et al.* (2011), Laverty *et al.* (2019), McGarvey *et al.* (2020), Grassa *et al.* (2021), van Velzen and Schranz (2021), Fulvio *et al.* (2021), Innes and Vergara (2023), and Fulvio *et al.* (2023) for an overview of the results in this area over the years]. A major achievement was the identification of the genomic location of the oxidocyclase genes synthesizing either CBDA or THCA within a long terminal repeat (LTR)-rich region on *C. sativa* chromosome 7 (26–31 Mb) (Grassa *et al.*, 2021). At this locus, the presence of either *CBDAS* or *THCAS* functional genes underlies the production of either CBDA or THCA (Laverty *et al.*, 2019; McKernan *et al.*, 2020, Preprint; Grassa *et al.*, 2021). Specifically, *CBDAS* functionality appears pivotal for regulating CBDA production (McKernan *et al.*, 2020, Preprint) and, even though recent studies showed that both *CBDAS* and *THCAS* loci display little sequence variation in a large panel of *C. sativa* accessions (Lynch *et al.*, 2024, Preprint), the relationship between *CBDAS* genetic diversity and CBDA levels deserves further attention, as induced *CBDAS* mutations can significantly affect *CBDAS* catalytic activity (Dai *et al.*, 2024). In addition to *CBDAS*, relevant genetic patterns were recently uncovered for some of the genes acting upstream of cannabinoid oxidocyclases within the cannabinoid pathway, including *OLS* and *OAC* (Innes and Vergara, 2023). Specifically, some *OLS* and *OAC* tandem paralogs display copy number variation across different hemp accessions, and future research could investigate functional redundancy of variable gene copies, as well as their possible gene dosage effect on the cannabinoid pathway (Innes and Vergara, 2023).

The greatest proportion of hemp cannabinoids is synthesized by stalked glandular trichomes located on hemp female flowers, which are the hemp plant structures richest in CBDA (Fig. 1) (Mahlberg and Kim, 2004; Livingston *et al.*, 2020). These trichomes are formed by multicellular, large stalks supporting 12–16 disk cells and a secretory cavity, and their occurrence is highest on calyces and bracts of hemp female flowers (Livingston *et al.*, 2020; Tanney *et al.*, 2021). The disk cells synthesize CBDA by forming polarized ‘supercells’ that contain non-photosynthetic metabolic plastids (Livingston *et al.*, 2022). CBDA is then accumulated in the secretory cavities of the trichomes (Livingston *et al.*, 2022; Xie *et al.*, 2023). Because of the importance of glandular trichomes for CBDA production, elucidation of the factors influencing their number, size, and metabolic activity is a relevant research area (Tanney *et al.*, 2021), particularly because these properties are genetically controlled in hemp (Punja *et al.*, 2023). Bassolino *et al.* (2020) recently identified three *C. sativa* MYB transcription factor genes likely to be involved in trichome development regulation based on homology with *Arabidopsis thaliana*. This study highlights the relevance of knowledge about glandular trichomes obtained from other dicot species, given the overall scarce information on the genetics of hemp glandular trichome development (Tanney *et al.*, 2021). In this context, both tomato (*Solanum lycopersicum*) and sweet wormwood (*Artemisia annua*) are models to study the genetics underlying trichome development, and two transcription factor gene families—*R2R3-MYB* and *HD-ZIP IV*—regulate the initiation of glandular trichome formation in these species (Chalvin *et al.*, 2020). Specifically, the genes *AaMYB1*, *AaMIXTA1*, and

SlSIMX1 from the *R2R3-MYB* family, as well as the genes *AaHD1*, *AaHD8*, *SICD2*, and *SIWO* from the *HD-ZIP IV* family all positively influence glandular trichome development in *A. annua* (Matías-Hernández *et al.*, 2017; Yan *et al.*, 2017; Shi *et al.*, 2018) and tomato (Nadakuduti *et al.*, 2012; Yang *et al.*, 2015; Ewas *et al.*, 2016, 2017). Moreover, their overexpression leads to increased yield of glandular trichome metabolites in *A. annua* (Yan *et al.*, 2017; Shi *et al.*, 2018). Conversely, jasmonate signaling genes such as *AaJAZ8* or *SJJAZ2* are known to repress glandular trichome formation by down-regulating *R2R3-MYB* and *HD-ZIP IV* transcription factor genes (Yan *et al.*, 2017; Yu *et al.*, 2018). Finally, quantitative trait loci (QTLs) associated with variability in shape and density of glandular trichomes in tomato have also been mapped (Momotaz *et al.*, 2010; Bennewitz *et al.*, 2018). Overall, these genetic resources can serve as a starting point for translational genomics studies aimed at studying the regulation of glandular trichome development in hemp, by identifying hemp gene orthologs or genomic regions that share conserved blocks of genes in the same relative positions as compared with known QTLs (syntenic QTL regions). Such gene orthologs and syntenic QTL regions can in turn be targeted for functional studies or screenings of favorable allelic diversity for both trichome- and metabolic-related traits. In this regard, the recent works of Kundan *et al.* (2022) and Haiden *et al.* (2022) appear relevant, as they respectively identified 99 *R2R3-MYB* genes in the hemp genome, and showed the functionality of a *C. sativa* *MIXTA* gene homolog to *A. annua* to increase trichome density upon overexpression in tobacco leaves.

Terpenes

After cannabinoids, terpenes are another important class of hemp secondary metabolites. Terpenes determine the typical *C. sativa* aroma and, like cannabinoids, are synthesized and accumulated in stalked glandular trichomes (Sommano *et al.*, 2020). Terpenes can be used for multiple applications, including as aroma regulators and bioactive compounds in cosmetics, flavor additives in food, or bioactive molecules in pharmaceuticals thanks their antimicrobial, antioxidant, anti-inflammatory, and antidiabetic properties (Nuutinen, 2018; Chen and Pan, 2021). Furthermore, terpenes can be blended with CBD to enhance the medicinal properties of CBD itself—a phenomenon known as the ‘entourage effect’ (Ferber *et al.*, 2020; Anand *et al.*, 2021). At a chemical level, terpenes are a diverse group of hydrocarbon molecules made up of isoprene chains (Hanuš and Hod, 2020; Sommano *et al.*, 2020). Hemp trichomes can synthesize monoterpenes (two isoprene subunits), sesquiterpenes (three isoprene subunits), diterpenes (four isoprene subunits), and triterpenes (six isoprene subunits), with the vast majority of *C. sativa* terpenes being mono- and sesquiterpenes (Booth *et al.*, 2020; Hanuš and Hod, 2020; Sommano *et al.*, 2020).

Terpene synthases (TPSs) are responsible for terpene synthesis in all land plants (Chen *et al.*, 2011). In hemp, TPSs

cyclize the last intermediates of the mevalonate (MEV) and non-mevalonate (MEP) pathways—geranyl diphosphate and farnesyl diphosphate, respectively—into the variety of mono- and sesquiterpenes found in hemp (Booth *et al.*, 2020). The *TPS* gene family is large and diversified across all land plants, accounting for eight known *TPS* subfamilies—*TPS-a* to *TPS-h* (Chen *et al.*, 2011; Jiang *et al.*, 2019). Among these, the *TPS-a* and *TPS-b* genes are of particular relevance for hemp, as they synthesize mono- and sesquiterpenes, respectively (Booth *et al.*, 2020). Research showed that *TPS* genes form a large gene family in hemp, with a variable total number of genes across different accessions. Specifically, Booth *et al.* (2017, 2020) identified 13 and 19 *TPS* genes in the transcriptome of the hemp cultivar ‘Finola’ and in the draft genome of the cultivar ‘Purple Kush’, respectively (Van Bakel *et al.*, 2011). More recently, Allen *et al.* (2019) and Xu *et al.* (2024) mapped 55 and 41 distinct *TPS* genes in the complete *C. sativa* genomes from cultivars CBDRx (Grassa *et al.*, 2021) and ‘Jamaica Lion’ (McKernan *et al.*, 2020, Preprint), respectively. Finally, McKernan *et al.* (2020, Preprint) found extensive copy number variation for *TPS* genes potentially associated with variability in terpene profiles across 40 re-sequenced *C. sativa* accessions. As *TPS* genes are organized in genomic clusters, (proximal), gene duplications have driven the expansion and diversification of these genes in hemp (Booth *et al.*, 2020; Xu *et al.*, 2024). Most of the hemp *TPS* genes belong to the *TPS-a* and *TPS-b* subfamilies, in line with the prevalence of mono- and sesquiterpenes among hemp terpenes (Booth *et al.*, 2017, 2020; Xu *et al.*, 2024). Nevertheless, the genomic differentiation of *TPS* genes, reflected in their expression variability across different hemp accessions, seems key to determine the specific blends of hemp terpenes—including α -pinene, limonene, myrcene, and β -caryophyllene (Zager *et al.*, 2019; Booth *et al.*, 2020). As such, a further sampling of the genomic and transcriptomic diversity of hemp *TPS* genes, coupled with functional analyses on individual *TPS* copies, seems a promising strategy to acquire the genetic knowledge needed to modify hemp terpenes toward specific blends.

In addition to *TPS* genes, the other genes within the MEV and MEP pathways are also important to understand the genetics of hemp terpene biosynthesis, as well as to modify the synthesis of these molecules. These genes encode multiple enzymes that globally catalyze the conversion of pyruvate into geranyl diphosphate (MEP pathway) and of acetoacetyl-CoA into farnesyl diphosphate (MEV pathway) [see Booth *et al.* (2020) for an overview]. Notably, most of the MEP and MEV pathway genes are present in multiple copies within the hemp genome, often arranged in tandem clusters (Booth *et al.*, 2020). Furthermore, MEP and MEV pathway genes display co-expression with specific *TPS* genes during terpene synthesis in hemp (Zager *et al.*, 2019). As such, it would be relevant to study the variability in copy number and in the coordinated regulation of the MEP and MEV pathway genes in panels of multiple hemp accessions, and to correlate the results with overall




Fig. 2. Representation of the plant oil biosynthesis pathway and the genes involved. Orange stars next to genes indicate that hemp orthologs have been identified in the seed hemp genome by [Wei et al. \(2024\)](#). Blue stars indicate genes that are interesting hemp targets for screenings of allelic diversity and copy number variation, or genetic engineering. ACCase, acetyl-CoA carboxylase; MT, malonyl-CoA-ACP transferase; FAS, fatty acid synthase; KAS, ketoacyl-ACP synthase; SAD, stearoyl-ACP desaturase; FATA, acyl-ACP thioesterase A; FATB, acyl-ACP thioesterase B; LACS, long-chain acyl-CoA synthetase; FAE, fatty acid elongase; FAD2, fatty acid desaturase 2; FAD3, fatty acid desaturase 3.

terpene production and profiles. This could identify targets to increase terpene production through biotechnology.

Oil

Seed oil is another important product of industrial hemp, with several high-value applications in nutraceutical, cosmetic, and pharmaceutical industries ([Devkota, 2022](#); [Smart et al., 2022](#)). This is due to the hemp seed oil composition, which includes 50–70% linoleic acid (18:2 n -6), 15–25% α -linolenic acid (18:3 n -3), and ~9% γ -linolenic (18:3 n -6) and stearidonic acids (18:4 n -3) ([Matthäus and Brühl, 2008](#); [Rezvankhah et al., 2019](#)). These are all polyunsaturated fatty acids (PUFAs), which deliver health benefits upon consumption or when used in medicines against diseases such as glaucoma, cancer, inflammation, arthritis, and allergies ([Rezvankhah et al., 2019](#)). Moreover, tocopherols (i.e. vitamin E) and phytosterols—which are antioxidants and can reduce low-density lipoprotein (LDL)-cholesterol—are also found in high amounts in hemp seed oil (up to 800 mg kg⁻¹ tocopherols and ~5 g kg⁻¹ phytosterols) ([Oomah et al., 2002](#); [Matthäus and Brühl, 2008](#)). Industrial hemp varieties have typically not been bred for oil content and composition, which is the reason why hemp oil yield is often unstable, while oil quality is heterogeneous, even within cultivars ([Matthäus and Brühl, 2008](#)). As such, pursuing an increased and stable oil yield, as well as an optimized and stable oil composition, is a major target for research on industrial hemp.

So far, improvement of hemp oil-related traits has followed conventional methods of selective breeding, as in the case of the development of the ‘Finola’ variety ([Callaway and Laakkonen, 1997](#)). As plant oil yield and quality are typically

highly heritable traits ([Schierholt and Becker, 2001](#); [Bachlava et al., 2008](#); [Khan et al., 2010](#)), selective breeding can lead to significant improvements. Still, research in different oilseed crops showed that genetic manipulation of the oil pathway is also very effective both to increase seed oil content and yield, and to tailor oil composition to specific uses ([Savadi et al., 2017](#); [Porokhvinova et al., 2022](#)). In this context, the work of [Bielecka et al. \(2014\)](#) pioneered this research area in hemp, by showing that the silencing of two *FATTY ACID DESATURASE (FAD)* genes identified through homology with other oilseed dicots greatly changed the oil fatty acid profile of hemp ([Fig. 2](#)). More recently, [Wei et al. \(2024\)](#) released the reference genome of a major Chinese seed hemp cultivar, ‘Yushe’, which was used to identify 36 genes underlying hemp oil synthesis and covering the conventional plant oil biosynthetic pathway ([Fig. 2](#)). Moreover, it was shown that seed hemp experienced an expansion of the *STEAROYL-ACP DESATURASE (SAD)* and *FAD* gene families relative to other species, which might underpin its high seed oil content and high amounts of oil PUFAs ([Wei et al., 2024](#)). Further research could explore variability in *SAD* and *FAD* copy number across diverse hemp accessions and verify correlations with variability in seed oil content and composition. Moreover, reverse genetics studies such as that of [Bielecka et al. \(2014\)](#) could be replicated for other hemp oil genes. In this regard, *FATTY ACID ELONGASE (FAE)* represents an interesting target, as six *FAE* copies have been found in seed hemp ([Wei et al., 2024](#)) and its down-regulation increased the levels of oleic acid in *Brassicaceae* ([Tian et al., 2011](#); [Shi et al., 2015](#)). This, coupled with the overexpression of endogenous *FAD* genes, could lead to

increased PUFAs levels (Liu *et al.*, 2024), which can be a valuable target for seed hemp improvement.

Despite being valuable for food and industrial applications, PUFA-rich plant oils are prone to lipid peroxidation by reactive oxygen species (Ali *et al.*, 2022), making the oil unstable. This issue can be controlled by decreasing the PUFA content and raising the content of the oxidatively stable oleic acid (C18:1) in the oil, for example by down-regulating *FAD* genes (Kinney, 1998; Bielecka *et al.*, 2014). However, this approach also decreases the yield of valuable PUFAs that are naturally prevalent in hemp oil. An alternative approach can aim at increasing tocopherols in hemp oil, which act as antioxidants against PUFA peroxidation (Mène-Saffrané and DellaPenna, 2010; Boonnoy *et al.*, 2017; Ali *et al.*, 2022), have several food and industrial applications, and are naturally present in relatively high amounts in hemp oil. Several key genes underlying tocopherol biosynthesis have been found in different plants, including *A. thaliana*, carrot, maize, and tobacco [see Fritsche *et al.* (2017) for a detailed review]. Specifically, the genes *p-HYDROXYPHENYLPYRUVATE DIOXYGENASE 1 (PSD1)* and several *VITAMIN E (VTE)* loci deeply affect the synthesis and content of tocopherols in multiple plant organs, including seeds, and are targets for engineering these plant traits (Fritsche *et al.*, 2017). Wei *et al.* (2024) identified 16 genes involved in tocopherol biosynthesis in the seed hemp genome, including six homologs of the *A. thaliana VTE* loci. This finding is a valuable starting point for reverse genetics research aimed at understanding the synthesis of hemp tocopherols, and modulating their content in seed oil.

Fibers

The utilization of hemp for the production of fibers is one of the most ancient uses of this crop (Ren *et al.*, 2021; Smart *et al.*, 2022). Hemp fibers are essentially made up of cell walls and are located in the stems of the plants (Petit, 2020; Smart *et al.*, 2022). They comprise bast fibers—originating from the cambium, rich in crystalline cellulose and low in lignin—and hurd fibers (or shives)—forming the woody core of the hemp stems, with a high content of lignin and xylan-rich hemicellulosic polysaccharides (Van der Werf *et al.*, 1994; Salentijn *et al.*, 2015). Bast fibers are the most valuable hemp fibers, and the maximization of their yield and quality are pre-eminent targets of industrial hemp breeding (Salentijn *et al.*, 2015). The quality of bast fibers involves the composition of the cell walls constituting the fibers themselves, and is maximized when fibers are rich in crystalline cellulose with a low angle of cellulose microfibril deposition, and display a reduced content of lignin and xylan hemicelluloses (Salentijn *et al.*, 2015; Petit, 2020).

The yield and quality of hemp bast fibers have a strong genetic basis, as shown by the high heritability of these traits (Hennink, 1994; Petit *et al.*, 2020c). Nevertheless, the study of their genetic architecture is difficult, as both fiber yield and fiber quality are complex traits, controlled by several genomic

loci and genes with intricately pleiotropic effects (Petit *et al.*, 2020b). The study of the genetics underlying complex traits typically involves genetic mapping to identify regions and genes associated with patterns of trait variability in the progenies of crossings of phenotypically contrasting lines, or in phenotypically diverse panels of accessions. In this context, the work of Petit *et al.* (2020b) pioneered genetic mapping in hemp aimed at uncovering the genetic architecture of fiber yield and quality. These analyses uncovered 16 QTLs associated with hemp fiber yield and quality variability across diverse environments, as well as multiple candidate genes underlying the QTLs (Petit *et al.*, 2020b). Notably, a large proportion of these candidate genes is involved in the synthesis of lignin and pectins (Petit *et al.*, 2020b). As such, selection to reduce the content of these cell wall constituents may represent a promising strategy to improve the quality of bast fibers. Interestingly, protocols to screen these phenotypes in hemp at the scale of breeding programs have been developed (Petit *et al.*, 2019), and can thus be used in selective breeding schemes.

In the absence of further studies on the genetic basis of fiber yield and quality in hemp, information on this trait from other plant species can be highly valuable. As hemp bast fibers are essentially constituted of cell walls, research on the genetics underlying variability in cell wall composition conducted in grasses, forage crops, and model species such as *A. thaliana* is particularly relevant. Notably, these studies have largely elucidated the biosynthetic pathway of lignin (Bonawitz and Chapple, 2010; Yoon *et al.*, 2015) and have shown its amenability to genetic engineering (Ralph *et al.*, 2019; Carpita and McCann, 2020). Specifically, knockouts of key lignin genes such as *CINNAMATE 4-HYDROXYLASE (C4H)*, *4-COUMARATE:CoA LIGASE (4CL)*, *CINNAMATE 3-HYDROXYLASE (C3H)*, *CINNAMOYL REDUCTASE (CCR)*, and *CAFFEYOYL O-METHYLTRANSFERASE (COMT)* all determine considerable reductions of lignin content in the cell walls and tissues of diverse plant species (Leple *et al.*, 2007; Sattler *et al.*, 2009, 2012; Bjurhager *et al.*, 2010; Ralph *et al.*, 2012; Saballos *et al.*, 2012; Xiang *et al.*, 2017). Therefore, the mining of these genes in the hemp genome and their knockout could represent valuable strategies to reduce lignin content in hemp bast fibers. However, tissue-specific patterns of gene expression should probably be studied, as lignin genes such as those just mentioned typically form multigene families whose members take part in lignin deposition in specific tissues or cells (Chantreau *et al.*, 2014; Le Roy *et al.*, 2017; MacMillan *et al.*, 2017). Accordingly, in hemp it would be important to ensure that reduction of lignin content is confined to bast fibers, while extensive crop growth sustained by the woody core of hemp stems can still take place.

Regarding the possibility of increasing the crystalline cellulose content in hemp bast fibers, research in other crops showed that the genetics underlying this trait is more complex and less understood than those of lignin biosynthesis. Given the current status of plant cell wall research, the most likely relevant

target genes to modify these traits are the *CELLULOSE SYNTHASE (CESA)* genes, whose down-regulation and overexpression affect both the total cellulose content and cellulose crystallinity in different plant species (Harris *et al.*, 2009; Joshi *et al.*, 2011; Jayawardhane *et al.*, 2020). Moreover, multiple *CESA* genes are specifically responsible for the deposition of plant fibers, for example in cotton (Li *et al.*, 2013, 2016), making these genes relevant in the context of fiber improvement. Research demonstrated that cotton-specific patterns in the genomic diversification and physical organization of multiple *CESA* gene copies probably underly the massive deposition of fibers observed in cotton (Pancaldi *et al.*, 2022a). In this respect, mining the hemp homologs of functionally studied *CESA* genes could identify gene targets for screenings of allelic diversity and copy number variation, or for genetic engineering approaches to modify the cellulose properties of fibers. To conclude, it is noteworthy that the genomic organization of cell wall genes has co-evolved with their differential functionalization across both different members of multigene families and different plant species (Pancaldi *et al.*, 2022a, 2023a). Therefore, it is important to consider the ‘genomic contexts’ (Dewey, 2011) of specific gene copies when mining homologs of target genes between diverse plant clades.

Flowering behavior and sex determination

The regulation of flowering time and the mechanisms underlying sex determination are crucial hemp traits affecting plant growth and the yield and quality of the major harvestable hemp products (Faux *et al.*, 2013; Salentijn *et al.*, 2019). Regarding flowering, hemp is a short-day crop with a quantitative flowering phenology resulting from the interaction of genetic and metabolic factors with environmental variables such as specific photoperiod length and temperature (Salentijn *et al.*, 2019; Petit *et al.*, 2020a). The onset of flowering marks the end of vegetative growth, the start of bast fiber formation, and an intensified lignin deposition in stems (Keller *et al.*, 2001; Faux *et al.*, 2013; Liu *et al.*, 2015). In parallel, the time between flowering onset and hemp harvest is crucial to optimize the content and composition of seed oil and proteins (Amaducci *et al.*, 2015). Hemp has an XY sex chromosomes system, which underlies the prevalence of dioecious female (XX) and male (XY) plants (Salentijn *et al.*, 2019; Smart *et al.*, 2022). However, monoecious accessions also exist and are generally genetic females (XX) with quantitative variation in the proportion of female and male flowers, indicating that sex regulation is not entirely controlled by sex chromosomes (Faux *et al.*, 2016; Petit *et al.*, 2020a). Notably, monoecious cultivars display more uniform plant height and total stem and seed production (which are all traits suitable for dual harvest of stems and seeds), while the total production and quality of bast fibers are generally higher in dioecious accessions (Salentijn *et al.*, 2019). Controlling sex is therefore crucial to obtain hemp cultivars fitted to specific applications (Moliterni *et al.*, 2004; Salentijn *et al.*, 2019).

Current understanding of the genetics underlying hemp flowering behavior is not complete. Nevertheless, it is clear that this trait has a strong genetic basis and is highly heritable (Petit *et al.*, 2020c). Moreover, studies on separate species identified several genes involved in flowering time regulation, highlighting the complex nature of this trait in plants [see Freytes *et al.* (2021) for an updated review]. Flowering-related genes belong to different, interacting, signaling pathways, including the photoperiod-dependent flowering pathway, the temperature-dependent pathway, and the endogenous pathway that incorporates stimuli from internal hormones, metabolites, and aging factors (Zhang *et al.*, 2014; Salentijn *et al.*, 2019) (Fig. 3). Some of the genes in these pathways are well studied and conserved across plant species, such as *FLOWERING LOCUS T (FT)* or *FLORIGEN*; a major flowering promoter gene responsive to photoperiod and highly conserved across plants (Wickland and Hanzawa, 2015), *FLOWERING LOCUS C (FLC)*; a central flowering repressor responsive to temperature and regulated by multiple transcription factors (Ruelens *et al.*, 2013; Cheng *et al.*, 2017), *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS1 (SOC)*; a conserved gene responsive to multiple flowering pathways and promoting floral meristem (Immink *et al.*, 2012), and *SQUAMOSA-PROMOTER BINDING PROTEIN-LIKE* genes (*SPL* genes; involved in the endogenous flowering pathway and conserved across angiosperms) (Chen *et al.*, 2010; Preston and Hileman, 2013). Moreover, these primary flowering genes interact with several other genes that can be extensively conserved across plants as well, including *APETALA1*, *LEAFY*, *AGAMOUS-LIKE PROTEIN*, *VERNALIZATION (VRN)*, and *FLOWERING LOCUS-D* (Thomson and Wellmer, 2019; Sharma *et al.*, 2020) (Fig. 3). The increased availability of hemp genomic resources allowed investigations of the genetics of flowering regulation also in hemp. Petit *et al.* (2020a) performed the first genome-wide association study (GWAS) on this trait, anchored to the draft hemp genome of Van Bakel *et al.* (2011). More recently, other genetic mapping studies were performed by Woods *et al.* (2021), Toth *et al.* (2022), and Dowling *et al.* (2024). Overall, these studies found several hemp QTLs associated with variability in flowering time, with candidate genes often corresponding to known critical flowering regulators. For example, the flowering-related QTLs of Petit *et al.* (2020a) contain homologs of *FT*, *FLC*, *FLD*, *SOC1*, *SPL*, and *VRN* genes, highlighting the likely importance of these flowering genes also in hemp (Fig. 3). More recently, Toth *et al.* (2022) successfully mapped on hemp chromosome 1 two major loci—*AUTOFLOWER1* and *EARLY1*—controlling, respectively, photoperiod insensitivity and flowering time across multiple hemp populations. Importantly, molecular markers were also developed for both loci (Toth *et al.*, 2022), allowing high-throughput screening of hemp material for key flowering alleles to tailor novel hemp varieties to specific latitudes and agricultural rotations. Another locus—*AUTOFLOWER2*—was identified by Dowling *et al.* (2024) on hemp chromosome 8, most probably caused by a




Fig. 3. Representation of the major hemp flowering pathways (photoperiod-dependent, temperature-dependent, and endogenous) and the main genes involved in those pathways, based on genetic knowledge on flowering regulation for both hemp and other plant species. These genes represent interesting targets for hemp improvement. Arrows between genes indicate known genetic interactions among them. AP1, *Apetala1*, LFY, *Leafy*, AGL, *Agamous-like protein*; FT, *Flowering locus T* (or *Florigen*); FLC, *Flowering Locus C*; VRN, *Vernalization*; FLD, *Flowering Locus D*; SPL, *Squamosa Protein-like*; SOC, *Suppressor of Overexpression of Constans 1*.

hemp *FT* gene. Finally, [Steel *et al.* \(2023\)](#) performed comparative mapping of flowering-related loci from several hemp mapping studies, finding ample co-localizations for some QTLs and genes identified across them. Similarly, co-localization between QTLs for flowering, agronomic, and morphological hemp traits was also found in a separate QTL mapping study ([Woods *et al.*, 2021](#)). Overall, these observations suggest that translating the ample knowledge on flowering time regulation available in model species to hemp can represent a promising strategy to unveil the complete genetic architecture of hemp flowering time. Moreover, the reported co-localizations between QTLs for flowering time and other traits indicate opportunities for the parallel improvement of multiple traits in hemp, by careful selection of the diversity in initial material.

Similarly to flowering behavior, the genetic factors underlying hemp sex determination are also not fully understood. Still, this trait is also highly heritable, meaning that it has a strong genetic basis, whose core lies in the XY hemp sex chromosomes ([Faux *et al.*, 2014](#); [Petit *et al.*, 2020c](#)). The hemp system of XY chromosomes is one of the oldest among plants, firstly evolved in the hemp–hop ancestor 20–25 million years ago ([Prentout *et al.*, 2020, 2021](#)). The X and Y chromosomes of hemp differ in size, with the Y chromosome being ~40 Mbp larger than the X chromosome due to a loading of long interspersed nuclear element (LINE)-like retrotransposons ([Sakamoto *et al.*, 1998](#); [Van Bakel *et al.*, 2011](#); [Divashuk *et al.*, 2014](#)). The accumulation of LINE-like retrotransposons in the

hemp Y chromosome underlies the suppression of recombination over a portion of the X–Y chromosomes, as well as their still-evolving structural heteromorphism ([Moliteri *et al.*, 2004](#); [Vergara *et al.*, 2016](#); [Kovalchuk *et al.*, 2020](#)), despite the occurrence of a relatively large pseudo-autosomal recombining region ([Peil *et al.*, 2003](#); [Prentout *et al.*, 2021](#)). Nevertheless, hemp sex determination seems to rely on the X-to-autosomes balance rather than on a purely Y-active mechanism ([Faux *et al.*, 2014](#)). Moreover, the occurrence of variable proportions of male and female flowers in female (XX) monoecious hemp cultivars suggests interaction between sex chromosomes and other regions of the hemp genome to determine the phenotypic expression of hemp sex ([Faux *et al.*, 2016](#); [Salentijn *et al.*, 2019](#); [Petit *et al.*, 2020a](#)). These observations explain why the mapping of stable male-linked sex markers is a notoriously difficult task in hemp ([Kovalchuk *et al.*, 2020](#)), even if high-throughput genotyping methods have recently allowed the development of effective assays ([Toth *et al.*, 2020](#); [Torres *et al.*, 2022](#)). Moreover, studying the genes located on the hemp sex chromosomes both on their own and in relation to other targets on the autosomal chromosomes could be promising to depict the mechanisms underlying hemp sex determination. In this direction, Prentout and co-workers applied comparative RNA-seq analysis to identify a common set of 112 sex-linked genes across hemp and hop ([Prentout *et al.*, 2020, 2021](#)), which appear particularly relevant to identify targets underlying the X and Y chromosome-dependent components of sex

determination in hemp. An RNA-seq approach was also used by Adal *et al.* (2021), who identified ~200 genes displaying up-regulation upon chemical induction of male flowers in genetically female *C. sativa* plants, including genes involved in anther and pollen development, as well as hormone signaling. Genes related to hormone signaling, including auxin- and gibberellin-responsive transcription factors and *bZIP* genes regulating auxin and gibberellin balances, were also found to be related to hemp sex determination by Petit *et al.* (2020a), who performed a GWAS on this trait in a hemp panel including monoecious cultivars. Overall, these studies show that integrating genomic, transcriptomic, and genetic mapping approaches at the whole-genome level retains the potential to further dissect the genetic architecture of sex determination in hemp. As such, the combined profiling of the expression of hemp genes from sex and autosomal chromosomes, along with their comparative functional analysis in monoecious male and female accessions, could be performed to reveal interactive networks underlying hemp sex determination.

The future of hemp improvement guided by novel tools

Following the resurgence in interest toward hemp as a multipurpose crop and the loosened restrictions to hemp cultivation in some countries (Devkota, 2022; Rathi *et al.*, 2022), the research conducted over the last decades have been of paramount importance in supporting hemp improvement. However, hemp currently remains underimproved, far from reaching its potential as a multipurpose crop (Salentijn *et al.*, 2015; Smart *et al.*, 2022). Specifically, great efforts in the development of genetic and genomic hemp resources have been made, but the genetic knowledge on major traits is still incomplete. Moreover, the translation of fundamental hemp research into superior marketed varieties has been limited so far (Salentijn *et al.*, 2015; Smart *et al.*, 2022).

Further study and improvement of different hemp traits will require different strategies, as the genetic factors underlying different characters differ significantly in complexity, genomic structure, and level of understanding. For example, the application of genomic selection coupled with high-throughput CBDA phenotyping could be explored to attain increased CBDA yields, by selecting favorable alleles at genomic loci involved in cannabinoid biosynthesis (McKernan *et al.*, 2020, Preprint; Innes and Vergara, 2023; Lynch *et al.*, 2024, Preprint). So far, genomic selection has not been performed in hemp, but the availability of multiple high-quality genome assemblies (Grassa *et al.*, 2021; Wei *et al.*, 2024) and of ever cheaper genotyping technologies allows for pursuing this goal. Alternatively, methods for targeted sequencing [see, for example, Scaglione *et al.* (2019)] could also be applied by restricting genotyping to CBDA-associated loci while increasing the accuracy of detecting and characterizing diverse alleles with a positive

effect on CBDA synthesis. Regarding phenotyping, near infrared spectroscopy is emerging as a valuable high-throughput methodology to analyze the content of different cannabinoids—including CBDA and THCA—in diverse types of hemp material (Callado *et al.*, 2018; Birenboim *et al.*, 2022; Tran *et al.*, 2023), allowing the screening of large populations for the content of these compounds. Finally, hemp polyploidization through *in vitro* approaches or crossing of genotypes with different ploidy levels can also be investigated further as a strategy to increase hemp CBDA yield. In principle, polyploid hemp accessions are expected to display increased CBDA concentrations and/or CBDA yield (Crawford *et al.*, 2021; Suchoff *et al.*, 2024). However, the studies performed are not conclusive in this respect, deserving further research (Mansouri and Bagheri, 2017; Crawford *et al.*, 2021). Nevertheless, the creation of sterile triploid hemp genotypes appears to be a good strategy to prevent reduction of CBDA yield due to uncontrolled cross-pollination (Kurtz *et al.*, 2020).

In addition to improving CBDA yield, the modulation of the content and fatty acid composition of hemp seed oil is also an essential trait, particularly to fit hemp oil to specific uses, such as food and cosmetic applications (for which high amounts of PUFAs, particularly linoleic and α -linolenic fatty acids, is desirable) or biofuel production (for which a high amount of oxidative stable fatty acids, such as oleic acid, represents an important target) (Callaway, 2004; Vogl *et al.*, 2004; Carlsson *et al.*, 2011). In this regard, similarly to what was discussed for cannabinoids, the screening of sequence and structural genetic variation through genotyping-based or whole-genome resequencing methods can also be applied to the loci underlying hemp seed oil synthesis (Wei *et al.*, 2024). Specifically, favorable alleles, copy number variation, or presence/absence variation could be searched for genes such as *SAD*, *FAD*, and *FAE*, which are key to regulate the oil content of oleic acid, PUFAs, and elongated fatty acids, respectively (Bates *et al.*, 2013; Bielecka *et al.*, 2014; Wei *et al.*, 2024). For example, copy number expansion or selection for more active alleles at the *SAD* loci should favor the synthesis of oleic acid for biofuel applications (Wang *et al.*, 2024). The same would be expected by selecting for reduced copy number and non-functional alleles at the hemp *FAD* and *FAE* loci (Carlsson *et al.*, 2011; Bielecka *et al.*, 2014). Conversely, boosting the activity of *FAD* genes can promote the accumulation of PUFAs, which would be beneficial for food or cosmetic applications (Wang *et al.*, 2021). The achievement of these goals will be facilitated by using high-throughput methods for phenotyping seed oil content and composition in large plant populations. For this purpose, both NMR and infrared spectroscopy are well-established methodologies across different plant species (Jasinski *et al.*, 2016; Melchinger *et al.*, 2018; Anderson *et al.*, 2019), and a protocol for their use in hemp has recently been developed (Siudem *et al.*, 2019). Finally, the generally high heritability of seed oil content and composition (Schierholt and Becker, 2001; Bachlava *et al.*, 2008; Khan *et al.*, 2010) allows

for crossing superior accessions identified through genetic screening and directly operates genomically informed selection of superior material from progenies. To this aim, it would also be worth developing seed hemp mapping populations both to evaluate genes in the oil pathway that are of particular relevance to determine desired patterns of oil content and quality variation, as well as to mine molecular markers that can be used in selective breeding to deal with linkage drag. These types of studies are currently missing for hemp seed oil traits.

Genomic engineering aimed at boosting or constraining specific branches of the hemp oil pathway is also a valid strategy to improve seed oil content or composition. This approach would be particularly suited to optimize hemp seed oil properties in plant material already improved for other traits such as CBDA profile, which could be difficult to keep genetically stable while breeding for other traits (Ingvarsen and Brinch-Pedersen, 2023). It is noteworthy that different tools to enable genomic engineering in hemp have recently been developed. These include the achievement of successful transformation and regeneration of hemp plants from hypocotyls (Galán-Ávila *et al.*, 2021), despite hemp's notorious recalcitrance to these techniques (Ingvarsen and Brinch-Pedersen, 2023). Moreover, Zhang *et al.* (2021) recently reported the first successful example of CRISPR/Cas9-mediated genome editing in hemp, consisting of the knockout of the *PHYTOENE DESATURASE* gene. Altogether, these methods could be applied in *cis*-genesis or gene editing experiments to overexpress one or more genes participating in oleic acid synthesis in plastids, from *ACETYL-CoA CARBOXYLASE* to *SAD* genes, as their overexpression typically translates to higher seed oil content. In parallel, *FAD* genes could be either down-regulated or overexpressed, to aim at higher oleic acid or PUFAs in hemp oil. This type of research could also make the oil pathway a model for the development of genome editing techniques in hemp.

Compared with cannabinoids and oil, knowledge of the genetics underlying fiber yield and quality, flowering behavior, and sex determination in hemp is more fragmented. In this sense, further genetic mapping studies should be performed, as they have proven to be valuable tools to identify genes and markers for these traits in hemp (Faux *et al.*, 2016; Petit *et al.*, 2020a, b; Toth *et al.*, 2022). Moreover, in parallel to 'conventional' QTL mapping and genome-wide association approaches, bulk segregant analysis (BSA) should also be considered as an approach for variable hemp traits that probably rely on a limited number of loci, such as sex determination in monoecious hemp accessions (Salentijn *et al.*, 2019). For this trait, BSA between pools of plants displaying extremely high proportions of male or female flowers, respectively, could reveal relevant underlying loci. In turn, comparative genomics could be used to further characterize those loci based on functional information from other species. More broadly, the translation of characterized genomic

loci from (model) plants to a crop such as hemp is also a promising approach to uncover the genetics of complex traits in *C. sativa*. In this respect, pipelines have recently been developed for projecting multiple genomic loci underlying quantitative traits in diverse model species to target crops, as well as for analyzing allelic variation of the projected loci in the target crops (Pancaldi *et al.*, 2022b, 2023b). These strategies exploit large-scale and high-throughput analysis of genome synteny (Zhao and Schranz, 2017) to translate entire syntenically conserved QTLs across species (Pancaldi *et al.*, 2023b) and identify underlying functionally conserved genes based on conserved 'genomic contexts' (Dewey, 2011; Zhao and Schranz, 2017; Pancaldi *et al.*, 2023b). Collectively, these methods can quickly identify conserved loci and genes underlying traits of interest in understudied crops, and characterize (favorable) variation at such loci by using, for example, whole-genome re-sequencing data. In turn, accessions carrying superior alleles can be selected for by targeted genotyping or sequencing of diverse populations and included in breeding programs, accelerating crop improvement. Breeding gain can be maximized if recurrent breeding schemes are applied, with the crossing of novel superior material with well-performing lines (Smart *et al.*, 2022). In this respect, speed breeding can also be considered to maximize the number of crosses and generations per year, and a method to apply speed breeding in hemp has recently been developed (Schilling *et al.*, 2023).

Conclusion

Hemp is a multipurpose crop with high potential to fuel diverse valuable industrial applications in a developing bio-based economy. From this perspective, the recent loosening of the restrictions on hemp cultivation in different countries has acted as a substantial boost to research and commercial interest in this crop. Accordingly, the last decades have already seen unprecedented development of genetic, genomic, and breeding resources for hemp. These developments have provided many new insights into important traits for successfully adopting hemp in agriculture and its products in industries. This review has highlighted future research areas, and proposed approaches in fundamental and applied genetics to further advance hemp. Unlocking the potential of this crop will require the integration of resources and tools in creative research efforts to deliver new insights in an effective, rapid, and cost-effective manner. For this purpose, all the diverse resources and tools available on key traits for hemp and other crops need to be collectively analyzed with innovative bioinformatic approaches, and the insights obtained should be translated into applied achievements by using the most cutting-edge breeding strategies and tools—from precise genomic engineering to speed breeding.

Author contributions

FP: writing, with input from EMJS and LMT. All authors read and approved the final manuscript.

Conflict of interest

The authors declare no conflict of interest.

Funding

This research was part of a project that received funding from the European Union's Horizon 2020 Research and Innovation Program MIDAS (Marginal lands, industrial crops and innovative bio-based value chains) project, under grant agreement no. 101082070.

References

- Adal AM, Doshi K, Holbrook L, Mahmoud SS. 2021. Comparative RNA-Seq analysis reveals genes associated with masculinization in female *Cannabis sativa*. *Planta* **253**, 1–17.
- Ali E, Hussain S, Hussain N, Kakar KU, Shah JM, Zaidi SHR, Jan M, Zhang K, Khan MA, Imtiaz M. 2022. Tocopherol as plant protector: an overview of tocopherol biosynthesis enzymes and their role as antioxidant and signaling molecules. *Acta Physiologicae Plantarum* **44**, 20.
- Allen KD, McKernan K, Pauli C, Roe J, Torres A, Gaudino R. 2019. Genomic characterization of the complete terpene synthase gene family from *Cannabis sativa*. *PLoS One* **14**, e0222363.
- Amaducci S, Scordia D, Liu F, Zhang Q, Guo H, Testa G, Cosentino S. 2015. Key cultivation techniques for hemp in Europe and China. *Industrial Crops and Products* **68**, 2–16.
- Anand U, Pacchetti B, Anand P, Sodergren MH. 2021. Cannabis-based medicines and pain: a review of potential synergistic and entourage effects. *Pain Management* **11**, 395–403.
- Anderson JV, Wittenberg A, Li H, Berti MT. 2019. High throughput phenotyping of *Camelina sativa* seeds for crude protein, total oil, and fatty acids profile by near infrared spectroscopy. *Industrial Crops and Products* **137**, 501–507.
- Andre CM, Hausman J-F, Guerriero G. 2016. *Cannabis sativa*: the plant of the thousand and one molecules. *Frontiers in Plant Science* **7**, 19.
- Bachlava E, Burton JW, Brownie C, Wang S, Auclair J, Cardinal AJ. 2008. Heritability of oleic acid content in soybean seed oil and its genetic correlation with fatty acid and agronomic traits. *Crop Science* **48**, 1764–1772.
- Bassolino L, Buti M, Fulvio F, Pennesi A, Mandolino G, Milc J, Francia E, Paris R. 2020. In silico identification of MYB and bHLH families reveals candidate transcription factors for secondary metabolic pathways in *Cannabis sativa* L. *Plants* **9**, 1540.
- Bates PD, Stymne S, Ohlrogge J. 2013. Biochemical pathways in seed oil synthesis. *Current Opinion in Plant Biology* **16**, 358–364.
- Bennewitz S, Bergau N, Tissier A. 2018. QTL mapping of the shape of type VI glandular trichomes in tomato. *Frontiers in Plant Science* **9**, 1421.
- Bielecka M, Kaminski F, Adams I, Poulson H, Sloan R, Li Y, Larson TR, Winzer T, Graham IA. 2014. Targeted mutation of $\Delta 12$ and $\Delta 15$ desaturase genes in hemp produce major alterations in seed fatty acid composition including a high oleic hemp oil. *Plant Biotechnology Journal* **12**, 613–623.
- Birenboim M, Kengisbuch D, Chalupowicz D, Maurer D, Barel S, Chen Y, Fallik E, Paz-Kagan T, Shimshoni JA. 2022. Use of near-infrared spectroscopy for the classification of medicinal cannabis cultivars and the prediction of their cannabinoid and terpene contents. *Phytochemistry* **204**, 113445.
- Bjurhager I, Olsson A-M, Zhang B, Gerber L, Kumar M, Berglund LA, Burgert I, Sundberg B, Salmén L. 2010. Ultrastructure and mechanical properties of *Populus* wood with reduced lignin content caused by transgenic down-regulation of cinnamate 4-hydroxylase. *Biomacromolecules* **11**, 2359–2365.
- Blessing EM, Steenkamp MM, Manzanares J, Marmar CR. 2015. Cannabidiol as a potential treatment for anxiety disorders. *Neurotherapeutics* **12**, 825–836.
- Bonawitz ND, Chapple C. 2010. The genetics of lignin biosynthesis: connecting genotype to phenotype. *Annual Review of Genetics* **44**, 337–363.
- Boonnoy P, Karttunen M, Wong-Ekkabut J. 2017. Alpha-tocopherol inhibits pore formation in oxidized bilayers. *Physical Chemistry Chemical Physics* **19**, 5699–5704.
- Booth JK, Page JE, Bohlmann J. 2017. Terpene synthases from *Cannabis sativa*. *PLoS One* **12**, e0173911.
- Booth JK, Yuen MM, Jancsik S, Madilao LL, Page JE, Bohlmann J. 2020. Terpene synthases and terpene variation in *Cannabis sativa*. *Plant Physiology* **184**, 130–147.
- Borin M, Palumbo F, Vannozi A, Scariolo F, Sacilotto GB, Gazzola M, Barcaccia G. 2021. Developing and testing molecular markers in *Cannabis sativa* (hemp) for their use in variety and dioecy assessments. *Plants* **10**, 2174.
- Braich S, Baillie RC, Jewell LS, Spangenberg GC, Cogan NO. 2019. Generation of a comprehensive transcriptome atlas and transcriptome dynamics in medicinal cannabis. *Scientific Reports* **9**, 16583.
- Braich S, Baillie RC, Spangenberg GC, Cogan NO. 2020. A new and improved genome sequence of *Cannabis sativa*. *Gigabyte* **2020**, gigabyte10.
- Callado CS-C, Núñez-Sánchez N, Casano S, Ferreira-Vera C. 2018. The potential of near infrared spectroscopy to estimate the content of cannabinoids in *Cannabis sativa* L.: a comparative study. *Talanta* **190**, 147–157.
- Callaway J. 2004. Hempseed as a nutritional resource: an overview. *Euphytica* **140**, 65–72.
- Callaway J, Laakkonen T. 1997. Cultivation of Cannabis oil seed varieties in Finland. *Journal of the International Hemp Association* **3**, 32–34.
- Carlsson AS, Yilmaz JL, Green AG, Stymne S, Hofvander P. 2011. Replacing fossil oil with fresh oil—with what and for what? *European Journal of Lipid Science and Technology* **113**, 812–831.
- Carpita NC, McCann MC. 2020. Redesigning plant cell walls for the biomass-based bioeconomy. *Journal of Biological Chemistry* **295**, 15144–15157.
- Cerino P, Buonerba C, Cannazza G, D'Auria J, Ottoni E, Fulgione A, Di Stasio A, Pierri B, Gallo A. 2021. A review of hemp as food and nutritional supplement. *Cannabis and Cannabinoid Research* **6**, 19–27.
- Chalvin C, Drevensek S, Dron M, Bendahmane A, Boualem A. 2020. Genetic control of glandular trichome development. *Trends in Plant Science* **25**, 477–487.
- Chantreau M, Portelette A, Dauwe R, et al. 2014. Ectopic lignification in the flax lignified bast fiber1 mutant stem is associated with tissue-specific modifications in gene expression and cell wall composition. *The Plant Cell* **26**, 4462–4482.
- Chen C, Pan Z. 2021. Cannabidiol and terpenes from hemp—ingredients for future foods and processing technologies. *Journal of Future Foods* **1**, 113–127.
- Chen F, Tholl D, Bohlmann J, Pichersky E. 2011. The family of terpene synthases in plants: a mid-size family of genes for specialized metabolism that is highly diversified throughout the kingdom. *The Plant Journal* **66**, 212–229.
- Chen X, Zhang Z, Liu D, Zhang K, Li A, Mao L. 2010. SQUAMOSA promoter-binding protein-like transcription factors: star players for plant growth and development. *Journal of Integrative Plant Biology* **52**, 946–951.

- Cheng J-Z, Zhou Y-P, Lv T-X, Xie C-P, Tian C-E.** 2017. Research progress on the autonomous flowering time pathway in Arabidopsis. *Physiology and Molecular Biology of Plants* **23**, 477–485.
- Crawford S, Rojas BM, Crawford E, Otten M, Schoenenberger TA, Garfinkel AR, Chen H.** 2021. Characteristics of the diploid, triploid, and tetraploid versions of a cannabigerol-dominant F1 hybrid industrial hemp cultivar, *Cannabis sativa* ‘Stem Cell CBG’. *Genes* **12**, 923.
- Dai L, Niu T, Luo R, et al.** 2024. Improvement of cannabidiolic acid synthetase activity through molecular docking and site-directed mutagenesis. *Industrial Crops and Products* **208**, 117860.
- Devkota HP.** 2022. Hemp (*Cannabis sativa* L.)—taxonomy, distribution and uses. In: Belwal T, Belwal NC, eds. *Revolutionizing the potential of hemp and its products in changing the global economy*. Cham: Springer, 1–10.
- Dewey CN.** 2011. Positional orthology: putting genomic evolutionary relationships into context. *Briefings in Bioinformatics* **12**, 401–412.
- Divashuk MG, Alexandrov OS, Razumova OV, Kirov IV, Karlov GI.** 2014. Molecular cytogenetic characterization of the dioecious *Cannabis sativa* with an XY chromosome sex determination system. *PLoS One* **9**, e85118.
- Dowling CA, Shi J, Toth JA, Quade MA, Smart LB, McCabe PF, Melzer R, Schilling S.** 2024. A FLOWERING LOCUS T ortholog is associated with photoperiod-insensitive flowering in hemp (*Cannabis sativa* L.). *The Plant Journal* **119**, 383–403.
- Ewas M, Gao Y, Ali F, Nishawy EM, Shahzad R, Subthain H, Amar M, Martin C, Luo J.** 2017. RNA-seq reveals mechanisms of SIMX1 for enhanced carotenoids and terpenoids accumulation along with stress resistance in tomato. *Science Bulletin* **62**, 476–485.
- Ewas M, Gao Y, Wang S, et al.** 2016. Manipulation of SIMX1 for enhanced carotenoids accumulation and drought resistance in tomato. *Science Bulletin* **61**, 1413–1418.
- Faux A-M, Berhin A, Dauguet N, Bertin P.** 2014. Sex chromosomes and quantitative sex expression in monoecious hemp (*Cannabis sativa* L.). *Euphytica* **196**, 183–197.
- Faux A-M, Draye X, Flamand M-C, Occre A, Bertin P.** 2016. Identification of QTLs for sex expression in dioecious and monoecious hemp (*Cannabis sativa* L.). *Euphytica* **209**, 357–376.
- Faux A-M, Draye X, Lambert R, D’Andrimont R, Raulier P, Bertin P.** 2013. The relationship of stem and seed yields to flowering phenology and sex expression in monoecious hemp (*Cannabis sativa* L.). *European Journal of Agronomy* **47**, 11–22.
- Ferber SG, Namdar D, Hen-Shoval D, Eger G, Koltai H, Shoval G, Shbiro L, Weller A.** 2020. The ‘entourage effect’: terpenes coupled with cannabinoids for the treatment of mood disorders and anxiety disorders. *Current Neuropharmacology* **18**, 87–96.
- Freytes SN, Canelo M, Cerdán PD.** 2021. Regulation of flowering time: when and where? *Current Opinion in Plant Biology* **63**, 102049.
- Fritsche S, Wang X, Jung C.** 2017. Recent advances in our understanding of tocopherol biosynthesis in plants: an overview of key genes, functions, and breeding of vitamin E improved crops. *Antioxidants* **6**, 99.
- Fulvio F, Paris R, Montanari M, et al.** 2021. Analysis of sequence variability and transcriptional profile of cannabinoid synthase genes in *Cannabis sativa* L. chemotypes with a focus on cannabichromenic acid synthase. *Plants* **10**, 1857.
- Fulvio F, Righetti L, Minervini M, Moschella A, Paris R.** 2023. The B1080/B1192 molecular marker identifies hemp plants with functional THCA synthase and total THC content above legal limit. *Gene* **858**, 147198.
- Galán-Ávila A, Gramazio P, Ron M, Prohens J, Herraiz FJ.** 2021. A novel and rapid method for *Agrobacterium*-mediated production of stably transformed *Cannabis sativa* L. plants. *Industrial Crops and Products* **170**, 113691.
- Galasso I, Russo R, Mapelli S, Ponzoni E, Brambilla IM, Battelli G, Reggiani R.** 2016. Variability in seed traits in a collection of *Cannabis sativa* L. genotypes. *Frontiers in Plant Science* **7**, 688.
- Grassa CJ, Weiblen GD, Wenger JP, Dabney C, Poplawski SG, Motley TS, Michael TP, Schwartz CJ.** 2021. A new cannabis genome assembly associates elevated cannabidiol (CBD) with hemp introgressed into marijuana. *New Phytologist* **230**, 1665–1679.
- Guerriero G, Behr M, Legay S, Mangeot-Peter L, Zorzan S, Ghoniem M, Hausman J-F.** 2017. Transcriptomic profiling of hemp bast fibres at different developmental stages. *Scientific Reports* **7**, 4961.
- Haiden SR, Apicella PV, Ma Y, Berkowitz GA.** 2022. Overexpression of CsMIXTA, a transcription factor from *Cannabis sativa*, increases glandular trichome density in tobacco leaves. *Plants* **11**, 1519.
- Hanuš LO, Hod Y.** 2020. Terpenes/terpenoids in cannabis: are they important? *Medical Cannabis and Cannabinoids* **3**, 25–60.
- Harris D, Stork J, Debolt S.** 2009. Genetic modification in cellulose-synthase reduces crystallinity and improves biochemical conversion to fermentable sugar. *GCB Bioenergy* **1**, 51–61.
- Hennink S.** 1994. Optimisation of breeding for agronomic traits in fibre hemp (*Cannabis sativa* L.) by study of parent–offspring relationships. *Euphytica* **78**, 69–76.
- Hillig KW.** 2005. Genetic evidence for speciation in *Cannabis* (Cannabaceae). *Genetic Resources and Crop Evolution* **52**, 161–180.
- Immink RG, Posé D, Ferrario S, et al.** 2012. Characterization of SOC1’s central role in flowering by the identification of its upstream and downstream regulators. *Plant Physiology* **160**, 433–449.
- Ingvarsdson CR, Brinch-Pedersen H.** 2023. Challenges and potentials of new breeding techniques in *Cannabis sativa*. *Frontiers in Plant Science* **14**, 1154332.
- Innes PA, Vergara D.** 2023. Genomic description of critical cannabinoid biosynthesis genes. *Botany* **101**, 270–283.
- Jasinski S, Lécureuil A, Durandet M, Bernard-Moulin P, Guerche P.** 2016. Arabidopsis seed content QTL mapping using high-throughput phenotyping: the assets of near infrared spectroscopy. *Frontiers in Plant Science* **7**, 1682.
- Jayawardhane KN, Singer SD, Ozga JA, Rizvi SM, Weselake RJ, Chen G.** 2020. Seed-specific down-regulation of Arabidopsis CELLULOSE SYNTHASE 1 or 9 reduces seed cellulose content and differentially affects carbon partitioning. *Plant Cell Reports* **39**, 953–969.
- Jiang S-Y, Jin J, Sarojam R, Ramachandran S.** 2019. A comprehensive survey on the terpene synthase gene family provides new insight into its evolutionary patterns. *Genome Biology and Evolution* **11**, 2078–2098.
- Johnson MS, Wallace JG.** 2021. Genomic and chemical diversity of commercially available high-CBD industrial hemp accessions. *Frontiers in Genetics* **12**, 682475.
- Johnson R.** 2019. *Defining hemp: a fact sheet*. Washington, DC: Congressional Research Service.
- Joshi CP, Thammannagowda S, Fujino T, et al.** 2011. Perturbation of wood cellulose synthesis causes pleiotropic effects in transgenic aspen. *Molecular Plant* **4**, 331–345.
- Keller A, Leupin M, Mediavilla V, Wintermantel E.** 2001. Influence of the growth stage of industrial hemp on chemical and physical properties of the fibres. *Industrial Crops and Products* **13**, 35–48.
- Khan NU, Marwat KB, Hassan G, Farhatullah SB, Makhdoom K, Ahmad W, Khan HU.** 2010. Genetic variation and heritability for cotton seed, fiber and oil traits in *Gossypium hirsutum* L. *Pakistan Journal of Botany* **42**, 615–625.
- Kinney A.** 1998. Production of specialised oils for industry. In: Harwood JL, ed. *Plant lipid biosynthesis: fundamentals and agricultural applications*. Cambridge: Cambridge University Press, 273–285.
- Kovalchuk I, Pellino M, Rigault P, Van Velzen R, Ebersbach J, Ashnest J, Mau M, Schranz M, Alcorn J, Laprairie R.** 2020. The genomics of cannabis and its close relatives. *Annual Review of Plant Biology* **71**, 713–739.
- Kriese U, Schumann E, Weber W, Beyer M, Brühl L, Matthäus B.** 2004. Oil content, tocopherol composition and fatty acid patterns of the seeds of 51 *Cannabis sativa* L. genotypes. *Euphytica* **137**, 339–351.
- Kundan M, Gani U, Fayaz M, Angmo T, Kesari R, Rahul VP, Gairola S, Misra P.** 2022. Two R2R3-MYB transcription factors, CsMYB33 and

- CsMYB78 are involved in the regulation of anthocyanin biosynthesis in *Cannabis sativa* L. *Industrial Crops and Products* **188**, 115546.
- Kurtz LE, Brand MH, Lubell-Brand JD.** 2020. Production of tetraploid and triploid hemp. *HortScience* **55**, 1703–1707.
- Lange BM, Zager JJ.** 2022. Comprehensive inventory of cannabinoids in *Cannabis sativa* L.: can we connect genotype and chemotype? *Phytochemistry Reviews* **21**, 1273–1313.
- Laverty KU, Stout JM, Sullivan MJ, et al.** 2019. A physical and genetic map of *Cannabis sativa* identifies extensive rearrangements at the THC/CBD acid synthase loci. *Genome Research* **29**, 146–156.
- Leple J-C, Dauwe R, Morreel K, et al.** 2007. Downregulation of cinnamoyl-coenzyme A reductase in poplar: multiple-level phenotyping reveals effects on cell wall polymer metabolism and structure. *The Plant Cell* **19**, 3669–3691.
- Le Roy J, Blervacq A-S, Créach A, Huss B, Hawkins S, Neutelings G.** 2017. Spatial regulation of monolignol biosynthesis and laccase genes control developmental and stress-related lignin in flax. *BMC Plant Biology* **17**, 124.
- Li A, Wang R, Li X, et al.** 2016. Proteomic profiling of cellulase-aid-extracted membrane proteins for functional identification of cellulose synthase complexes and their potential associated-components in cotton fibers. *Scientific Reports* **6**, 26356.
- Li A, Xia T, Xu W, et al.** 2013. An integrative analysis of four CESA isoforms specific for fiber cellulose production between *Gossypium hirsutum* and *Gossypium barbadense*. *Planta* **237**, 1585–1597.
- Liu F-H, Hu H-R, Du G-H, Deng G, Yang Y.** 2017. Ethnobotanical research on origin, cultivation, distribution and utilization of hemp (*Cannabis sativa* L.) in China. *Indian Journal of Traditional Knowledge* **16**, 235–242.
- Liu M, Fernando D, Daniel G, Madsen B, Meyer AS, Ale MT, Thygesen A.** 2015. Effect of harvest time and field retting duration on the chemical composition, morphology and mechanical properties of hemp fibers. *Industrial Crops and Products* **69**, 29–39.
- Liu Y, Du Z, Li Y, Lu S, Tang S, Guo L.** 2024. Improving linolenic acid content in rapeseed oil by overexpression of CsFAD2 and CsFAD3 genes. *Molecular Breeding* **44**, 9.
- Livingston SJ, Quilichini TD, Booth JK, Wong DC, Rensing KH, Laflamme-Yonkman J, Castellarin SD, Bohlmann J, Page JE, Samuels AL.** 2020. Cannabis glandular trichomes alter morphology and metabolite content during flower maturation. *The Plant Journal* **101**, 37–56.
- Livingston SJ, Rensing KH, Page JE, Samuels AL.** 2022. A polarized supercell produces specialized metabolites in cannabis trichomes. *Current Biology* **32**, 4040–4047.e4.
- Lynch RC, Padgitt-Cobb LK, Garfinkel AR, Knaus BJ, Hartwick NT, Allsing N, Aylward A, Mamerto A, Kitony JK, Colt K.** 2024. Domesticated cannabinoid synthases amid a wild mosaic cannabis pangenome. *bioRxiv* 2024.05. 21.595196. [Preprint].
- MacMillan CP, Birke H, Chuah A, Brill E, Tsuji Y, Ralph J, Dennis ES, Llewellyn D, Pettolino FA.** 2017. Tissue and cell-specific transcriptomes in cotton reveal the subtleties of gene regulation underlying the diversity of plant secondary cell walls. *BMC Genomics* **18**, 539.
- Mahlberg PG, Kim ES.** 2004. Accumulation of cannabinoids in glandular trichomes of *Cannabis* (Cannabaceae). *Journal of Industrial Hemp* **9**, 15–36.
- Mansouri H, Bagheri M.** 2017. Induction of polyploidy and its effect on *Cannabis sativa* L. In: Chandra S, Lata H, ElSohly M, eds. *Cannabis sativa* L.—Botany and biotechnology. Cham: Springer, 365–383.
- Matías-Hernández L, Jiang W, Yang K, Tang K, Brodelius PE, Pelaz S.** 2017. AaMYB 1 and its orthologue AtMYB 61 affect terpene metabolism and trichome development in *Artemisia annua* and *Arabidopsis thaliana*. *The Plant Journal* **90**, 520–534.
- Matthäus B, Brühl L.** 2008. Virgin hemp seed oil: an interesting niche product. *European Journal of Lipid Science and Technology* **110**, 655–661.
- McGarvey P, Huang J, McCoy M, et al.** 2020. De novo assembly and annotation of transcriptomes from two cultivars of *Cannabis sativa* with different cannabinoid profiles. *Gene* **762**, 145026.
- McKernan KJ, Helbert Y, Kane LT, Ebling H, Zhang L, Liu B, Eaton Z, McLaughlin S, Kingan S, Baybayan P.** 2020. Sequence and annotation of 42 cannabis genomes reveals extensive copy number variation in cannabinoid synthesis and pathogen resistance genes. *bioRxiv* 2020.01.03.894428. [Preprint].
- Melchinger A, Böhm J, Utz H, Müller J, Munder S, Mauch F.** 2018. High-throughput precision phenotyping of the oil content of single seeds of various oilseed crops. *Crop Science* **58**, 670–678.
- Mène-Saffrané L, Dellapenna D.** 2010. Biosynthesis, regulation and functions of tocopherols in plants. *Plant Physiology and Biochemistry* **48**, 301–309.
- Moliterni VC, Cattivelli L, Ranalli P, Mandolino G.** 2004. The sexual differentiation of *Cannabis sativa* L.: a morphological and molecular study. *Euphytica* **140**, 95–106.
- Momotaz A, Scott JW, Schuster DJ.** 2010. Identification of quantitative trait loci conferring resistance to *Bemisia tabaci* in an F2 population of *Solanum lycopersicum* × *Solanum habrochaites* accession LA1777. *Journal of the American Society for Horticultural Science* **135**, 134–142.
- Morel A, Lebard P, Dereux A, Azuar J, Questel F, Bellivier F, Marie-Claire C, Fatséas M, Vorspan F, Bloch V.** 2021. Clinical trials of cannabidiol for substance use disorders: outcome measures, surrogate endpoints, and biomarkers. *Frontiers in Psychiatry* **12**, 565617.
- Nadakuduti SS, Pollard M, Kosma DK, Allen C Jr, Ohlrogge JB, Barry CS.** 2012. Pleiotropic phenotypes of the sticky peel mutant provide new insight into the role of CUTIN DEFICIENT2 in epidermal cell function in tomato. *Plant Physiology* **159**, 945–960.
- Novakova P, Sal J.** 2019. Use of technical hemp for concrete—hempcrete. IOP Conference Series: Materials Science and Engineering **603**, 052095.
- Nuutinen T.** 2018. Medicinal properties of terpenes found in *Cannabis sativa* and *Humulus lupulus*. *European Journal of Medicinal Chemistry* **157**, 198–228.
- O'Brien K.** 2022. Cannabidiol (CBD) in cancer management. *Cancers* **14**, 885.
- Oomah BD, Busson M, Godfrey DV, Drover JC.** 2002. Characteristics of hemp (*Cannabis sativa* L.) seed oil. *Food Chemistry* **76**, 33–43.
- Pancaldi F, Schranz ME, Van Loo EN, Trindade LM.** 2023a. Highly differentiated genomic properties underpin the different cell walls of Poaceae and eudicots. *Plant Physiology* **194**, 274–295.
- Pancaldi F, Van Loo EN, Schranz ME, Trindade LM.** 2022a. Genomic architecture and evolution of the cellulose synthase gene superfamily as revealed by phylogenomic analysis. *Frontiers in Plant Science* **13**, 870818.
- Pancaldi F, Van Loo EN, Senio S, Al Hassan M, Van Der Crujisen K, Paulo M-J, Dolstra O, Schranz ME, Trindade LM.** 2023b. Syntenic cell wall QTLs as versatile breeding tools: intraspecific allelic variability and predictability of biomass quality loci in target plant species. *Plants* **12**, 779.
- Pancaldi F, Vlegels D, Rijken H, Van Loo EN, Trindade LM.** 2022b. Detection and analysis of syntenic quantitative trait loci controlling cell wall quality in angiosperms. *Frontiers in Plant Science* **13**, 855093.
- Peil A, Flachowsky H, Schumann E, Weber W.** 2003. Sex-linked AFLP markers indicate a pseudoautosomal region in hemp (*Cannabis sativa* L.). *Theoretical and Applied Genetics* **107**, 102–109.
- Petit J.** 2020. Novel molecular tools to uncover the genetic architecture of hemp fibre quality. PhD thesis. Wageningen University.
- Petit J, Gulisano A, Dechesne A, Trindade LM.** 2019. Phenotypic variation of cell wall composition and stem morphology in hemp (*Cannabis sativa* L.): optimization of methods. *Frontiers in Plant Science* **10**, 959.
- Petit J, Salentijn EM, Paulo M-J, Denneboom C, Trindade LM.** 2020a. Genetic architecture of flowering time and sex determination in hemp (*Cannabis sativa* L.): a genome-wide association study. *Frontiers in Plant Science* **11**, 569958.
- Petit J, Salentijn EM, Paulo M-J, Denneboom C, Van Loo EN, Trindade LM.** 2020b. Elucidating the genetic architecture of fiber quality in hemp (*Cannabis sativa* L.) using a genome-wide association study. *Frontiers in Genetics* **11**, 566314.

- Petit J, Salentijn EM, Paulo M-J, Thouminot C, Van Dinter BJ, Magagnini G, Gusovius H-J, Tang K, Amaducci S, Wang S. 2020c. Genetic variability of morphological, flowering, and biomass quality traits in hemp (*Cannabis sativa* L.). *Frontiers in Plant Science* **11**, 102.
- Porokhvinova E, Matveeva T, Khafizova G, Bemova V, Doubovskaya A, Kishlyan N, Podolnaya L, GavriloVA V. 2022. Fatty acid composition of oil crops: genetics and genetic engineering. *Genetic Resources and Crop Evolution* **69**, 2029–2045.
- Prentout D, Razumova O, Rhoné B, Badouin H, Henri H, Feng C, Käfer J, Karlov G, Marais GA. 2020. An efficient RNA-seq-based segregation analysis identifies the sex chromosomes of *Cannabis sativa*. *Genome Research* **30**, 164–172.
- Prentout D, Stajner N, Cerenak A, Tricou T, Brochier-Armanet C, Jakse J, Käfer J, Marais GA. 2021. Plant genera *Cannabis* and *Humulus* share the same pair of well-differentiated sex chromosomes. *New Phytologist* **231**, 1599–1611.
- Preston JC, Hileman LC. 2013. Functional evolution in the plant SQUAMOSA-PROMOTER BINDING PROTEIN-LIKE (SPL) gene family. *Frontiers in Plant Science* **4**, 80.
- Punja ZK, Sutton DB, Kim T. 2023. Glandular trichome development, morphology, and maturation are influenced by plant age and genotype in high THC-containing cannabis (*Cannabis sativa* L.) inflorescences. *Journal of Cannabis Research* **5**, 12.
- Ralph J, Akiyama T, Coleman HD, Mansfield SD. 2012. Effects on lignin structure of coumarate 3-hydroxylase downregulation in poplar. *Bioenergy Research* **5**, 1009–1019.
- Ralph J, Lapierre C, Boerjan W. 2019. Lignin structure and its engineering. *Current Opinion in Biotechnology* **56**, 240–249.
- Ranalli P, Venturi G. 2004. Hemp as a raw material for industrial applications. *Euphytica* **140**, 1–6.
- Rathi V, Singh G, Kumar P, Chaudhary M, Singh P, Mishra M. 2022. Legality of worldwide cannabis use and associated economic benefits. In: Belwal T, Belwal NC, eds. *Revolutionizing the potential of hemp and its products in changing the global economy*. Cham: Springer, 27–68.
- Ren G, Zhang X, Li Y, *et al.* 2021. Large-scale whole-genome resequencing unravels the domestication history of *Cannabis sativa*. *Science Advances* **7**, eabg2286.
- Rezvankhah A, Emam-Djomeh Z, Safari M, Askari G, Salami M. 2019. Microwave-assisted extraction of hempseed oil: studying and comparing of fatty acid composition, antioxidant activity, physicochemical and thermal properties with Soxhlet extraction. *Journal of Food Science and Technology* **56**, 4198–4210.
- Ruelens P, De Maagd RA, Proost S, Theißen G, Geuten K, Kaufmann K. 2013. FLOWERING LOCUS C in monocots and the tandem origin of angiosperm-specific MADS-box genes. *Nature Communications* **4**, 2280.
- Saballos A, Sattler SE, Sanchez E, Foster TP, Xin Z, Kang C, Pedersen JF, Vermerris W. 2012. *Brown midrib2 (Bmr2)* encodes the major 4-coumarate:coenzyme A ligase involved in lignin biosynthesis in sorghum (*Sorghum bicolor* L. Moench). *The Plant Journal* **70**, 818–830.
- Sakamoto K, Akiyama Y, Fukui K, Kamada H, Satoh S. 1998. Characterization; genome sizes and morphology of sex chromosomes in hemp (*Cannabis sativa* L.). *Cytologia* **63**, 459–464.
- Salentijn EM, Petit J, Trindade LM. 2019. The complex interactions between flowering behavior and fiber quality in hemp. *Frontiers in Plant Science* **10**, 614.
- Salentijn EM, Zhang Q, Amaducci S, Yang M, Trindade LM. 2015. New developments in fiber hemp (*Cannabis sativa* L.) breeding. *Industrial Crops and Products* **68**, 32–41.
- Sattler SE, Palmer NA, Saballos A, Greene AM, Xin Z, Sarath G, Vermerris W, Pedersen JF. 2012. Identification and characterization of four missense mutations in *brown midrib 12 (Bmr12)*, the caffeic O-methyltransferase (COMT) of sorghum. *Bioenergy Research* **5**, 855–865.
- Sattler SE, Saathoff AJ, Haas EJ, Palmer NA, Funnell-Harris DL, Sarath G, Pedersen JF. 2009. A nonsense mutation in a cinnamyl alcohol dehydrogenase gene is responsible for the sorghum *brown midrib6* phenotype. *Plant Physiology* **150**, 584–595.
- Savadi S, Lambani N, Kashyap PL, Bisht DS. 2017. Genetic engineering approaches to enhance oil content in oilseed crops. *Plant Growth Regulation* **83**, 207–222.
- Scaglione D, Pinosio S, Marroni F, Di Centa E, Fornasiero A, Magris G, Scalabrin S, Cattonaro F, Taylor G, Morgante M. 2019. Single primer enrichment technology as a tool for massive genotyping: a benchmark on black poplar and maize. *Annals of Botany* **124**, 543–552.
- Schierholt A, Becker H. 2001. Environmental variability and heritability of high oleic acid content in winter oilseed rape. *Plant Breeding* **120**, 63–66.
- Schilling S, Melzer R, Dowling CA, Shi J, Muldoon S, McCabe PF. 2023. A protocol for rapid generation cycling (speed breeding) of hemp (*Cannabis sativa*) for research and agriculture. *The Plant Journal* **113**, 437–445.
- Schluttenhofer C, Yuan L. 2017. Challenges towards revitalizing hemp: a multifaceted crop. *Trends in Plant Science* **22**, 917–929.
- Sharma N, Geuten K, Giri BS, Varma A. 2020. The molecular mechanism of vernalization in Arabidopsis and cereals: role of Flowering Locus C and its homologs. *Physiologia Plantarum* **170**, 373–383.
- Shi J, Lang C, Wu X, Liu R, Zheng T, Zhang D, Chen J, Wu G. 2015. RNAi knockdown of fatty acid elongase1 alters fatty acid composition in *Brassica napus*. *Biochemical and Biophysical Research Communications* **466**, 518–522.
- Shi P, Fu X, Shen Q, Liu M, Pan Q, Tang Y, Jiang W, Lv Z, Yan T, Ma Y. 2018. The roles of AaMIXTA 1 in regulating the initiation of glandular trichomes and cuticle biosynthesis in *Artemisia annua*. *New Phytologist* **217**, 261–276.
- Sirikantaramas S, Morimoto S, Shoyama Y, Ishikawa Y, Wada Y, Shoyama Y, Taura F. 2004. The gene controlling marijuana psychoactivity: molecular cloning and heterologous expression of Δ^1 -tetrahydrocannabinolic acid synthase from *Cannabis sativa* L. *Journal of Biological Chemistry* **279**, 39767–39774.
- Siudem P, Wawer I, Paradowska K. 2019. Rapid evaluation of edible hemp oil quality using NMR and FT-IR spectroscopy. *Journal of Molecular Structure* **1177**, 204–208.
- Small E. 2015. Evolution and classification of *Cannabis sativa* (marijuana, hemp) in relation to human utilization. *The Botanical Review* **81**, 189–294.
- Smart LB, Toth JA, Stack GM, Monserrate LA, Smart CD. 2022. Breeding of hemp (*Cannabis sativa*). *Plant Breeding Reviews* **46**, 239–288.
- Sommano SR, Chittasupho C, Ruksiriwanich W, Jantrawut P. 2020. The cannabis terpenes. *Molecules* **25**, 5792.
- Stack GM, Snyder SI, Toth JA, *et al.* 2023. Cannabinoids function in defense against chewing herbivores in *Cannabis sativa* L. *Horticulture Research* **10**, uhad207.
- Stack GM, Toth JA, Carlson CH, Cala AR, Marrero-González MI, Wilk RL, Gentner DR, Crawford JL, Philippe G, Rose JK. 2021. Season-long characterization of high-cannabinoid hemp (*Cannabis sativa* L.) reveals variation in cannabinoid accumulation, flowering time, and disease resistance. *GCB Bioenergy* **13**, 546–561.
- Steel L, Welling M, Ristevski N, Johnson K, Gendall A. 2023. Comparative genomics of flowering behavior in *Cannabis sativa*. *Frontiers in Plant Science* **14**, 1227898.
- Suchoff DH, Inoa SH, Stack GM, Wares AJ, Snyder SI, Murdock MJ, Rose JK, Smart LB, Caton TA, Pearce RC. 2024. Characterization of agronomic performance and sterility in triploid and diploid cannabinoid hemp. *Agronomy Journal* **116**, 2470–2482.
- Tahir MN, Shahbazi F, Rondeau-Gagné S, Trant JF. 2021. The biosynthesis of the cannabinoids. *Journal of Cannabis Research* **3**, 7.
- Tang Q, Xu Y, Gao F, Cheng C, Deng C, Chen J, Yuan X, Zhang X, Su J. 2023. Transcriptomic and metabolomic analyses reveal the differential accumulation of phenylpropanoids and terpenoids in hemp autotetraploid and its diploid progenitor. *BMC Plant Biology* **23**, 616.

- Tanney CA, Backer R, Geitmann A, Smith DL.** 2021. Cannabis glandular trichomes: a cellular metabolite factory. *Frontiers in Plant Science* **12**, 721986.
- Taura F, Sirikantaramas S, Shoyama Y, Yoshikai K, Shoyama Y, Morimoto S.** 2007. Cannabidiolic-acid synthase, the chemotype-determining enzyme in the fiber-type *Cannabis sativa*. *FEBS Letters* **581**, 2929–2934.
- Thomson B, Wellmer F.** 2019. Molecular regulation of flower development. *Current Topics in Developmental Biology* **131**, 185–210.
- Tian B, Wei F, Shu H, Zhang Q, Zang X, Lian Y.** 2011. Decreasing erucic acid level by RNAi-mediated silencing of fatty acid elongase 1 (BnFAE1.1) in rapeseeds (*Brassica napus* L.). *African Journal of Biotechnology* **10**, 13194–13201.
- Torres A, Pauli C, Givens R, Argyris J, Allen K, Monfort A, Gaudin RJ.** 2022. High-throughput methods to identify male *Cannabis sativa* using various genotyping methods. *Journal of Cannabis Research* **4**, 57.
- Toth JA, Stack GM, Cala AR, Carlson CH, Wilk RL, Crawford JL, Viands DR, Philippe G, Smart CD, Rose JK.** 2020. Development and validation of genetic markers for sex and cannabinoid chemotype in *Cannabis sativa* L. *GCB Bioenergy* **12**, 213–222.
- Toth JA, Stack GM, Carlson CH, Smart LB.** 2022. Identification and mapping of major-effect flowering time loci *Autoflower1* and *Early1* in *Cannabis sativa* L. *Frontiers in Plant Science* **13**, 991680.
- Tran J, Vassiliadis S, Elkins AC, Cogan NO, Rochfort SJ.** 2023. Developing prediction models using near-infrared spectroscopy to quantify cannabinoid content in *Cannabis sativa*. *Sensors* **23**, 2607.
- Überall MA.** 2020. A review of scientific evidence for THC: CBD oromucosal spray (nabiximols) in the management of chronic pain. *Journal of Pain Research* **13**, 399–410.
- Van Bakel H, Stout JM, Cote AG, Tallon CM, Sharpe AG, Hughes TR, Page JE.** 2011. The draft genome and transcriptome of *Cannabis sativa*. *Genome Biology* **12**, R102–R118.
- Van der Werf HM, Van Der Veen JH, Bouma A, Ten Cate M.** 1994. Quality of hemp (*Cannabis sativa* L.) stems as a raw material for paper. *Industrial Crops and Products* **2**, 219–227.
- van Velzen R, Schranz ME.** 2021. Origin and evolution of the cannabinoid oxidocyclase gene family. *Genome Biology and Evolution* **13**, ewab130.
- Vergara D, Baker H, Clancy K, Keepers KG, Mendieta JP, Pauli CS, Tittes SB, White KH, Kane NC.** 2016. Genetic and genomic tools for *Cannabis sativa*. *Critical Reviews in Plant Sciences* **35**, 364–377.
- Vogl CR, Mölleken H, Lissek-Wolf G, Surböck A, Kobert J.** 2004. Hemp (*Cannabis sativa* L.) as a resource for green cosmetics: yield of seed and fatty acid compositions of 20 varieties under the growing conditions of organic farming in Austria. *Journal of Industrial Hemp* **9**, 51–68.
- Wang J, Liu Z, Liu H, Peng D, Zhang J, Chen M.** 2021. *Linum usitatissimum* FAD2A and FAD3A enhance seed polyunsaturated fatty acid accumulation and seedling cold tolerance in *Arabidopsis thaliana*. *Plant Science* **311**, 111014.
- Wang J-J, Shao Y-N, Xin Y, Zhang C, Yuan G, Liu Z-J, Chen M-X.** 2024. Heterogeneous expression of stearyl-acyl carrier protein desaturase genes SAD1 and SAD2 from *Linum usitatissimum* enhances seed oleic acid accumulation and seedling cold and drought tolerance capability in *Brassica napus*. *Journal of Integrative Agriculture* **23**, 1864–1878.
- Wei H, Yang Z, Niyitanga S, et al.** 2024. The reference genome of seed hemp (*Cannabis sativa*) provides new insights into fatty acid and vitamin E synthesis. *Plant Communications* **5**, 100718.
- Wickland DP, Hanzawa Y.** 2015. The FLOWERING LOCUS T/TERMINAL FLOWER 1 gene family: functional evolution and molecular mechanisms. *Molecular Plant* **8**, 983–997.
- Woods P, Campbell BJ, Nicodemus TJ, Cahoon EB, Mullen JL, McKay JK.** 2021. Quantitative trait loci controlling agronomic and biochemical traits in *Cannabis sativa*. *Genetics* **219**, iyab099.
- Xiang Z, Sen SK, Min D, Savithri D, Lu F, Jameel H, Chiang V, Chang H-M.** 2017. Field-grown transgenic hybrid poplar with modified lignin biosynthesis to improve enzymatic saccharification efficiency. *ACS Sustainable Chemistry & Engineering* **5**, 2407–2414.
- Xie Z, Mi Y, Kong L, Gao M, Chen S, Chen W, Meng X, Sun W, Chen S, Xu Z.** 2023. *Cannabis sativa*: origin and history, glandular trichome development, and cannabinoid biosynthesis. *Horticulture Research* **10**, uhad150.
- Xu J, Kong L, Ren W, Wang Z, Tang L, Wu W, Liu X, Ma W, Zhang S.** 2024. Identification and expression analysis of TPS family gene in *Cannabis sativa* L. *Heliyon* **10**, e27817.
- Yan T, Chen M, Shen Q, et al.** 2017. HOMEODOMAIN PROTEIN 1 is required for jasmonate-mediated glandular trichome initiation in *Artemisia annua*. *New Phytologist* **213**, 1145–1155.
- Yang C, Gao Y, Gao S, Yu G, Xiong C, Chang J, Li H, Ye Z.** 2015. Transcriptome profile analysis of cell proliferation molecular processes during multicellular trichome formation induced by tomato *Wov* gene in tobacco. *BMC Genomics* **6**, 173–174.
- Yoon J, Choi H, An G.** 2015. Roles of lignin biosynthesis and regulatory genes in plant development. *Journal of Integrative Plant Biology* **57**, 902–912.
- Yu X, Chen G, Tang B, Zhang J, Zhou S, Hu Z.** 2018. The jasmonate ZIM-domain protein gene *SIJAZ2* regulates plant morphology and accelerates flower initiation in *Solanum lycopersicum* plants. *Plant Science* **267**, 65–73.
- Zager JJ, Lange I, Srividya N, Smith A, Lange BM.** 2019. Gene networks underlying cannabinoid and terpenoid accumulation in cannabis. *Plant Physiology* **180**, 1877–1897.
- Zhang X, Xu G, Cheng C, et al.** 2021. Establishment of an *Agrobacterium*-mediated genetic transformation and CRISPR/Cas9-mediated targeted mutagenesis in hemp (*Cannabis sativa* L.). *Plant Biotechnology Journal* **19**, 1979–1987.
- Zhang Y, Zhou Y, Chen Q, Huang X, Tian CE.** 2014. Molecular basis of flowering time regulation in Arabidopsis. *Chinese Bulletin of Botany* **49**, 469.
- Zhao T, Schranz ME.** 2017. Network approaches for plant phylogenomic synteny analysis. *Current Opinion in Plant Biology* **36**, 129–134.