



HHS Public Access

Author manuscript

Nat Microbiol. Author manuscript; available in PMC 2024 December 20.

Published in final edited form as:

Nat Microbiol. 2024 November ; 9(11): 2765–2773. doi:10.1038/s41564-024-01827-2.

A cross-systems primer for synthetic microbial communities

Elijah C. Mehlferber^{1,2,*}, Gontran Arnault³, Bishnu Joshi⁴, Laila P. Partida-Martinez⁵, Kathryn A. Patras^{4,6}, Marie Simonin³, Britt Koskella^{7,8,*}

¹School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA, USA

²Center for Microbial Dynamics and Infection, Georgia Institute of Technology, Atlanta, GA, USA

³Univ Angers, Institut Agro, INRAE, IRHS, SFR QUASAV, Angers, France

⁴Department of Molecular Virology and Microbiology, Baylor College of Medicine, Houston, Texas, USA

⁵Departamento de Ingeniería Genética, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional (Cinvestav), Irapuato 36824, México

⁶Alkek Center for Metagenomics and Microbiome Research, Baylor College of Medicine, Houston, Texas, USA

⁷Department of Integrative Biology, University of California, Berkeley, CA, United States

⁸San Francisco Chan Zuckerberg Biohub, San Francisco, CA, United States

Abstract

The design and use of Synthetic Communities, or SynComs, represents one of the most promising strategies for disentangling the complex interactions within microbial communities, and between these communities and their hosts. Compared to natural communities, these simplified consortia provide the opportunity to study ecological interactions at tractable scales, as well as facilitating reproducibility and fostering interdisciplinary science. However, the effective implementation of the SynCom approach requires several important considerations regarding the development and application of these model systems. There are also emerging ethical considerations when both designing and deploying SynComs in clinical, agricultural, or environmental settings. Here, we outline current best practices in developing, implementing and evaluating SynComs across different systems, including a focus on important ethical considerations for SynCom research.

Introduction

Microbial organisms represent the bulk of the diversity present on Earth and as sequencing technologies and computational tools have advanced, we have become increasingly aware

*Corresponding Authors.
Author Contributions

EM and BK developed the framework for the manuscript, with inputs from all authors. EM lead the writing of the manuscript, with contributions by GA, BJ, LPM, KP, MS and BK. Authors GA, BJ, LPM, KP contributed equally to the manuscript, these authors are presented in alphabetical order by surname.

Competing Interests

The authors declare no competing interests.

of the important role they play across systems. As these organisms live in complex and often highly dynamic communities, there is a clear need to understand when and why microbial taxa coexist, how they interact with one another, and how these interactions translate to function – especially given that these outcomes often cannot be predicted based on knowledge of individual taxa. One of the most promising strategies to disentangle these complex relationships within communities and between communities and their hosts is the design of model consortia, generally referred to as synthetic communities, or SynComs.

SynComs can be defined as “consortia of microorganisms designed to mimic, at some scale, the observed functions and structure of the microbiome in natural conditions”¹. This approach was first pioneered in 1965 when Russel Schaedler colonized germ-free mice with defined bacterial isolates², although the term “Synthetic Community” was first used, to the best of our knowledge, by Kim et al. in 2008 to describe a three species community comprised of soil bacteria³. The approach has since gained popularity in both plant and human systems^{4–6}. Historically, SynComs have been composed of bacterial species and have primarily focused on coexistence, competition, cross-feeding and functions encoded on bacterial genomes or plasmids. Though less common, researchers can also include fungal, protist, archaeal, and viral taxa within these experimental communities^{7,8}. Given their popularity, most of the examples included throughout this piece will focus on bacterial SynComs, though we acknowledge the importance of these multi-kingdom community approaches. It is further important to delineate between the types of synthetic communities discussed in this piece: those composed of naturally sourced organisms meant to model some functions of their originating communities^{9,10}, versus those that represent a group of Synthetic Organisms designed to perform a certain function^{11,12}, usually via genetic engineering, with the latter being more typically used in Synthetic Biology.

Compared to natural communities, SynComs provide several advantages to researchers. Their defined membership enables the reconstitution of identical communities across experiments, allowing reproducibility across time and labs¹³. Like the development of model systems in biology, this approach allows researchers to integrate knowledge of a given system to accelerate progress and foster interdisciplinary science. While most research on microbiomes relies on destructive sampling, SynComs allow for repeated manipulation of the community to dissect the role of individual species (and their abundances) in its assembly and function.

The SynCom approach can be used to facilitate answers to fundamental research questions, as well as for specific applications, and is equally suitable for host-associated and free-living microbial communities (Figure 1). In both cases, SynComs provide excellent opportunities to model ecological interactions at tractable scales and can offer key insights into community dynamics. As applied systems, non-host associated (environmentally-derived) SynComs can be harnessed for bioremediation, chemical engineering, and biofuel production. Host-associated SynComs can be used to increase crop production in agricultural settings, through both growth promotion and disease protection, as well as to understand and treat microbiome-associated animal diseases. In the context of human health, SynComs can be designed to treat disease and dysbiosis, such as enrichment of opportunistic pathogens, which are associated with infectious (e.g. *Clostridioides difficile* infection¹⁴) or

metabolic (e.g. *diabetes mellitus*¹⁵) origins. A better fundamental understanding of how within-microbiome interactions affect the balance of microbial taxa and ability of resident communities to resist pathogen invasion will support the development of targeted microbial interventions to reduce such community disturbances. In all cases, SynComs address the need to understand the host-microbe and microbe-microbe interactions underlying these phenotypes, and act as potential interventions for re-establishing stable microbial communities.

In the following sections we outline current best practices in developing, implementing and evaluating SynComs, including a focus on important ethical considerations for SynCom research and application (Figure 2). This piece is not a comprehensive review of SynCom-associated studies, nor is it the first to outline approaches to developing SynComs^{1,4,5,11,16,17}, but rather serves as a cross-systems primer for those hoping to develop a SynCom for research or application, or for evaluating SynCom-associated work. The information provided herein represents a comprehensive starting point for those unfamiliar with the field, and citations have been chosen carefully to give readers the opportunity to follow up on any points, or explore specific systems, in greater detail.

Designing the community

Many strategies exist to design SynComs depending on the objectives and research system of interest. These can be summarized as a continuum from bottom-up to top-down designs (Figure 3). The bottom-up approach relies on the assembly of a specific set of microbial strains of interest, chosen due to their suitability to some criteria of the study (including the feasibility of isolating and culturing them) (Figure 3a). In this case, phenotypically and genomically defined strains are typically combined to characterize microbial interaction dynamics and mechanism, community functions, and emergent properties of known strain assemblages. These simple SynComs have facilitated the discovery of inter-microbial antagonism pathways between microorganisms¹⁸, as well as microbial cross-feeding and degradative synergies that are critical for ecosystem functioning¹⁹. For example, the OMM mouse community, shows how microbial interactions and cross-feeding can impact their host by shaping their exposure to certain metabolic by-products²⁰. This approach is crucial for identifying the molecular mechanisms driving microbial interactions, but it relies on simplification of the microbial diversity and environmental conditions. Strains are often selected because of the extensive knowledge available on their genetic and phenotypic attributes (i.e. model strains) and not because they co-exist in nature (i.e. from different sources of isolation). Moreover, they are selected because they can grow alone, leading to bias in the types of strains being included. Additionally, recent work has demonstrated that strains coexisting within a stable complex community might fail to coexist in pairwise co-cultures, showing that multi-species coexistence is an emergent phenomenon²¹.

In contrast to building SynComs from characterized strains, a top-down design relies on assembling a large diversity of strains (for example, by sampling of a natural source) and reducing their complexity in a stepwise fashion to gain insight into the sub-components of the community (Figure 3b). The objectives for this approach can be wide-ranging, from mimicking the natural phylogenetic diversity, to identifying core taxa or core functions of a

microbiome, to understanding the specific role of taxa of interest in complex assemblages. Simplification of the initial community can be achieved through natural or knowledge-driven filtering approaches or bottlenecks. One such straightforward approach is to let the environment or host ‘filter’ or select for strains capable of colonizing and surviving in/on it from the initial strain pool²². Complementary approaches include performing experimental evolution to enrich taxa or functions of interest over multiple cycles of reinoculation²³ or applying random filtering such as serial dilution²⁴ to create random subsets of the wider community for subsequent exploration. Alternatively, knowledge-driven filtering can be performed based on existing data that informs add-in/drop-out of taxonomic or functional groups of interest, which can be identified from functional assays or metagenomes²⁵, and/or microbial hubs identified via co-occurrence networks²⁶. These bottom-up and top-down approaches for SynCom development are complementary and necessary to eventually “meet in the middle”, allowing researchers to understand and predict microbiota assembly and functions across scales of complexity.

A crucial aspect in SynCom design is the meticulous sourcing and selection of the strains. Depending on the objectives of the study, strains can be sourced from the study system (e.g. same soil or individual), from across environments, or even from (inter)national strain collections. It is important to acknowledge that the sourcing of strains poses a significant limitation to the SynCom approach, as even complex SynComs may lack certain keystone taxa (i.e. those with outsized impact on community stability or function²⁷) that might be essential for realistic community dynamics⁴. During the selection process, careful consideration must be given to the number of strains that align with the project’s goals and system complexity²⁸. Factors like ease of cultivation and growth rates play a role in determining the feasibility of incorporating specific strains, but it is critical to appreciate the varying growth capabilities of strains under different conditions (i.e. acknowledging that selection based on one set of criteria likely reduces success or function of the SynCom under different conditions). Mitigating these effects can be achieved by aligning media and culture conditions with the specific requirements of the studied system²⁹. Additionally, it is important to note that the convenience of handling specific strains does not always correspond directly to their significance within the system.

Strain preparation and Inoculation

Various factors must be considered to standardize the use of SynComs across experiments and studies (Box 1). First, the choice between *in vitro* and *in vivo systems* is fundamental, necessitating consideration of the ecological relevance and applicability of the chosen system. When studying host-associated communities, an *in vitro* approach may be most appropriate (at initially for hypothesis generation and when the relevant interactions are solely inter-microbial, but if the interactions of interest are host-microbial, an *in vivo* approach is necessary. Next, the impact of inoculum concentration must be considered given the density- and frequency-dependent nature of many microbial interactions³⁰. This can be achieved by inoculating at ecologically relevant densities or inoculating at lower densities and allowing the community to establish *in situ*. There may be compelling reasons to increase the concentration, especially when the SynCom is required to outcompete the resident community (host or environment) within a coalescence framework³¹. Establishing

standardized protocols for SynCom inoculation is essential, including the timing and frequency of inoculation³², growth prior to inoculation (i.e. physiological state of the strains³³ and media composition), and subsequent sampling of community dynamics. Additionally, it is necessary to determine if these communities can be stored throughout the duration of the study³⁴, or if they need to be remade each time to ensure that they have the same concentration, evenness, and physiological state.

Evaluating a SynCom

When designing a SynCom it is important to remember the oft cited quote by George Box, “all models are wrong, some are useful”. SynComs, after all, are meant to be tractable models of natural systems. The question is not whether they represent those systems perfectly, but rather if they represent the features of the system that the researchers aim to study. It is therefore of critical importance that the SynCom is designed with specific questions and context in mind, that these questions are well articulated, and that the features selected to be represented in that model are relevant to the system in its natural or applied environment. For fundamental questions, employing a simplified SynCom can prove highly advantageous in demonstrating the feasibility or existence of specific functions or interactions (Fig. 1). For example, Yang et al.³⁵ employed a community consisting of 6 species from the same genus, and although highly simplified compared to natural communities, this SynCom allowed them to test the role of community diversity in robustness against invasion, though of course a more complex community might reveal additional contributing factors. More complex questions might require more complex communities. To identify a conserved set of host (*Arabidopsis thaliana*) genes that are upregulated in response to colonization by bacteria, Maier et al.³⁶ constructed a SynCom consisting of 38 strains representing the breadth of phyla naturally associated with the plant. Establishing that this was a general plant response required a SynCom that captured more of the natural diversity that is found to associate with their host. While these two communities are quite different, each is sufficient to represent models of the interactions of interest.

When evaluating the effectiveness of a SynCom, it is important to focus on the system being modeled rather than the composition or complexity compared to other established communities (Fig. 3). For every SynCom study, there will be a tradeoff between tractability and relevance. Simple communities are easier to work with, but less representative of real systems. They run the risk of missing emergent properties and context-specific outcomes, such as higher order competitive interactions, priority effects, or the impact of rare keystone members, making them potentially less generalizable. In contrast, while more complex models might capture these effects, they can be harder to implement, show lower reproducibility, and offer significant challenges when it comes to data interpretation.

Researchers must think critically about their questions and ensure that their community is sufficiently designed to answer them, while also transparently communicating the limitations of their model. To ensure your SynCom aligns with the questions being asked, methods can be implemented to validate community performance. This includes understanding what features of your community you need to validate (sequencing depth, growth, survival, interactions, productivity, host phenotypes, ecosystem function etc.) and identifying methods

to do so (see Box 1 for suggested computational resources). In light of new methods for barcoding/labeling strains³⁷ and multi-omics approaches, both validating composition (or change in composition) of the SynCom and evaluating the functions of the SynCom can be done simultaneously, but depending on if the study is focusing on ecological (validate composition) or functional (validate function) properties of the community both approaches might not be needed.

The classic approach to understanding bacterial community composition is through 16S rRNA gene sequencing (though other targets such as *gyrB* and *rpoB* can be used). However, this method only resolves relative abundance of the bacteria present, and can run into issues with copy number variation, primer bias, as well as difficulties delineating at the species level and the inability to distinguish between living and dead bacteria. When employing a well-defined SynCom, many of these issues can be addressed or avoided by employing alternative/additional methods. Plating, if community members can be morphologically distinguished, allows for a relatively cheap and effective determination of living bacterial numbers and diversity. For more complex communities, absolute abundance can also be approximated through qPCR^{38,39} or ddPCR⁴⁰ using general or species-specific primers. Further corrections for copy number can be employed if the SynCom member genomes have been sequenced⁴¹, and primer bias can be addressed through comparisons to known mock communities⁴², for example, using the Zymo community standards. When addressing strain resolution, methods like DADA2⁴³ can distinguish between strains if they differ by at least one base pair in the sequenced region (though this is not the case for all taxonomic groups). Other approaches such as long read sequencing⁴⁴ or metagenomic barcoding can be used to distinguish more closely related strains. Finally, live/dead PCR using PMA has been employed to remove relic DNA from sequencing samples, limiting quantification to cells that are still intact (and therefore likely alive) at the time of sequencing⁴⁵.

Perhaps more complicated is determining that a SynCom is performing the functions of interest. When evaluating these functions, more complex “Omics” enabled methods can be employed. Metagenomics can be used to quantify the genetic and functional composition of the community, including both gene presence and relative abundance⁴⁶, and could be used to determine if these are representative of the natural system. Further approaches could be applied to approximate a community metagenome by normalizing genome assemblies (for gene content) against 16S rRNA gene sequencing data. RNAseq can be applied to determine whether the host is responding to the SynCom under the conditions of interest, as well as evaluating the microbial responses through community-wide RNAseq⁴⁷. Additionally, both host and community can be evaluated simultaneously using dual RNA-seq⁴⁸. The extent to which host and community responses are reflective of the actual system can be further quantified by comparing gene expression in the natural and model systems⁴⁹.

Ethical considerations in SynCom development and application

When developing SynComs to model or treat human disease it is important that these systems are representative of diverse human groups (i.e. those living across rural and urban settings, from different geographic areas, and across the socio-economic continuum) and focus on both well-studied and typically neglected diseases⁵⁰. Their design should consider

that microbiome composition can vary across geographic and economic boundaries^{51,52}, either by including these groups when defining important strains or by designing communities to specifically represent them. When a SynCom is intended for clinical application it must be composed of known, culturable, and reproducible communities that are verified to be free from harmful pathogens or virulence factors that could pose risks to human health. This is now possible because the genomes and features of the community (e.g. metatranscriptome, metaproteome) can be more easily characterized. Even after this, however, rigorous multi-center, longitudinal cohort studies are required to identify SynCom off-target effects, such as unintentional transfer of pathobionts from donor to recipient⁵³, inadvertent propagation of genes (e.g., antimicrobial resistance genes), unintended impacts on the endogenous microbiota such as competitive enrichment of other pathogens⁵⁴, or unanticipated effects on host metabolism or susceptibility to disease. Finally, it is important to consider the long-term effects on the resident community, as well as the potential for transmission beyond the original recipient, including horizontal transmission within the household, as well as vertical transmission from parent to child⁵⁵.

Likewise, there are several critical factors that must be considered when SynComs are to be used in natural or agricultural settings. We are optimistic that the field can learn from, and avoid, past mistakes made with novel biological technologies (i.e. antibiotics or species introduction biocontrol) as SynComs become more widely applied. Given the high densities at which microbial amendments are typically introduced, these impacts could be more disruptive than ‘natural’ microbial dispersal⁵⁶. Moreover, since many members of SynComs have been specifically selected to grow well (often across diverse habitats), they should be considered to pose a risk for invasion, possibly leading to loss of natural microbial diversity. As such, the development and deployment of SynComs outside of the laboratory should adhere to the four principles of ethics (do good, don’t harm, respect, and act justly) and the eleven guiding principles for microbiome research⁵⁷. Practically, studies should be undertaken to assess the associated risks under the conditions that these SynComs might be applied. Further, while biocontainment has been a long-standing focus for engineered microbial organisms⁵⁸, it has received far less attention for Syncoms and probiotics more generally. As use of Syncoms in medical, agricultural, and environmental settings becomes more prominent, this will need more thorough assessment.

Beyond SynCom release, there are also ethical considerations around their use in research. These include embracing FAIR data principles to ensure that all data underlying published findings are findable, accessible, interoperable and reusable⁵⁹. There are numerous data and strain repositories that can be used to achieve this goal, but true reusability requires useful metadata^{59–61} (Box 1). The National Microbiome Data Collaborative (<https://microbiomedata.org/>) offers many examples of how this can be done and is itself an exemplary effort of how to bring these issues to the attention of the research community.

Future Perspectives

In summary, while SynComs represent an important resource for increasing our fundamental knowledge of microbial systems, as well as a valuable applied tool, there are critical considerations when designing and implementing them. While this piece provides a high-

level overview of those considerations, more system-specific reading will certainly be useful as the reader begins to construct or evaluate a SynCom. We suggest the following reviews as excellent next steps depending on the specific system in question; plant^{5,17} (including the review by Northen et al., in this issue⁶²) animal⁶³, agriculture^{1,64}, human^{4,16,65}.

Despite the progress made in developing the SynCom approach, the field is still in its infancy and researchers must continue to collaboratively establish and share best practices. As research becomes more collaborative and more standardized, the field may move towards “model” SynComs for use across research groups. However, it is essential to first identify the most effective systems and communities, and in doing so we will likely need to expand our efforts to include less culturable organisms, as well as increasing diversity across kingdoms and trophic levels. Additionally, as more research groups begin working with SynComs, it is imperative to explore methods for integrating findings across different models to uncover common principles and patterns, including standardizing the reporting of metadata associated with these studies. Likewise, the field should develop best practices for calibrating and testing the effectiveness of communities as models for specific research questions. Looking ahead, the potential role of artificial intelligence⁶⁶ in advancing the development and study of SynComs should also be considered, as tools to accomplish this are beginning to be implemented.

Acknowledgements

This manuscript was inspired by an ASM Microbe panel discussion involving many of the authors and organized by the ASM EEB track in 2023, we would like to specifically thank Benjamin Callahan for early discussions on this topic. EM would like to acknowledge funding from the NSF EAGER award # 1838299 as well as support from the NSF Postdoctoral Research Fellowships in Biology award # 2209151. BK is a Chan Zuckerberg San Francisco Biohub investigator. BJ and KP would like to acknowledge funding from the National Institutes of Health award # U19 AI157981. GA and MS were supported by the 3rd Programme for Future Investments (France 2030) and operated by the SUCSEED project (ANR- 20- PCPA-0009) funded by the ‘Growing and Protecting crops Differently’ French Priority Research Program (PPR-CPA), part of the national investment plan operated by the French National Research Agency (ANR). LPM would like to acknowledge funding from Conahcyt (Consejo Nacional de Humanidades, Ciencias y Tecnologías) with award #'s A1-S-9889 and CBF2023-2024-2642.

References

1. de Souza RSC, Armanhi JSL & Arruda P. From Microbiome to Traits: Designing Synthetic Microbial Communities for Improved Crop Resiliency. *Frontiers in Plant Science* 11, (2020).
2. Schaedler RW, Dubs R & Costello R ASSOCIATION OF GERMFREE MICE WITH BACTERIA ISOLATED FROM NORMAL MICE. *J Exp Med* 122, 77–82 (1965). [PubMed: 14325475]
3. Kim HJ, Boedicker JQ, Choi JW & Ismagilov RF Defined spatial structure stabilizes a synthetic multispecies bacterial community. *Proceedings of the National Academy of Sciences* 105, 18188–18193 (2008).
4. van Leeuwen PT, Brul S, Zhang J & Wortel MT Synthetic microbial communities (SynComs) of the human gut: design, assembly, and applications. *FEMS Microbiol Rev* 47, fuad012 (2023). [PubMed: 36931888]
5. Vorholt JA, Vogel C, Carlström CI & Müller DB Establishing Causality: Opportunities of Synthetic Communities for Plant Microbiome Research. *Cell Host & Microbe* 22, 142–155 (2017). [PubMed: 28799900]
6. Lebeis SL et al. Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa. *Science* 349, 860–864 (2015). [PubMed: 26184915]
7. Thonar C, Frossard E, Šmilauer P. & Jansa J Competition and facilitation in synthetic communities of arbuscular mycorrhizal fungi. *Molecular Ecology* 23, 733–746 (2014). [PubMed: 24330316]

8. Durán P et al. Microbial Interkingdom Interactions in Roots Promote Arabidopsis Survival. *Cell* 175, 973–983.e14 (2018). [PubMed: 30388454]
9. Mehlferber EC et al. Phyllosphere microbial associations improve plant reproductive success. *Front Plant Sci* 14, 1273330 (2023). [PubMed: 38143578]
10. Flores-Núñez VM et al. Synthetic Communities Increase Microbial Diversity and Productivity of Agave tequilana Plants in the Field. *Phytobiomes Journal* 7, 435–448 (2023).
11. Johns NI, Blazejewski T, Gomes AL & Wang HH Principles for designing synthetic microbial communities. *Current Opinion in Microbiology* 31, 146–153 (2016). [PubMed: 27084981]
12. McCarty NS & Ledesma-Amaro R Synthetic Biology Tools to Engineer Microbial Communities for Biotechnology. *Trends Biotechnol* 37, 181–197 (2019). [PubMed: 30497870]
13. Sasse J et al. Multilab EcoFAB study shows highly reproducible physiology and depletion of soil metabolites by a model grass. *New Phytologist* 222, 1149–1160 (2019). [PubMed: 30585637]
14. Theriot CM & Young VB Interactions Between the Gastrointestinal Microbiome and *Clostridium difficile*. *Annual Review of Microbiology* 69, 445–461 (2015).
15. Singer-Englar T, Barlow G & Mathur R Obesity, diabetes, and the gut microbiome: an updated review. *Expert Review of Gastroenterology & Hepatology* 13, 3–15 (2019). [PubMed: 30791839]
16. Vázquez-Castellanos JF, Biclou A, Vrancken G, Huys GR & Raes J Design of synthetic microbial consortia for gut microbiota modulation. *Current Opinion in Pharmacology* 49, 52–59 (2019). [PubMed: 31430629]
17. Marín O, González B & Poupin MJ From Microbial Dynamics to Functionality in the Rhizosphere: A Systematic Review of the Opportunities With Synthetic Microbial Communities. *Front. Plant Sci* 12, (2021).
18. Peterson SB, Bertolli SK & Mougous JD Interbacterial antagonism: at the center of bacterial life. *Curr Biol* 30, R1203–R1214 (2020). [PubMed: 33022265]
19. D'Souza G et al. Ecology and evolution of metabolic cross-feeding interactions in bacteria. *Natural Product Reports* 35, 455–488 (2018). [PubMed: 29799048]
20. Pérez Escrivá P, Fuhrer T & Sauer U Distinct N and C Cross-Feeding Networks in a Synthetic Mouse Gut Consortium. *mSystems* 7, e01484–21 (2022). [PubMed: 35357218]
21. Chang C-Y, Baji D, Vila JCC, Estrela S & Sanchez A Emergent coexistence in multispecies microbial communities. *Science* 381, 343–348 (2023). [PubMed: 37471535]
22. Finkel OM et al. A single bacterial genus maintains root growth in a complex microbiome. *Nature* 587, 103–108 (2020). [PubMed: 32999461]
23. Anand G, Goel V, Dubey S & Sharma S Tailoring the rhizospheric microbiome of *Vigna radiata* by adaptation to salt stress. *Plant Growth Regul* 93, 79–88 (2021).
24. Auchtung JM, Preisner EC, Collins J, Lerma AI & Britton RA Identification of Simplified Microbial Communities That Inhibit *Clostridioides difficile* Infection through Dilution/Extinction. *mSphere* 5, e00387–20 (2020). [PubMed: 32727857]
25. Kumar N, Hitch TCA, Haller D, Lagkouvardos I & Clavel T MiMiC: a bioinformatic approach for generation of synthetic communities from metagenomes. *Microbial Biotechnology* 14, 1757–1770 (2021). [PubMed: 34081399]
26. Caballero-Flores G, Pickard JM & Núñez G Microbiota-mediated colonization resistance: mechanisms and regulation. *Nat Rev Microbiol* 21, 347–360 (2023). [PubMed: 36539611]
27. Banerjee S, Schlaeppi K & van der Heijden MGA Keystone taxa as drivers of microbiome structure and functioning. *Nat Rev Microbiol* 16, 567–576 (2018). [PubMed: 29789680]
28. Emmenegger B et al. Identifying microbiota community patterns important for plant protection using synthetic communities and machine learning. *Nat Commun* 14, 7983 (2023). [PubMed: 38042924]
29. Gerna D, Clara D, Allwardt D, Mitter B & Roach T Tailored Media Are Key to Unlocking the Diversity of Endophytic Bacteria in Distinct Compartments of Germinating Seeds. *Microbiology Spectrum* 10, e00172–22 (2022). [PubMed: 35867396]
30. Abisado RG, Benomar S, Klaus JR, Dandekar AA & Chandler JR Bacterial Quorum Sensing and Microbial Community Interactions. *mBio* 9, 10.1128/mbio.02331-17 (2018).

31. Rocca JD, Muscarella ME, Peralta AL, Izabel-Shen D & Simonin M Guided by Microbes: Applying Community Coalescence Principles for Predictive Microbiome Engineering. *mSystems* 6, 10.1128/msystems.00538-21 (2021).
32. Debray R et al. Priority effects in microbiome assembly. *Nat Rev Microbiol* 20, 109–121 (2022). [PubMed: 34453137]
33. Mutlu A, Kaspar C, Becker N & Bischofs IB A spore quality-quantity tradeoff favors diverse sporulation strategies in *Bacillus subtilis*. *ISME J* 14, 2703–2714 (2020). [PubMed: 32724142]
34. Parnell JJ, Vintila S, Tang C, Wagner MR & Kleiner M Evaluation of ready-to-use freezer stocks of a synthetic microbial community for maize root colonization. *Microbiology Spectrum* 12, e02401–23 (2023). [PubMed: 38084978]
35. Yang T et al. Resource availability modulates biodiversity-invasion relationships by altering competitive interactions: Resource availability modulates biodiversity. *Environmental Microbiology* 19, 2984–2991 (2017). [PubMed: 28229529]
36. Maier BA et al. A general non-self response as part of plant immunity. *Nat. Plants* 7, 696–705 (2021). [PubMed: 34007033]
37. Ordon J et al. Chromosomal barcodes for simultaneous tracking of near-isogenic bacterial strains in plant microbiota. *Nat Microbiol* 9, 1117–1129 (2024). [PubMed: 38503974]
38. Jian C, Luukkonen P, Yki-Järvinen H, Salonen A & Korpela K Quantitative PCR provides a simple and accessible method for quantitative microbiota profiling. *PLOS ONE* 15, e0227285 (2020). [PubMed: 31940382]
39. Zemb O et al. Absolute quantitation of microbes using 16S rRNA gene metabarcoding: A rapid normalization of relative abundances by quantitative PCR targeting a 16S rRNA gene spike-in standard. *Microbiologyopen* 9, e977 (2020). [PubMed: 31927795]
40. Morella NM, Yang SC, Hernandez CA & Koskella B Rapid quantification of bacteriophages and their bacterial hosts in vitro and in vivo using droplet digital PCR. *Journal of Virological Methods* 259, 18–24 (2018). [PubMed: 29859196]
41. Kembel SW, Wu M, Eisen JA & Green JL Incorporating 16S Gene Copy Number Information Improves Estimates of Microbial Diversity and Abundance. *PLOS Computational Biology* 8, e1002743 (2012). [PubMed: 23133348]
42. Brooks JP et al. The truth about metagenomics: quantifying and counteracting bias in 16S rRNA studies. *BMC Microbiology* 15, 66 (2015). [PubMed: 25880246]
43. Callahan BJ et al. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods* 13, 581–583 (2016). [PubMed: 27214047]
44. Callahan BJ et al. High-throughput amplicon sequencing of the full-length 16S rRNA gene with single-nucleotide resolution. *Nucleic Acids Research* 47, e103 (2019). [PubMed: 31269198]
45. Carini P et al. Effects of Spatial Variability and Relic DNA Removal on the Detection of Temporal Dynamics in Soil Microbial Communities. *mBio* 11, (2020).
46. de Souza RSC, Armanhi JSL, Damasceno N de B, Imperial J. & Arruda P. Genome Sequences of a Plant Beneficial Synthetic Bacterial Community Reveal Genetic Features for Successful Plant Colonization. *Frontiers in Microbiology* 10, (2019).
47. Giannoukos G et al. Efficient and robust RNA-seq process for cultured bacteria and complex community transcriptomes. *Genome Biol* 13, r23 (2012). [PubMed: 22455878]
48. Westermann AJ, Gorski SA & Vogel J Dual RNA-seq of pathogen and host. *Nat Rev Microbiol* 10, 618–630 (2012). [PubMed: 22890146]
49. Lewin GR et al. Application of a quantitative framework to improve the accuracy of a bacterial infection model. *Proceedings of the National Academy of Sciences* 120, e2221542120 (2023).
50. Mohajeri MH et al. The role of the microbiome for human health: from basic science to clinical applications. *Eur J Nutr* 57, 1–14 (2018).
51. Mallott EK et al. Human microbiome variation associated with race and ethnicity emerges as early as 3 months of age. *PLoS Biol* 21, e3002230 (2023). [PubMed: 37590208]
52. Gupta VK, Paul S & Dutta C Geography, Ethnicity or Subsistence-Specific Variations in Human Microbiome Composition and Diversity. *Front. Microbiol* 8, (2017).

53. DeFilipp Z et al. Drug-Resistant *E. coli* Bacteremia Transmitted by Fecal Microbiota Transplant. *N Engl J Med* 381, 2043–2050 (2019). [PubMed: 31665575]
54. Varga JJ et al. Antibiotics Drive Expansion of Rare Pathogens in a Chronic Infection Microbiome Model. *mSphere* 7, e00318–22 (2022). [PubMed: 35972133]
55. Bogaert D et al. Mother-to-infant microbiota transmission and infant microbiota development across multiple body sites. *Cell Host Microbe* 31, 447–460.e6 (2023). [PubMed: 36893737]
56. Ladau J et al. Microbial invasions and inoculants: a call to action. (2023).
57. Lange L et al. Microbiome ethics, guiding principles for microbiome research, use and knowledge management. *Environmental Microbiome* 17, 50 (2022). [PubMed: 36180931]
58. Pantoja Angles A, Valle-Pérez AU, Hauser C & Mahfouz MM Microbial Biocontainment Systems for Clinical, Agricultural, and Industrial Applications. *Frontiers in Bioengineering and Biotechnology* 10, (2022).
59. Huttenhower C, Finn RD & McHardy AC Challenges and opportunities in sharing microbiome data and analyses. *Nat Microbiol* 8, 1960–1970 (2023). [PubMed: 37783751]
60. Yilmaz P et al. Minimum information about a marker gene sequence (MIMARKS) and minimum information about any (x) sequence (MIXS) specifications. *Nat Biotechnol* 29, 415–420 (2011). [PubMed: 21552244]
61. Mirzayi C et al. Reporting guidelines for human microbiome research: the STORMS checklist. *Nat Med* 27, 1885–1892 (2021). [PubMed: 34789871]
62. Northen T PLACEHOLDER.
63. Jennings SAV & Clavel T Synthetic Communities of Gut Microbes for Basic Research and Translational Approaches in Animal Health and Nutrition. *Annual Review of Animal Biosciences* 12, 283–300 (2024). [PubMed: 37963399]
64. Shayanthan A, Ordoñez PAC & Oresnik IJ The Role of Synthetic Microbial Communities (SynCom) in Sustainable Agriculture. *Front. Agron* 4, (2022).
65. Cheng AG et al. Design, construction, and in vivo augmentation of a complex gut microbiome. *Cell* 185, 3617–3636.e19 (2022). [PubMed: 36070752]
66. Emmenegger B et al. Identifying microbiota community patterns important for plant protection using synthetic communities and machine learning. *Nat Commun* 14, 7983 (2023). [PubMed: 38042924]
67. Schloss PD et al. Introducing mothur: Open-Source, Platform-Independent, Community-Supported Software for Describing and Comparing Microbial Communities. *Applied and Environmental Microbiology* 75, 7537–7541 (2009). [PubMed: 19801464]
68. Edgar RC UNOISE2: improved error-correction for Illumina 16S and ITS amplicon sequencing. Preprint at 10.1101/081257 (2016).
69. Edgar RC, Haas BJ, Clemente JC, Quince C & Knight R UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27, 2194–2200 (2011). [PubMed: 21700674]
70. Davis NM, Proctor DM, Holmes SP, Reiman DA & Callahan BJ Simple statistical identification and removal of contaminant sequences in marker-gene and metagenomics data. *Microbiome* 6, 226 (2018). [PubMed: 30558668]
71. Frøslev TG et al. Algorithm for post-clustering curation of DNA amplicon data yields reliable biodiversity estimates. *Nat Commun* 8, 1188 (2017). [PubMed: 29084957]
72. Dixon P VEGAN, a package of R functions for community ecology. *Journal of Vegetation Science* 14, 927–930 (2003).
73. McMurdie PJ & Holmes S phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. *PLOS ONE* 8, e61217 (2013). [PubMed: 23630581]
74. Barnett D. J. m, Arts I. C. w & Penders J. microViz: an R package for microbiome data visualization and statistics. *Journal of Open Source Software* 6, 3201 (2021).
75. Lahti L & Shetty S microbiome R package. (2017) doi: 10.18129/B9.bioc.microbiome.
76. Bankevich A et al. SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. *Journal of Computational Biology* 19, 455–477 (2012). [PubMed: 22506599]

77. Wick RR, Judd LM, Gorrie CL & Holt KE Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. *PLOS Computational Biology* 13, e1005595 (2017). [PubMed: 28594827]
78. Schwengers O et al. Bakta: rapid and standardized annotation of bacterial genomes via alignment-free sequence identification. *Microb Genom* 7, 000685 (2021). [PubMed: 34739369]
79. Shade A & Stopnisek N Abundance-occupancy distributions to prioritize plant core microbiome membership. *Current Opinion in Microbiology* 49, 50–58 (2019). [PubMed: 31715441]
80. Fisher CK & Mehta P Identifying Keystone Species in the Human Gut Microbiome from Metagenomic Timeseries Using Sparse Linear Regression. *PLoS One* 9, e102451 (2014). [PubMed: 25054627]
81. Wang X-W et al. Identifying keystone species in microbial communities using deep learning. *Nat Ecol Evol* 8, 22–31 (2024). [PubMed: 37974003]
82. Kurtz ZD et al. Sparse and Compositionally Robust Inference of Microbial Ecological Networks. *PLOS Computational Biology* 11, e1004226 (2015). [PubMed: 25950956]
83. Karkaria BD, Fedorec AJH & Barnes CP Automated design of synthetic microbial communities. *Nat Commun* 12, 672 (2021). [PubMed: 33510148]
84. Toju H et al. Scoring Species for Synthetic Community Design: Network Analyses of Functional Core Microbiomes. *Front. Microbiol* 11, (2020).
85. Paredes SH et al. Design of synthetic bacterial communities for predictable plant phenotypes. *PLOS Biology* 16, e2003962 (2018). [PubMed: 29462153]

Box 1.**Resources and Best Practices for SynCom Design****Resources**

1. Computational resources for processing and preparing amplicon sequencing data (short or long reads): dada2⁴³, Mothur⁶⁷, UNOISE3⁶⁸, UCHIME3⁶⁹, Decontam⁷⁰, LULU⁷¹
2. Resources for tracking and characterizing SynCom/resident microbiota: vegan⁷², Phyloseq⁷³, MicroViz⁷⁴, Microbiome R Package⁷⁵.
3. Sequencing pipelines for assembling and annotating strain genomes: SPADES⁷⁶ or Unicycler⁷⁷ are considered best practices for assembly, while BAKTA⁷⁸ is the defacto option for annotation.
4. Methods for identifying keystone species, core or key functional taxa: Abundance Occupancy curves⁷⁹, LIMITS⁸⁰, the DKI machine learning framework⁸¹, SPIEC-EASI⁸²
5. Resources for the automated design and predictive effects of synthetic communities: used by Karkaria et al.⁸³, Toju et al.⁸⁴ and Paredes et al.⁸⁵

Best Practices

1. Methods used to select strains should be documented and published in work referencing the SynCom.
2. Strains should undergo whole-genome sequencing and these data should be made publicly available.
3. Strains within SynComs should be made available to other researchers via deposition in a public collection such as ATCC, DSM, CBS, CIRM etc.
4. Within fields, methods for strain preparation and inoculation should be standardized (i.e. growth conditions prior to experimentation, strain inoculation density, sampling methods during experiment).
5. Integrity of strain freezer stocks should be maintained, and best practices followed, as it is vital to prevent these strains from becoming lab adapted. Freeze thaw cycles and the number of passages of a strain in non-native conditions should be minimized and documented. Strains should periodically be validated for identity and redundant copies of the library should be stored in separate locations.




Figure 1. Dual continuums of “question” and “system” for SynCom research.

Research questions using SynComs can range from fundamental questions or basic science, that is, trying to understand the rules and functioning underpinning different systems, to applied questions. Here communities are designed to fulfill certain purposes, for example, [AU: please complete this sentence using a brief example from the figure]. Likewise, the system being used can be placed on a continuum from environmental to free living and host-associated microbial communities.




Figure 2. Flow diagram of approaches used when designing, evaluating and deploying a SynCom.

(A) All studies begin by designing the community (green). SynCom design can proceed from either Bottom-up (increase complexity through iterations) or Top-down (reduce complexity through iterations) approaches. When designing communities it is important to consider the number of strains needed to be relevant, as well as the sourcing of those strains. (B) Strains are then prepared and used for inoculation (yellow). Important considerations include the strain growth conditions, applied concentration, experimental system and methods of inoculation. (C) After a SynCom has been implemented, it is critical to evaluate if it provides relevant information about the system being modeled. To do so, the questions must first be well defined, after which the relevant features can be assessed by tracking the composition and functioning of the community. (D) When designing and

applying SynComs across both human and environmental systems, there are important ethical considerations to take into account. In human systems, these communities should be representative of diversity seen across geographic, cultural and economic boundaries, and communities applied to patients should be tested for off target effects. When applying a SynCom to a natural system, care must be taken to ensure that these species do not spread and become invasive.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 3. Examples of bottom-up and top-down design approaches for SynComs.

(A) Bottom-up approaches can include selecting strains that represent the phylogenetic diversity of the natural community at some level, identifying strains that perform some functions of interest in the natural community, or through the prediction of key interactions in the community that a researcher might want to model. (B) Top-down designs can employ host or environmental filtering. This is where a larger community is applied into the study environment and only those strains that pass some growth or persistence metrics are included. It can also be achieved through the recapitulation of key features in community interaction networks or through a sequential drop out, where strains are sequentially removed in order to select the minimal complexity required to model the interactions of interest. In practice these approaches are not mutually exclusive, and researchers can choose to employ a combination of bottom-up or top-down strain selection approaches to define their communities.