



OPEN ACCESS

APPROVED BY
Frontiers Editorial Office,
Frontiers Media SA, Switzerland

*CORRESPONDENCE
Takuya Takahashi
✉ takahast@yokohama-cu.ac.jp

RECEIVED 22 November 2024
ACCEPTED 25 November 2024
PUBLISHED 10 December 2024

CITATION

Yatomi T, Tomasi D, Tani H, Nakajima S,
Tsugawa S, Nagai N, Koizumi T, Nakajima W,
Hatano M, Uchida H and Takahashi T (2024)
Corrigendum:
 α -amino-3-hydroxy-5-methyl-4-isoxazole
propionic acid (AMPA) receptor density
underlies intraregional and interregional
functional centrality.
Front. Neural Circuits 18:1533008.
doi: 10.3389/fncir.2024.1533008

COPYRIGHT

© 2024 Yatomi, Tomasi, Tani, Nakajima,
Tsugawa, Nagai, Koizumi, Nakajima, Hatano,
Uchida and Takahashi. This is an open-access
article distributed under the terms of the
[Creative Commons Attribution License \(CC
BY\)](#). The use, distribution or reproduction in
other forums is permitted, provided the
original author(s) and the copyright owner(s)
are credited and that the original publication
in this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Corrigendum: α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) receptor density underlies intraregional and interregional functional centrality

Taisuke Yatomi^{1,2}, Dardo Tomasi³, Hideaki Tani¹,
Shinichiro Nakajima¹, Sakiko Tsugawa¹, Nobuhiro Nagai¹,
Teruki Koizumi¹, Waki Nakajima², Mai Hatano², Hiroyuki Uchida¹
and Takuya Takahashi^{2*}

¹Department of Neuropsychiatry, Keio University School of Medicine, Tokyo, Japan, ²Department of Physiology, Yokohama City University Graduate School of Medicine, Yokohama, Japan, ³Laboratory of Neuroimaging (LNI), National Institute on Alcohol Abuse and Alcoholism, NIH, Bethesda, MD, United States

KEYWORDS

α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) receptor, [¹¹C]K-2, positron emission tomography, synaptic plasticity, resting-state functional MRI (rsfMRI), functional connectivity density mapping, functional network, functional centrality

A Corrigendum on

[\$\alpha\$ -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid \(AMPA\) receptor density underlies intraregional and interregional functional centrality](#)

by Yatomi, T., Tomasi, D., Tani, H., Nakajima, S., Tsugawa, S., Nagai, N., Koizumi, T., Nakajima, W., Hatano, M., Uchida, H., and Takahashi, T. (2024). *Front. Neural Circuits*. 18:1497897.
doi: 10.3389/fncir.2024.1497897

In the published article, there was an error in the order of affiliation 2 and 3. Instead of “²Laboratory of Neuroimaging (LNI), National Institute on Alcohol Abuse and Alcoholism, NIH, Bethesda, MD, United States, ³Department of Physiology, Yokohama City University Graduate School of Medicine, Yokohama, Japan”, affiliation 2 and 3 should have been correctly written as “²Department of Physiology, Yokohama City University Graduate School of Medicine, Yokohama, Japan, ³Laboratory of Neuroimaging (LNI), National Institute on Alcohol Abuse and Alcoholism, NIH, Bethesda, MD, United States”.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.