



OPEN ACCESS

EDITED AND REVIEWED BY
Marijn Lijffijt,
IonTX, Inc, United States

*CORRESPONDENCE

Kevin T. Mintz
✉ kmintz@stanford.edu

RECEIVED 05 November 2024

ACCEPTED 22 November 2024

PUBLISHED 23 December 2024

CITATION

Mintz KT, Gammer B, Khan AJ, Shaub G,
Levine S and Sisti D (2024) Corrigendum:
Physical disability and psychedelic therapies:
an agenda for inclusive research and practice.
Front. Psychiatry 15:1523364.
doi: 10.3389/fpsy.2024.1523364

COPYRIGHT

© 2024 Mintz, Gammer, Khan, Shaub, Levine
and Sisti. This is an open-access article
distributed under the terms of the [Creative
Commons Attribution License \(CC BY\)](#). The
use, distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication in
this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Corrigendum: Physical disability and psychedelic therapies: an agenda for inclusive research and practice

Kevin T. Mintz^{1*}, Brinn Gammer², Amanda J. Khan^{3,4},
Gretchen Shaub⁵, Steven Levine⁵ and Dominic Sisti²

¹Stanford University, Center for Biomedical Ethics, Stanford, CA, United States, ²Department of Medical Ethics & Health Policy, University of Pennsylvania, Philadelphia, PA, United States, ³Department of Psychiatry, University of California, San Francisco, San Francisco, CA, United States, ⁴Sage Integrative Health, Berkeley, CA, United States, ⁵COMPASS Pathways, London, United Kingdom

KEYWORDS

physical disability, sensory disability, psychedelic therapies, ethics, structural ableism in healthcare, justice, inclusion in clinical research

A Corrigendum on

Physical disability and psychedelic therapies: an agenda for inclusive research and practice

By Mintz KT, Gammer B, Khan AJ, Shaub G, Levine S and Sisti D (2022) *Front. Psychiatry* 13:914458. doi: 10.3389/fpsy.2022.914458

In the published article, there were errors in **Table 2**. The authors wish to correct the errors and clarify data presented in **Table 2**, pertaining to the prevalence of mental illness in several populations.

Under “Mild/Moderate Cerebral Palsy”, the data in column 2 refer to men only. A footnote has been added to clarify this.

Under “Spinal cord injury”, the data in column 2 were reported as 7.1–26.6% and 13.2–40% for Anxiety and PTSD, respectively. The correct values are 13.2–40% and 7.1–26.6%. The table has been updated to reflect this.

Under “Deafness”, the data for Substance Use Disorder in column 2 reflect a population in outpatient psychiatric care. A footnote has been added to clarify this.

Also under “Deafness”, the data in column 2 were reported as 30.1%, 14.2%, 5.2%, 48.4% and 4.9% for Anxiety, Bipolar, Impulse control disorders, Substance Use Disorders and ADHD, respectively. The correct values are 18.7%, 3.7%, 15.8%, 27.8% and 11.2%. The table has been updated to reflect this.

Under “Blindness”, the data in column 2 refer to patients older than 65. A footnote has been added to clarify this.

The corrected **Table 2** and its caption appear below.

The authors apologize for these errors and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

TABLE 2 | Disability community prevalence of mental disorders.

Disabling condition	Prevalence within disability communities	Source
MS	MDD - 46% Anxiety - 16.5% Bipolar - 2.4% Schizophrenia - 2%	Marrie RA, Horwitz R, Cutter G, Tyry T, Campagnolo D, Vollmer T, 2009 (33)
ALS	MDD - 17%	Zucchi E, Ticozzi N, Mandrioli J, 2019 (34)
Mild/Moderate Cerebral Palsy ^a	Anxiety Disorders - 19.5% Mood Disorders - 19.5% Schizophrenia - 2.8%	Whitney DG, Warschausky SA, Ng S, Hurvitz EA, Kamdar NS, Peterson MD, 2019 (35) ^a
Spinal cord injury	MDD - 20-30% Anxiety - 13.2-40% PTSD - 7.1–26.6%	Post M, van Leeuwen, C, 2012 (36)
Traumatic brain injury	MDD - 24.5% Anxiety - 24.5% Bipolar - -1.6% Schizophrenia - 4.0% PTSD - 16.5%	Rogers JM, Read CA, 2007 (37)
Deafness ^b	Substance Use Disorder - 11.5% Anxiety Disorders – 18.7% Bipolar – 3.7% Impulse control disorders – 15.8% Substance Use Disorders – 27.8% ADHD – 11.2%	Diaz DR, Landsberger SA, Povlinski J, Sheward J, Sculley C, 2013 (38) ^b
Blindness ^c	MDD - 18.2% Anxiety Disorders - 14.3% Schizophrenia - 1.4% Alcohol Misuse - 3.3% Psychoactive Substance Misuse - 12.9% Anorexia or Bulimia -.5%	Court H, McLean G, Guthrie B, Mercer SW, Smith DJ, 2014 (39) ^c

^aStatistics in this table reflect the statistics for men with CP in this article.

^bThis article is focused on a population in outpatient psychiatric care.

^cThis article is focused on patients 65 and older in the UK.

The authors thank Jacob Kramer, an MSW student at Rhode Island College for helping bring these errors to our attention, particularly with regard to the D/deaf community.