

CONTROL OF THE HYPERPOLARIZATION-ACTIVATED CATION CURRENT BY EXTERNAL ANIONS IN RABBIT SINO-ATRIAL NODE CELLS

By A. M. FRACE, F. MARUOKA AND A. NOMA

From the Department of Physiology, Faculty of Medicine, Kyushu University, Fukuoka, 812, Japan

(Received 17 October 1991)

SUMMARY

1. Effects of varying concentrations of anions on the hyperpolarization-activated current (I_f) were studied in myocytes isolated from the rabbit sino-atrial node. Substituting Cs^+ for the intracellular K^+ clearly separated I_f from the delayed rectifier K^+ current. Control properties, including gating kinetics and ion selectivity, similar to previous studies were obtained.

2. Substitution of extracellular Cl^- with larger anions including isethionate, glutamate, acetate, and aspartate, reduced the amplitude of I_f without changing the reversal potential. Substitution with small anions such as iodide or nitrate supported an intact I_f . These effects were reproduced in the excised outside-out patch conformation.

3. The conductance for I_f was a saturating function of the extracellular Cl^- concentration ($[\text{Cl}^-]_o$) with an equilibrium binding constant ($K_{1/2}$) of 11 mM and a slope factor of about 1 when substituted with large anions. Total removal of small anions completely abolished I_f .

4. The voltage-dependent gating of I_f was not affected by changing ($[\text{Cl}^-]_o$), suggesting that Cl^- modulates conductance properties of I_f .

5. The results indicate that I_f conductance is unique in that it is dependent on an extracellular anion (Cl^-), yet it is carried exclusively by cations, K^+ and Na^+ . These effects are independent of any measurable voltage-dependent gating parameters.

INTRODUCTION

In various types of cells having a relatively low resting potential, hyperpolarization of the membrane induces an activation of an inward current (I_f or I_h). The cell types include cardiac pacemaker cells (Seyama, 1979; Yanagihara & Irisawa, 1980; DiFrancesco, Ferroni, Mazzanti & Tromba, 1986), atrio-ventricular node cells (Noma, Irisawa, Kokobun, Kotake, Nishimura & Watanabe, 1980) and Purkinje cells (DiFrancesco, 1981*a, b*, 1982), photoreceptors (Bader & Bertrand, 1984), and neurones in spinal sensory ganglia, and central nervous system (Mayer & Westbrook, 1983; McCormick & Pape, 1990). As a common characteristic among different cells,

I_f shows a very slow time course of activation on hyperpolarization, relative to its quick deactivation on depolarization. The conductance of the I_f channel is characterized by its multi-ionic nature. The amplitude of I_f is increased by raising the extracellular K^+ concentration ($[K^+]_o$) and decreased by reducing $[Na^+]_o$. The measurement of the reversal potential suggests that the current is carried by both K^+ and Na^+ (DiFrancesco, 1981*a*). However, since the early examination of I_f (Seyama, 1979), several studies have demonstrated that the current is also sensitive to alteration of $[Cl^-]_o$. Substitution of Cl^- with large anions reduces the amplitude of I_f (Yanagihara & Irisawa, 1980; Mayer & Westbrook, 1983; McCormick & Pape, 1990). The finding may suggest that Cl^- also carries current through the I_f channel under an assumption that the intracellular Cl^- is depleted in the absence of external Cl^- (Seyama, 1979). Alternatively, the finding may be explained by assuming a non-specific blocking action of impermeable anions (McCormick & Pape, 1990), or by assuming an activation of the channel by Cl^- (present study).

In recent years several groups have refined the isolation technique of single sinoatrial (SA) node cells and are compiling more data on I_f characteristics as well as its role in the pacemaker mechanism (DiFrancesco *et al.* 1986; Yatani & Brown, 1990; Denyer & Brown, 1990; van Ginneken & Giles, 1991). We have also developed an improved procedure for isolating SA node cells, in general agreement with these groups. We have undertaken a whole-cell voltage clamp analysis on the activation of I_f and its dependence on anions by performing systematic ion-replacement experiments on isolated SA node cells.

METHODS

SA nodal cell isolation

Albino rabbits weighing 1–2 kg were anaesthetized with intravenous injection of sodium pentobarbitone (30–50 mg/kg). The heart was dissected out and mounted on a modified Langendorff perfusion system and perfused with Tyrode solution followed by a Ca^{2+} -free Tyrode solution for 3–5 min. After contractions ceased 300 ml of a Ca^{2+} -free Tyrode solution containing collagenase (Yakult, Tokyo, Japan; 0.13–0.03 mg/ml, depending on lot number) was perfused followed by 150 ml of Ca^{2+} -free Tyrode solution containing protease (Sigma, 0.1 mg/ml). The SA node region was then dissected out, trimmed, and cut into strips. A second Ca^{2+} -free collagenase digestion (0.6–0.4 mg/ml) was then performed for 10–30 min. The digested SA node tissue was then washed in a 'KB' solution (Isenberg & Klöckner, 1982) and dispersed by triturating with a fire-polished pipette. Dispersed cells were stored in 'KB' solution at 4 °C for up to 1 day. Cells used in this study were identified visually as long, spindle-shaped cells approximately 50–80 μm in length and 10–15 μm in width in normal Tyrode solution. These cells have faint striations and a prominent, centrally located nucleus (similar to type A in Denyer & Brown, 1990). Other cell types were generally larger, rod shaped, and striated, or of rounded shape. The present cell-isolation procedure and solutions used provide an I_f that exhibits very little run-down in normal Na^+ - or K^+ -Tyrode solution.

Solutions

Normal Tyrode solution contained (mM): NaCl, 140; KCl, 5.4; $MgCl_2$, 0.5; $CaCl_2$, 1.8; NaH_2PO_4 , 0.33; glucose, 5.5 and HEPES, 5.0 (pH = 7.4). In recording I_f , $BaCl_2$ (2 mM) and $NiCl_2$ (1 mM) were also added to block other currents. In experiments which examined exchanging external anions, a sodium salt of aspartate, glutamate, acetate, isethionate, iodide, or nitrate was substituted for NaCl on an equimolar basis. In solutions requiring low Cl^- concentrations, $MgCl_2$, $NiCl_2$ and $BaCl_2$ were omitted and calcium gluconate was used. Variations in $[Na^+]_o$ or $[K^+]_o$ were made by substituting them with Tris. All solutions were used at 37 °C.

The internal solution contained (mM): potassium aspartate or caesium aspartate, 110; KCl, 20; CaCl₂, 2.0; EGTA, 5.0; MgCl₂, 5.0; K₂ATP, 5.0, and HEPES, 5.0 (pH = 7.2). The free Ca²⁺ is calculated to be 0.1 μM; 100 μM-GTP and 50 μM-cyclic AMP were included. In some instances 130 mM-CsCl was used as the major intracellular salt.

Voltage clamp recording and analysis

Whole-cell and isolated patch recordings were performed using the original methods of Hamill, Marty, Neher, Sakmann & Sigworth (1981). Currents were recorded with a patch clamp amplifier (List EPC-7, Darmstadt, Germany). The current, voltage, and trigger signals were stored on video tape using a PCM converter system (NF Electronic Instruments, RP-882, Tokyo, Japan) for subsequent computer analysis (NEC PC98 XL, Tokyo, Japan).

The liquid junction potentials of the aspartate-rich K⁺ and Cs⁺ internal solutions were -11.4 ± 0.3 mV ($n = 10$) and -12.3 ± 0.7 mV ($n = 7$), respectively, when in contact with the control Tyrode solution. The measurement of junction potential drifted by a few millivolts when the electrode pipette was kept in the Tyrode solution, most probably due to diffusion at the pipette tip. We do not know exactly how large the junction potential was between the pipette solution and the intracellular solution, although it should be small. Therefore, voltage recordings measured in the aspartate-rich internal solution were corrected by -10 mV for practical purposes. To minimize junction-potential error during Cl⁻ variation a 3 M-KCl leakage electrode was used as a bath reference electrode. This reference system maintains a small, continuous leak of 3 M-KCl solution from a reservoir. The reference electrode was put downstream from the preparation. In other instances a large 3 M-KCl-agar bridge was used. Data are presented as means \pm standard deviation.

RESULTS

Isolation of the hyperpolarization-activated current using the Cs⁺-rich internal solution

In nearly every spindle-shaped cell examined a large, inward current was activated on hyperpolarization negative to approximately -70 mV. After the initial break-in and voltage clamp, the current often decreased or increased to a steady-state level within minutes, presumably due to equilibration with the pipette solution, but the threshold potential for activation was not obviously changed. Rapid run-down of the current, as described in DiFrancesco *et al.* (1986), was not evident in normal Tyrode solution or in high-K⁺ solutions, presumably due to a higher intracellular Ca²⁺ of pCa 7 (Hagiwara & Irisawa, 1989) and inclusion of cyclic AMP and GTP (Yatani & Brown, 1990).

The activation of I_f on hyperpolarization is usually overlapped with a time-dependent deactivation of the delayed rectifier K⁺ current. To isolate I_f, the delayed rectifier K⁺ current was blocked by adding 2 mM-Ba²⁺ and 1 mM-Ni²⁺ in the external solution, which also partially inhibited the time-independent background conductance. For complete suppression of the outward current of the delayed rectifier K⁺ channel, Cs⁺-rich internal solution was used. Figure 1A shows a family of I_f recordings in response to single hyperpolarizing steps. It is evident that the current trace at -60 mV shows no obvious time-dependent change and very slow activation of I_f is visible at -70 mV. At stronger hyperpolarizations I_f showed exponential activation at the onset of the pulse. With the Cs⁺-rich solution the sigmoidal time course of the initial current was not significant.

The amplitude of the inward I_f tail, which was recorded on terminating the hyperpolarizing pulse, increased with increasing conditional hyperpolarization and showed a trend of saturation. Voltage-dependent activation of I_f was assessed by

measuring the tail current. Two different current parameters including integration of the area of the tail current or measurement of the peak amplitude of tail current were used. The values were normalized to the maximum value in each experiment. Data obtained using the K^+ -rich solution in the pipette are also shown for comparison. We




Fig. 1. Control current properties of I_t recorded with the Cs^+ -rich internal solution. *A*, original current recordings are superimposed. The pulse protocol is shown in the lower part: the holding potential was -50 mV and a 4.5 s pulse is applied in 10 mV increments. Cs^+ pipette solution; Na^+ Tyrode solution with 2 mM- Ba^{2+} and 1 mM- Ni^{2+} . *B*, voltage-dependent activation of I_t , determined in Na^+ Tyrode solution using K^+ (\circ) or Cs^+ (\bullet) pipette solution. The tail current values are normalized and plotted against membrane potential for six experiments. The continuous line is a fit with eqn (1).

conclude that the activation curve is not affected by replacing internal K^+ with Cs^+ . All data points in Fig. 1*B* were then fitted with the following Boltzmann relation:

$$K_{act} = \frac{1}{1 + \exp\{QF(V_m - V_{\frac{1}{2}})/RT\}}, \quad (1)$$

where K_{act} is the activation parameter, Q is valency of gating charge, V_m is the membrane potential, $V_{\frac{1}{2}}$ is the potential at half-maximum activation, and F , R and T have their usual meanings in thermodynamics. Regardless of the major intracellular cation or means of measuring the current amplitude, the activation profile in Na^+ Tyrode solution displays a threshold near -70 mV, a $V_{\frac{1}{2}}$ of 93.1 ± 8.4 mV, $Q = 3.03 \pm 0.55$ ($n = 6$; 4 intracellular Cs^+ , 2 intracellular K^+). These data are shown in Fig. 1*B* with a representative fitted line.

The 'instantaneous' $I-V$ relations were measured from a double-pulse tail current protocol as indicated in the insets of Fig. 2. Under physiological ionic conditions the tail current reversed from inward to outward with depolarization at -32.7 ± 1.4 mV ($n = 19$), which is slightly less negative than the intersection of the instantaneous $I-V$ curve. This is because the instantaneous $I-V$ relation is also determined by other background conductances. The value of the reversal potential is in good agreement with previous studies if corrected for the junction potential (DiFrancesco, 1981*b*; van Ginneken & Giles, 1991). The value of the Na^+ conductance relative to the K^+ conductance (pNa/pK) is 0.27 according to the Goldman-Hodgkin-Katz equation. With the Cs^+ -rich solution the tail current was always inward and the instantaneous

I-V relation was a smooth curve (Fig. 2*B*). Thus, it is concluded that internal Cs⁺ neither carries outward current, nor affects the inward flux through the I_t channel. In the following experiments, data obtained with the K⁺-rich solution were also used when the separation of potassium current (I_K) from I_t was not critical in interpreting the experimental data.




Fig. 2. Late current (●) and 'instantaneous' current-voltage relationships (○) recorded in potassium aspartate (*A*) and caesium aspartate pipette solutions (*B*) with the control Na⁺ Tyrode solution. The late current was measured near the end of the hyperpolarizing pulse of the single-pulse protocol as shown in Fig. 1*A*. Current traces and the voltage protocols for the instantaneous *I-V* relationships are shown in the insets.

Effects of anion replacement

Substitution of the majority of extracellular Cl⁻ in Na⁺ Tyrode solution by several organic anions was found to greatly reduce the amplitude of I_t, whereas substitution with small anions such as I⁻ or NO₃⁻ supported a full-amplitude I_t with properties indistinguishable from control. Figure 3*A* shows representative records comparing the currents between control external Cl⁻ and NO₃⁻ (upper panel) or control Cl⁻ and isethionate (lower). The amplitude of I_t was reduced to about 30% of the control in the isethionate solution, but the time course of the current was not changed. The effects of different anions on the current amplitude were examined by normalizing the *I-V* relation referring to the amplitude of I_t at -120 mV in the control. It was found, regardless of species, that all the organic anions tested reduced I_t to nearly the same extent, and that the current amplitude was almost identical with Cl⁻, I⁻, and NO₃⁻ (Fig. 3*B*). The amplitude of the current in the organic anion solutions is about 30% of the control over the potential range -120 to -90 mV, indicating no obvious voltage dependence of the effect of Cl⁻ replacement. These findings are difficult to explain by assuming a blocking action of organic anions (McCormick & Pape, 1990), since the molecular structure of these ions is different.

No measurable Cl⁻ permeability was evident from experiments of replacing external Cl⁻. The reversal potential of I_t was measured by plotting the instantaneous




Fig. 3. *A*, current traces at -110 mV before and after substituting NO_3^- (upper panel) and isethionate (lower panel) for Cl^- in Tyrode solution are superimposed. *B*, late current-voltage relationships in Na^+ Tyrode solution using different extracellular anions, aspartate (■), glutamate (▽), acetate (◆), isethionate (◇), Cl^- (△), I^- (□), or NO_3^- (●). The Cl^- from CaCl_2 , MgCl_2 , NiCl_2 and BaCl_2 remains in the extracellular solution (10.6 mM). Currents were normalized to the peak inward current at the maximum hyperpolarized potential. The instantaneous current jumps at the pulse onset were assumed to be due to leak current systems and were subtracted.




Fig. 4. The instantaneous *I-V* relations in K^+ Tyrode solution containing 150 (●) and 13 (○) mM- Cl^- aspartate substitution in *A*, and those obtained at a constant $[\text{Cl}^-]_o$ of 150 mM in 24 mM $[\text{K}^+]_o$ (○; Tris substitution) and 150 mM $[\text{K}^+]_o$ solutions (●) in *B*. Pulse protocol is essentially the same as shown in Fig. 2.

I-V relation before and after the reduction of $[\text{Cl}^-]_o$ from 150 to 13 mM (aspartate replacement) in 150 mM-external Na^+ (Fig. 4*A*). The slope of the instantaneous *I-V* curve is reduced at all potentials at low $[\text{Cl}^-]_o$. Although the Cl^- reversal potential was expected to change by about 64 mV from -39 mV in the control to $+25$ mV in 13 mM- Cl^- , the reversal potential remained constant at around -33 mV, which is determined by both Na^+ and K^+ . This finding may also indicate that the ion

selectivity, pNa/pK, is not affected by lowering [Cl⁻]_o. When [K⁺]_o was changed from 150 to 24 mM at a constant [Cl⁻]_o in the Na⁺-free solution (Tris-HCl replacement), the reversal potential shifted as expected from a change in the K⁺ equilibrium




Fig. 5. Effects of varying [Cl⁻]_o in the whole cell (*A* and *B*) and the outside-out patch (*C*) in K⁺ Tyrode solution. *A*, current traces obtained in control 150 mM-Cl⁻ and after total replacement of Cl⁻ with aspartate are superimposed. The pulse protocol is indicated above the current. *B*, currents recorded at [Cl⁻]_o of 2, 4, 14 and 150 mM are superimposed. *C*, the control records at 150 mM [Cl⁻]_o obtained before (150, lower trace) and after (arrow-head) the application of 10 mM-Cl⁻ are superimposed with the test run (10).

potential (Fig. 4*B*). Thus we conclude that Cl⁻ is not permeating the I_f channel, which is a cation channel.

Dose-dependent potentiation of I_f by small anions

An activating action by Cl⁻, rather than I_f channel block by organic anions, was strongly supported by recording I_f at very low [Cl⁻]_o. The K⁺ Tyrode solution was used to obtain the maximum current in these experiments. On complete substitution of Cl⁻ with aspartate virtually no I_f was recorded, even with large voltage steps (Fig. 5*A*). Addition of only 2, 4, or 14 mM-Cl⁻ to the external solution induced an I_f of progressively larger amplitudes during hyperpolarizing pulses (Fig. 5*B*). In these solutions changes in the concentration of organic anions is less than 10% of the control concentration. It is difficult to assume that decreasing the concentration of the organic anion from 148 to 136 mM progressively released the channel block.

The cell-free patch recording of channel current may be free from complications arising from cell-induced mechanisms, such as changes in cell volume or intracellular ionic concentrations and pH, stimulated by the Cl⁻ replacement. Recording of I_f from a cell-free patch has already been reported, initially by Yatani & Brown (1990). Therefore, the protocol was also performed using isolated outside-out patches. Although the I_f in this conformation underwent run-down, currents were rapidly reduced in 10 mM-Cl⁻ and recovered in control Cl⁻ solution (Fig. 5*C*). Also, experiments were conducted with Cl⁻ replacing aspartate as the major intracellular anion ([Cl⁻]_i = 140 vs. 24 mM in the normal control) with no change in the effect.

The findings described above strongly suggest that the external presence of small anions is essential for activation of the I_f channel by hyperpolarization. Dose dependence of this effect was quantified by measuring the steady-state current at




Fig. 6. Dose-response relationship for peak I_t vs. $[Cl^-]_o$. The amplitudes of I_t in each experiment ($n = 32$) are normalized to the peak current obtained at -120 mV in control 150 mM-KCl Tyrode solution which preceded the reduced Cl^- application. The continuous line is the fit to the Hill equation. The arrow-head points to the $K_{1/2}$ concentration.




Fig. 7. Voltage-dependent activation in various $[Cl^-]_o$ (Δ) compared to those obtained in the 150 mM $[Cl^-]_o$ in K^+ Tyrode solution (\blacklozenge). Data in each experiment are normalized to the peak tail current. Continuous line, fitting eqn (1), is for the control K^+ data.

a single, large, hyperpolarizing test potential (-120 mV) over a wide range of $[Cl^-]_o$ (substituting with glutamate or aspartate) and a dose-response relation was compiled ($n = 32$). The concentration dependence was fitted with a single-site binding relation with an equilibrium binding constant ($K_{1/2}$) of 11.5 mM and a Hill coefficient of 1.13 (Fig. 6).

Mechanisms of I_t depression in low- Cl^- solution

The reduction in the amplitude of I_t might be caused by a change in the voltage-operated gating kinetics, or by a reduction in the single-channel conductance in the low- Cl^- solution. We failed to record single-channel events in the cell-attached patch recording or in the excised outside-out patch, most probably due to extremely small

single-channel conductance (DiFrancesco, 1986). However, we could test the gating kinetics at low [Cl⁻]_o. Figure 7 compares the quasi-steady-state activation-voltage relationship between I_t in the lower [Cl⁻]_o solutions and that measured at 140 mM [Cl⁻]_o in the K⁺ Tyrode solution. In six experiments, there was little effect




Fig. 8. Semilog plots of the time-dependent activation of I_t . The upper panel shows those for control 150 mM [Cl⁻]_o in K⁺ Tyrode solution, and the lower panel at 14 mM [Cl⁻]_o (aspartate substitution). The total time-dependent current (upper point plots) and the subtracted fast exponential component (lower plots) are shown. The theoretical curve, a sum of the slow and fast components, was superimposed on the original current in the insets, although the good fit made the two traces indistinguishable. The horizontal bar in the inset indicates the duration of the test pulse (-100 mV) of 5 s, and the vertical bar 1 nA. See text for detail.

on the voltage-dependent activation parameters over this concentration range. The $V_{\frac{1}{2}}$ in low Cl⁻ was 94.4 ± 3.7 mV and Q was 2.53 ± 0.56 .

The rate of I_t activation was measured by plotting the current on a logarithmic scale. In Fig. 8, the ordinate indicates the current amplitude measured in reference to an assumed steady-state level. Since the current did not reach a steady level

during the pulse, it was necessary to assume a steady-state level. To determine an appropriate value, the reference level was arbitrarily shifted until the semi-logarithmic plot became linear near the end of the pulse as shown in the figure. By calculating the regression line over the late linear portion, the slow component was determined first. Then this slow component was subtracted from the original current, and the difference was replotted to measure the fast component. The sum of these two exponential components almost perfectly fitted the original current record as shown in the inset of Fig. 8. It should be noted that at small hyperpolarizations a single exponential often gave a good fit to the currents. On reduction of the current in 14 mM $[Cl^-]_o$ the current kinetics are nearly unaffected with rate constants (τ) of 0.38 and 1.67 as compared with the control values of 0.36 and 1.56 s. Essentially the same results were obtained in three experiments.

DISCUSSION

The results of this paper address the question of how substitution of extracellular Cl^- by large anions reduces the size of I_f . This effect of Cl^- substitution has been described, initially by Seyama (1979), followed by Yanagihara & Irisawa (1980) in cardiac SA node. More recently, a similar phenomenon has been described for the very similar I_f in spinal sensory ganglion neurones (Mayer & Westbrook, 1983) and in thalamic relay neurones (McCormick & Pape, 1990). These latter reports suggested a non-specific blocking activity for the large anions. On further Cl^- removal or on addition of only a few millimoles of Cl^- in the presence of high concentrations of organic anions (Fig. 3), it is clear that I_f is directly dependent on small anions in the extracellular solution. Anions of < 0.25 nm seem to be able to support a normal current while larger, organic anions allow none. The response is a membrane-delimited one in that it is sustained in an excised-patch conformation. Also, intracellular Cl^- depletion was avoided in experiments containing high Cl^- in the pipette solution.

Dual involvement of anions and cations in a single-conductance system is not a novel result, having been described for anion permeability of gramicidin channels (Neher, 1975) and for cation permeability of an anion channel in hippocampal neurones (Franciolini & Nonner, 1987). However, each of these systems is found to have an experimentally measurable permeability of both anion and cation. We can find no measurable Cl^- permeability in the I_f system. No shifts in reversal potential are found after large changes in $[Cl^-]_o$. Furthermore, removing Cl^- from the bath solution for the cardiac and neuronal preparations would lead to an increase of inward current rather than a decrease, if anion permeation is a factor.

A model for binding of impermeant anions has been proposed to interpret changes in thallos ion conductance in the gramicidin channel (Eisenman, Sandbloom & Neher, 1978). The model proposes that small anions perform a screening role for cations bound at external sites of a multi-ion pore. The effects on gramicidin channel conductance are quite small in comparison with the present result. The cardiac I_f conductance was totally eliminated by completely omitting external Cl^- . We postulate that such a type of positive charge screening, whether it be a fixed charge at a recessed site on the protein or a bound cation is a necessary step in I_f channel permeation by cations, such as Na^+ and K^+ . Access by the large organic anions to the screening sites may be sterically blocked by a small tunnel structure of the channel

protein. The voltage-independent nature of this Cl⁻ effect suggests that these interactions should be out of the membrane potential field in the case of the cardiac I_f channel. This view is consistent with the finding that the voltage-operated gating properties were not modified by varying the Cl⁻ concentration. The voltage-dependent gating should depend on a molecular structure within the membrane potential field. The fact that voltage-dependent activation and I_f kinetics are unaffected by Cl⁻ variation suggests that the manner by which Cl⁻ exerts its effect is by varying channel conductance. While a change in either unit conductance or open probability could produce such a macroscopic result, it seems likely that such a large change in open probability would also be reflected in either gating kinetics or activation of I_f, as it is in other ion-gated channels (for example the Ca²⁺-activated K⁺ channel, Barrett, Magleby, & Pallotta, 1982). Although a novel Cl⁻ binding scheme which produces a functional channel in an all-or-none manner, without subsequent effects on channel properties, could be argued, we reason that a Cl⁻ effect on I_f channel permeation is the most likely interpretation. Direct single-channel measurement of these parameters is required to test these hypotheses.

The present study succeeded in separating the activation of I_f from the deactivation of the I_K by using Cs⁺ in the internal solution. It should be noted that the deactivation of I_K causes a time-dependent change of the whole-cell current in the same direction as the activation of I_f over the potential range positive to the K⁺ equilibrium potential. In the present study, the outward current through the delayed rectifier K⁺ channel was suppressed by replacing the intracellular K⁺ with Cs⁺, but the inward current through I_f channel was not blocked by the Cs⁺. The finding is in contrast to the block of the I_f channel with Cs⁺ from the external side (DiFrancesco, 1982; Noma, Morad & Irisawa, 1983). It may be speculated that the I_f channel does not allow access of Cs⁺ from the internal mouth of the channel. The asymmetrical effect of Cs⁺ across the membrane is well established for the inward rectifier K⁺ channel (see, for example, Matsuda & Noma, 1984). If the ion selectivity of the I_f channel in the presence of 5.4 mM-K⁺ in the external solution is determined by the Goldman-Hodgkin-Katz equation using the reversal potential of about -33 mV in the present study, the pNa/pK ratio is only 0.27. Thus, the I_f channel is also relatively selective for K⁺.

We thank Professor W. Trautwein for his valuable suggestions during his stay at Kyushu University. Thanks are also due to Fumiko Katsuda and Miho Fukushima for secretarial services. This work was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Science and Culture, Japan. M.F. was supported by the Japan Society for the Promotion of Science.

REFERENCES

- BADER, C. R. & BERTRAND, D. (1984). Effect of changes in intra- and extracellular sodium on inward (anomalous) rectification in salamander photoreceptors. *Journal of Physiology* **347**, 611-631.
- BARRETT, J. N., MAGLEBY, K. L. & PALLOTTA, B. S. (1982). Properties of single calcium-activated potassium channels in cultured rat muscle. *Journal of Physiology* **331**, 211-230.
- DENYER, J. C. & BROWN, H. F. (1990). Rabbit sino-atrial node cells: isolation and electrophysiological properties. *Journal of Physiology* **428**, 405-424.
- DI FRANCESCO, D. (1981a). A new interpretation of the pace-maker current in calf Purkinje fibres. *Journal of Physiology* **314**, 359-376.

- DiFRANCESCO, D. (1981b). A study of the ionic nature of the pace-maker current in calf Purkinje fibres. *Journal of Physiology* **314**, 377–393.
- DiFRANCESCO, D. (1982). Block and activation of the pace-maker channel in calf Purkinje fibres: effects of potassium, caesium, and rubidium. *Journal of Physiology* **329**, 485–507.
- DiFRANCESCO, D. (1986). Characterization of single channels in cardiac sino-atrial node cells. *Nature* **324**, 470–473.
- DiFRANCESCO, D., FERRONI, A., MAZZANTI, M. & TROMBA, C. (1986). Properties of the hyperpolarizing-activated current (i_t) in cells isolated from the rabbit sino-atrial node. *Journal of Physiology* **377**, 61–88.
- EISENMAN, G., SANDBLOOM, J. & NEHER, E. (1978). Interactions in cation permeation through the gramicidin channel: Cs, Rb, K, Na, Li, Tl, H, and effects of anion binding. *Biophysical Journal* **22**, 307–340.
- FRANCIOLINI, F. & NONNER, W. (1987). Anion and cation permeability of a chloride channel in rat hippocampal neurons. *Journal of General Physiology* **90**, 453–478.
- HAGIWARA, N. & IRISAWA, H. (1989). Modulation by intracellular Ca^{2+} of the hyperpolarization-activated inward current in rabbit single sino-atrial node cells. *Journal of Physiology* **409**, 121–141.
- HAMILL, O. P., MARTY, A., NEHER, E., SAKMANN, B. & SIGWORTH, F. J. (1981). Improved patch-clamp techniques for high resolution current recording from cells and cell-free membrane patches. *Pflügers Archiv* **391**, 85–100.
- ISENBERG, G. & KLÖCKNER, U. (1982). Calcium tolerant ventricular myocytes prepared by preincubation in a 'KB medium'. *Pflügers Archiv* **395**, 6–18.
- MCCORMICK, D. A. & PAPE, H.-C. (1990). Properties of a hyperpolarization-activated cation current and its role in rhythmic oscillation in thalamic relay neurones. *Journal of Physiology* **431**, 291–318.
- MATSUDA, H. & NOMA, A. (1984). Isolation of calcium current and its sensitivity to monovalent cations in dialysed ventricular cells of guinea-pig. *Journal of Physiology* **357**, 553–573.
- MAYER, M. L. & WESTBROOK, G. L. (1983). A voltage-clamp analysis of inward (anomalous) rectification in mouse spinal sensory ganglion neurones. *Journal of Physiology* **340**, 19–45.
- NEHER, E. (1975). Ionic specificity of the gramicidin channel and thallos ion. *Biochimica et Biophysica Acta* **401**, 540.
- NOMA, A., IRISAWA, H., KOKOBUN, S., KOTAKE, H., NISHIMURA, M. & WATANABE, Y. (1980). Slow current systems in the A-V node of the rabbit heart. *Nature* **285**, 228–229.
- NOMA, A., MORAD, M. & IRISAWA, H. (1983). Does the 'pacemaker current' generate the diastolic depolarization in the rabbit SA node cells? *Pflügers Archiv* **397**, 190–194.
- SEYAMA, I. (1979). Characteristics of the anion channel in the sino-atrial node cell of the rabbit. *Journal of Physiology* **294**, 447–460.
- VAN GINNEKEN, A. C. G. & GILES, W. (1991). Voltage clamp measurements of the hyperpolarization-activated inward current I_t in single cells from rabbit sino-atrial node. *Journal of Physiology* **434**, 57–83.
- YANAGIHARA, K. & IRISAWA, H. (1980). Inward current activated during hyperpolarization in the rabbit sinoatrial node cell. *Pflügers Archiv* **385**, 11–19.
- YATANI, A. & BROWN, A. M. (1990). Regulation of cardiac pacemaker current I_t in excised membranes from sinoatrial node cells. *American Journal of Physiology* **285**, H1947–1951.