

Commentary

Narcotic-based sedation regimens for critically ill mechanically ventilated patientsPratik Pandharipande¹ and E Wesley Ely²¹Assistant Professor, Department of Anesthesiology/Division of Critical Care, Vanderbilt University School of Medicine, Nashville, Tennessee, USA²Associate Professor, Department of Medicine, Center for Health Services Research and Division of Allergy/Pulmonary/Critical Care Medicine, Vanderbilt University School of Medicine and the VA Tennessee Valley Geriatric Research, Education and Clinical Center (GRECC), Nashville, Tennessee, USACorresponding author: Pratik Pandharipande, pratik.pandharipande@vanderbilt.edu

Published online: 18 April 2005

This article is online at <http://ccforum.com/content/9/3/247>

© 2005 BioMed Central Ltd

Critical Care 2005, **9**:247-248 (DOI 10.1186/cc3523)See related research by Breen *et al.* in this issue [<http://ccforum.com/content/9/3/R200>]**Abstract**

Sedatives and analgesics are routinely used in the intensive care unit to relieve pain and anxiety. These agents have numerous side effects and may contribute to poor outcomes such as increased length of mechanical ventilation, longer ICU stays and acute and long-term cognitive dysfunction. Modifying sedation paradigms utilizing either narcotic-based regimens with remifentanyl or fentanyl, or by using α_2 agonists such as dexmedetomidine may help in improving these outcomes in critically ill patients.

Benzodiazepines and narcotics are an integral component of the pharmacological treatment of millions of critically ill mechanically ventilated patients. This month in *Critical Care*, Breen and coworkers [1] report that narcotic-based sedation strategies, compared with the traditional approach involving benzodiazepines with supplemental analgesics for pain, might have improved patient outcomes. In this prospective, randomized, unblinded trial, target-based sedation with remifentanyl and rescue midazolam was associated with decreased duration of mechanical ventilation and shorter times to extubation compared with sedation with benzodiazepines, with rescue fentanyl or morphine. The study has important ramifications in the light of recent investigations that have questioned whether currently used strategies for providing sedation and analgesia in mechanically ventilated patients are optimal.

The Society of Critical Care Medicine, in its 2002 clinical practice guidelines [2], recommended protocolized target-based sedation. Those guidelines recommend the use of lorazepam for long-term sedation and midazolam or propofol for short-term sedation, with fentanyl or morphine for analgesia. Unfortunately, benzodiazepines have numerous adverse effects, including the potential for prolonged ventilation, development of delirium and contribution to

chronic cognitive dysfunction [3-7]. Hence, modification to delivery patterns of sedatives or changing sedation paradigms could alter patient outcomes.

Prior studies [6,8-10] have demonstrated reductions in duration of mechanical ventilation and intensive care unit (ICU) length of stay with protocolized, target-based sedation and daily wake-up trials, and by modifying the route of administration of these drugs (intermittent versus continuous infusion). Although sedatives and analgesics act on the central nervous system, studies comparing sedative and analgesic regimens have tended not to report neurological outcomes. This was not possible until recently because of the lack of monitoring tools that could measure outcomes such as delirium in nonverbal patients. Newer reliable and validated sedation scales such as the Sedation Agitation Scale, Richmond Agitation Sedation Scale, and the Confusion Assessment Method in the ICU for detecting delirium in mechanically ventilated patients permit measurement of these outcomes [11-14]. This is important because delirium is an independent predictor of prolonged mechanical ventilation, longer ICU stay and mortality [15].

Benzodiazepines and propofol act primarily as γ -aminobutyric acid (GABA) agonists to exert their sedative effects. Along similar lines, alcohol is an agonist at the GABA receptor, which is believed to be the probable mechanism underlying the cognitive impairment seen in alcoholic persons [16,17]. Use of novel agents that are GABA receptor sparing (e.g. remifentanyl at the μ opioid receptor or dexmedetomidine at the α_2 receptors) may help to reduce exposure to benzodiazepines and consequently the degree of brain dysfunction, and may improve outcomes such as duration of mechanical ventilation and ICU length of stay.

Remifentanyl has been shown to be efficacious and safe in critically ill patients [18]. It is an attractive agent for sedation in the ICU because of its favorable pharmacokinetic and pharmacodynamic profiles. The drug is metabolized by non-specific plasma esterases and is therefore independent of the function of organs such as the liver and kidneys. Studies in critically ill patients [19,20] have demonstrated a predictable and constant offset time, with little or no accumulation of drug over time. However, there are conflicting reports on the development of tolerance as well as withdrawal after discontinuation of remifentanyl [20-22]. Additionally, remifentanyl does not possess amnestic or anxiolytic properties.

The study reported in the present issue by Breen and coworkers [1] was an unblinded study conducted in 10 countries and in 15 medical centers. Although the titration of remifentanyl was based on a fixed protocol, titration of the comparator benzodiazepine infusion was left to the discretion of the individual centers, based on local clinical practice. Furthermore, even though there was no statistical difference in the time from start of study drug to the beginning of the weaning process, this process was begun in the remifentanyl group an average of 15 hours earlier. In addition, it is not clear why the patients in the comparator benzodiazepine arm took longer to be extubated once weaning was instituted. It would be interesting to know whether the benzodiazepine group took much longer to return to an acceptable level of arousal for extubation or whether there were other confounding factors, such as delirium, that delayed extubation in this group. The authors do mention that the sedation levels were matched in both groups, during treatment and in the post-treatment period, and that the differences were due to the drug *per se* and not the level of sedation. However, a physiological basis for this difference is not offered. This is important in the context of an unblinded trial, in which there is always the potential for bias.

What this field now needs is a new guard of studies focusing on acute and long-term cognitive outcomes of sedative regimens in addition to outcomes such as days on mechanical ventilation and ICU length of stay. Modifying sedation paradigms by utilizing agents such as remifentanyl or dexmedetomidine may provide clinicians with alternatives to the 'gold standard' benzodiazepines. However, for that to happen we need rigorously conducted, blinded, randomized controlled trials.

Competing interests

The author(s) declare that they have no competing interests.

Acknowledgements

Dr Pandharipande is the recipient of the Vanderbilt Physician Scientist Development Award. Dr Ely is the Associate Director of Research for the VA Tennessee Valley Geriatric Research and Education Clinical Center (GRECC). He is a recipient of the Paul Beeson Faculty Scholar Award from the Alliance for Aging Research and is a recipient of a K23 from the National Institute of Health (#AG01023-01A1).

References

1. Breen D, Karabinis A, Malbrain M, Morais R, Albrecht S, Jarnvig I-L, Parkinson P, Kirkham AJT: **Decreased time on mechanical ventilation when comparing analgesia-based sedation using remifentanyl versus standard hypnotic-based sedation for up to 10**

- days in ICU patients: a randomised trial. [ISRCTN47583497]. *Crit Care* 2005, **9**:R200-R210.
2. Jacobi J, Fraser GL, Coursin DB, Riker R, Fontaine D, Wittbrodt ET, Chalfin DB, Masica MF, Bjerke S, Coplin WM, et al.: **Clinical practice guidelines for the sustained use of sedatives and analgesics in the critically ill adult.** *Crit Care Med* 2002, **30**:119-141.
3. Dubois MJ, Bergeron N, Dumont M, Dial S, Skrobik Y: **Delirium in an intensive care unit: a study of risk factors.** *Intensive Care Med* 2001, **27**:1297-304.
4. Pandharipande P, Shintani A, Peterson J, Ely EW: **Sedative and analgesic medications are independent risk factors in icu patients for transitioning into delirium [abstract].** *Crit Care Med* 2004, **32**:A19.
5. Marcantonio ER, Juarez G, Goldman L, Mangione CM, Ludwig LE, Lind L, Katz N, Cook F, Orav J, Lee TH: **The relationship of post-operative delirium with psychoactive medications.** *JAMA* 1994, **272**:1518-1522.
6. Kollef MH, Levy NT, Ahrens T, Schaiff R, Prentice D, Sherman G: **The use of continuous IV sedation is associated with prolongation of mechanical ventilation.** *Chest* 1999, **114**:541-548.
7. Inouye SK, Schlesinger MJ, Lyndon TJ: **Delirium: a symptom of how hospital care is failing older persons and a window to improve quality of hospital care.** *Am J Med* 1999, **106**:565-573.
8. Kress JP, Pohlman AS, O'Connor MF, Hall JB: **Daily interruption of sedative infusions in critically ill patients undergoing mechanical ventilation.** *N Engl J Med* 2000, **342**:1471-1477.
9. Brook AD, Ahrens TS, Schaiff R, Prentice D, Sherman G, Shannon W, Kollef MH: **Effect of a nursing implemented sedation protocol on the duration of mechanical ventilation.** *Crit Care Med* 1999, **27**:2609-2615.
10. Brattebo G, Hofoss D, Flaatten H, Muri AK, Gjerde S, Plsek PE: **Effect of a scoring system and protocol for sedation on duration of patients' need for ventilator support in a surgical intensive care unit.** *BMJ* 2002, **324**:1386-1389.
11. Sessler CN, Gosnell M, Grap MJ, Brophy GT, O'Neal PV, Keane KA, Tesoro EP, Elswick RK: **The Richmond Agitation-Sedation Scale: validity and reliability in adult intensive care patients.** *Am J Respir Crit Care Med* 2002, **166**:1338-1344.
12. Riker R, Picard JT, Fraser G: **Prospective evaluation of the sedation-agitation scale for adult critically ill patients.** *Crit Care Med* 1999, **27**:1325-1329.
13. Ely EW, Inouye SK, Bernard GR, Gordon S, Francis J, May L, Truman B, Speroff T, Gautam S, Margolin R, et al.: **Delirium in mechanically ventilated patients: validity and reliability of the confusion assessment method for the intensive care unit (CAM-ICU).** *JAMA* 2001, **286**:2703-2710.
14. Ely EW, Truman B, Shintani A, Thomson JWW, Wheeler AP, Gordon S, Francis J, Speroff T, Gautam S, Margolin R, et al.: **Monitoring sedation status over time in ICU patients: reliability and validity of the Richmond Agitation-Sedation Scale (RASS).** *JAMA* 2003, **289**:2983-2991.
15. Ely EW, Shintani A, Truman B, Speroff T, Gordon S, Harrell FE, Inouye S, Bernard G, Dittus R: **Delirium as a predictor of mortality in mechanically ventilated patients in the intensive care unit.** *JAMA* 2004, **291**:1753-1762.
16. Samson HH, Harris RA: **Neurobiology of alcohol abuse.** *Trends Pharmacol Sci* 1992, **13**:206-211.
17. Martz A, Deitrich RA, Harris RA: **Behavioral evidence for the involvement of gamma-aminobutyric acid in the actions of ethanol.** *Eur J Pharmacol* 1983, **89**:53-62.
18. Cavaliere F, Antonelli M, Arcangeli A, Conti G, Costa R, Pennisi MA, Proietti R: **A low-dose remifentanyl infusion is well tolerated for sedation in mechanically ventilated, critically-ill patients.** *Can J Anaesth* 2002, **49**:1088-1094.
19. De Bellis P, Gerbi G, Bacigalupo P, Buscaglia G, Massobrio B, Montagnani L, Servidei L: **Experience with remifentanyl in the ICU [in Italian].** *Minerva Anestesiol* 2002, **68**:765-773.
20. Karabinis A, Mandragos K, Stergiopoulos S, Komnos A, Soukup J, Speelberg B, Kirkham AJ: **Safety and efficacy of analgesia-based sedation with remifentanyl versus standard hypnotic-based regimens in intensive care unit patients with brain injuries: a randomised, controlled trial [ISRCTN50308308].** *Crit Care* 2004, **8**:R268-R280.
21. Vinik HR, Kissin I: **Rapid development of tolerance to analgesia during remifentanyl infusion in humans.** *Anesth Analg* 1998, **86**:1307-1311.
22. Apitzsch H, Olthoff D, Thieme V, Wiegel M, Bohne V, Vetter B: **Remifentanyl and alfentanil: sympathetic-adrenergic effect in the first postoperative phase in patients at cardiovascular risk [in German].** *Anaesthesist* 1999, **48**:301-309.