

What is already known on this topic

Socioeconomic position in childhood is associated with future insulin resistance and coronary heart disease

What this study adds

Among Danish children, those with the most educated and highest earning parents were the least insulin resistant

The opposite was true for children from Estonia and Portugal; those from the most educated and highest earning parents were the most insulin resistant

These results are a reminder that socioeconomic inequalities are dynamic and vary between countries, over time, and between generations within the same country

Contributors: See bmj.com

Funding: This study was supported by the following grants. Denmark: Danish Heart Foundation, Danish Medical Research Council, Health Foundation, Danish Council for Sports Research, Foundation of 17-12-1981, Foundation in Memory of Asta Florida Bolding nee Andersen, and Faculty of Health Sciences, University of Southern Denmark. Estonia: Estonian Science Foundation grant numbers 3277 and 5209. Portugal: FEDER funding from the Government of Madeira. DAL is funded by a UK Department of Health career scientist award. The views expressed in this paper are those of the authors and not necessarily any funding body.

Competing interests: None declared.

Ethical approval: All study protocols conformed to the international guidelines on biomedical research and each research team complied with the ethical procedures of that country.

- 1 Wannamethee SG, Whincup PH, Shaper G, Walker M. Influence of fathers' social class on cardiovascular disease in middle-aged men. *Lancet* 1996;348:1259-63.
- 2 Lawlor DA, Ebrahim S, Davey Smith G. Socioeconomic position in childhood and adulthood and insulin resistance: cross sectional survey using data from the British women's heart and health study. *BMJ* 2002;325:805-7.
- 3 Davey Smith G, Lynch J. Life course approaches to socioeconomic differentials in health. In: Kuh D, Ben-Shlomo Y, eds. *A life course approach to chronic disease epidemiology*. Second ed. Oxford: Oxford University Press, 2004:77-115.

- 4 Kunst AE, Groenohof F, Andersen O, Borgan JK, Costa G, Desplanques G, et al. Occupational class and ischemic heart disease mortality in the United States and 11 European countries. *Am J Public Health* 1999;89:47-53.
- 5 Leon DA. Commentary: Unequal inequalities across Europe. *BMJ* 1998;316:1642.
- 6 Leclerc A, Lert F, Fabien C. Differential mortality: some comparisons between England and Wales, Finland and France, based on inequality measures. *Int J Epidemiol* 1990;19:1001-10.
- 7 Kunst AE, Geurts JJM, van den Berg J. International variation in socioeconomic inequalities in self reported health. *J Epidemiol Community Health* 1995;49:117-23.
- 8 Kunst AE, Mackenbach JP. The size of mortality differences associated with educational level in nine industrialized countries. *Am J Public Health* 1994;84:932-7.
- 9 Mackenbach JP, Kunst AE, Cavelaars AEJM, Groenohof F, Geurts JJM, EU Working Group on Socioeconomic Inequalities in Health. Socio-economic inequalities in morbidity and mortality in western Europe. *Lancet* 1997;349:1655-9.
- 10 Van DE, Koolman X. Explaining the differences in income-related health inequalities across European countries. *Health Econ* 2004;13:609-28.
- 11 Riddoch C, Edwards D, Page A, Froberg K, Anderssen SA, Wedderkopp N, et al. The European youth heart study—cardiovascular disease risk factors in children: rationale, aims, study design and validation of methods. *Journal of Physical Activity and Health* 2005;2:115-29.
- 12 Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985;28:412-9.
- 13 Gunnell D. Commentary: Early insights into height, leg length, proportionate growth and health. *Int J Epidemiol* 2001;30:221-2.
- 14 Galobardes B, Lynch JW, Davey Smith G. Childhood socioeconomic circumstances and cause-specific mortality in adulthood: systematic review and interpretation. *Epidemiol Rev* 2004;26:7-21.
- 15 Leinsalu M, Vagero D, Kunst AE. Estonia 1989-2000: enormous increase in mortality differences by education. *Int J Epidemiol* 2003;32:1081-7.
- 16 Kunst A. *Cross-national comparisons of socio-economic differences in mortality*. Rotterdam: Erasmus University, 1997.
- 17 Shkolnikov VM, Leon DA, Adamets S, Andreev E, Deev A. Educational level and adult mortality in Russia: an analysis of routine data 1979 to 1994. *Soc Sci Med* 1998;47:357-69.
- 18 Vodopivec M. Worker reallocation during Estonia's transition to market. *International Journal of Manpower* 2002;23:77-97.
- 19 Brandou F, Brun J-F, Mercier J. Limited accuracy of surrogates of insulin resistance during puberty in obese and lean children at risk for altered glucoregulation. *J Clin Endocrinol Metab* 2005;90:761-7.
- 20 Kunst AE, Groenohof F, Mackenbach JP, EU Working Group on Socio-economic Inequalities in Health. Occupational class and cause specific mortality in middle aged men in 11 European countries: comparison of population based studies. *BMJ* 1998;316:1636-42.
- 21 Harro M, Villa I, Liiv K, Aru J, Alep J. Nutrition-related health indicators and their major determinants in the new member states: case of Estonia. *J Public Health* 2005; in press.
- 22 Davey Smith G. Socioeconomic differentials. In: Kuh D, ed. *A life course approach to chronic disease epidemiology*. Oxford: Oxford Medical Publications, 1997:242-73.

(Accepted 31 March 2005)

Commentary: Health inequalities and social dynamics in Europe

Denny Vågerö, Mall Leinsalu

Centre for Health Equity Studies (CHESS), Stockholm University/Karolinska Institutet, SE-106 91 Stockholm, Sweden

Denny Vågerö
professor

Stockholm Centre on Health of Societies in Transition, Södertörn University College, SE-141 89 Huddinge, Sweden
Mall Leinsalu
lecturer

Correspondence to:
D Vågerö
denny.vagero@chess.su.se

Lawlor and colleagues make the valid point that health inequalities are dynamic and change over time and between countries.¹ Unexpectedly, in Estonia and Portugal they found that a high level of insulin resistance is more common among children of more highly educated parents. They ask whether this is because of the new wealth of these families, perhaps a preference for Western style “junk” food?

A well known, but often ignored, fact is that the social distribution of risk factors, disease, and mortality varies by disease entity, time period, and country. “Anomalies,” or deviations from the standard pattern of poorer people having poorer health, include breast cancer and malignant melanoma. Certain risk factors for heart disease, such as smoking or obesity, may previously have been more common among people who are wealthy; in some countries this is still so. Studies from several countries suggest that as coronary heart

disease became more common, its inverse class pattern among men became more pronounced, perhaps even preceded by a reversal some decades ago.² Similarly, the generally higher mortality rate ratios for circulatory diseases (comparing manual and non-manual workers) in northwestern Europe than in southern Europe may be a result of a “phase difference,” implying that those ratios will soon increase in the south.³ Is what the authors report from Portugal, Estonia, and Denmark consistent with such a phase transition? We feel that this is more likely to be the case for Portugal (Madeira) than for Estonia (Tartu).

Health inequalities are indeed constantly changing, driven by the social dynamics of a country. Many paradoxes are hidden in this truth. For Russia, the standard pattern of mortality by education applies to both men and women, in spite of contrasting risk factor distributions, with many “anomalies”—for instance, in obesity

and lipid profiles among men.⁴ For Estonia, highly educated adults, compared with those with less education, gained a considerable advantage during the 1990s, in terms of mortality from circulatory diseases and total mortality.⁵ According to Lawlor and colleagues, the children of these highly educated parents may paradoxically have the poorest insulin resistance profile. We cannot say how much of a paradox this really is, however, as we know nothing about whether their parents as children also had a poor insulin profile.

Genetic, fetal, and early childhood factors should all be relevant in determining insulin resistance. In Lawlor and colleagues' study, parental education was important for insulin resistance among prepubertal and postpubertal children. Parental education can be taken as a measure of social circumstances when their children were born—that is, before the collapse of the Soviet system. Income was measured in 1997-2000; for Estonia this means when new food markets had opened up. In mutual adjustments, education but not income had an independent effect; thus it seems unlikely that it is consumption of "burgers, crisps, and processed food" that is creating the pattern of high insulin resistance among children of highly educated parents. We also noted that, in this study, children of highly educated fathers in Estonia had a 200 g lower birth weight than others, consistent with their higher insulin resistance.

Anomalies such as those reported for Estonia and Portugal may be of special significance, as they point towards gaps in our understanding and warn against too simplistic a view of health inequalities. Correctly understanding the development of health and mortality in the formerly communist led countries of central and eastern Europe is likely to challenge (and has already challenged) many cherished epidemiological "truths."

We thank Ilona Koupil for important comments to a previous version of this commentary.

Contributors: DV wrote the first version. ML commented and suggested changes. Both authors have read and accepted the final version.

Funding: DV is financed through the Swedish Council for Working Life and Social Research. ML is financed through the Swedish Foundation for Baltic and East European Studies.

Competing interests: None declared.

- 1 Lawlor DA, Harro M, Wedderkopp N, Andersen LB, Sardinha LB, Riddoch CJ, et al Association of socioeconomic position with insulin resistance among children from Denmark, Estonia, and Portugal: cross sectional study. *BMJ* 2005;331:183-6.
- 2 Marmot MG, Adelstein AM, Robinson N, Rose GA. Changing social class distribution of heart disease. *BMJ*:1978;iii:109-12.
- 3 Mackenbach JP, Bakker MJ, Kunst AE, Diderichsen F. Socioeconomic inequalities in health in Europe: an overview. In: Mackenbach JP, Bakker MJ, eds. *Reducing inequalities in health: a European perspective*. London and New York: Routledge, 2002.
- 4 Vågerö D, Koupil I, Plavinskaya S, Shestov D. Educational differences in mortality 1975-2000 are largely explained by known risk factors among men, but not among women—the St Petersburg cohort study. CHES, Stockholm University/Karolinska Institute, 2005. [Submitted for publication.]
- 5 Leinsalu M, Vågerö D, Kunst A. Estonia 1989-2000: enormous increase in mortality differences by education. *Int J Epidemiol* 2003;32:1081-7.

Changes in dietary fat and declining coronary heart disease in Poland: population based study

Witold A Zatonski, Walter Willett

We previously described a rapid decline in mortality due to coronary heart disease in Poland between 1991 and 1994, corresponding with increases in the ratio of polyunsaturated fat to saturated fat in people's diet and fruit consumption.¹ The changes in food consumption followed changes in economic policy, including reductions in subsidies for dairy and other animal fats. We describe subsequent trends and use data from cohort studies to estimate the contributions from smoking and diet to these changes.

Methods and results

Mortality due to coronary heart disease has continued to fall in Poland in both sexes and across educational levels. Compared with 1990, by 2002 for the age band 45-64 years it had fallen by 38% in men (340 per 100 000 to 212/100 000) and by 42% in women (76/100 000 to 44/100 000). By 1999 (the latest year with comparable data), consumption of saturated fat had fallen by 7% (44.8 g/day to 41.5 g/day), consumption of polyunsaturated fat had risen by 57% (14.8 g/day to 23.3 g/day), and the ratio of the two had increased by 70%. Per head, consumption of imported

fruit rose from 2.8 kg/year in 1990 to 8.8 kg/year in 1991 and 10.4 kg/year in 1999.

From 1990 to 2004, the prevalence of smoking in Poland fell among people younger than 40 but increased from 23.1% to 35.2% among women aged 40-60. Among men, it fell from 51.1% to 46.1% at age 40-49, from 51.1% to 46.1% at 50-59, and from 34.3% to 28.2% over 60. If we assume a relative risk of 3 for current smoking then a reduction in prevalence of 5% (from 55% to 50%) would reduce the risk of coronary heart disease by about 5%. This, and similar declines in rates of coronary heart disease among men and women, indicates that changes in smoking contributed little to falling rates of coronary heart disease.

In the prospective nurses' health study, the ratio of dietary saturated and unsaturated fats was inversely related to the incidence of coronary heart disease.² The change in coronary mortality in Poland was similar to that predicted by the slope relating this ratio to the risk of coronary heart disease in the study (figure). The increased consumption of imported fruit between

Cancer Centre and Institute of Oncology, 5 Roentgena Str, 02-781 Warsaw, Poland

Witold A Zatonski
director of the Cancer Epidemiology and Prevention Division

Harvard School of Public Health, 665 Huntington Avenue, Boston, MA 02115, USA
Walter Willett
professor

Correspondence to: W Willett
walter.willett@channing.harvard.edu

BMJ 2005;331:187-9



Two additional references (w1 and w2) are on bmj.com