

## *p*-Cresol and 3,5-Xylenol Methylhydroxylases in *Pseudomonas putida* N.C.I.B. 9869

By MALCOLM J. KEAT and DAVID J. HOPPER

Department of Biochemistry, University College of Wales, Aberystwyth, Dyfed SY23 3DD, Wales, U.K.

(Received 7 April 1978)

*Pseudomonas putida* N.C.I.B. 9869, when grown on 3,5-xylenol, hydroxylates the methyl groups on 3,5-xylenol and on *p*-cresol by two different enzymes. 3,5-Xylenol methylhydroxylase, studied only in relatively crude extracts, requires NADH, is not active with *p*-cresol and is inhibited by cyanide, but not by CO. The *p*-cresol methylhydroxylase requires an electron acceptor and will act under anaerobic conditions. It was purified and is a flavocytochrome *c* of mol.wt. approx. 114000 consisting of two subunits of equal size. The enzyme catalyses the hydroxylation of *p*-cresol ( $K_m$  16  $\mu$ M) and the further oxidation of product, *p*-hydroxybenzyl alcohol ( $K_m$  27  $\mu$ M) to *p*-hydroxybenzaldehyde. A different *p*-cresol methylhydroxylase of the flavocytochrome *c* type is induced by growth on *p*-cresol. It too was purified and has mol.wt. approx. 100000, and again consisted of two equal-size subunits. The  $K_m$  for *p*-cresol is 3.6  $\mu$ M and for *p*-hydroxybenzyl alcohol, 15  $\mu$ M.

Although the bacterial oxidation of methyl-substituted aromatic compounds may proceed with retention of the methyl group intact until after ring fission, often the initial attack involves oxidation of the methyl group to carboxyl (Chapman, 1972). The first step of hydroxylation to yield an alcohol is followed by oxidation to an aldehyde and then an acid, in a manner analogous to the conversion of alkanes into carboxylic acids or the  $\omega$ -oxidation of fatty acids (Gholson *et al.*, 1963). Detailed studies have been made on the enzymes used by some bacteria for alkane hydroxylation (McKenna & Coon, 1970; Jurtshuk & Cardini, 1971; Ruettinger *et al.*, 1977). These use molecular oxygen with a requirement for reduced nicotinamide nucleotide and are multi-component systems. Less is known about the enzymes that oxidize methyl groups attached to the benzene ring, although the *p*-xylene hydroxylase from *Pseudomonas aeruginosa* appears to be a mixed-function oxygenase and has been resolved into two components by DEAE-cellulose chromatography (Nozaka & Kusunose, 1968). An enzyme that hydroxylates the methyl group of *p*-cresol has been purified from a strain of *Pseudomonas putida* (Hopper, 1976; Hopper & Taylor, 1977). It is a flavocytochrome and will act under anaerobic conditions with a requirement for an electron acceptor.

The degradation of both *p*-cresol and 3,5-xylenol by *Ps. putida* N.C.I.B. 9869 is initiated by oxidation of a methyl group (Hopper & Chapman, 1971; Hopper & Taylor, 1973), and the results of a study of the hydroxylases involved are presented in the present paper.

### Materials and Methods

#### *Maintenance and growth of organism*

The organism, *Ps. putida* N.C.I.B. 9869, was maintained by growth on nutrient-agar slants for 24 h at 30°C, followed by storage at 4°C. It was subcultured monthly.

Cells were grown at 30°C in medium containing (per litre): Na<sub>2</sub>HPO<sub>4</sub>, 4.33 g; KH<sub>2</sub>PO<sub>4</sub>, 2.65 g; NH<sub>4</sub>Cl, 2.0 g; nitrilotriacetic acid, 0.1 g; salts solution (Rosenberger & Elsdon, 1960), 4.0 ml; 3,5-xylenol or *p*-cresol, 0.3 g. For whole-cell studies cells were grown in 1 litre of medium in 2-litre Erlenmeyer flasks at 30°C in an orbital shaker (Gallenkamp and Co., Widnes, Lancs., U.K.). Larger quantities of cells for preparation of extracts and for enzyme purification were grown in a Microferm MF-114 fermenter (New Brunswick Scientific Co., New Brunswick, NJ, U.S.A.) which was aerated at a rate of 4 litres/min and stirred at 150 rev./min. Growth was followed by measuring turbidity at 540 nm in a Unicam SP.500 spectrophotometer. Further additions of substrate were made when necessary to give higher yields. Cells were harvested in an Alfa Laval LAB 102B continuous-flow centrifuge (Alfa Laval Co., Brentford, Middx., U.K.) and stored at -20°C.

#### *Enzyme assays*

The procedures for assaying *p*-cresol methylhydroxylases A and B were those described by Hopper & Taylor (1977) modified by using 50 mm-glycine/NaOH buffer, pH 9.6, the pH optimum for both enzymes.

### Preparation of extracts

Extracts were prepared by suspending each 1 g of wet cells in 2 ml of ice-cold 50 mM-KH<sub>2</sub>PO<sub>4</sub>/NaOH buffer, pH 7.0, cooling the suspension in an ice/water bath and disrupting the bacteria with a Dawe Soniprobe, type 7530 A, fitted with a micro-tip for volumes up to 20 ml or in a 500 W MSE ultrasonic disintegrator fitted with a 3 cm probe for larger volumes. The smaller volumes were disrupted for a total of 4 min in ½ min periods and the larger volumes for 8 min also in ½ min periods. The extracts were centrifuged at 2°C for 10 min at 40000g and the supernatants, referred to as crude extracts, retained. Such extracts were red-brown in colour and contained 35–45 mg of protein/ml. Ice-cold saturated (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> solution was added slowly, with stirring, to crude extract to adjust it to 70% (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> saturation. After 1 h at 0°C the precipitate was removed by centrifuging at 25000g for 20 min. The pellet was dissolved in a volume of 50 mM-phosphate buffer, pH 7.0, equal to that of the original crude extract and dialysed with stirring for 3 h against two changes of 100 vol. of the same buffer. This extract is referred to as (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>-treated extract.

### Polyacrylamide-gel electrophoresis

Electrophoresis of enzymes was carried out on 6.4% (w/v) polyacrylamide gels with an average pore radius of 1.5 nm (Gordon, 1969) by the method of Ornstein & Davis (1964) at pH 8.9. Sodium dodecyl sulphate/polyacrylamide-gel electrophoresis was performed as described by Weber & Osborn (1969). Disaggregation of the enzymes required incubation for 2 h at 37°C in 10 mM-phosphate buffer, pH 7.0, containing 1% (v/v) sodium dodecyl sulphate and 1% (w/v) mercaptoethanol. The incubation mixture was then dialysed overnight against 10 mM-phosphate buffer, pH 7.0, containing 0.1% (w/v) sodium dodecyl sulphate and 0.1% (w/v) mercaptoethanol. Gels were stained for protein with either Amido Black 10B (Colour Index 20470) or Coomassie Brilliant Blue G-250 (Colour Index 42655) and cleared by washing with 7% (v/v) acetic acid in water or acetic acid/ethanol/water (2:9:9, by vol.) respectively. Gels were stained for enzyme activity by placing them in reaction mixtures containing, in 1.5 ml of 50 mM-glycine/NaOH buffer, pH 9.6, 2 μmol of either *p*-cresol or *p*-hydroxybenzyl alcohol, 2 μmol of phenazine methosulphate and 1 mg of either Nitro Blue Tetrazolium or 2-(4-iodophenyl)-3-(4-nitrophenyl)-5-phenyltetrazolium chloride. These were incubated at 20°C in the dark. The Nitro Blue Tetrazolium gave a blue colour with the enzyme and the latter compound gave a red colour. Gels, stained for protein, were scanned at 600 nm in a Gilford 2400S spectrophotometer (Gilford Instruments, Ober-

lin, OH, U.S.A.) fitted with a 2410S gel-transport attachment.

### Ultracentrifuge studies

These were carried out with a Beckman Spinco model E ultracentrifuge equipped with Rayleigh interference optics. Samples were previously dialysed for 24 h against 16.7 mM-phosphate buffer, pH 7.0, at 2°C. Sedimentation-equilibrium experiments were conducted by using the short-column meniscus-depletion method of Yphantis (1964) and were performed at 20410 rev./min for 24 h at 20°C for hydroxylase A and at 17890 rev./min for 24 h at 9°C for hydroxylase B. The initial protein concentration was 0.5 mg/ml. In these runs the solution compartment in the double-sector cells, equipped with sapphire windows, contained 100 μl of enzyme solution on 50 μl of fluorocarbon oil (Beckman, Palo Alto, CA, U.S.A.); the reference compartment contained 120 μl of diffusate on 40 μl of fluorocarbon oil. The attainment of equilibrium was determined experimentally by measuring fringe displacements of given radial positions. A value of 0.725 cm<sup>3</sup>/g was assumed for the partial specific volume,  $\bar{v}$ , of the proteins.

### Aerobic incubations

Oxidation of *p*-cresol by crude extracts was followed at 30°C in Warburg respirometers containing, in a 2.8 ml reaction volume, 33 mg of protein, 125 μmol of KH<sub>2</sub>PO<sub>4</sub>/NaOH buffer, pH 7.0, 2 μmol of phenazine methosulphate and 6 μmol of substrate, with 0.2 ml of 20% (w/v) KOH in the centre well. For oxidation of *p*-cresol or *p*-hydroxybenzyl alcohol by partially purified hydroxylase A flasks contained, in 2.8 ml of 50 mM-glycine/NaOH buffer, pH 9.6, 2 μmol of phenazine methosulphate, 3.2 mg of protein and 6 μmol of substrate, with 0.2 ml of 20% (w/v) KOH in the centre well.

### Anaerobic incubations

Reaction mixtures in Thunberg tubes contained 5 ml of purified enzyme (1.2 mg and 0.8 mg of protein/ml for hydroxylase A and B respectively), 5 ml of 50 mM-glycine/NaOH buffer, pH 9.6, 50 μmol of phenazine methosulphate, 60 μmol of 2,6-dichlorophenol-indophenol and 30 μmol of either *p*-cresol or *p*-hydroxybenzyl alcohol in the side arms. The tubes were alternately evacuated and flushed with N<sub>2</sub> five times and, after tipping of substrate, were incubated at 30°C for 3 h. The reaction was stopped by addition of 5 ml of 20% (w/v) orthophosphoric acid.

### Isolation of products

Protein from acidified reaction mixtures was removed by centrifuging at 10000g for 10 min and

the supernatants were extracted twice with an equal volume of diethyl ether. The pooled extracts were dried over anhydrous  $\text{Na}_2\text{SO}_4$  and then evaporated to dryness.

#### Chromatography

T.l.c. was carried out on Kieselgel GF<sub>254</sub> (E. Merck A.-G., Darmstadt, Germany) coated on glass plates to a thickness of 0.25 mm and activated at 110°C for 30 min. Phenolic compounds were separated by using solvent A [benzene/dioxan/acetic acid (90:25:4, by vol.) (Pastuska, 1961)]. For 2,4-dinitrophenylhydrazones solvent B [benzene/tetrahydrofuran (19:1, v/v) (Byrne, 1965)] was used. Descending paper chromatography of phenolic compounds was carried out on Whatman no. 1 paper with solvent C [benzene/acetic acid/water (125:72:3, by vol.) (Smith, 1960)] and solvent D [5% (w/v) sodium formate/formic acid (200:1, v/v) (Smith, 1960)].

#### Preparation of 2,4-dinitrophenylhydrazones

Reaction mixtures or 5 ml of solutions of standards were incubated at 30°C for 30 min with 0.1% (w/v) 2,4-dinitrophenylhydrazine in 2M-HCl. The precipitated 2,4-dinitrophenylhydrazones were collected by centrifuging at 10000g for 10 min. The pellet was washed with 2M-HCl and dissolved in 3 ml of ethyl acetate. The solution was dried over anhydrous  $\text{Na}_2\text{SO}_4$ .

#### Absorption spectra

Visible-absorption spectra were measured in a Unicam SP.1800 double-beam recording spectrophotometer. For difference spectra of the *p*-cresol methylhydroxylases 2 ml of enzyme solution containing 0.5–1.0 mg of protein/ml were first oxidized with 2  $\mu\text{l}$  of 50% (w/w)  $\text{H}_2\text{O}_2$ , and the baseline spectrum recorded with oxidized enzyme in both sample and reference cuvettes (1 cm light-path). The enzyme in the sample cuvette was then reduced by addition of a few crystals of sodium dithionite or 0.1 ml of 20 mM substrate, and the difference spectrum recorded.

#### Spectrophotofluorimetric studies

Fluorimetric measurements were made in 1 cm light-path cuvettes in an Aminco-Bowman spectrophotofluorimeter (American Instrument Co., Silver Spring, MD, U.S.A.) fitted with a xenon light-source and with slit arrangement no. 3 (Aminco instruction manual).

#### NADH- and NADPH-generating systems

NADH was generated in Warburg flasks by including 0.1 ml of 10 mM-NAD<sup>+</sup>, 0.1 ml of 0.2M-

ethanol and 0.1 ml of alcohol dehydrogenase (1 mg/ml, 300 units/mg) in the reaction mixture. NADPH was generated by inclusion of 0.1 ml of 10 mM-NADP<sup>+</sup>, 0.1 ml of 0.2M-glucose 6-phosphate and 0.1 ml of glucose 6-phosphate dehydrogenase (30 units/ml). One unit is defined as the conversion of 1  $\mu\text{mol}$  of substrate/min under the manufacturers' recommended conditions.

#### Protein determinations

The biuret method of Gornall *et al.* (1949) was used to determine protein, except where very low concentrations were present, when the method of Lowry *et al.* (1951) was used. Standard curves were prepared by using bovine serum albumin. The elution of protein from columns was followed by measuring  $A_{280}$  of the fractions.

#### Purification of *p*-cresol methylhydroxylase A

Crude extract from 3,5-xylene-grown cells was heated rapidly, with continuous stirring, to 70°C, kept at this temperature for 5 min and then quickly cooled in ice/water. The precipitated protein was removed by centrifuging at 18000g for 10 min. The supernatant was loaded on a DEAE-cellulose column (18.5 cm  $\times$  2.5 cm), which was washed with 200 ml of 16.7 mM- $\text{KH}_2\text{PO}_4$ /NaOH buffer, pH 7.0. The enzyme was then eluted with a linear gradient from 0 to 0.3 M-KCl in 1 litre of the buffer and 10 ml fractions were collected. Fractions 52–58 inclusive were pooled and dialysed for 4 h against 2 litres of 16.7 mM- $\text{KH}_2\text{PO}_4$ /NaOH buffer, pH 7.0, with three changes of buffer. The enzyme was concentrated by loading on a small DEAE-cellulose column (3 cm  $\times$  2 cm) and eluting with the phosphate buffer containing 0.5 M-KCl, and then loaded on a Sephadex G-200 column (45 cm  $\times$  2.5 cm). It was eluted with 50 mM- $\text{KH}_2\text{PO}_4$ /NaOH buffer, pH 7.0. Fractions of 5 ml were collected and fractions 33–38 were pooled. A summary of the purification is shown in Table 1.

#### Purification of *p*-cresol methylhydroxylase B

Crude extract from *p*-cresol-grown cells was heat-treated as described for *p*-cresol methylhydroxylase A. The extract was then loaded on a DEAE-cellulose column (12 cm  $\times$  2.5 cm) and the enzyme washed straight through with 16.7 mM- $\text{KH}_2\text{PO}_4$ /NaOH buffer, pH 7.0. The enzyme was concentrated by the gradual addition at 2°C of 0.48 g of  $(\text{NH}_4)_2\text{SO}_4$  per ml of extract followed by centrifuging at 25000g for 20 min. The pellet of precipitated protein was redissolved in a small volume of 50 mM- $\text{KH}_2\text{PO}_4$ /NaOH buffer, pH 7.0, and applied to a Sephadex G-200 column (45 cm  $\times$  2.5 cm). The enzyme was eluted with the same buffer and 5 ml fractions were collected.

Table 1. Summary of the purification of *p*-cresol methylhydroxylase from *Ps. putida* N.C.I.B. 9869 grown on 3,5-xylenol (hydroxylase A)

The starting material was 90 g wet wt. of cells. One unit of enzyme activity is the amount that catalyses the uptake of  $\frac{1}{2}$   $\mu$ mol of O<sub>2</sub>/min at 30°C in the O<sub>2</sub>-monitor assay.

Purification step	Volume (ml)	Total protein (mg)	Total activity (units)	Specific activity (units/mg)	Recovery (%)	Purification
(1) Crude extract	240	7900	1529	0.19	100	1
(2) Heat-treatment	210	1125	446	0.40	29.2	2.1
(3) DEAE-cellulose chromatography	70	77	407	5.28	26.6	27.7
(4) Sephadex G-200 gel filtration	30	19	189	9.95	12.4	52.3

Table 2. Summary of the purification of *p*-cresol methylhydroxylase from *Ps. putida* N.C.I.B. 9869 grown on *p*-cresol (hydroxylase B)

The starting material was 45 g wet wt. of cells. One unit of enzyme activity is the amount that catalyses the uptake of  $\frac{1}{2}$   $\mu$ mol of O<sub>2</sub>/min at 30°C in the O<sub>2</sub>-monitor assay.

Purification step	Volume (ml)	Total protein (mg)	Total activity (units)	Specific activity (units/mg)	Recovery (%)	Purification
(1) Crude extract	130	5410	1929	0.36	100	1
(2) Heat-treatment	110	908	470	0.52	24.4	1.4
(3) DEAE-cellulose chromatography	120	540	432	0.80	22.4	2.2
(4) Sephadex G-200 gel filtration	55	106	190	1.79	9.9	5.0
(5) Hydroxyapatite chromatography	40	5.6	38	6.78	2.0	18.8

Fractions 27–37 were pooled and dialysed against 1 litre of 40mM-KH<sub>2</sub>PO<sub>4</sub>/NaOH buffer, pH6.8, at 2°C for 4h with three changes of buffer. The extract was loaded on a hydroxyapatite column (13cm × 2.5cm) and the enzyme eluted with a linear gradient constructed from 350ml of 40mM-KH<sub>2</sub>PO<sub>4</sub>/NaOH buffer, pH6.8, and 350ml of 220mM-KH<sub>2</sub>PO<sub>4</sub>/NaOH buffer, pH6.8. Fractions of 5ml were collected and the protein in pooled fractions 75–80 was concentrated by addition of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> as described before. The protein pellet was redissolved in 2ml of 50mM-KH<sub>2</sub>PO<sub>4</sub>/NaOH buffer, pH7.0, and dialysed for 3h at 2°C against 1 litre of the same buffer, which was changed twice. A summary of the purification is shown in Table 2.

### Chemicals

CO was prepared by the controlled addition of formic acid to conc. H<sub>2</sub>SO<sub>4</sub> (Vogel, 1956).

*o*-Cresol, *m*-cresol, *p*-cresol, 2,3-xyleneol, 2,5-xyleneol, 2,6-xyleneol, 3,4-xyleneol, *o*-hydroxybenzyl alcohol, DL-1-phenylethanol, Nitro Blue Tetrazolium, 2-(4-iodophenyl)-3-(4-nitrophenyl)-5-phenyl-tetrazolium chloride and phenazine methosulphate were from BDH, Poole, Dorset, U.K. 3,5-Xyleneol, *p*-methoxyphenol, *p*-ethylphenol and 2,4-xyleneol (redistilled before use) were from Aldrich Chemical Co., Milwaukee, WI, U.S.A. Amido Black 10B and

2,6-dichlorophenol-indophenol were from Hopkin and Williams, Chadwell Heath, Essex, U.K. Coomassie Brilliant Blue G-250 was from Serva Feinbiochemica G.m.b.H., Heidelberg, Germany. 4-Hydroxybenzyl alcohol was from K & K Laboratories, Plainview, NY, U.S.A. *p*-Methylbenzyl alcohol was from NIPA Laboratories, Pontypridd, Glam., U.K. Bovine serum albumin (Cohn fraction V) was from Koch-Light Laboratories, Colnbrook, Bucks., U.K. Sephadex G-200 was from Pharmacia Fine Chemicals, Uppsala, Sweden. Preswollen micro-granular DEAE-cellulose DE-52 was from W. R. Balston, Maidstone, Kent, U.K. Bio-Gel HTP hydroxyapatite was from Bio-Rad Laboratories, Richmond, CA, U.S.A. Fumarase, glyceraldehyde 3-phosphate dehydrogenase, phosphorylase *a*, glucose 6-phosphate dehydrogenase, alcohol dehydrogenase, NAD<sup>+</sup>, NADP<sup>+</sup> and glucose 6-phosphate were from Boehringer Corp., London W.5, U.K.

### Results

#### *Oxidation of compounds by whole cells*

The oxidation of a number of compounds by washed-cell suspensions of *Ps. putida* N.C.I.B. 9869 grown on either 3,5-xyleneol or *p*-cresol was followed in Warburg respirometers (Fig. 1). Although 3,5-




Fig. 1. Oxidation of compounds by washed cells grown with (a) 3,5-xyleneol and (b) *p*-cresol

Each flask contained 1.5 ml of bacterial suspension (5 mg dry wt.) in 50 mM-KH<sub>2</sub>PO<sub>4</sub>/NaOH buffer, pH 7.0, 0.3 ml of 10 mM-substrate, tipped from the side arm and 0.2 ml of 20% (w/v) KOH in the centre well. Substrates were: ●, 3,5-xyleneol; ▲, *p*-cresol; ■, *p*-hydroxybenzoic acid. The temperature was 30°C. The gas phase was air. Oxygen uptakes in the absence of substrate [74 μl/h for (a) and 54 μl/h for (b)] have been subtracted.

xyleneol-grown cells showed initial rapid oxidation of *p*-cresol as well as of growth substrate, oxidation of 3,5-xyleneol by *p*-cresol-grown cells was very slow. Neither of these compounds was oxidized by succinate-grown cells.

#### *p*-Cresol methylhydroxylase activity in cell extracts

Crude cell-free extracts of the organism grown on 3,5-xyleneol or *p*-cresol were centrifuged at 150000g for 60 min at 4°C and tested in the Warburg apparatus for ability to oxidize *p*-cresol. There was only very slow oxidation of *p*-cresol above endogenous and the rate was not enhanced by inclusion of 6 μmol of NAD<sup>+</sup>, 6 μmol of NADP<sup>+</sup>, an NADH-generating system or an NADPH-generating system. Inclusion of 2 μmol of phenazine methosulphate, however, resulted in very rapid oxidation of substrate. This phenazine methosulphate-stimulated oxidation of *p*-cresol, similar to that described by Hopper (1976), could be followed in the O<sub>2</sub> monitor and formed the basis of the routine enzyme assay. Similar activity was also found in the extract from *p*-cresol-grown cells, but not in that from succinate-grown cells.

#### Purification of *p*-cresol methylhydroxylase from 3,5-xyleneol- and *p*-cresol-grown cells

Schemes for the purification of *p*-cresol methylhydroxylase from 3,5-xyleneol-grown cells, referred to as hydroxylase A, and from *p*-cresol-grown cells, referred to as hydroxylase B, were developed and are described in the Materials and Methods section. Both procedures yielded red-coloured proteins active with both *p*-cresol and *p*-hydroxybenzyl alcohol.

Enzyme activity was stimulated approx. 2-fold in each case by the particles obtained from centrifuging crude extract at 150000g for 60 min at 4°C or, in the case of *p*-cresol methylhydroxylase A, by the early fractions from DEAE-cellulose chromatography of crude extract. Heat treatment of the stimulatory material at 50°C for 1 min resulted in complete loss of its activity, which accounts for the apparent high loss of enzyme units on heat treatment of crude extracts in the purifications.

#### Polyacrylamide-gel electrophoresis of *p*-cresol methylhydroxylases

After electrophoresis on polyacrylamide gels, hydroxylase B (100 μg) gave a single band when stained for protein (Fig. 2) or when stained for enzyme activity with either *p*-cresol or *p*-hydroxybenzyl alcohol as substrate. Hydroxylase A (100 μg), however, gave one major band followed by two other bands of decreasing intensity when gels were stained for protein (Fig. 2). None of these corresponded to hydroxylase B and all three bands appeared, with the same pattern of intensities, when gels were stained for enzyme activity, again with either *p*-cresol or *p*-hydroxybenzyl alcohol as substrates. A red-coloured band could be seen on electrophoresis of hydroxylase A corresponding to the major protein band. When this band was excised from a number of gels and extracted with 16.7 mM-phosphate buffer, pH 7.0, electrophoresis of the preparation on fresh gels again gave three bands. Similar treatment of the portion of gel 0.5 cm behind the red band gave the same pattern. A mixture of hydroxylase A and B preparations gave a pattern equivalent to the sum of the separate proteins. A similar mixed pattern was seen when 10 μl of heat-treated extract (5.4 mg of protein/ml) from cells grown on a mixture of 3,5-xyleneol and *p*-cresol (0.015%, w/v, of each) was subjected to electrophoresis and gels stained for enzyme activity.

#### Substrate specificity of *p*-cresol methylhydroxylases

Hydroxylases A and B showed similar substrate specificities and were active with a number of compounds (Table 3). Neither enzyme was active under these conditions with *m*-cresol, *o*-cresol, 2,3-xyleneol,




Fig. 2. Scans of polyacrylamide gels after electrophoresis of purified *p*-cresol methylhydroxylases. Gels were loaded with 100  $\mu$ g of purified enzyme, hydroxylase A (a) and hydroxylase B (b), and after electrophoresis were stained for protein with Amido Black. They were then scanned at 600 nm.

2,5-xyleneol, 2,6-xyleneol, 3,5-xyleneol, *m*-hydroxybenzyl alcohol, *o*-hydroxybenzyl alcohol, *p*-methoxyphenol, DL-1-phenylethanol, *p*-xylene, *p*-hydroxyacetophenone, *o*-hydroxyacetophenone or *p*-methylbenzyl alcohol.

#### Identification of products from aerobic and anaerobic incubations

*p*-Cresol and *p*-hydroxybenzyl alcohol were incubated aerobically in Warburg flasks with hydroxylase A in the presence of phenazine methosulphate. After incubation for 50 min at 30°C the contents of the flasks were acidified to pH 1.0 with 5M-HCl, and the products were isolated. The products were chromatographed by t.l.c. on silica gel in

Table 3. Relative activities of *p*-cresol methylhydroxylases A and B with various substrates

Substrates were tested by the O<sub>2</sub>-monitor assay with 100  $\mu$ g of hydroxylase A or 80  $\mu$ g of hydroxylase B and 1.0  $\mu$ mol of substrate. The values are given as a percentage of the rate with *p*-cresol.

Substrate	Hydroxylase activity	
	A	B
<i>p</i> -Cresol	100	100
2,4-Xyleneol	71	74
3,4-Xyleneol	70	78
<i>p</i> -Hydroxybenzyl alcohol	69	81
<i>p</i> -Ethylphenol	66	55

solvent A and on paper in solvents C and D. In all systems the product from both *p*-cresol and *p*-hydroxybenzyl alcohol corresponded to authentic *p*-hydroxybenzaldehyde, with no trace of starting material.

Both hydroxylases A and B were active under anaerobic conditions. Addition of *p*-cresol or *p*-hydroxybenzyl alcohol from the side arm of a Thunberg cuvette, made anaerobic by alternate evacuation and filling with N<sub>2</sub> four times, to the reaction mixture for spectrophotometric assay resulted in a rapid decrease in A<sub>600</sub> due to reduction of 2,6-dichlorophenol-indophenol. The presence of both enzyme and phenazine methosulphate were essential for this activity.

The anaerobic reaction mixtures were scaled up to obtain sufficient products for identification. Chromatography by t.l.c. and on paper in the same systems as before showed *p*-hydroxybenzaldehyde as the major product from *p*-cresol and *p*-hydroxybenzyl alcohol with both hydroxylases. A minor product from *p*-cresol corresponded to *p*-hydroxybenzyl alcohol. Further confirmation of *p*-hydroxybenzaldehyde as a product came from t.l.c. of the 2,4-dinitrophenylhydrazone of the product in solvent B, where it corresponded to the derivative from authentic material.

#### Kinetic studies

Spectrophotometric assays were performed with 14  $\mu$ g of pure hydroxylase A or 17  $\mu$ g of pure hydroxylase B per assay and a range of substrate concentrations from 3.3 to 13.3  $\mu$ M. Double-reciprocal plots of the rate obtained and the substrate concentrations gave straight lines from which the *K<sub>m</sub>* values for *p*-cresol and *p*-hydroxybenzyl alcohol were calculated. Hydroxylase A gave a *K<sub>m</sub>* of 16  $\mu$ M for *p*-cresol and 27  $\mu$ M for *p*-hydroxybenzyl alcohol. The corresponding values for hydroxylase B were 3.6 and 15  $\mu$ M.




Fig. 3. Absorption spectrum of hydroxylase A. The spectrum of purified *p*-cresol methylhydroxylase A (1.3 mg/ml) was recorded against a blank of 50 mM-KH<sub>2</sub>PO<sub>4</sub>/NaOH buffer, pH 7.0.

#### Molecular-weight determinations of *p*-cresol methylhydroxylases

A value of 114000 was obtained for the molecular weight of hydroxylase A by the sedimentation-equilibrium (meniscus-depletion) method. The value for hydroxylase B, from a similar experiment, but run at a lower temperature, was 99000. The respective values for hydroxylase A and B from their relative elution volumes ( $V_e/V_0$ ) from a Sephadex G-200 column calibrated by plotting relative elution volumes of known proteins against log(molecular weight) (Andrews, 1965) were 108000 and 100000.

Electrophoresis of the enzymes, after incubation with sodium dodecyl sulphate and mercaptoethanol, on polyacrylamide gels containing sodium dodecyl sulphate gave single bands when stained for protein with Coomassie Brilliant Blue. Comparison of their mobilities relative to Bromophenol Blue with the mobilities of standard proteins gave mol.wts. of approx. 58000 for hydroxylase A and 56000 for hydroxylase B. The standard proteins were glyceraldehyde 3-phosphate dehydrogenase (mol.wt. 37000), fumarase (mol.wt. 48500), bovine serum albumin (mol.wt. 68000) and phosphorylase *a* (subunit mol.wt. 94000).

#### Absorption spectra of *p*-cresol methylhydroxylases

The visible-absorption spectra of the two enzymes are very similar and typical of haemoproteins. The spectrum of hydroxylase A, shown in Fig. 3, has  $\lambda_{max}$  for  $\alpha$ -,  $\beta$ - and  $\delta$ -peaks of 553, 523 and 418 nm

respectively. The equivalent peaks for hydroxylase B are at 551, 523 and 418 nm. These peaks also appeared in difference spectra of reduced against oxidized enzymes and the same spectra were produced by reducing oxidized enzyme with sodium dithionite or by adding the substrates *p*-cresol or *p*-hydroxybenzyl alcohol. No reduction occurred with 3,5-xyleneol.

The enzymes behaved differently when attempts were made to prepare their alkaline pyridine haemochromagens. A solution of hydroxylase A (1 ml containing 0.83 mg of protein) was mixed with 1 ml of 0.2M-NaOH/pyridine (3:1, v/v), and a few crystals of sodium dithionite were added. The  $\alpha$ -,  $\beta$ - and  $\delta$ -peaks appeared at 550, 520 and 414 nm respectively with an  $A_{550}$  of 0.12. The spectrum is characteristic of a *c*-type cytochrome (Bartsch, 1971) and from the millimolar absorption coefficient of 31.18 litre·mmol<sup>-1</sup>·cm<sup>-1</sup> for the  $\alpha$ -peak of the pyridine haemochrome of cytochrome *c* and a mol.wt. of 114000, the ratio of haem to protein for hydroxylase A was calculated as 1.06:1. When the same treatment was applied to hydroxylase B, reoxidation of the ferrohaemochromagen was too rapid for the spectrum to be recorded. However, a spectrum with an  $\alpha$ -peak at 550 nm was obtained when 1.5 ml of purified enzyme (0.8 mg of protein/ml) was mixed with 1.5 ml of the NaOH/pyridine mixture and placed, with a few crystals of sodium dithionite, in a Thunberg cuvette which was alternately evacuated and flushed with argon three times.

#### Flavin content of *p*-cresol methylhydroxylase

A solution of hydroxylase A (6 ml containing 0.71 mg of protein/ml) was acidified to pH 2.0 with 2M-HCl at 4°C and 3 vol. of ice-cold acetone was added. The solution was centrifuged at 18000g for 15 min at 4°C, yielding a yellow pellet that contained approx. half of the protein. The pellet did not dissolve in 1.0 ml of 50 mM-phosphate buffer, pH 7.0, but, after acidification of the suspension to pH 2.0 with 2M-HCl and incubation in the dark at 100°C for 3 min followed by centrifugation at 18000g for 15 min, some yellow colour remained in the supernatant. The spectrum (Fig. 4) suggests the presence of flavin and some haem. Addition of sodium dithionite resulted in the disappearance of the peak at 450 nm, which was then restored by oxidation by bubbling air through the solution.

Attempts to dissociate flavin from pure enzyme, by incubating at 100°C for 5 min or by adding (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> to 90% saturation followed by acidification to pH 3, were unsuccessful. However, further evidence for flavin came from fluorescence studies on pure enzyme or on a tryptic digest of the yellow protein precipitated by acid/acetone. The latter was prepared by resuspending the pellet in 2 ml of 50 mM-phosphate buffer, pH 7.0, and incubating at 38°C for




Fig. 4. Absorption spectrum of yellow component of hydroxylase A after acetone/HCl extraction. The spectrum of the material described in the text was recorded against a blank of 50mM-KH<sub>2</sub>PO<sub>4</sub>/NaOH buffer, pH7.0.

3h with 3mg of trypsin. Undigested protein was removed by heating at 100°C for 3min followed by centrifuging. The preparations gave a fluorescence emission peak at 523nm with maximum fluorescence when excited at 450nm and for the tryptic digest the excitation spectrum for emission at 520nm gave peaks at 465, 450, 380 and 295nm.

The peak at 465nm was much larger in the spectrum obtained with pure enzyme and is probably due to the presence of the haem group.

The acetone from the supernatant of the original extraction was evaporated with a stream of N<sub>2</sub>, and the pyridine haemochromagen was prepared, as before, with the residue. This gave a spectrum with peaks at 550, 523 and 418nm.

A similar acid/acetone extraction of hydroxylase B again resulted in a yellow precipitate. Attempts to dissociate flavin were again unsuccessful, although the pure enzyme fluoresced at 525nm with a maximum when excited at 450nm and the excitation spectrum gave peaks at 465, 450, 380 and 295nm. Although the supernatant from acid/acetone extraction was red-brown in colour, attempts to prepare the alkaline pyridine haemochrome failed.

#### Oxidation of 3,5-xyleneol by cell extracts

Measurements of the oxidation of 3,5-xyleneol by extracts of cells grown on 3,5-xyleneol were made in a Warburg apparatus. There was no increase in the




Fig. 5. Oxidation of 3,5-xyleneol by (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>-treated extracts.

Warburg flasks contained (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>-treated extract (40mg of protein) from 3,5-xyleneol-grown cells and 6μmol of 3,5-xyleneol in 2.8ml of 50mM-phosphate buffer, pH7.0. The centre well contained 0.2ml of 20% (w/v) KOH. Flasks also contained an NADH-generating system (●), NAD<sup>+</sup> (3μmol) (○), an NADPH-generating system (▲) or NADP<sup>+</sup> (3μmol) (△).

rate of oxygen uptake above the endogenous rate when 6μmol of 3,5-xyleneol was added to a reaction mixture (40mg of crude-extract protein in 2.8ml of 50mM-phosphate buffer, pH7.0) unless NAD<sup>+</sup> (3μmol) was included, when a rate of 600μl of O<sub>2</sub>/h was measured compared with 260μl of O<sub>2</sub>/h for a flask without substrate. This stimulation was not given by NADP<sup>+</sup> or an NADPH-generating system. When an NADH-generating system was used there was no significant oxidation above endogenous until after a lag of around 70min. However, when (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>-treated extracts were used, an NADH-generating system was required for oxidation of 3,5-xyleneol, and NAD<sup>+</sup> alone was not effective (Fig. 5). The rate after such a treatment was much less than that with crude extract. Attempts to assay the 3,5-xyleneol hydroxylase in the oxygen monitor were unsuccessful.

#### Substrate specificity of 3,5-xyleneol hydroxylase

The relative rates of oxygen uptake when four phenols were tested as substrates are shown in Table 4.



Table 4. *Relative rates of oxidation of phenols by (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>-treated extracts*

Oxidation of substrates by (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>-treated extracts from 3,5-xylenol-grown cells was followed in a Warburg apparatus. Flasks contained approx. 40 mg of protein in 50 mm-phosphate buffer, pH 7.0, an NADH-generating system and 6 μmol of substrate in a total volume of 2.8 ml with 0.2 ml of 20% (w/v) KOH in the centre well. The endogenous rate was subtracted from each and the rates relative to that for 3,5-xylenol are given.

Substrate	Relative rate
3,5-Xylenol	100
2,5-Xylenol	100
<i>m</i> -Cresol	18
<i>p</i> -Cresol	0

#### *Inhibition of 3,5-xylenol hydroxylase*

The oxidation of 3,5-xylenol by crude extracts supplemented with NAD<sup>+</sup> was completely inhibited by NaCN at a concentration of 0.5 mM. No inhibition was observed when the atmosphere in the flask consisted of approx. 50% CO in air.

#### **Discussion**

The oxidation of both 3,5-xylenol and *p*-cresol by cells grown on 3,5-xylenol (Fig. 1) suggested the presence of a methylhydroxylase capable of attacking groups both *meta* and *para* to the hydroxy group. However, the more detailed study of these activities has shown that the initial hydroxylations are due to two very different enzymes.

The requirement of the 3,5-xylenol hydroxylase for NADH, as shown by experiments with (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>-treated extracts, is typical of an enzyme of the mono-oxygenase type, and inhibition by cyanide but not CO suggests a system more like the octane hydroxylase of McKenna & Coon (1970) than the cytochrome *P*-450 system described by Jurtschuk & Cardini (1971). The apparent lag in 3,5-xylenol oxidation by crude extracts supplied with NADH may be an artefact and reflect high activities of alternative NADH-oxidizing systems. A similar observation was reported for a bacterial methane-oxidizing system and it was shown that addition of methane to extracts caused a redirection of electron flow to O<sub>2</sub> rather than any overall stimulation of the rate of O<sub>2</sub> consumption (Tonge *et al.*, 1974).

Extracts also oxidized other *m*-methyl-substituted phenols (Table 4), which is in accord with the partial oxidation of 2,5-xylenol to 3-hydroxy-4-methylbenzoate by this organism and the oxidation of *m*-cresol by 3,5-xylenol-grown cells (Hopper & Chapman, 1971). However, there was no activity

with *p*-cresol and clearly this enzyme does not account for its oxidation by 3,5-xylenol-grown cells (Fig. 1).

Oxidation of the methyl group of *p*-cresol is initiated by a second hydroxylase with very different properties, closely resembling the enzyme described by Hopper & Taylor (1977). The requirement for an electron acceptor and the activity under anaerobic conditions suggests hydroxylation by dehydrogenation followed by hydration, by the mechanism proposed by Hopper (1976) rather than by an oxygenase enzyme. Such a mechanism requires the appropriate groups *para* to each other and all of the substrates for the enzyme have this orientation (Table 3). A similar scheme has been postulated for hydroxylation of  $\alpha$ -conidindrin in its metabolism by *Pseudomonas multivorans*, although no enzymes have been demonstrated (Toms & Wood, 1970). *p*-Hydroxybenzyl alcohol was identified as a product, but a second dehydrogenation by the enzyme can convert this into *p*-hydroxybenzaldehyde, which was the major product under the conditions used. This second reaction is not necessarily of physiological importance, since cells also contain NAD<sup>+</sup>-linked alcohol dehydrogenases (Keat & Hopper, 1976).

The *p*-cresol methylhydroxylase in 3,5-xylenol-grown cells (hydroxylase A) was purified, although electrophoresis on polyacrylamide gels gave three bands (Fig. 2). These were probably due to aggregation, since all three were active and a re-run of protein eluted from one band again gave three bands. Also, only one band was seen after electrophoresis on gels containing sodium dodecyl sulphate. A similar phenomenon has been reported by Pujar & Ribbons (1976) for 4,5-dihydroxyphthalate decarboxylase. The mol.wt. of 58000 for the hydroxylase from electrophoresis on sodium dodecyl sulphate-containing gels compared with 114000 by ultracentrifuge studies suggests there are two subunits of about equal molecular weight. One of these is a *c*-type cytochrome as judged by the spectrum of the alkaline pyridine haemochromogen (Bartsch, 1971) and a value of one haem per molecule of enzyme can be calculated. The cytochrome *c* subunit can be extracted with acid/acetone, although the peak at 413 nm in the absorption spectrum of the yellow protein remaining after extraction indicates some residual haem. This also accounts for one of the peaks in the fluorescence excitation spectrum for this yellow protein. The remaining peaks in both spectra suggest the presence of flavin and, as is typical for a flavin, the peak at 450 nm is lost on reduction. Attempts to dissociate the flavin were unsuccessful and possibly it is attached covalently to the second subunit. The involvement of the haem in the enzyme reaction was demonstrated by its reduction by substrate, and the enzyme is one of the class of proteins known as flavocytochrome *c* (Bartsch *et al.*, 1968).

A similar *p*-cresol methylhydroxylase (hydroxylase B) with the same specificity was induced by growth on *p*-cresol itself. It too converted *p*-cresol into the alcohol and aldehyde under anaerobic conditions with a requirement for an electron acceptor and again contained haem and flavin in two subunits of equal size. However, this does not appear to be the same enzyme as that in 3,5-xyleneol-grown cells. The patterns of bands after electrophoresis of the purified enzymes are clearly different (Fig. 2), as also were the bands of activity after electrophoresis of crude extracts, which showed that both enzymes were induced by growth on a mixture of *p*-cresol and 3,5-xyleneol. Hydroxylase B, unlike hydroxylase A, was not retained on the DEAE-cellulose column during purification and gave a lower molecular weight and slightly lower  $K_m$  values for *p*-cresol and *p*-hydroxybenzyl alcohol than hydroxylase A. The pyridine haemochromogen of hydroxylase B was reoxidized much more rapidly than that of hydroxylase A, making the recording of its spectrum more difficult.

Thus oxidation of *p*-cresol by 3,5-xyleneol-grown cells is the result of gratuitous induction of a *p*-cresol methylhydroxylase. However, the physiological role of this enzyme is not known, since it is not present in *p*-cresol-grown cells, which contain a similar, but not identical, hydroxylase.

We thank Mr. D. G. Taylor for skilled technical assistance and the Science Research Council for a Research Studentship to M. J. K.

## References

- Andrews, P. (1965) *Biochem. J.* **96**, 595–606
- Bartsch, R. G. (1971) *Methods Enzymol.* **23**, 344–363
- Bartsch, R. G., Meyer, T. E. & Robinson, A. B. (1968) in *Structure and Function of Cytochromes* (Okunuki, K., Kamen, M. D. & Sekuu, I., eds.), pp. 443–451, University of Tokyo Press, Tokyo
- Byrne, G. A. (1965) *J. Chromatogr.* **20**, 528–540
- Chapman, P. J. (1972) in *Degradation of Synthetic Organic Molecules in the Biosphere*, pp. 17–55, Printing and Publishing Office, National Academy of Sciences, Washington
- Gholson, R. K., Baptist, J. N. & Coon, M. J. (1963) *Biochemistry* **2**, 1155–1159
- Gordon, A. H. (1969) in *Electrophoresis of Proteins in Polyacrylamide and Starch Gels* (Work, T. S. & Work, E., eds.), pp. 9–13, North-Holland, Amsterdam and London
- Gornall, A. G., Bardawill, C. J. & David, M. M. (1949) *J. Biol. Chem.* **177**, 751–766
- Hopper, D. J. (1976) *Biochem. Biophys. Res. Commun.* **69**, 462–468
- Hopper, D. J. & Chapman, P. J. (1971) *Biochem. J.* **122**, 19–28
- Hopper, D. J. & Taylor, D. G. (1973) *J. Bacteriol.* **122**, 1–6
- Hopper, D. J. & Taylor, D. G. (1977) *Biochem. J.* **167**, 155–162
- Jurtshuk, P. & Cardini, G. E. (1971) *C.R.C. Crit. Rev. Microbiol.* **1**, 239–289
- Keat, M. J. & Hopper, D. J. (1976) *Biochem. Soc. Trans.* **4**, 494–495
- Lowry, O. H., Rosebrough, N. J., Farr, A. L. & Randall, R. J. (1951) *J. Biol. Chem.* **193**, 265–275
- McKenna, E. J. & Coon, M. J. (1970) *J. Biol. Chem.* **245**, 3882–3889
- Nozaka, J. & Kusunose, M. (1968) *Agric. Biol. Chem.* **32**, 1033–1039
- Ornstein, L. & Davis, B. J. (1964) *Ann. N.Y. Acad. Sci.* **121**, 305–650
- Pastuska, G. (1961) *Z. Anal. Chem.* **179**, 355–358
- Pujar, B. G. & Ribbons, D. W. (1976) *Abstr. Am. Soc. Microbiol. Annu. Meet.* **76th** 145
- Rosenberger, R. F. & Elsdon, S. R. (1960) *J. Gen. Microbiol.* **22**, 726–739
- Ruettinger, R. T., Griffith, G. R. & Coon, M. J. (1977) *Arch. Biochem. Biophys.* **183**, 528–537
- Smith, I. (1960) *Chromatographic and Electrophoretic Techniques*, 2nd edn., vol. 1, pp. 291–298, W. Heinemann, London
- Toms, A. & Wood, J. M. (1970) *Biochemistry* **9**, 733–740
- Tonge, G. M., Harrison, D. E. F., Knowles, C. J. & Higgins, I. J. (1974) *FEBS Lett.* **58**, 293–299
- Vogel, A. I. (1956) *Practical Organic Chemistry*, 3rd edn., p. 185, Longman, Green and Co., London
- Weber, K. & Osborn, M. (1969) *J. Biol. Chem.* **244**, 4406–4412
- Yphantis, D. A. (1964) *Biochemistry* **3**, 297–317