

# PCR amplification introduces errors into mononucleotide and dinucleotide repeat sequences

L A Clarke, C S Rebelo, J Gonçalves, M G Boavida, P Jordan

## Abstract

**The polymerase chain reaction (PCR) is used universally for accurate exponential amplification of DNA. We describe a high error rate at mononucleotide and dinucleotide repeat sequence motifs. Subcloning of PCR products allowed sequence analysis of individual DNA molecules from the product pool and revealed that: (1) monothymidine repeats longer than 11 bp are amplified with decreasing accuracy, (2) repeats generally contract during PCR because of the loss of repeat units, (3) Taq and proofreading polymerase Pfu generate similar errors at mononucleotide and dinucleotide repeats, and (4) unlike the parent PCR product pool, individual clones containing a single repeat length produce no “shadow bands”. These data demonstrate that routine PCR amplification alters mononucleotide and dinucleotide repeat lengths. Such sequences are common components of genetic markers, disease genes, and intronic splicing motifs, and the amplification errors described here can be mistaken for polymorphisms or mutations.**

(J Clin Pathol: Mol Pathol 2001;54:351–353)

Keywords: polymerase chain reaction; nucleotide repeat sequences; shadow bands

The polymerase chain reaction (PCR) is one of the most widely used techniques in molecular biology and has made possible a great variety of both diagnostic and research applications. Examples are the detection of gene mutations, the analysis of polymorphic markers and microsatellite loci, analysis of gene expression, DNA cloning, and site directed mutagenesis. Because PCR involves the exponential amplification of target sequences, a high degree of polymerase fidelity is essential if the introduction of a large number of replication errors during the PCR reaction is to be avoided. Most commercially available Taq polymerases introduce errors at the rate of approximately  $10^{-5}$  to  $10^{-6}$  point mutations/bp/duplication, with higher fidelity polymerases such as Pfu and Deep Vent generating up to eight times fewer errors.<sup>1</sup> In contrast, we found that mononucleotide and dinucleotide repeats were not faithfully reproduced during PCR.

## Methods and results

A stretch of 26 A nucleotides in intron 5 of the hMSH2 gene, termed Bat-26,<sup>2</sup> was amplified from the genomic DNA of several healthy individuals under routine PCR conditions (200 ng

genomic DNA and 1.25 U AmpliTaq (Perkin Elmer, Branchburg, New Jersey, USA) in 50  $\mu$ l containing 1.5mM MgCl<sub>2</sub>, 300 ng each primer,<sup>2</sup> and 200 $\mu$ M dNTPs). Direct sequencing of the PCR product produced an illegible sequence after the mononucleotide repeat, indicating a possible difference between two alleles. Such allele differences, along with any polymerase slippage incurred during the PCR reaction, can both be visualised using gene scanning technology. However, for our study a strategy of subcloning and sequencing of individual clones of the PCR product was used for the qualitative determination of the composition of the PCR product pool, by identifying the sequences of individual DNA molecules. This revealed differences in the length of the Bat-26 poly-A stretch, which varied from 19 to 28 bp, which is incompatible with the concept of simple polymorphism or mutation (table 1). The sequencing of more than 30 individual clones revealed that the repeat was predominantly shortened, with only 35% of the clones containing the predicted sequence of (A)<sub>26</sub> (table 1). These data suggested that a systematic polymerase error had taken place during the PCR reaction, which was specific to the mononucleotide repeat.

To determine in more detail the performance of PCR amplification at similar sequences, repeats of 21, 13, 11, and nine monothymidines were amplified and subcloned. The polypyrimidine tract of the hMLH1 intron 11 splice acceptor site contains a (T)<sub>21</sub> monothymidine repeat, termed Bat-21 (AcNb U40971),<sup>3</sup> with an adjacent (TA)<sub>11</sub> dinucleotide repeat. The polypyrimidine tract preceding exon 2 in the gene hMSH2 contains a (T)<sub>13</sub> stretch, termed Bat-13 (AcNb U41207),<sup>3</sup> and intron 4 of the human RAC1 gene contains both (T)<sub>9</sub> and (T)<sub>11</sub> runs (AcNb AJ132695).<sup>4</sup> As shown in table 1, sequencing of individual cloned PCR products revealed incorrect amplification of the (T)<sub>21</sub> and (T)<sub>13</sub> repeats, whereas (T)<sub>9</sub> was replicated faithfully. The limit for correct amplification was reached with (T)<sub>11</sub>, where 90% of the cloned products contained the predicted number of thymidines. The predominant observation was of repeat contraction; the (T)<sub>21</sub> repeat appeared to expand, but this tendency was accounted for upon inspection of individual Bat-21 clones by A-T transversions in the adjacent TA repeat. Two different amplification errors at this combined repeat therefore led to an overall expansion of the poly-T stretch.

To determine the performance of the high fidelity Pfu polymerase at such sequences, the

Centro de Genética Humana, Instituto Nacional de Saúde ‘Dr. Ricardo Jorge’, Avenida Padre Cruz, 1649–016 Lisboa, Portugal  
L A Clarke  
C S Rebelo  
J Gonçalves  
M G Boavida  
P Jordan

Correspondence to:  
Dr Jordan  
peter.jordan@insa.min-saude.pt

Accepted for publication  
27 March 2001



forming a size ladder according to the length of the poly-A stretch. This demonstrates that the phenomenon of shadow bands commonly observed in PCR based genetic analysis of microsatellite loci can be attributed at least in part to artefactual PCR amplification errors. In addition, others have shown that improper annealing of PCR products during non-denaturing gel electrophoresis can further compound the occurrence of multiple band patterns.<sup>13</sup>

### Discussion

In our study, we describe a high error rate in PCR amplification of mononucleotide and dinucleotide repeats from genomic DNA. Our findings relate mainly to monothymidine/monoadenosine repeats, and suggest that the longer the repeat, the greater the errors made during amplification. Although our additional data include only TA and CA dinucleotide repeats, we believe it is possible to extrapolate the general rule that PCR amplification of mononucleotide or dinucleotide repeats results in error. Use of the high fidelity proofreading polymerase Pfu in place of Taq restricted the occurrence of such errors only in the case of short monothymidine repeats. We think that contraction of the repeat itself is the most common type of error occurring during PCR amplification. This type of error is commonly referred to as polymerase "slippage" and is probably caused by slipped strand mispairing.<sup>14,15</sup> Alternatively, the loss or gain of nucleotide repeat units without affecting the surrounding sequence<sup>10,12</sup> can be explained by mega-priming. In this case, fragments that were either incompletely synthesised or broken in their repetitive element during PCR can anneal under formation of mismatches, then extend, resulting in repeat length variation, independent of polymerase type or PCR conditions. This conclusion was drawn from work using synthetic oligonucleotides in the absence of genomic DNA.<sup>16,17</sup> Previous work has suggested that such polymerase error can be reduced by optimisation of PCR reaction conditions and buffer composition.<sup>1,13,18-20</sup> In our present study, a fixed set of routine PCR conditions were used to determine the effect that repeat size and length alone would have on polymerase error rate. Our data lead to the conclusion that polymerase performance itself imposes considerable constraints upon PCR amplification fidelity independent of PCR conditions, primer composition, or PCR product length.

Although the molecular mechanism of the described errors during amplification of repetitive DNA sequence motifs remains a matter of discussion, such errors are frequently encountered during cloning or genetic analysis of introns, polypyrimidine tracts, or microsatellite markers. In addition, they are of great interest during analysis and diagnosis of various diseases. For example, mononucleotide and dinucleotide microsatellite sequences are hotspots for mammalian polymerase error during *in vivo* DNA replication,<sup>21</sup> and are highly unstable in tumours resulting from mismatch repair deficiency.<sup>22,23</sup> On this basis, they are

widely used as markers for certain cancer syndromes,<sup>24</sup> including hereditary non-polyposis colorectal cancer.<sup>25</sup> Furthermore, dinucleotide repeats are used as polymorphic markers to determine heterozygosity. With the increased use of automated DNA sequencing, the described *in vitro* amplification errors at repeat motifs can easily be mistaken, in research and diagnostics, for polymorphism or mutation.

We thank C Caldas for comments on the manuscript, L Vieira for donating D15S128 primers, S Pedro for running the ABI sequencing apparatus, and S Beck and S Vieira for assistance with high performance liquid chromatography analysis. This study was supported by PRAXIS XXI grant 2/2.1/SAU/1397/95 from the Fundação para a Ciência e a Tecnologia, and PRAXIS XXI postdoctoral fellowship BPD/4140/96 to LAC.

- Cline J, Braman JC, Hogrefe HH. PCR fidelity of Pfu DNA polymerase and other thermostable DNA polymerases. *Nucleic Acids Res* 1996;24:3546-51.
- Hoang JM, Cottu PH, Thuille B, et al. Bat-26, an indicator of the replication error phenotype in colorectal cancers and cell lines. *Cancer Res* 1997;57:300-3.
- Zhou XP, Hoang JM, Cottu P, et al. Allelic profiles of mononucleotide repeat microsatellites in control individuals and in colorectal tumours with or without replication errors. *Oncogene* 1997;15:1713-18.
- Matos P, Skaug J, Márques B, et al. Small GTPase Rac1: structure, localisation and expression of the human gene. *Biochem Biophys Res Commun* 2000;277:741-51.
- Weissenbach J, Gyapay G, Dib C, et al. A second-generation linkage map of the human genome. *Nature* 1992;359:794-801.
- Lubahn DB, Brown TR, Simental JA, et al. Sequence of the intron/exon junctions of the coding region of the human androgen receptor gene and identification of a point mutation in a family with complete androgen insensitivity. *Proc Natl Acad Sci U S A* 1989;86:9534-8.
- Tsuji S. Molecular genetics of triplet repeats: unstable expansions of triplet repeats as a new mechanism for neurodegenerative diseases. *Intern Med* 1997;36:3-8.
- Zoghbi HY, Orr HT. Glutamine repeats and neurodegeneration. *Annu Rev Neurosci* 2000;23:217-47.
- Walsh PS, Fildes NJ, Reynolds R. Sequence analysis and characterization of stutter products of the tetranucleotide repeat locus vWA. *Nucleic Acids Res* 1996;24:2807-12.
- Hauge XY, Litt M. A study of the origin of "shadow bands" seen when typing dinucleotide repeat polymorphisms by the PCR. *Hum Mol Genet* 1993;2:411-15.
- Litt M, Hauge XY, Sharma V. Shadow bands seen when typing polymorphic dinucleotide repeats: some causes and cures. *Biotechniques* 1993;15:280-4.
- Murray V, Monchawin C, England PR. The determination of the sequences present in the shadow bands of a dinucleotide repeat PCR. *Nucleic Acids Res* 1993;21:2395-8.
- Bovo D, Ruge M, Shiao YH. Origin of spurious multiple bands in the amplification of microsatellite sequences. *J Clin Pathol: Mol Pathol* 1999;52:50-1.
- Levinson G, Gutman GA. Slipped-strand mispairing: a major mechanism for DNA sequence evolution. *Mol Biol Evol* 1987;4:203-21.
- Krawczak M, Cooper DN. Gene deletions causing human genetic disease: mechanisms of mutagenesis and the role of the local DNA sequence environment. *Hum Genet* 1991;86:425-41.
- Hamdan H, Tynan JA, Fenwick RA, et al. Automated detection of trinucleotide repeats in fragile X syndrome. *Mol Diagn* 1997;2:259-69.
- Behn-Krappa A, Doerfler W. Enzymatic amplification of synthetic oligodeoxyribonucleotides: implications for triplet expansions in the human genome. *Hum Mutat* 1994;3:19-24.
- Brail L, Fan E, Levin DB, et al. Improved polymerase fidelity in PCR-SSCP. *Mutat Res* 1993;303:171-5.
- Lyons-Darden T, Topal MD. Effects of temperature, Mg<sup>2+</sup> concentration and mismatches on triplet-repeat expansion during DNA replication *in vitro*. *Nucleic Acids Res* 1999;27:2235-40.
- Wu MJ, Chow LW, Hsieh M. Amplification of GAA/TTC triplet repeat *in vitro*: preferential expansion of (TTC)<sub>n</sub> strand. *Biochim Biophys Acta* 1998;1407:155-62.
- Kunkel TA. The mutational specificity of DNA polymerase- $\beta$  during *in-vitro* DNA synthesis. *J Biol Chem* 1985;260:5787-96.
- Liu B, Nicolaidis NC, Markowitz S, et al. Mismatch repair gene defects in sporadic colorectal cancers with microsatellite instability. *Nat Genet* 1995;9:48-55.
- Marra G, Schär P. Recognition of DNA alterations by the mismatch repair system. *Biochem J* 1999;388:1-13.
- Rodriguez-Bigas MA, Boland CR, Hamilton SR, et al. National Cancer Institute workshop on hereditary non-polyposis colorectal cancer syndrome: meeting highlights and Bethesda guidelines. *J Natl Cancer Inst* 1997;89:1758-62.
- Lynch HT. Hereditary non-polyposis colorectal cancer (HNPCC). *Cytogenet Cell Genet* 1999;86:130-5.