

The 14–Amino Acid CLV3, CLE19, and CLE40 Peptides Trigger Consumption of the Root Meristem in *Arabidopsis* through a CLAVATA2-Dependent Pathway^W

Martijn Fiers,^{a,b} Elzbieta Golemic,^c Jian Xu,^d Lonneke van der Geest,^a Renze Heidstra,^{b,d} Willem Stiekema,^b and Chun-Ming Liu^{a,b,1}

^aPlant Research International, 6700 AA Wageningen, The Netherlands

^bCentre for BioSystems Genomics, 6700 AB Wageningen, The Netherlands

^cInstitute of Plant Physiology, Polish Academy of Sciences, 30-239 Kraków, Poland

^dDepartment of Molecular Genetics, Utrecht University, 3584 CH, Utrecht, The Netherlands

CLAVATA3 (CLV3), CLV3/ESR19 (CLE19), and CLE40 belong to a family of 26 genes in *Arabidopsis thaliana* that encode putative peptide ligands with unknown identity. It has been shown previously that ectopic expression of any of these three genes leads to a consumption of the root meristem. Here, we show that in vitro application of synthetic 14–amino acid peptides, CLV3p, CLE19p, and CLE40p, corresponding to the conserved CLE motif, mimics the overexpression phenotype. The same result was observed when CLE19 protein was applied externally. Interestingly, *clv2* failed to respond to the peptide treatment, suggesting that CLV2 is involved in the CLE peptide signaling. Crossing of the CLE19 overexpression line with *clv* mutants confirms the involvement of CLV2. Analyses using tissue-specific marker lines revealed that the peptide treatments led to a premature differentiation of the ground tissue daughter cells and misspecification of cell identity in the pericycle and endodermis layers. We propose that these 14–amino acid peptides represent the major active domain of the corresponding CLE proteins, which interact with or saturate an unknown cell identity-maintaining CLV2 receptor complex in roots, leading to consumption of the root meristem.

INTRODUCTION

In multicellular organisms, cell-to-cell communication is essential for coordinating growth and differentiation. In animals, peptides are known to be the major players in cell-to-cell communication (for review, see Alberts et al., 1994). This is in contrast with plants in which most intercellular communication is mediated by phytohormones, such as auxin, cytokinin, gibberellic acid, abscisic acid, ethylene, and brassinosteroids (Mandave, 1988; Kende and Zeevaart, 1997). However, in recent years, several putative peptide ligands have been identified in plants (for review, see Lindsey et al., 2002; Ryan et al., 2002) and have been shown to mediate signaling events during plant–pathogen interactions (Pearce et al., 1991), cell division (Matsubayashi and Sakagami, 1996), and anther–stigma interactions (Schopfer et al., 1999; Kim et al., 2003).

CLAVATA3 (CLV3) is a putative peptide ligand of *Arabidopsis thaliana* that interacts with a disulphide-linked CLV1/CLV2 re-

ceptor complex to restrict the stem cell population in the shoot apical meristem (SAM) in a non-cell-autonomous manner (Fletcher et al., 1999). CLV1 is a membrane-bound leucine-rich repeat receptor kinase, while CLV2 is a leucine-rich repeat receptor-like protein lacking a kinase domain (Clark et al., 1997; Jeong et al., 1999). The stem cells are marked by CLV3 expression, while the SAM organizing center is marked by the expression of the WUSCHEL (WUS) stem cell-promoting transcription factor (Laux, 2003). A feedback regulation loop between CLV3 and WUS maintains the number of stem cells in the SAM (Brand et al., 2000; Schoof et al., 2000). As such, *clv1*, *clv2*, and *clv3* mutants have enlarged SAMs, while the *wus* mutant or CLV3 overexpression terminates SAM development (Laux et al., 1996; Hobe et al., 2003). Although biochemical studies showed that CLV3 is required for the formation of a 450-kD functional CLV1/CLV2 receptor complex with several associated proteins (Trotochaud et al., 1999), the biochemical nature of the active ligand encoded by CLV3 has not yet been elucidated.

CLV3 belongs to the CLV3/ESR (CLE) family of 26 genes in *Arabidopsis*, of which 25 are transcribed in one or more tissues (Cock and McCormick, 2001; Hobe et al., 2003; Sharma et al., 2003; Fiers et al., 2004). These genes encode small proteins that contain a putative secretion signal at their N termini and a conserved 14–amino acid motif (CLE motif) at or near their C termini (Cock and McCormick, 2001). In the *clv3-1* and *clv3-5* mutants, a single amino acid change (from G to A) in the CLE motif is enough to disrupt the function of CLV3 (Fletcher et al., 1999), indicating the importance of this motif. First identified in maize

¹To whom correspondence should be addressed. E-mail chunming.liu@wur.nl; fax 31-317-418094.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Chun-Ming Liu (chunming.liu@wur.nl).

^WOnline version contains Web-only data.

Article, publication date, and citation information can be found at www.plantcell.org/cgi/doi/10.1105/tpc.105.034009.

(*Zea mays*) as being expressed in endosperm regions surrounding the embryo, *ESR* genes encode extracellular proteins with unknown functions (Bonello et al., 2002). Such *CLE* genes have not only been found in many plant species, but also in parasitic nematodes (Wang et al., 2005).

One interesting feature of these *CLE* proteins is that they share very little sequence similarity outside the *CLE* motif (Cock and McCormick, 2001). Remarkably, *CLE40* from *Arabidopsis* and *Hg-SYV46* from nematode can fully complement the *clv3* mutant phenotype when expressed under the control of the *CLV3* and cauliflower mosaic virus 35S (*35S*) promoter, respectively, illustrating the functional redundancy of these genes (Hobe et al., 2003; Wang et al., 2005). The redundancy could also be seen from the T-DNA insertional mutants of *CLE19* and *CLE40*, which gave no phenotype and a very subtle phenotype, respectively (Hobe et al., 2003; Fiers et al., 2004). By contrast, overexpression of several *CLE* genes, such as *CLV3*, *CLE19*, *CLE40*, or *Hg-SYV46*, under the control of the *35S* promoter induce striking developmental phenotypes in root and shoot development in *Arabidopsis* (Hobe et al., 2003; Fiers et al., 2004; Wang et al., 2005). Termination of root meristem development has been observed in transgenic plants overexpressing any of these four genes (Hobe et al., 2003; Fiers et al., 2004; Wang et al., 2005). These studies point to the existence of common and redundant signaling machinery in roots that responds to different *CLE* genes to trigger the differentiation of the root meristem.

Here, we report the development of a novel in vitro root assay to elucidate how *CLE* genes trigger the consumption of the root meristem. Application of chemically synthesized 14-amino acid peptides, *CLV3p*, *CLE19p*, and *CLE40p* (hereafter referred to as *CLE* peptides), corresponding to the conserved *CLE* motif of the related *CLE* proteins, led to consumption of the root meristem in a manner that closely resembles the overexpression phenotypes mentioned above. The same effect was achieved by application of heterologously produced full-length *CLE19* protein. The *clv2* mutation abolished the sensitivity of roots to these peptides, suggesting that *CLV2* is involved in the *CLE* signaling pathway. Using different marker lines, we demonstrate that the peptide treatments led to an inward shifting of cell identity in several cell layers and to a premature differentiation of the ground tissue daughter cells. We propose that the *CLE* peptides represent the major functional domain of the *CLE* proteins and that this domain interacts with *CLV2* to trigger the termination of the root meristem.

RESULTS

In Vitro Application of *CLE* Peptides Causes a Short Root Phenotype

It has been reported previously that overexpression of *CLV3*, *CLE19*, *CLE40*, or *Hg-SYV46* in *Arabidopsis* under the control of *35S* promoter resulted in termination of root meristem development (Hobe et al., 2003; Fiers et al., 2004; Wang et al., 2005). The same phenotype was observed when *CLE19* was expressed under the control of the root meristem-specific promoter *RCH1* (Casamitjana-Martinez et al., 2003). The amino acid sequences

of *CLV3*, *CLE19*, and *CLE40* were compared to identify common motifs. The only conserved sequence among these proteins resides in the *CLE* motif (Figure 1A). We therefore examined whether a chemically synthesized 14-amino acid peptide of *CLE19p* (Figure 1B), corresponding to the conserved *CLE* motif of *CLE19*, is also able to trigger the consumption of the root meristem. As a control, we used a 16-amino acid peptide corresponding to the C terminus of *AGAMOUS* that has no similarity with the *CLE* peptides (*AGp*, Figure 1B). *Arabidopsis* seeds were germinated on vertical plates with media containing different concentrations of the individual peptides. The length of the primary root was measured after 7 d (Figure 2A). Although no clear effect on root growth was observed when *CLE19p* was applied at the 0.1 to 1 μ M concentration range, a dramatic inhibitive effect was observed at 10 μ M or higher (Figure 2A). Further increase of the *CLE19p* concentration from 10 to 100 μ M gave only a slight decrease in root length (Figure 2A). *AGp* had no significant effect on root length. As such, we used a concentration of 10 μ M for the subsequent analyses. These results gave us the first indication that the conserved motif of *CLE19* is sufficient to mimic the short root phenotype generated by overexpression of the *CLE19* gene.

Furthermore, *CLV3p* and *CLE40p* (Figure 1B), corresponding to the *CLE* motifs of *CLV3* and *CLE40*, respectively, were tested for their effect on root growth. Unlike *CLV3*, *CLE40* is expressed in roots (Hobe et al., 2003). Several peptides, including a mutant *CLV3* peptide (*CLV3m*, with a G-to-A conversion as in *clv3-1* and *clv3-5* mutants), two truncated peptides (*CLV3t* and *CLE19t*, with five amino acids removed the C termini of *CLV3p* and *CLE19p*, respectively), and a shuffled *CLV3* peptide (*CLV3s*, the same amino acid composition as *CLV3p* but with a shuffled sequence), were chemically synthesized as controls (Figure 1B). The peptides were added individually to medium at a concentration of 10 μ M. Seedlings treated with *CLV3p* and *CLE40p*, similar to those treated with *CLE19p*, showed a clear reduction in root length, as compared with the controls with either no peptide or treated with *CLV3m*, *CLV3s*, *CLV3t*, or *CLE19t*, which did not have a significant effect on root length (Figure 2B). Statistical analysis showed no significant differences among treatments with *CLV3p*, *CLE19p*, and *CLE40p*.

To investigate if all peptides derived from the *CLE* motif of the *CLE* proteins are able to produce the short root phenotype, we synthesized a 14-amino acid peptide (*CLE5p*) of *CLE5*. The *CLE* motif of *CLE5* differs from that of *CLV3*, *CLE19*, and *CLE40* (Figure 1B). *CLE5* is expressed in roots (Sharma et al., 2003). *CLE5p* shares between 43 and 50% identity with the *CLE* peptides. Treatment with *CLE5p* did not cause any reduction in root length (Figure 2B), indicating that the sequence of the peptide is critical for the function of the *CLE* peptides in triggering the consumption of root meristem, and not all peptides with a *CLE* motif from one of the 26 *CLE* genes results in a similar response.

To examine if *CLV3p* application in solid media could complement the *clv3* phenotype, we measured the size of the *clv3* SAMs using Nomarski optics after 4, 8, and 14 d of treatment. We failed to detect any changes, suggesting that either *CLV3p* cannot be transported from the root to the SAM, or the peptide is not able to function in the SAM.




Figure 1. The CLE Motif and the Peptides Used in the Root Assays.

(A) Alignment of CLV3, CLE5, CLE19, *Bn* CLE19, and CLE40. The signal sequences are underlined, and the CLE motif is framed.

(B) Chemically synthesized peptides used in the root assays.

Identical amino acids in five or more proteins/peptides are shaded in black, and amino acids that are similar or identical in at least three proteins are highlighted in gray.

The CLE Peptides Trigger the Consumption of the Root Meristem in a Manner Similar to *CLE19* Overexpression

We next investigated if the short root phenotype generated after the CLE peptide treatment resembles the phenotype observed after overexpression of *CLE19* (Fiers et al., 2004). Using Nomarski optics, we observed that roots treated for 14 d with CLV3s, CLV3m, and CLE5p were morphologically indistinguishable from roots grown on plates without peptide (Figures 3A to 3D). These roots have a cell division zone that consists of a population of cytoplasm-dense cells, followed by gradually enlarged elongated cells. By contrast, roots treated with the active CLE peptides were much thinner, with a significantly decreased number of meristematic cells (Figures 3E to 3G). These roots seem to have an equal number of cell layers along the radial axis, as in the wild type, except for a region above the quiescent center (QC) where the formation of ground tissue appeared to be delayed. The thinner root phenotype seems to be caused by reduction in cell expansion. In CLE peptide-treated roots, only a few cytoplasm-dense cells could be recognized, which were immediately followed by elongated and highly vacuolated cells above this region. This consumption of the root meristem closely resembles the phenotype observed in roots of *CLE19* overexpression lines (Casamitjana-Martinez et al., 2003; Fiers et al., 2004).

The numbers of meristematic cells in the primary root of wild-type and CLE peptide-treated roots were quantified by counting the number of nonelongated cytoplasm-dense cells along the

cortex layer, starting from the QC after 14 d of treatment. In wild-type seedlings, the number of meristematic cells was on average 81, while only one to four such cells could be found from seedlings germinated in the presence of a CLE peptide ($n = 10$; Figures 3E to 3H).

Previous reports showed that ectopic expression of *CLV3*, *CLE19*, or *CLE40* caused consumption of the root meristem without immediately disturbing auxin distribution in the columella initial cells or QC function (Casamitjana-Martinez et al., 2003; Hobe et al., 2003; Fiers et al., 2004). To determine if this was also the case after the treatments with CLE peptides, seedlings of *DR5:β-glucuronidase (GUS)* (a reporter line showing auxin distribution; Sabatini et al., 1999) and *QC25* (QC-specific marker; Casamitjana-Martinez et al., 2003) marker lines were examined. Although a clear reduction in root length and a decreased number of meristematic cells were observed, no visible difference was observed in the *DR5:GUS* and *QC25* (see Supplemental Figure 1 online) expression patterns. This observation revealed that neither the peptide treatment nor the overexpression acts primarily on the QC.

CLV2 Is Involved in Perception of the CLE Peptides in Roots

To identify components of the signal transduction pathway involved in perception of the CLE peptide signal in roots, we determined whether the CLV signaling pathway is involved in the short root phenotype by treating *clv1*, *clv2*, and *clv3* mutants




Figure 2. Treatment with CLE Peptides Gave a Short Root Phenotype in *Arabidopsis*.

(A) Effect of different concentrations of CLE19p and AGp (control) on wild-type (*Col-0*) root development. The lengths of the main roots were measured after 7 d of growth on peptide-containing media ($n = 8$ for each treatment).

(B) Effects of different peptides on the root lengths of *Arabidopsis* (*Ler*). The roots were measured after 14 d of growth on media with 10 μ M various peptides. Note that CLE5p, CLV3m, CLV3t, and CLE19t did not inhibit root growth ($n = 10$ for each treatment). Data and error bars represent mean \pm SD.

with the CLE peptides. Seedlings of the wild type (*Landsberg erecta* [*Ler*]), *clv1-1*, and *clv3-2* grown on plates with any of the CLE peptides showed a significantly shorter root length and decreased number of meristematic cells (Figures 4A and 4B). For example, the number of meristematic cells in *clv1-1* roots grown on peptide-free media was 47 ± 12 mm, which was reduced to 2 ± 2 mm after treatment with CLE19p (Figure 4B).

clv2-1, however, showed significantly less sensitivity to the CLE peptides (Figures 4A and 4B). In the absence of peptide application, we did not detect any defects in the root of the *clv2*, using both Nomarski optics and confocal microscopy (see Supplemental Figure 2 online). The primary roots of *clv2* mutants treated with CLE peptides also maintained a normal meristem morphology (Figure 4E), in contrast with fully differentiated root meristems of the wild type, *clv1-1*, and *clv3-2* after the peptide

treatments (Figures 4C, 4D, and 4F). Seedlings of *clv2-1* treated with CLV3p, CLE19p, or CLE40p had 51 to 83 meristematic cells, as compared with two to four such cells for *clv1-1* seedlings and one to three for *clv3-2* (Figure 4B).

Heterologously Produced CLE19 Protein Also Triggers a Consumption of the Root Meristem

To examine if the full-length CLE protein functions like the CLE peptides, a construct was made in which the sequence encoding the *Brassica napus* (*Bn*) CLE19 protein without the putative secretion signal peptide was fused to the C terminus of glutathione *S*-transferase (GST) and expressed in *Escherichia coli*. After cleaving with thrombin and purification (Figure 5A), the *Bn* CLE19 protein was evaluated for its effect on *Arabidopsis* roots. Application of the *Bn* CLE19 protein at a concentration of 10 μ M strongly inhibited root growth of *Arabidopsis* seedlings, as compared with the control GST protein, which did not have any detectable effect (Figure 5B). After 10 d of treatment, the average root length in the GST-treated seedlings was ~ 34 mm, while for the *Bn* CLE19-treated seedlings it was ~ 7 mm (Figure 5B). Nomarski-based observations showed that *Bn* CLE19 protein-treated roots exhibited a similar phenotype as the CLE peptide-treated or *CLE19* overexpression roots. No additional phenotypes were observed as compared with the treatment with CLE19p, demonstrating a functional similarity between the CLE peptides and the full-length protein.

Genetic Analysis Confirms That CLV2 Is Involved in Perceiving the CLE19 Signal in Roots

After establishing the role of CLV2 in perception of CLE peptides in roots, we studied if CLV2 is also needed for the perception of the ectopically expressed *CLE19* in transgenic *Arabidopsis*. We crossed a $P_{35S}::Bn$ *CLE19* overexpression line (as described in Fiers et al., 2004) with *Ler*, *clv1-1*, *clv2-1*, and *clv3-2*. In the F1 generation, we identified seedlings with short roots and transferred them to the greenhouse to obtain F2 seeds. Plants with long roots were selected from each family of F2 seedlings. We reasoned that plants with long roots should not carry the $P_{35S}::Bn$ *CLE19* transgene, unless the line carries a mutation that can suppress the function of the transgene. As shown in Table 1, all plants from the F2 populations of $P_{35S}::Bn$ *CLE19* \times *Ler*, $P_{35S}::Bn$ *CLE19* \times *clv1-1*, and $P_{35S}::Bn$ *CLE19* \times *clv3-2* with a long-root phenotype did not carry the transgene. Plants with both long roots and the transgene were only identified in the F2 population of $P_{35S}::Bn$ *CLE19* \times *clv2-1* (Table 1). All these 11 plants also displayed the *clv2* mutant phenotype in siliques. A similar result was obtained with the $P_{35S}::At$ *CLE19* transgenic line (data not shown), demonstrating that the *clv2* mutation also suppressed the short-root phenotype produced by the *At* *CLE19* transgene. Consistent with this conclusion, none of the short-root plants obtained from the F2 population of $P_{35S}::Bn$ *CLE19* \times *clv2-1* were homozygous for the *clv2* mutation (data not shown), further confirming that the homozygous *clv2* mutation was needed to suppress the short-root phenotype induced by *Bn* *CLE19* overexpression.





Figure 3. Effect of CLE Peptides on Root Meristems.

(A) to (G) The morphology of the primary roots of *Arabidopsis* (*Ler*) 14 d after treatments with different peptides. Note the consumption of the root meristem (E) to (G) after treatments with CLE peptides. The bar in (G) = 50 μ m for (A) to (G).

(H) Number of meristematic cells in roots of *Ler* seedlings after 14 d of treatment with different peptides. Data and error bars represent mean \pm SD ($n = 10$).

Treatment with CLE Peptides Leads to a Misspecification of Cell Identities

To study the effect of CLE peptides on root development, three tissue-specific marker lines were selected, treated, and inspected by confocal microscopy. These lines were (1) *J0571*, which shows green fluorescent protein (*GFP*) expression in cortex and endodermis layers, including the ground tissue initials and their daughter cells, and occasionally in the QC; (2) *P_{SCR}:GFP*, which reflects the expression of the *SCARECROW* (*SCR*) transcription factor (Wysocka-Diller et al., 2000) and shows *GFP* expression in the endodermis layer, QC, and ground tissue initials; and (3) promoter cortex2-driven nuclear-localized Histone 2B:yellow fluorescent protein (*P_{CO2}:YFP-H2B*), which shows *YFP* expression in cortex cells, but not in their initials, nor in the QC (Heidstra et al., 2004). These reporter lines were grown on media containing either CLE peptides or controls (CLV3s, CLV3m, or without peptide).

In all these lines, *GFP* expression was localized in the expected cell layer in the absence of peptides (Figures 6A, 6E, and 6I) or with control peptides (Figures 6B, 6F, and 6H). In all samples examined, a maximum of one ground tissue daughter cell was observed above the single ground tissue initial cell (Figure 6A, labeled with an arrowhead). In a number of samples, neither

visible ground tissue daughter cell nor ground tissue initial cell could be detected (Figure 6I). When seedlings of *J0571* were treated with CLE peptides for 4 d, a single-file, with up to four ground tissue daughter cells, was observed (Figures 6C and 6D, labeled with arrowheads). More than 50% of the plants treated with CLE peptides exhibited this phenotype. This is an early defect, since at this stage the reduced root growth was not yet evident. After an 8-d treatment with CLE peptides, these single-file ground tissue daughter cells were enlarged and elongated, but the number of cells in the file did not increase further (see Supplemental Figure 3 online), suggesting that the division of these cells was impaired.

What is the identity of this single file of cells located at the position of the ground tissue daughter cell? The *GFP* expression in the *J0571* marker line (Figures 6C and 6D, indicated by arrowheads) suggested that they still maintained their ground tissue identity, despite the fact that the division pattern was altered. Similarly, when *P_{SCR}:GFP* seedlings were treated with CLE peptides, *GFP* expression was observed in this single file, suggesting that it still maintains an endodermis and ground tissue identity (inset in Figure 6G, marked by arrowheads). Since the *SCR* promoter is not active in the cortex layer (Figure 6E), we could exclude the possibility that these cells obtained a full




Figure 4. Effect of Peptides on the Root Growth in Different Mutants.

All roots were measured after 2 weeks growing on media containing different peptides.

(A) The length of the primary roots of *clv1-1*, *clv2-1*, and *clv3-2* grown on media with different peptides. Note that the CLE peptides did not give a significant inhibition in *clv2-1* ($n = 10$).

(B) Number of meristematic cells in roots of *clv1-1*, *clv2-1*, and *clv3-2* grown on media with different peptides ($n = 10$). Data and error bars in **(A)** and **(B)** represent mean \pm SD.

(C) to **(F)** The morphology of roots of *Ler* and *clv* mutants treated with CLE19p, showing that *clv2-1* is not sensitive to the peptide. Bar in **(C)** = 50 μ m for **(C)** to **(F)**.

cortex identity. To further clarify the identity of these cells, the cortex-specific marker line $P_{CO_2}:YFP-H2B$ (Figure 6I) was examined upon peptide treatment. Interestingly, the single file of cells showed *YFP* expression (Figures 6J to 6M, indicated by arrowheads). We also noticed that the *YFP* expression was excluded from the ground tissue initial cells (Figures 6J and 6M, marked by arrows). As such, we concluded that the ground tissue daughter cells had obtained a cortex/endodermis double identity before the asymmetrical periclinal division occurs. This phenotype partially resembles the phenotype of the *scr* mutant in which $P_{CO_2}:YFP-H2B$ and *SCR* are expressed in the single-layered ground tissue (Heidstra et al., 2004).

Examination of the cell layers across the root meristem showed that, beside the changes in the ground tissue, all other cell layers were morphologically recognizable. However, in both the *J0571* and $P_{SCR}:GFP$ marker lines treated with CLE peptides for 4 d, we observed the expression of *GFP* in the pericycle layer (Figures 6C, 6D, 6G, and 6H, indicated by asterisks). This ectopic *GFP* expression was observed in >70% of roots treated with CLE19p and ~40% treated with CLV3p and CLE40p. The pericycle initial cells (located next to QC) and the adjacent few cells in the same layer often showed strong *GFP* signal, while *GFP* levels decreased in the above cells (Figures 6C, 6D, 6G, and 6H). In a few cases, noncontinuous *GFP* expression was




Figure 5. Heterologous Production of CLE19 Protein and Its Effect on Root Development.

(A) SDS-PAGE gel of thrombin-cleaved GST-Bn CLE19 and GST protein produced in *E. coli* with a mass of ~5 and 24 kD, respectively. The molecular mass (in kD) is shown at the right.

(B) Effect of purified the *Bn* CLE19 or GST proteins on root development. Wild-type (*Col-0*) seedlings were grown for 10 d on plates containing 10 μ M *Bn* CLE19 or GST. Data and error bars represent mean \pm SD ($n = 8$).

observed in this layer (Figure 6C). The number of pericycle cells with *GFP* expression at one focal plane ranged from 1 to 50. These cells were often distributed asymmetrically along the two sides of the root (Figures 6C, 6D, 6G, and 6H). Interestingly, in 8-d-old seedlings treated with CLE peptides, *GFP* expression in the pericycle cells was no longer detectable (see Supplemental Figure 3 online).

Similarly, upon the treatment with CLE peptides, we observed the expression of the cortex marker in the endodermis layer (Figures 6J to 6L, indicated by asterisks). Approximately 70% of roots treated with the CLE19p showed *YFP* expression in the endodermal layer, while the frequency of *YFP* expression was ~40% for CLV3p- and CLE40p-treated roots.

Penetration of CLE19p into Roots

To determine whether the CLE peptides enter the root or function on the root surface, a labeled CLE19p (R-CLE19p) was synthesized with a lissamine rhodamine fluorophore coupled to the N-terminal Lys. Wild-type (*Columbia-0* [*Col-0*]) seedlings were incubated with 10 μ M R-CLE19p, free lissamine rhodamine, or propidium iodide (PI) and analyzed by confocal microscopy.

PI is a vital dye that is widely used for staining of the cell wall. It can only enter the cells if the membrane integrity is disrupted (Robinson et al., 2002). R-CLE19p was able to enter the root. R-CLE19p fluorescence was observed in all cell layers of the roots within 1 min after peptide application. The fluorescence was located predominantly in the intercellular spaces (Figure 7A). After 4 min the signal was more intense, but still only located in the intercellular spaces (Figure 7D). Interestingly, the free lissamine rhodamine entered the roots with a lower efficiency, as indicated by the weak fluorescence (Figures 7B and 7E). PI entered the roots with the highest efficiency (Figures 7C and 7F), taking only 1 min to reach the saturation level (Figures 7C and 7D).

DISCUSSION

Previous studies suggested that CLV3 acts as a secreted protein that is required for the formation of the active CLV1/CLV2 signaling complex (Fletcher et al., 1999; Trotochaud et al., 1999; Rojo et al., 2002). However, the precise molecular identity of the functional CLV3 protein is still unknown. Several attempts have been made to identify the mature CLV3 protein using antibodies (Rojo et al., 2002), but so far only a CLV3-T7 fusion protein could be detected (Rojo et al., 2002). Recently, we and other labs have demonstrated that ectopic expression of *CLV3*, *CLE19*, *CLE40*, or *HgSYV46* leads to a consumption of the root meristem (Casamitjana-Martinez et al., 2003; Hobe et al., 2003; Fiers et al., 2004; Wang et al., 2005). Sequence alignment showed that the similarity between the putative proteins encoded by these genes was restricted to the CLE motif. Both *CLV3* and *CLE19* appeared to be not expressed in root meristems (Fletcher et al., 1999; Fiers et al., 2004), suggesting that the phenotype of ectopic expression represents a gain-of-function phenotype. We developed an in vitro root assay in *Arabidopsis* to examine how these proteins might generate the same phenotype in roots. The in vitro assay showed that three chemically synthesized 14-amino acid CLE peptides, corresponding to the conserved CLE motif of CLV3, CLE19, and CLE40, were able to mimic the overexpression phenotype when applied in vitro.

Three CLE peptides share only 50% (between CLV19p and CLE3p or CLE40p) to 64% (between CLV3p and CLE40p) sequence identity. The similar phenotype observed suggests that a common signaling pathway is involved and that the receptor(s) is able to recognize multiple ligands with certain

Table 1. Analysis of the F2 Plants with Long Roots from Crosses between *P*_{35S}:*Bn* CLE19 and *Ler* or Different *clv* Mutants (*clv1*, *clv2*, and *clv3*)

Cross Combinations	F2 Plants with Long Roots Examined	Plants with Multiple Carpels ^a		Plants with Wild-Type Carpels	
		Without <i>P</i> _{35S} : <i>Bn</i> CLE19	With <i>P</i> _{35S} : <i>Bn</i> CLE19	Without <i>P</i> _{35S} : <i>Bn</i> CLE19	With <i>P</i> _{35S} : <i>Bn</i> CLE19
<i>P</i> _{35S} : <i>Bn</i> CLE19 \times <i>Ler</i>	12	0	0	12	0
<i>P</i> _{35S} : <i>Bn</i> CLE19 \times <i>clv1-1</i>	12	2	0	10	0
<i>P</i> _{35S} : <i>Bn</i> CLE19 \times <i>clv2-1</i> ^b	60	5	11	33	0
<i>P</i> _{35S} : <i>Bn</i> CLE19 \times <i>clv3-2</i>	13	3	0	10	0

^a Multiple carpels in the pistil is the typical phenotype for all *clv* mutants.

^b Two independent experiments were performed, and the data were combined in sum.




Figure 6. Effects of CLE Peptides on the Cell Identity of Roots.

The ground tissue initial cells are marked by arrows, and the ground tissue daughter cells, including cells at this position, are labeled with arrowheads. Abnormal expression of *GFP* or *YFP* in other cell layers is marked by asterisks.

(A) to (D) Confocal analysis after 4-d treatment of the roots of *J0571*. Note that treatments with CLV3p (C) or CLE19p (D) lead to a delayed separation of cortex and endodermis and expression of *GFP* in pericycle cells. No peptide (A), 10 μ M CLVm (B), CLV3p (C), and CLE19p (D).

(E) to (H) Confocal analyses of *P_{SCR}::GFP* seedlings after 4-d treatments. No peptide (E), 10 μ M CLV3s (F), 10 μ M CLV3p (G), and 10 μ M CLE19p (H). The inset in (G) shows the delayed separation of the cortex and endodermis cells (with *GFP* expression).

(I) to (M) *P_{CO2}::YFP-H2B* marker line treated with different peptides. Note the *YFP* expression in the ground tissue daughter cells and the pericycle cells, but not in ground tissue initial cells.

(I) No peptide, 4 d after germination. The inset shows the absence of *YFP* expression in the QC, ground tissue initial, and ground daughter cells.

(J) 10 μ M CLE19p, 4 d after treatment.

(K) 10 μ M CLE40p, 5 d after treatment.

(L) and (M) 10 μ M CLE19p, 5 d after treatment.

Bars = 30 μ m.




Figure 7. Penetration of Fluorescence-Labeled CLE19p in Roots.

The roots of *Arabidopsis* (*Col-0*) were incubated for 1 min (**[A]** to **[C]**) or 4 min (**[D]** to **[F]**) with 10 μ M R-CLE19p (**[A]** and **[D]**), 10 μ M lissamine rhodamine (**[B]** and **[E]**), or 10 μ M PI (**[C]** and **[F]**).

Bar in **(F)** = 30 μ m for **(A)** to **(F)**.

sequence variations. It is possible that in the endogenous situation the specificity of *CLV3*, *CLE19*, and *CLE40* is defined by the regulation of expression. The control peptides, including *CLV3s*, *CLV3m*, *CLV3t*, and *CLE19t*, did not affect root development. *CLE5p*, a peptide corresponding to a distant member of the *CLE* genes expressed in roots, was also unable to trigger the consumption of the root meristem. These observations suggest that the *CLE* peptides act in a sequence-specific manner.

Experiments with a labeled peptide showed that this peptide is able to efficiently enter the intercellular spaces of the root. The fluorescence signal was located predominately in the cell wall, suggesting that active transport is not likely to be involved. Although the labeled peptide penetrated into roots faster than the free fluorescence dye, PI, which cannot enter the intact plasma membrane (Robinson et al., 2002), enters into the roots even faster. The differences in penetration efficiency can be explained either by the differences in hydrophobicity or molecular masses.

Using a QC marker and the *DR5:GUS* line, we observed that application of the *CLE* peptides resulted in a consumption of the root meristem without directly interfering with the QC function or auxin distribution in roots. This conclusion is further supported by the observation that the ground tissue initial cells located next to the QC and tightly regulated by the QC (van den Berg et al., 1995, 1997) were not affected. These data are in agreement with previous observations that overexpression of *CLV3*, *CLE19*, or *CLE40* in *Arabidopsis* causes a termination of root meristems without acting primarily on the QC (Casamitjana-Martinez et al., 2003; Hobe et al., 2003; Fiers et al., 2004).

To determine if the sequence before the *CLE* motif could influence the activity, *Bn CLE19* protein (with its N-terminal secretion signal removed) was produced in *E. coli* and tested in the root assay. The full-length *CLE19* protein appeared to be functional in *Arabidopsis* in a similar fashion as *CLE19p*, sug-

gesting that the extra sequence before the *CLE* motif does not affect its function. This is consistent with what is known for other peptide ligands in plants, such as phytosulphokine and flagellin (Felix et al., 1999; Yang et al., 1999). Although the pre-proteins for phytosulphokine in rice (*Oryza sativa*), *Arabidopsis*, and asparagus (*Asparagus officinalis*) are different in sequence, the functional peptides are believed to be the same (Yang et al., 1999). Since no endogenous *CLE* ligand has been identified yet, we cannot exclude the possibility that the endogenous peptides are longer or shorter than the *CLE* peptides tested.

For the identification of signaling components involved in perception of these *CLE* peptides, we examined whether components of the *Arabidopsis* *CLV* signaling pathways, specifically *CLV1*, *CLV2*, and *CLV3*, are involved in *CLE* peptide signaling. Interestingly, the *clv2* mutant was insensitive to *CLE* peptide treatment, suggesting that *CLV2* is functionally involved in perception of the *CLE* peptides in roots. Genetic analysis demonstrated that *CLV2* is also involved in the perception of the *P_{35S}:Bn CLE19* transgene, suggesting that the same machinery is involved in the perception of the transgenic *Bn CLE19* and *CLE* peptides. Among these *CLV* genes, *CLV2* is the only one expressed in *Arabidopsis* roots (Birbaum et al., 2003). It seems that *CLV2* not only participates in the *CLV1/CLV2* receptor complex to transmit the *CLV3* signal in the SAM (Jeong et al., 1999), but also functions in an unidentified *CLV1*-like receptor kinase complex in roots to perceive the *CLE* peptides. The fact that *clv2* does not show any root phenotype suggests that either *CLV2* is redundant in roots while the redundant protein(s) cannot sense the *CLE* peptides, or *CLV2* is not functional in roots and the *CLE* peptides act ectopically to generate a gain-of-function phenotype.

Three cell layer identity markers were examined to further study the effects of the peptide treatment. First, we observed that the treatments led to the accumulation of multiple cells at the

position of the ground tissue daughter cell. This is striking since in the wild-type situation an asymmetrical division always occurs immediately after the ground tissue daughter cell is produced, to form an outer cortex cell and an inner endodermis cell (van den Berg et al., 1995, 1997). This tightly regulated cell division pattern is controlled by SCR in a cell autonomous manner, and the function of SHORT ROOT (SHR) is required for the activation of SCR, which in turn induces the rotation of the division plane (Helariutta et al., 2000; Heidstra et al., 2004).

Detailed analyses showed that two ground tissue markers (*J0571* and *P_{SCR}::GFP*) were expressed in the row of ground tissue daughter cells, suggesting that these cells still maintain their ground tissue identity. In the absence of CLE peptide application, *YFP* expression in the cortex-specific marker line (*P_{CO2}::YFP-H2B*) was excluded from the ground tissue initials and the ground tissue daughter cells (see inset in Figure 6I; Heidstra et al., 2004). However, upon treatment with CLE peptides, *YFP* expression was observed in the single file of cells located at the position of the ground tissue daughter cell (Figures 6J to 6M), indicating that these cells obtained a cortex identity prior to the asymmetrical division. Since *SCR* was expressed in these cells, we conclude that they have obtained a cortex/endodermis double identity. This single cell layer of ground tissue with a double identity is similar to that in *scr* but different from the *shr* mutant, in which the single layer has only cortex identity (Di Laurenzio et al., 1996; Helariutta et al., 2000). Interestingly, both *scr* and *shr* mutants have a short root phenotype in which the root meristem gradually differentiates after germination, suggesting that interrupted cell layer formation could have a direct effect on the maintenance of meristematic activity in roots. Whether CLE peptides interfere with SCR signaling components remains to be studied. Additionally, it is also not known if the CLE peptides first inhibit the division of the ground tissue daughter cells, which then leads to the differentiation of these cells, or vice versa.

Another striking effect observed after treatment with CLE peptides is the confusion in cell layer identity. The ground tissue genes (*P_{SCR}::GFP* and *J0571*) were transiently activated in the pericycle layer, and a cortex gene was active in the endodermal layer. The common feature is that the inner cell layers take the identity of the outer cell layer. Such a confusion or misspecification of layer identity could be a consequence of the failure of cells to sense the presence of neighboring cells. It is well known that positional signals are used by plant cells to define their identity (van den Berg et al., 1995, 1997). As such, we believe that these CLE peptides and the *CLV3*, *CLE19*, and *CLE40* transgenes act in a dominant-negative fashion to interact with or saturate an unknown cell identity-maintaining *CLV2* receptor complex. This in turn blocks intercellular communication among different cells and cell layers in the root, which leads to consumption of the root meristem.

METHODS

Plant Growth Conditions and Plant Strains

The marker lines *J0571* (made by Jim Haseloff) and *P_{SCR}::GFP* (Wysocka-Diller et al., 2000) and the mutants of *clv1-1*, *clv2-1*, and *clv3-2* were provided by the Nottingham Arabidopsis Stock Centre. *P_{CO2}::YFP-H2B*,

DR5::GUS, and *QC25* were used as previously described (Casamitjana-Martinez et al., 2003; Heidstra et al., 2004).

Root Assay

Seeds were gas-sterilized in a desiccator for 1 h with 100 mL of bleach (4% NaClO) mixed with 3 mL of HCl in a beaker. For peptide treatments, the sterilized seeds were plated at a distance of 0.5 cm on media containing different concentrations of peptides, half-strength Murashige and Skooge microelements and macroelements (Duchefa), 1% (w/v) sucrose, and 0.5 g/L MES, pH 5.8, with 1.5% (w/v) agar. Peptides and proteins were added to the sterilized media before the medium was solidified. Plates were first incubated at 4°C in the dark for 2 d and then transferred to a room with a temperature of 23°C, 16 h light per day, and cultured nearly vertically. The root length was measured from the base of the hypocotyl to the tip of the primary root. GUS analysis was as described by Fiers et al. (2004), and GFP analyses were performed after 4, 6, and 8 d of growth. Peptides were ordered from Mimopopes with a purity of >70% and dissolved in a filter-sterilized sodium phosphate buffer (50 mM, pH 6).

To examine the penetration of the peptide into roots, *CLE19p* with a lissamine rhodamine fluorophore coupled to the N-terminal Lys (*R-CLE19p*) was ordered (Mimopopes). Four-day-old seedlings of *Arabidopsis thaliana* (*Col-0*) were incubated in either 10 μM *R-CLE19p*, 10 μM lissamine rhodamine, or 10 μg/mL PI by incubating the roots for different time periods, after which they were examined using confocal microscopy.

Microscopy

Roots and dissected SAMs were cleared following the protocol of Sabatini et al. (1999) and analyzed using a Nikon microscope equipped with Nomarski optics. For confocal microscopy, roots were counterstained with 10 μg/mL PI (Sigma-Aldrich) and analyzed with a Leica SP2 inverted confocal microscope following the protocol of Heidstra et al. (2004).

Protein Purification

Two primers were designed (5'-TATGGATCCGCTTCATTTCGGAGT-TTG-3' and 5'-ATACTCGAGTTACCTGTTGTGAAGTGA-3') to amplify *Bn CLE19* (without the signal sequence) from the cDNA. The PCR fragment was digested with *Bam*HI and *Hind*III (Invitrogen) and cloned into the *pGEX4T-2* vector, and the fusion protein was purified as described by the manufacturer (Amersham Biosciences). The *Bn CLE19* and the GST control were tested using SDS-PAGE and quantified with Coomassie Brilliant Blue plus protein assay reagent (Pierce).

Genetic Analysis

The F1 of the crosses between a *P_{35S}::Bn CLE19* plant and individual *clv* mutants (*clv1-1*, *clv2-1*, and *clv3-2*) or the wild type (*Ler*) were examined using the root assay mentioned above for a normal or short root phenotype. F2 seeds were harvested from plants with short roots and assayed again for individuals with a long or short root phenotype. These two groups were transplanted separately to soil and checked for the presence of the transgene with a PCR for the *NPTII* gene (5'-TGGGCA-CAACAGACAATCGGCTGC-3' and 5'-TGCGAATCGGGAGCGCGGATA-CCG-3'). Homozygous *clv* mutants were identified from each group by their carpel phenotype.

Accession Numbers

Sequence data from this article can be found in the EMBL/GenBank data libraries under accession numbers AT2G27250 (*CLV3*), AT2G31082

(*CLE5*), At3g24225 (*CLE19*), AF343656 (*Bn CLE19*), and AT5G12990 (*CLE40*).

ACKNOWLEDGMENTS

We thank Kim Boutilier and Gerco Angenent for critical reading of the manuscript. We also thank the Wageningen University MicroSpectroscopy Centre for technical support, the Wisconsin Arabidopsis Knockout Center, the Salk Institute, the Torrey Mesa Research Institute (Syngenta), and the Nottingham Arabidopsis Stock Centre for providing the T-DNA insertion lines. This work was financially supported in part by the Dutch Ministry of Agriculture, Nature Management, and Fisheries (DWK281/392) and by the Centre for BioSystems Genomics, which is part of the Netherlands Genomics Initiative/Netherlands Organization for Scientific Research. E.G. was supported by the European Union CROP-STRESS project (QLAM 2001-00424).

Received May 4, 2005; revised July 1, 2005; accepted July 1, 2005; published July 29, 2005.

REFERENCES

- Alberts, B., Bray, D., Lewis, J., Raff, M., Roberts, K., and Watson, J.D. (1994). *Molecular Biology of the Cell*. (New York: Garland).
- Birnbaum, K., Shasha, D.E., Wang, J.Y., Jung, J.W., Lambert, G.M., Galbraith, D.W., and Benfey, P.N. (2003). A gene expression map of the Arabidopsis root. *Science* **302**, 1956–1960.
- Bonello, J.F., Sevilla-Lecoq, S., Berne, A., Risueno, M.C., Dumas, C., and Rogowsky, P.M. (2002). ESR proteins are secreted by the cells of the embryo surrounding region. *J. Exp. Bot.* **53**, 1559–1568.
- Brand, U., Fletcher, J.C., Hobe, M., Meyerowitz, E.M., and Simon, R. (2000). Dependence of stem cell fate in Arabidopsis on a feedback loop regulated by *CLV3* activity. *Science* **289**, 617–619.
- Casamitjana-Martinez, E., Hofhuis, H.F., Xu, J., Liu, C.M., Heidstra, R., and Scheres, B. (2003). Root-specific *CLE19* over-expression and the *sol1* and *sol2* suppressors implicate a CLV-like pathway in the control of Arabidopsis root meristem maintenances. *Curr. Biol.* **13**, 1435–1441.
- Clark, S.E., Williams, R.W., and Meyerowitz, E.M. (1997). The *CLAVATA1* gene encodes a putative receptor kinase that controls shoot and floral meristem size in Arabidopsis. *Cell* **89**, 575–585.
- Cock, J.M., and McCormick, S. (2001). A large family of genes that share homology with *CLAVATA3*. *Plant Physiol.* **126**, 939–942.
- Di Laurenzio, L., Wysocka-Diller, J., Malamy, J.E., Pysh, L., Helariutta, Y., Freshour, G., Hahn, M.G., Feldmann, K.A., and Benfey, P.N. (1996). The *SCARECROW* gene regulates an asymmetric cell division that is essential for generating the radial organization of the Arabidopsis root. *Cell* **86**, 423–433.
- Felix, G., Duran, J.D., Volko, S., and Boller, T. (1999). Plants have a sensitive perception system for the most conserved domain of bacterial flagellin. *Plant J.* **18**, 265–276.
- Fiers, M., Hause, G., Boutilier, K., Casamitjana-Martinez, E., Weijers, D., Offringa, D., van der Geest, L., van Lookeren Campagne, M., and Liu, C.M. (2004). Mis-expression of the *CLV3/ESR*-like gene *CLE19* in Arabidopsis leads to a consumption of root meristem. *Gene* **327**, 37–49.
- Fletcher, J.C., Brand, U., Running, M.P., Simon, R., and Meyerowitz, E.M. (1999). Signaling of cell fate decisions by *CLAVATA3* in Arabidopsis shoot meristems. *Science* **283**, 1911–1914.
- Heidstra, R., Welch, D., and Scheres, B. (2004). Mosaic analyses using marked activation and deletion clones dissect Arabidopsis SCARECROW action in asymmetric cell division. *Genes Dev.* **18**, 1964–1969.
- Helariutta, Y., Fukaki, H., Wysocka-Diller, J., Nakajima, K., Jung, J., Sena, G., Hauser, M.T., and Benfey, P.N. (2000). The *SHORT-ROOT* gene controls radial patterning of the Arabidopsis root through radial signaling. *Cell* **101**, 555–567.
- Hobe, M., Muller, R., Grunewald, M., Brand, U., and Simon, R. (2003). Loss of *CLE40*, a protein functionally equivalent to the stem cell restricting signal *CLV3*, enhances root waving in Arabidopsis. *Dev. Genes Evol.* **213**, 371–381.
- Jeong, S., Trotochaud, A.E., and Clark, E. (1999). The Arabidopsis *CLAVATA2* gene encodes a receptor-like protein required for the stability of the *CLAVATA1* receptor-like kinase. *Plant Cell* **11**, 1925–1933.
- Kende, H., and Zeevaert, J.A.D. (1997). The five “classical” plant hormones. *Plant Cell* **9**, 1197–1210.
- Kim, S., Mollet, J.C., Dong, J., Zhang, K., Park, S.Y., and Lord, E.M. (2003). Chemocyanin, a small basic protein from the lily stigma, induces pollen tube chemotropism. *Proc. Natl. Acad. Sci. USA* **100**, 16125–16130.
- Laux, T. (2003). The stem cell concept in plants: A matter of debate. *Cell* **113**, 281–283.
- Laux, T., Mayer, K.F., Berger, J., and Jürgens, G. (1996). The *WUSCHEL* gene is required for shoot and floral meristem integrity in Arabidopsis. *Development* **122**, 87–96.
- Lindsey, K., Casson, S., and Chille, P. (2002). Peptides: New signaling molecules in plants. *Trends Plant Sci.* **7**, 78–83.
- Mandave, N.B. (1988). Plant growth-promoting brassinosteroids. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **39**, 23–52.
- Matsubayashi, Y., and Sakagami, Y. (1996). Phytosulfokine, sulfated peptides that induce the proliferation of single mesophyll cells of *Asparagus officinalis* L. *Proc. Natl. Acad. Sci. USA* **93**, 7623–7627.
- Pearce, G., Strydom, D., Johnson, S., and Ryan, C.A. (1991). A polypeptide from tomato leaves induces wound-inducible inhibitor proteins. *Science* **253**, 895–898.
- Robinson, J.P., Darzynkiewicz, Z., Dean, P.N., Hibbs, A.R., Orfao, A., Rabinovitch, P.S., and Wheelless, L.L. (2002). *Current Protocols in Cytometry*. (New York: John Wiley & Sons).
- Rojo, E., Sharma, V.K., Kovaleva, V., Raikhel, N.V., and Fletcher, J.C. (2002). *CLV3* is localized to the extracellular space, where it activates the Arabidopsis *CLAVATA* stem cell signaling pathway. *Plant Cell* **14**, 969–977.
- Ryan, C.A., Pearce, G., Scheer, J., and Moura, D.S. (2002). Polypeptide hormones. *Plant Cell* **14** (suppl.), S251–S264.
- Sabatini, S., Beis, D., Wolkenfelt, H., Murfett, J., Guilfoyle, T., Malamy, J., Benfey, P., Leyser, O., Bechtold, N., and Weisbeek, P. (1999). An auxin-dependent distal organizer of pattern and polarity in the Arabidopsis root. *Cell* **99**, 463–472.
- Schoof, H., Lenhard, M., Haecker, A., Mayer, K.F.X., Jürgens, G., and Laux, T. (2000). The stem cell population of Arabidopsis shoot meristems is maintained by a regulatory loop between the *CLAVATA* and *WUSCHEL* genes. *Cell* **100**, 635–644.
- Schopfer, C.R., Nasrallah, M.E., and Nasrallah, J.B. (1999). The male determinant of self-incompatibility in *Brassica*. *Science* **286**, 1697–1700.
- Sharma, V.K., Ramirez, J., and Fletcher, J.C. (2003). The Arabidopsis *CLV3*-like (*CLE*) genes are expressed in diverse tissues and encode secreted proteins. *Plant Mol. Biol.* **51**, 415–425.
- Trotochaud, A.E., Hao, T., Wu, G., Yang, Z., and Clark, S.E. (1999). The *CLAVATA1* receptor-like kinase requires *CLAVATA3* for its

- assembly into a signalling complex that includes KAPP and a Rho-related protein. *Plant Cell* **11**, 393–405.
- van den Berg, C., Willemsen, V., Hage, W., Weisbeek, P., and Scheres, B.** (1995). Cell fate in the *Arabidopsis* root meristem determined by directional signalling. *Nature* **378**, 62–65.
- van den Berg, C., Willemsen, V., Hendriks, G., Weisbeek, P., and Scheres, B.** (1997). Short-range control of cell differentiation in the *Arabidopsis* root meristem. *Nature* **390**, 287–289.
- Wang, X., Mitchum, M.G., Gao, B., Li, C., Diab, H., Baum, T.J., Hussey, R.S., and Davis, E.L.** (2005). A parasitism gene from a plant-parasitic nematode with function similar to *CLAVATA3/ESR (CLE)* of *Arabidopsis thaliana*. *Mol. Plant Pathol.* **6**, 187–191.
- Wysocka-Diller, J.W., Helariutta, Y., Fukaki, H., Malamy, J.E., and Benfey, P.N.** (2000). Molecular analysis of SCARECROW function reveals a radial patterning mechanism common to root and shoot. *Development* **127**, 595–603.
- Yang, H., Matsubayashi, Y., Nakamura, K., and Sakagami, Y.** (1999). *Oryza sativa* PSK gene encodes a precursor of phytosulfokine- α , a sulfated peptide growth factor found in plants. *Proc. Natl. Acad. Sci. USA* **96**, 13560–13565.