

Implications for policy makers and future research
Nurse led hospital at home or early discharge schemes for patients with COPD should be prioritised over the type of nurse led models of chronic disease management that have been studied to date. There is little evidence available at present to support the models that have been evaluated. Existing services providing this sort of care should be robustly evaluated against the aims of the particular service.

The evidence around long term or intensive case management and hospital readmission is currently equivocal and requires further study. The potential benefits of schemes for chronic disease management in patients with COPD receiving long term oxygen therapy should also be explored further. Several potentially important outcomes have not been evaluated, including patients' satisfaction, self management, patients' coping and adherence, smoking cessation, and the effects on carers.

We thank Brian Schirn, Gene Feder, Martin Underwood, Gill Foster, Jill Goddard, Hazel Kilvington, Marie Montague, Sarah Cotter, Melanie Wakelin, Anne Spencer, Linda Stephenson, Lynette Edwards, staff at the Royal College of Nursing, members of the British Lung Foundation Breathe Easy groups, the International Primary Care Respiratory Group, and the Association of Respiratory Nurses.

Contributors: See bmj.com

Funding: NHS research and development service delivery and organisation programme. SJCT is funded by a Department of Health career scientist award in public health.

Competing interests: None declared.

Ethical approval: Not required.

- Ram FS, Wedzicha JA, Wright J, Greenstone M. Hospital at home for patients with acute exacerbations of chronic obstructive pulmonary disease: systematic review of the evidence. *BMJ* 2004;329:315-9.
- Candy B, Taylor SJ, Griffiths CJ, Ramsay J, Wedzicha JA, Schirn B, et al. *Evaluating the effectiveness of innovations involving nurses for people in the community with chronic obstructive airways disease*. London: Centre of General Practice and Primary Care, Institute of Community Health Science, Barts and The London, Queen Mary's School of Medicine and Dentistry, 2004.

- Verhagen AP, de Vet HC, de Bie RA, Kessels AG, Boers M, Bouter LM, et al. The Delphi list: a criteria list for quality assessment of randomized clinical trials for conducting systematic reviews developed by Delphi consensus. *J Clin Epidemiol* 1998;51:1235-41.
- Jadad AR, Moore RA, Carroll D. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996; 17:1-12.
- Oxford Centre for Evidence-Based Medicine. *Levels of evidence and grades of recommendation*. www.cebm.net/levels_of_evidence.asp#levels (accessed Mar 2005).
- Cockcroft A, Bagnall P, Heslop A, Andersson N, Heaton R, Batstone J, et al. Controlled trial of respiratory health worker visiting patients with chronic respiratory disability. *BMJ* 1987;294:225-8.
- Bergner M, Hudson LD, Conrad DA, Patmont CM, McDonald CJ, Perrin EB, et al. The cost and efficacy of home care for patients with chronic lung disease. *Med Care* 1988;26:566-79.
- Littlejohns P, Baveystock CM, Parnell H, Jones PW. Randomised controlled trial of the effectiveness of a respiratory health worker in reducing impairment, disability, and handicap due to chronic airflow limitation. *Thorax* 1991;46:559-64.
- Smith BJ, Appleton SL, Bennett PW, Roberts GC, Del Fante P, Adams R, et al. The effect of a respiratory home nurse intervention in patients with chronic obstructive pulmonary disease (COPD). *Aust N Z J Med* 1999;29:718-25.
- Farrero E, Escarrabill J, Prats E, Maderal M, Manresa, F. Impact of a hospital-based home-care program on the management of COPD patients receiving long-term oxygen therapy. *Chest* 2001;119:364-9.
- Egan E, Clavarino A, Burrigide L, Teuwen M, White, E. A randomized control trial of nursing-based case management for patients with chronic obstructive pulmonary disease. *Lippincott's Case Manag* 2002;7:170-9.
- Hermiz O, Comino E, Marks G, Daffurn K, Wilson S, Harris, M. Randomised controlled trial of home based care of patients with chronic obstructive pulmonary disease. *BMJ* 2003;325:938-40.
- Bourbeau J, Julien M, Maltais F, Rouleau M, Beaupre A, Begin R, et al. Reduction of hospital utilization in patients with chronic obstructive pulmonary disease: a disease-specific self-management intervention. *Arch Intern Med* 2003;163:585-91.
- Monnikhof E, van der Valk P, van der Palen J, van Herwaarden C, Zielhuis G. Effects of a comprehensive self-management programme in patients with chronic obstructive pulmonary disease. *Eur Respir J* 2003;22:815-20.
- Department of Health. *Improving chronic disease management*. www.dh.gov.uk/assetRoot/04/07/52/13/04075213.pdf (accessed Oct 2004).
- Weingarten SR, Henning JM, Badamgarav E, Knight K, Hasselblad V, Gano A, et al. Interventions used in disease management programmes for patients with chronic illness—which ones work? Meta-analysis of published reports. *BMJ* 2002;325:925-8.
- Feachem RGA, Sekhri NK, White KL. Getting more for their dollar: a comparison of the NHS with California's Kaiser Permanente. *BMJ* 2002;324:135-43.
- Elphick HE, Tan A, Ashby D, Smyth RL. Systematic reviews and lifelong diseases. *BMJ* 2002;325:381-4.

(Accepted 27 May 2005)

doi 10.1136/bmj.38512.664167.8F

The partial smoking ban in licensed establishments and health inequalities in England: modelling study

Alan A Woodall, Emma J Sandbach, Catherine M Woodward, Paul Aveyard, Graham Merrington

Telford and Wrekin Primary Care Trust, Telford TF1 5RY

Alan A Woodall
specialist registrar in public health

Emma J Sandbach
health intelligence analyst

Catherine M Woodward
director of public health

continued over

The UK government's white paper *Choosing Health* proposes prohibiting smoking in public places in England, but exempts public houses (pubs) not serving catered food and licensed establishments that require membership.¹ However, passive inhalation of smoke at work may cause 600 deaths per year in the United Kingdom and increases morbidity and mortality among bar workers.² Furthermore, people attempting to quit smoking find that socialising with other smokers makes quitting difficult, and lapses in quitting are more likely in premises where smoking is permitted.³

Concerns exist that exempt establishments are located primarily in deprived areas with the highest smoking prevalence and that a partial ban worsens health inequalities.⁴ We examined if exempt establishments were located predominantly in deprived areas in the borough of Telford and Wrekin.

Methods and results

We determined the catering status of pubs from regularly updated records of local authority licensing and environmental health. These allowed us to identify premises preparing catered food, those serving manufactured snacks, and those not serving food. *Choosing Health* proposes prohibiting smoking only in pubs serving catered food; all others can be exempt. All licensed members' clubs may choose exemption, which we assumed for this study.

 A statistical appendix in on bmj.com

This article was first posted on bmj.com on 19 August 2005: <http://bmj.com/cgi/doi/10.1136/bmj.38576.467292.EB>

Modelled and observed percentage of pubs and all licensed establishments exempt from the smoking ban proposed by *Choosing Health* by socioeconomic status of an area

Fifth of multiple deprivation index 2004	Median super-output area score	Predicted % (95% CI) in England		Observed % (95% CI) in Telford and Wrekin	
		Pubs*	Pubs, nightclubs, or licensed establishments requiring membership†	Pubs‡	Pubs, nightclubs, or licensed establishments requiring membership§
1 (least deprived)	5.74	26.9 (23.7 to 29.8)	39.3 (35.6 to 43.1)	31 (9 to 61)	47 (23 to 72)
2	10.96	31.7 (25.2 to 38.0)	44.7 (37.5 to 52.2)	21 (10 to 35)	30 (18 to 45)
3	17.02	37.9 (27.0 to 48.5)	51.2 (39.7 to 62.6)	56 (38 to 73)	67 (52 to 80)
4	26.61	48.4 (30.0 to 64.9)	61.3 (43.3 to 76.7)	51 (38 to 64)	63 (52 to 73)
5 (most deprived)	45.22	68.5 (36.4 to 87.3)	77.9 (50.5 to 92.4)	55 (32 to 77)	69 (49 to 85)

* χ^2 test for trend 9.10, df=1, P=0.003. † χ^2 test for trend 10.73, df=1, P<0.001. ‡ χ^2 test for trend 9.44, df=1, P=0.002. § χ^2 test for trend 11.93, df=1, P<0.001.

Postcode mapping of establishments to each super-output area allowed extraction of scores on the index of multiple deprivation scores by using the software MapInfo, version 8.0 (MapInfo Limited, Windsor, UK). The super-output area is a contiguous aggregate of census output areas containing some 1500 people.

We used logistic regression to examine the relation between the proportion of exempt premises and deprivation score for pubs and licensed members' establishments in Telford and Wrekin borough. With the regression coefficients generated, we used median deprivation scores for each English fifth to estimate the probability of a typical English establishment being exempt.

Out of 174 pubs in the borough, 99 (57%) served catered food. Hence, 75 (43%) would be exempt from smoking prohibition. Including all licensed members' establishments in the model showed that 127 (56%) would be exempt. The model predicted that two thirds of English pubs in deprived areas would be exempt, whereas only a quarter would be exempt in affluent areas. Including members' clubs in the model showed that two fifths of establishments in affluent areas and four fifths of establishments in deprived areas would be exempt (table). A significant linear trend between deprivation and probability of exemption existed in all analyses.

Comment

Prohibiting smoking only in pubs that serve catered food and allowing exemptions for other licensed drinking establishments may worsen health inequalities.

What is already known on this topic

Passive smoking is a serious risk to health

Smoking in public places is banned completely in many countries, but the current proposals for England will allow for some establishments licensed to sell alcohol to be exempt from a smoking ban

What this study adds

Most licensed establishments in the poorest areas would be exempt from the workplace ban on smoking, while most in the more affluent areas would be subject to the ban. This is likely to worsen socioeconomic inequalities in health and smoking prevalence

Choosing Health estimates that only 10-30% of pubs could be smoking (p100), but our data indicate that the proportion of exempt pubs is higher (43%). This is a small study in one borough, so care must be taken extrapolating the findings. However, Telford and Wrekin is similar to England in terms of demographics and socioeconomic profile. Higher exemption rates were also observed in a survey of 29 local authorities, but no data on deprivation were obtained.⁵ Our results show that people in deprived areas are more likely to live near licensed establishments exempt from legislation to protect them against smoking. It is possible that people from deprived neighbourhoods may visit establishments in affluent areas, whereas those living in affluent neighbourhoods make the reverse journey. It is more likely that the poorest people with the worst health and highest smoking prevalence would be those most likely to be harmed by passive smoking either working in pubs or as customers, and would be those most likely to have their attempt to stop smoking undermined. We urge the UK government to ban smoking in all enclosed public places, similar to the ban proposed in Scotland and enacted in Ireland, to prevent worsening health inequalities.

We thank Andrea Roalfe for statistical advice. AAW dedicates this work to the memory of Thomas Edward Kennedy.

Contributors: AAW initiated the study. AAW, PA, and CW contributed to the design of the study. GM provided data from the local authority and assisted with study design. AAW, PA, and ES carried out the statistical analyses. AAW wrote the paper. All authors contributed to interpretation of findings and contributed critical comments to the paper. AAW is the principal guarantor of the paper.

Funding: None.

Competing interests: None declared.

Ethical approval: Ethical approval was not sought as the data is in the public domain and no issues regarding breach of confidentiality were identified.

- 1 Department of Health. *Choosing health: making healthy choices easier*. London: Stationery Office, 2004.
- 2 Jamrozik K. Estimate of deaths attributable to passive smoking among UK adults: database analysis. *BMJ* 2005;330:812.
- 3 Wiltshire S, Bancroft A, Parry O, Amos A. "I came back here and started smoking again": perceptions and experiences of quitting among disadvantaged smokers. *Health Educ Res* 2003;18:292-303.
- 4 Raine R, Walt G, Basnett I. The white paper on public health *BMJ* 2004;329:1247.
- 5 British Medical Association. *Booze, fags and food*. London: BMA 2005. www.bma.org.uk/ap.nsf/Content/boozefagsandfood (accessed 22 Jun 2005).

(Accepted 22 July 2005)

doi 10.1136/bmj.38576.467292.EB

Department of Primary Care and General Practice, University of Birmingham, Edgbaston, Birmingham B15 2TT

Paul Aveyard
senior lecturer

Borough of Telford and Wrekin Council, Social Care, Telford TF3 4LE

Graham Merrington
environmental health manager

Correspondence to: A A Woodall (alan.woodall@telfordpct.nhs.uk)