

Environmental Occurrence of *Madurella mycetomatis*, the Major Agent of Human Eumycetoma in Sudan

Abdalla Ahmed,^{1,2} Daniel Adelmann,³ Ahmed Fahal,⁴ Henri Verbrugh,² Alex van Belkum,^{2*} and Sybren de Hoog³

Institute for Endemic Diseases and Faculty of Medical Laboratory Sciences,¹ and Department of Surgery, Faculty of Medicine,⁴ University of Khartoum, Khartoum, Sudan, and Department of Medical Microbiology & Infectious Diseases, Erasmus University Medical Center Rotterdam (EMCR), Dr. Molewaterplein 40, 3015 GD Rotterdam,² and Centraalbureau voor Schimmelcultures, Uppsalaalaan 8, 3584 CT Utrecht,³ The Netherlands

Received 13 July 2001/Returned for modification 30 September 2001/Accepted 14 December 2001

***Madurella mycetomatis* is the main causative agent of human eumycetoma, a severe debilitating disease endemic in Sudan. It has been suggested that eumycetoma has a soil-borne or thorn prick-mediated origin. For this reason, efforts were undertaken to culture *M. mycetomatis* from soil samples ($n = 43$) and thorn collections ($n = 35$) derived from areas in which it is endemic. However, ribosomal sequencing data revealed that the black fungi obtained all belonged to other fungal species. In addition, we performed PCR-mediated detection followed by restriction fragment length polymorphism (RFLP) analysis for the identification of *M. mycetomatis* DNA from the environmental samples as well as biopsies from patients with mycetoma. In the case of the Sudanese soil samples, 17 out of 74 (23%) samples were positive for *M. mycetomatis* DNA. Among the thorn collections, 1 out of 22 (5%) was positive in the PCR. All PCR RFLP patterns clearly indicated the presence of *M. mycetomatis*. In contrast, 15 Dutch and English control soil samples were all negative. Clinically and environmentally obtained fungal PCR products share the same PCR RFLP patterns, suggesting identity, at least at the species level. These observations support the hypothesis that eumycetoma is primarily environmentally acquired and suggest that *M. mycetomatis* needs special conditions for growth, as direct isolation from the environment seems to be impossible.**

Eumycetoma is a subcutaneous fungal infection in which the etiologic agent occurs in the form of more or less compact mycelial grains. The immune response is a granulomatous foreign body reaction characterized by impressive local necrosis and the formation of extensive fistula (8, 9, 18). Areas where eumycetoma is endemic are arid and hot climate zones with limited rainfall. A genuine “mycetoma belt” is found in tropical and subtropical regions, with foci in southern India, Somalia, Senegal, Argentina, and Sudan (12, 17). Cases are particularly common in Sudan, where, for example, Abbott (1) reported 1,231 cases in a 2.5-year period. The main causative agent of eumycetoma in Sudan is *Madurella mycetomatis* (4, 15). The disorder is supposed to originate from traumatic inoculation of plant material or soil contaminated by these fungi. Abbott (1) noted a history of thorn pricks in 10 out of 62 cases of mycetoma, and in two of these, there were actually thorns found embedded in tissue. Basset et al. (3) reported two cases with thorns inside lesions from which only non-*Madurella* fungi could be isolated. It is unlikely that *M. mycetomatis* represents a primary pathogen for humans. Rather, the fungi involved probably are saprophytes, which are able to tolerate uncharacteristic conditions when inoculated by coincidence. However, direct evidence for this hypothesis is still lacking. Thirumalachar and Padhye (22) were barely able to recover *M.*

mycetomatis from soil. This is rather unexpected for a saprophytic fungus.

The present paper reports efforts to find *M. mycetomatis* in the environment in Sudan. Isolating fungi by culturing soils and thorns, which are preconditioned for traumatic inoculation, was attempted. Isolated strains were compared to patient strains from Sudan that were identified as *M. mycetomatis* (2). Environmental fungal isolates with a colony appearance resembling that of *Madurella* were identified and compared by molecular methods. In addition, direct PCR-mediated testing of environmental material was performed.

MATERIALS AND METHODS

Strains and culture conditions. Strains presented in the current study are listed in Table 1. Sudanese *Madurella* strains used for comparison were those of Ahmed et al. (2), isolated by direct culture of black grains obtained from deep biopsies of patient lesions at the Mycetoma Research Center, Khartoum, Sudan. They were maintained on malt extract agar (MEA) slants at 24°C and transferred to fresh tubes regularly to preserve viability. Reference strains, selected from genera being reported as agents of eumycetoma, were retrieved from the collection of the Centraalbureau voor Schimmelcultures (Utrecht, The Netherlands).

Cultivation of environmental pigmented fungi. Strains were cultured from soil and thorn samples collected in the Gezira area along the Blue Nile southeast of Khartoum, Sudan (Fig. 1). From 43 samples (up to 20 cm in depth), 4 g of soil was suspended in 30 ml of sterile 0.9% saline by high-speed vortexing for 5 min. After settling, the supernatant was diluted 1:5 and 1:50 for inoculation of two or three MEA culture plates with 0.5 ml of the suspension.

Thirty-five thorn samples were collected and divided into two groups: fresh ones from green bushes (15 samples) and old ones lying on the ground or from old dry bushes (20 samples). Thorns primarily derived from the abundant species *Acacia mellifera*. Two or more thorns, depending on their size, were crushed in 2 ml of 0.9% sodium chloride solution. Two dilutions were made to inoculate two plates each (fresh thorns, 1:1 and 1:10; old thorns, 1:10 and 1:100). To decrease

* Corresponding author. Mailing address: Erasmus University Medical Center Rotterdam, Department of Medical Microbiology & Infectious Diseases, Dr. Molewaterplein 40, 3015 GD Rotterdam, The Netherlands. Phone: 31-10-4635813. Fax: 31-10-4633875. E-mail: vanbelkum@bacl.azr.nl.

TABLE 1. Survey of strains

Strain	Species	Source	Similarity with reference <i>M. mycetomatis</i> rRNA seq. ^a
Reference strains			
CBS 247.48	<i>M. mycetomatis</i>	United States (Gammel)	100%
CBS 201.38	<i>Madurella</i> sp. 1	Indonesia	Confident alignment ^b
CBS 248.48	<i>Madurella</i> sp. 1	New Mexico	Confident alignment
CBS 217.55	<i>Madurella</i> sp. 1	Argentina	Confident alignment
CBS 216.29	<i>Madurella</i> sp. 2	Italy	Distantly related
Clinical isolates			
DH11849	<i>M. mycetomatis</i>	Sudan, patient	0-bp difference
DH11850	<i>M. mycetomatis</i>	Sudan, patient	0-bp difference
DH11854	<i>M. mycetomatis</i>	Sudan, patient	0-bp difference
P2	<i>M. mycetomatis</i>	Sudan, patient	1-bp difference
DH11853	<i>M. mycetomatis</i>	Sudan, patient	2-bp difference
DH11862	<i>M. mycetomatis</i>	Sudan, patient	3-bp difference
Environmental isolates			
DH11963	<i>Exserohilum mcginnisii</i>	Sudan, soil	Nonrelevant
dH11959	<i>Bipolaris papendorfi</i>	Sudan, soil	Nonrelevant
dH11984	<i>Bipolaris spicifera</i>	Sudan, soil	Nonrelevant
4	<i>Ulospora bilgrami</i>	Sudan, thorn	Nonrelevant
5	<i>Ulospora bilgrami</i>	Sudan, thorn	Nonrelevant
6	<i>Ulospora bilgrami</i>	Sudan, thorn	Nonrelevant
7	<i>Ulospora bilgrami</i>	Sudan, thorn	Nonrelevant
8	<i>Ulospora bilgrami</i>	Sudan, thorn	Nonrelevant
9	<i>Chaetomium brasiliense</i>	Sudan, thorn	Nonrelevant
10	<i>Chaetomium murorum</i>	Sudan, soil	Nonrelevant
dH11971	<i>Phoma</i>	Sudan, thorn	Nonrelevant
dH11983	— ^c	Sudan, soil	Nonrelevant
dH11975	—	Sudan, soil	Nonrelevant

^a CBS 247.48 is considered to be the *M. mycetomatis* reference strain to which the ribosomal sequences for the other strains were matched for reasons of comparison.

^b "Confident alignment" indicates that difference was less than 10%.

^c —, isolates belong to currently undefined fungal species.

the number of contaminants in the case of several samples, two thorns were briefly rubbed with a cotton swab with 70% ethanol before dilution (fresh thorns, 1:1; old thorns, 1:10) and used to inoculate two plates. Aliquots of 0.5 ml were used per culture plate with Sabouraud's dextrose agar (SDA) containing 100 mg of chloramphenicol/liter. Plates were incubated at 37°C and checked for appropriate colonies at 3-day intervals. Strains with a colony appearance similar to that of *M. mycetomatis* and exuding a rusty brown, diffusible pigment into the agar were selected (16, 22).

DNA isolation of fungi isolated from soil. For DNA extraction, about 1 cm² of mycelium was transferred to a 2:1 mixture of silica gel and Celite 545 with 300 µl of cetyltrimethylammonium bromide (CTAB) buffer (200 mM Tris-HCl, [pH 7.5], 200 mM Na-EDTA, 8.2% NaCl, 2% CTAB). The material was ground with a micro-pestle (Eppendorf). After adding an additional 200 µl of CTAB buffer and vigorous shaking, the sample was incubated for 10 min at 65°C in a water bath. An equal volume of chloroform was added, vortexed briefly, and centrifuged for 5 min at 14,000 rpm (model 5915C centrifuge; Eppendorf, Hamburg, Germany). After transferring the aqueous supernatant to a new Eppendorf tube, 2 volumes (≈800 µl) of 96% ethanol were added and mixed gently. DNA was precipitated at -20°C for at least 30 min. The pellet, obtained by centrifugation, was washed twice with 500 µl of 70% ethanol. DNA was dried overnight at room temperature and suspended in 97.5 µl of TE buffer (10 mM Tris, 10 mM Na-EDTA, pH 8.0) with 2.5 µl of RNase solution (200 U of pancreatic RNase A/ml in 0.01 M sodium acetate). Samples were incubated for 5 to 30 min at 37°C and stored at -20°C.

Ribosomal PCR and sequencing for identification of soil organisms. DNA samples isolated from unknown strains isolated from soil or thorns was analyzed by a general primer-mediated ribosomal PCR test. The rDNA internal transcribed spacer (ITS) domain was amplified with primers V9G and LS266 (11). Other primer combinations (V9G/ITS4, LS266/ITS5) were used in case of negative results. The near-complete SSU rDNA gene was amplified with primers NS1 and NS24. Of several programs used, optimal results were obtained with the following: 94°C for 5 min followed by 30 cycles of 94°C for 1 min, 48°C for 1 min, and 72°C for 2 min, followed by 72°C for 3 min. One unit of Super-Taq polymerase (Sphaero Q, Leiden, The Netherlands) was used for a 50-µl reaction mixture

with 2 µl of DNA. In some cases, 6% dimethylsulfoxide (Merck, Darmstadt, Germany) was added. Amplicons were purified using the GFX PCR DNA and Gel Band Purification Kit according to the manufacturer's instructions (Amersham Pharmacia, Roosendaal, The Netherlands). Concentrations of amplicons were estimated by comparison with a marker (SmartLadder; Eurogentec, Seraing, Belgium) on a 1% agarose gel.

The sequencing reactions were carried out with 15 to 50 ng of DNA for a 10-µl reaction mixture, including 4 pmol of primer and 4 µl of BigDye RR Mix (Applied Biosystems) using primers ITS4 and ITS1 (ITS domain) or Oli1, Oli5, Oli9-11, Oli13, and Oli14 (reference 21 and Table 2) and NS24 (SSU gene) with 25 cycles, as follows: 96°C for 10 s, 50°C for 5 s, and 60°C for 4 min. Sequence reaction products were analyzed on an automated capillary sequencer (ABI, Gouda, The Netherlands). The sequences were adjusted using SeqMan of Lasergene software (DNASTAR Inc., Madison, Wis.) and analyzed using BioNumerics (Applied Maths, Kortrijk, Belgium). Sequences were compared to a set of dedicated sequences available at CBS (ITS fragment ITS 1+2, including 5.8S and the nearly complete SSU). Sequences of putative agents of eumycetoma were included (e.g., *Acremonium* spp., *Chaetomium*, *Neotestudina* spp., *Pseudallescheria* spp., *Bipolaris*, *Leptosphaeria*, and *Madurella* spp.). The sequences alignment was considered confident when less than a 10% difference was observed. In case no match was found, sequences were run against the EMBL databank using BLAST.

Processing of soil and thorn samples for direct DNA testing. Two grams of soil (74 Sudanese samples, 14 Dutch controls, and 1 United Kingdom control) or four or five thorns ($n = 22$) were suspended in 5 or 2 ml of samples, sterile physiological saline containing 2% sodium dodecyl sulfate (SDS) (Merck, Darmstadt, Germany) and 5 mg of proteinase K (Merck). The samples were incubated at room temperature on a horizontal shaker for 10 min and then incubated at 37°C (1 h). One milliliter of the clear upper layer of the suspensions was extracted with an equal volume of phenol/chloroform/isoamyl alcohol (25:24:1). The DNA was precipitated from the aqueous layer with cold absolute ethanol (Merck). The pellet was washed once with 70% ethanol, air dried, and dissolved in 100 µl of sterile distilled water. This crude extract was further purified with the




FIG. 1. Detailed map of the region in Sudan where the thorns were collected and the soil was sampled. Main sampling sites were in the encircled region comprising the region southeastern to Khartoum (Wad Madani and Sinnar). Other regions where mycetoma is highly endemic are highlighted by the elliptic red-dot regions. The red bars indicate the boundaries of the Sudanese mycetoma belt.

QIAmp Tissue kit (Qiagen, Hilden, Germany) according to the manufacturer's recommendations. The extracted DNA was then immediately tested by PCR.

PCR RFLP analysis of DNA extracted from thorns and soil samples. This analysis aimed at the broad-spectrum amplification of fungal DNA, followed by an *M. mycetomatis*-specific nested PCR. In the first PCR, ITS4 and ITS5 primers were used: 5'-TCCTCCGCTTATTGATATGC-3' and 5'-GGAAGTAAAAGT CGTAACAAGG-3', respectively (23). For the nested PCR, *M. mycetomatis* specific primers 26.1A (5'-AATGAGTTGGGCTTTAACGG-3) and (AATGA GTTGGGCTTTAACGG-3) and 28.3A (5'-TCCCGTAGT'GTAGTGT-CCC T'-3) were employed (2). The PCRs were performed in 50- μ l reaction volumes containing 0.2 U of SuperTaq polymerase (HT Biotech, Cambridge, United Kingdom), PCR buffer, deoxynucleotide triphosphate mix, 50 pmol of each primer, and 10 μ l of extracted DNA (approximately 100 ng of DNA) in the first PCR round and 5 μ l of the amplified material in the nested PCR. Cycling was performed in a model 60 thermocycler (Biomed, Theres, Germany) for 40 cycles

(94°C for 1 min, 58°C for 1 min, 72°C for 3 min). PCR products were examined by electrophoresis in 1% agarose gels containing ethidium bromide (0.3 μ g/ml). The electrophoresis was carried out in 0.5 \times Tris-borate-EDTA (TBE) buffer at a constant current of 100 mA for 1 h.

Restriction fragment length polymorphism (RFLP) in the *M. mycetomatis* ITS regions was studied by analysis of the PCR products generated by primers 26.1A and 28.3A by cleavage with *Cfo*I, *Msp*I (Boehringer Mannheim GmbH, Mannheim, Germany), and *Hae*III (New England Biolabs, Boston, Mass.) restriction enzymes. The restriction enzymes were used as recommended by the manufacturers. Fifteen microliters of PCR product was incubated overnight at 37°C with the restriction enzyme mixtures, which contained enzyme's buffer and 5 U of each enzyme per reaction. RFLP was determined by electrophoresis in 3% Nusieve GTG agarose gels (FMC Bioproducts, Rockville, Md.). The electrophoresis was performed in 0.5 \times TBE buffer at a constant current of 100 mA for 3 h.

TABLE 2. Universal and *Madurella*-specific primers used for PCR and sequencing^a

Primer	Position (gene/nucleotides)	Reference	Primer sequence (5' → 3') ^b
ITS4	LSU/41–60	23	←TCCTCCGCTTATTGATATGC
ITS5	SSU/1745–1767	23	GGAAGTAAAAGTCGTAACAAGG→
V9D	SSU/1609–1627	6	TAAAGTCCCTGCCCTTTGTA→
LS266	LSU/287–266	19	←GCATTCCCAAACAACCTCGACTC
NS1	SSU/20–38	23	GTAGTCATATGCTTGTCT→
NS24	SSU/1769–1750	10	←AAACCTTGTTACGACTTTTA
Oli5	SSU/83–102	14	GAAACTGCGAATGGCTCATT→
Oli9	SSU/573–591	14	CGCGGTAATTCAGCTCCA→
Oli10	SSU/951–935	14	←TGGYRAATGCTTTTCGC
Oli11	SSU/962–980	14	TTRATCAAGAACGAAAAGT→
Oli13	SSU/1418–1438	14	←ATAACAGGTCTGTGATGCC
Oli14	SSU/1418–1438	14	ATAACAGGTCTGTGATGCC→
26.1A ^c	ITS2/111–130	2	←AATGAGTTGGGCTTTAACGG
28.3A ^c	ITS1/474–494	2	TCCCGTAGTGTAGTGTCCCT→

^a Positions correspond to the *Saccharomyces cerevisiae* consensus sequence.

^b R = purine A or G; Y = pyrimidine C or T.

^c Primer sequences and positions are based on the *M. mycetomatis* ITS sequence (GenBank accession no. AF162133).

RESULTS

Cultivation and identification of soil- and thorn-borne fungi. Overall, 43 soil samples and 35 batches of thorns were analyzed for the presence of black-pigmented fungi resembling *M. mycetomatis*. Ultimately, 52 colonies from 24 samples were subcultured in a pure form on SDA. Strains were studied microscopically and by molecular analysis. None of the strains turned out to represent *M. mycetomatis* based on either morphological criteria or ribosomal sequencing. Some of the strains were excluded prior to sequencing based on ribosomal gene RFLP analyses (result not shown). Sequence results are listed in Table 1, presenting data for 13 environmental, 6 clinical, and 5 reference isolates. *M. mycetomatis* CBS 247.48 was found to be practically identical (0- to 3-bp difference) to six strains of cases of mycetoma from Sudan. CBS 247.48 might be Gammel's strain from Cleveland, Ohio, dating from 1927, which is the type strain of *Madurella americana*. Three further clinical strains putatively representing *M. mycetomatis* (CBS 201.38 from Indonesia, CBS 248.48 from New Mexico, and CBS 217.55 from Argentina) were found at some distance but could be aligned with confidence. *M. mycetomatis* CBS 216.29 from Italy was found to be different from both clusters. About 105 out of 190 positions of ITS1 and about 140 out of 160 positions of ITS2 of *M. mycetomatis* could be aligned with environmental *Chaetomium* species from Sudan, which were identified down to the species level by morphology and by comparison with sequences from the public domain. Three strains from Sudanese soil were found to be members of Pleosporales with distoseptate conidia, viz. *Exserohilummcginnisii*, *Bipolaris papendorffii*, and *Bipolaris spicifera*. Their ITS sequences showed no homology with *Leptosphaeria*. Five strains from Sudanese thorns and showing superficial resemblance to one of the known agents of mycetoma, *Neotestudina rosatii*, were morphologically identified as *Ulospora bilgramii* (13). Five reference strains of *N. rosatii* were clearly different but could be aligned with confidence. The type strain of *N. rosatii* CBS 427.62 originated from a case of mycetoma, whereas the remaining four strains were environmental (not shown in Table 1).

PCR-mediated detection of *M. mycetomatis* from various

sources. The first round of PCR, involving the ITS4-ITS5 primer pair, done directly on nucleic acid extracts from soil and thorn samples produced clearly visible products in all instances. Frequently mixtures of various PCR products were produced, indicating the presence of DNA from various fungal species. These results demonstrated that the PCR tests were not inhibited by contaminants from soil of thorns. The combination of primers V9G and LS266 generated the expected DNA amplicons of 1,100 bp as well.

Out of 22 thorn batches tested in this study, only one reacted with the *Madurella*-specific primers in the nested PCR and produced the characteristic 424-bp fragment matching the positive control (Fig. 2). The *M. mycetomatis* DNA was detected in significantly higher rates in the soil samples, since 17 positives (23%) were observed among the 74 soil samples from endemic foci. All the fourteen Dutch and the single British control soil samples remained negative in all PCR experiments. All the PCR products thus obtained ($n = 18$) displayed the same RFLP pattern, which exactly matched those of the clinical *M. mycetomatis* strains (Fig. 3). Patterns generated with all different restriction enzymes were identical and differed from those of the other species mentioned in Table 1.

DISCUSSION

The classical etiology of human eumycetoma involves a relatively limited set of agents as established pathogens typically causing mycetoma (20). This view is corroborated by the overrepresentation of a small number of species involved in the reported cases of eumycetoma, such as *Acremonium* spp., *Pseudallescheria boydii*, and *Madurella* spp. An environmental source has been suggested for the latter microorganism, but direct proof thereof is largely lacking. Presumably, when more ecological data become available, it could become clear that *M. mycetomatis* actually has another important ecological niche, rendering subcutaneous growth in humans insignificant from a fungal perspective (7). In addition, there are large problems with the correct identification of *M. mycetomatis*. It shows poor morphological differentiation, is sterile, and easily causes confusion among clinicians and microbiologists (5). Hence, reli-




FIG. 2. Molecular detection of *M. mycetomatis* in soils and thorn samples. (A) Results of the broad-spectrum ITS PCR employing primers ITS4 and ITS5. Note that amplicons can be observed in all lanes. (B) Result of the nested PCR using *M. mycetomatis*-specific primers. Lanes 1 to 6, amplicons derived from Dutch control samples; lanes 7 to 14, amplified material from Sudanese samples; lane 15, results obtained with purified *M. mycetomatis* DNA; lanes 16 and 17, negative controls; lanes M, 100-bp molecule sizing ladder, and the 600-bp fragment is indicated (arrow).

able identification of *M. mycetomatis* is still problematic. In order to solve this problem, Ahmed et al. (2) undertook a molecular study into *Madurella*-like agents of mycetoma in Sudan. These authors found morphological dissimilarity among strains originating from cases of mycetoma but noted that all clinical isolates proved to be identical. A single ribosomal spacer genotype was predominant, and this species was identified as *M. mycetomatis*. These findings now enable the targeted search, by molecular techniques, for the environmental source of this important cause of eumycetoma.

Although it is unlikely that *M. mycetomatis* in nature is present in the same form it is in patients, it is suggested that the isolation of this species directly from the environment is very difficult using the current protocols. Despite failed attempts to culture the organism, we now present data showing that it is possible to detect *M. mycetomatis* in the environment of areas in Sudan where it is endemic by molecular methods. In addition, thorny *A. mallifera* bushes, which are very common in the region sampled and are responsible for a high number of thorn pricks, were shown to occasionally harbor the same species of fungus. This supports the observation of Abbott (1) that the frequency of mycetoma is associated, albeit weakly, with the presence of thorny trees. The low recovery rate suggests, however, that *M. mycetomatis* is not an ordinary saprobe on the thorns.




FIG. 3. Molecular identification of *M. mycetomatis* from the environment and clinical biopsies. Shown are digests for the specific 26.1A/28.3A PCR products obtained for two clinical isolates (lanes 2 and 3, 5 and 6, and 8 and 9) and a single environmental isolate (lanes 1, 4, and 7). Note that per restriction digest, the patterns are identical. On the left (lane M), the 10-bp molecule sizing ladder is shown, and the 100- and 1,000-bp markers are indicated (arrow).

Human subcutaneous inoculation apparently is an efficient selective means for recovery of the fungus. A comparable situation is known in black yeast-like fungi, where *Cladophialophora bantiana* (11) and *Fonsecaea pedrosoi* (De Hoog et al., unpublished data) are homologous examples. These fungi are thus far known only from mammals and can be obtained from the environment after mouse passage only. Interestingly, *F. pedrosoi* is one of the agents of chromoblastomycosis, which is also a mycosis associated with (minor) trauma.

Other fungal species (anamorph genera *Bipolaris*, *Drechslera*, and *Exserohilum*) were frequently isolated from soil in Sudan. Among these was *E. mcginnisii*, which thus far is only known from a single strain from a human patient and hence has been regarded as a true pathogen. However, its occurrence in the environment suggests that this is also one of the saprobes which only occasionally is able to infect a human host.

In conclusion, we have demonstrated the presence of *M. mycetomatis* DNA on or in *Acacia* thorns and in soil samples. The fungal prevalence in the soil especially suggests that this may be the prime reservoir from which infections originate. This is further corroborated by the fact that clinical isolates and environmental strains seem to share the same ribosomal genotype, confirming that at least the same fungal species is concerned. The challenge for current research is to elucidate the processes that underlie the culture-refractory nature of environmental *M. mycetomatis* isolates. Whether or not the fungus readily adapts to humans provides another interesting and important research topic.

ACKNOWLEDGMENTS

Part of this research was financed through a Young Investigators award provided to A.V. by the European Society of Clinical Microbiology and Infectious Diseases. D.A. thanks Boehringer-Ingelheim for financial support.

A. Aptroot and G. Verkleij are acknowledged for their help with identification of fungi. R. C. Summerbell and A. A. Padhye are thanked for strains and for comments on the manuscript. K. Luijster-

burg and A. H. G. Gerrits van den Ende are thanked for technical assistance.

REFERENCES

- Abbott, P. 1956. Mycetoma in the Sudan. *Trans. R. Soc. Trop. Med. Hyg.* **50**:11–30.
- Ahmed, A. O. A., M. M. Mukhtar, M. Kool-Sijmons, A. H. Fahal, G. S. De Hoog, A. H. G. Gerrits van den Ende, E. E. Zijlstra, H. Verbrugh, A. M. El Sir Abugroun, A. M. Elhassan, and A. Van Belkum. 1999. Development of a species-specific PCR-restriction fragment length polymorphism analysis procedure for identification of *Madurella mycetomatis*. *J. Clin. Microbiol.* **37**:3175–3178.
- Basset, A., R. Camain, R. Baylet, and D. Lambert. 1965. Role des épines de mimosacées dans l'inoculation des mycétomes (à propos de deux observations). *Bull. Soc. Pathol. Exot.* **58**:22–24.
- Brumpt, E. 1906. Les mycétomes. Asselin & Houzeau, Paris, France.
- De Hoog, G. S., A. Buiting, C. S. Tan, A. B. Stroebel, C. Ketterings, E. J. De Boer, B. Naafs, R. Brimicombe, M. K. E. Nohlmans-Paulssen, G. T. J. Fabius, A. H. Klokke, and L. G. Visser. 1993. Diagnostic problems with imported cases of mycetoma in The Netherlands. *Mycoses* **36**:81–87.
- De Hoog, G. S., and G. van den Ende. 1998. Molecular diagnostic of clinical strain of filamentous Basidiomycetes. *Mycoses* **41**(5–6):183–189.
- De Hoog, G. S., F. D. Marvin-Sikkema, G. A. Lahpor, J. C. Gottschall, R. A. Prins, and E. Gueho. 1994. Ecology and physiology of the emerging opportunistic fungi *Pseudallescheria boydii* and *Scedosporium prolificans*. *Mycoses* **37**:71–78.
- El Hassan, A. M., A. H. Fahal, I. A. El Hag, and E. A. G. Khalil. 1994. The pathology of mycetoma. *Sudan Med. J.* **32**(Suppl. 1):23–45.
- Fahal, A. H., and S. H. Suliman. 1994. Clinical presentation of mycetoma. *Sudan Med. J.* **32**:46–66.
- Gargas, A., and J. W. Taylor. 1995. Phylogeny of Discomycetes and early radiation of apothecial Ascomycotina inferred from SSU rDNA sequence data. *Exp. Mycol.* **19**:7–15.
- Gerrits van den Ende, A. H. G., and G. S. De Hoog. 1999. Variability and molecular diagnostics of the neurotropic species *Cladophialophora bantiana*. *Stud. Mycol.* **43**:152–162.
- Gumaa, S. A. 1994. The aetiology and epidemiology of mycetoma. *Sudan Med. J.* **32**(Suppl. 1):14–22.
- Hawksworth, D. L. 1979. Ascospore sculpturing and generic concepts in the *Testudinaceae* (syn. *Zopfiaceae*). *Can. J. Bot.* **57**:91–99.
- Hendriks, L., Y. van de Peer, M. van Herck, J. M. Neefs, and R. De Wachter. 1990. The 18S ribosomal RNA sequence of the sea anemone *Anemonia sulcata* and its evolutionary position among other eukaryotes. *FEBS Lett.* **269**:445–449.
- Laveran, M. A. 1902. Au sujet d'un cas de mycetoma à grains noirs. *Bull. Acad. Med. Paris* **47**:773–776.
- Mackinnon, J. E. 1954. A contribution to the causal organisms of maduro-mycosis. *Trans. R. Soc. Trop. Med. Hyg.* **48**:470–480.
- Mahgoub, E. S., and I. G. Murray. 1973. Mycetoma. William Heinemann, London, United Kingdom.
- Mariat, F., P. Destombes, and G. Segretain. 1977. The mycetomas: clinical features, pathology, etiology, and epidemiology. *Contrib. Microbiol. Immunol.* **4**:1–39.
- Masclaux, F., E. Gueho, G. S. de Hood, and R. Christen. 1995. Phylogenetic relationship of human-pathogenic *Cladosporium* (Xylohypha) species inferred from partial LS rRNA sequences. *J. Med. Vet. Mycol.* **33**:327–338.
- Rippon, J. W. 1988. Medical mycology. The pathogenic fungi and the pathogenic actinomycetes, 3rd ed. Saunders, Philadelphia, Pa.
- Sterflinger, K., R. De Baere, G. S. De Hoog, R. De Wachter, W. E. Krumbein, and G. Haase. 1997. *Contiosporium perforans* and *C. apollinis*, two new rock-inhabiting fungi isolated from marble in the Sanctuary of Delos (Cyclades, Greece). *Antonie Leeuwenhoek* **72**:349–363.
- Thirumalachar, M. J., and A. A. Padhye. 1968. Isolation of *M. mycetomi* from soil in India. *Hindustan Antibiot. Bull.* **10**:314–318.
- White, T. J., T. Burns, S. Lee, and J. Taylor. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics, p. 315–322. *In* M. A. Innis, D. H. Gelfand, J. J. Sninsky, and T. J. White (ed.), *PCR protocols: a guide to methods and applications*. Academic Press, San Diego, Calif.