

# Anterior chamber depth changes after retinal detachment

R. E. HARTLEY AND R. J. MARSH

*From Moorfields Eye Hospital and the Western Ophthalmic Hospital, London*

Glaucoma is a rare complication of retinal detachment surgery. Smith (1967) reported postoperative angle closure in 4 per cent. of 1,000 consecutive scleral buckling procedures. He believed the mechanism to be a forward rotation of the ciliary body with shallowing of the anterior chamber and angle closure. Fiore and Newton (1970) investigated 34 patients after scleral buckling with encirclement and found that four were shallow after 2 months; the remainder became shallow temporarily but had returned to preoperative levels in 2 months. One of these had a choroidal detachment which recovered. We have set out to pursue these findings further: firstly, to compare anterior depth changes after various types of detachment surgery, and secondly, to investigate the early postoperative period.

## Methods

Patients suffering from untreated retinal detachments presenting at Moorfields Eye Hospital and the Western Ophthalmic Hospital over a period of 9 months were included in the survey. The patients were assessed preoperatively as follows: a history was recorded with emphasis on any previous family incidence of glaucoma. Visual acuities and refractive errors were noted. Applanation values, with pupils dilated, were taken. Gonioscopy was performed, recording the angles as deep, medium, or shallow (a short note was added on the trabecular appearance). The anterior chamber depths were measured with the Haag-Streit Type 2 Pachometer (Lowe, 1966) when the pupils were dilated. The corneal thickness was measured and then the distance from the anterior corneal surface to the anterior lens capsule. The instrument provides a reliable comparative reading without further correction. Operative procedures were documented to include the use of osmotic agents, the type and size of plomb or encircling strap, its average distance from the limbus, any muscles detached, any drainage of subretinal fluid, the number of quadrants treated with cryotherapy, and operative complications, such as vitreous or choroidal haemorrhages, paracentesis, and, lastly, in some cases, applanation on completion.

Postoperative readings were made of anterior chamber depth and applanation. Gonioscopy was carried out if any marked shallowing occurred. These readings were taken on the first postoperative day, if possible, and at least twice in the first week. Readings were continued weekly until the preoperative depth was regained or for a follow-up period of at least 3 months in all other cases.

## Results

A total of 56 patients was examined and they were divided into four main groups (Table I).

The amount of shallowing of the anterior chamber, the time that this occurred, and any special comments are set out in Tables II to V.

**Table I** Four main groups totalling 56 patients

A	Plombs with drainage	5
B	Plombs without drainage	28
C	Encircling straps with drainage	19
D	Encircling straps without drainage	4

GROUP A Plomb with drainage. Five patients in four of whom the chambers became shallow (Table II).

**Table II** Plombs with drainage

Patient no.	Amount of shallowing of anterior chamber (mm.)			Comments on retina
	First 3 days	End of first week	Late	
1	0.6	0.6	nil	Flat
2	0.9	0.7	0.6	Detached
3	0.1	nil	nil	Flat
4	0.5	0.25	nil	Flat

#### Special remarks on Group A

**Case 2** Corneal oedema was present from the second day. The retina remained detached. The eye was encircled 1 week later with no success and strap pain was experienced.

GROUP B Plomb without drainage. 28 patients, in thirteen of whom the anterior chamber became shallow (Table III, overleaf).

#### Special remarks on Group B

**Case 1** At the first week small choroidal detachments were seen nasally. At Day 12 the angle was closed for 180°. At 4 weeks iris atrophy was seen and the retina re-detached. The tension was not raised at any time.

**Case 2** At 10 days ring choroidal detachments were present and the angle closed nasally and temporally.

**Case 7** Had unchanging residual fluid which could well be due to a previous schisis.

**Case 13** Vitreous haemorrhage occurred during surgery. The retina re-detached on Day 7. The patient was re-operated on twice but the retina remained detached.

GROUP C Encirclement with drainage. Nineteen patients of whom seven shallowed, one deepened, and eleven remained unchanged (Table IV, overleaf).

#### Special Remarks on Group C

**Case 2** On Day 2 there was iridocorneal contact with glaukomflecken; tension was uncontrolled medically and on Day 4 the strap was cut. The glaucoma resolved but sectoral iris atrophy ensued.

**Case 3** Raised tension occurred on Day 1; 2.6 per cent. of the angle was closed. The tension was

**Table III** *Plombs without drainage*

Patient no.	Amount of shallowing of anterior chamber (mm.)			Comments on retina
	First 3 days	End of first week	Late	
1	—	0·5	0·1	Detached
2	—	0·5	nil	Flat
3	—	1·2	nil	Flat
4	—	0·8	0·5	Flat
5	0·2	0·2	0·1	Flat
7	0·3	nil	nil	Residual fluid
8	0·15	nil	nil	Flat
9	0·2	0·15	0·4	Flat
10	0·15	0·1	nil	Flat
11	0·2	0·2	nil	Flat
12	0·1	0·2	0·2	Flat
13	0·5	0·4	—	Detached

**Table IV** *Encirclement with drainage*

Patient no.	Amount of shallowing of anterior chamber (mm.)			Comments on retina
	First 3 days	End of first week	Late	
1	0·4	0·6	—	Flat
2	1·3	1·1	nil	Flat
3	1·3	0·8	0·8	Flat
4	0·4	0·3	nil	Flat
5	0·15	1·2	—	Re-detached
6	0·5	0·9	0·3	Flat
7	0·3	1·0	0·5	Re-detached

uncontrolled medically and on Day 7 a peripheral iridectomy was carried out. The glaucoma resolved over the next 3 weeks with a slow initial response.

**Case 4** A giant tear was present.

**Case 5** There was an inferior coloboma of iris ciliary body and choroid. The retina was flat on Day 3 but then slowly detached. The strap was tightened 2 months later with only temporary success.

**Case 6** Choroidal detachments were seen on Day 5, but slowly resolved.

**Case 7** There was a total detachment with incipient massive vitreous retraction. On Day 5 the retina remained detached and there was strap pain. Massive vitreous retraction occurred.

**Table V** *Encirclement without drainage*

Patient no.	Amount of shallowing of anterior chamber (mm.)			Comments on retina
	First 3 days	End of first week	Late	
1	0.3	nil	nil	Flat
2	0.1	nil	—	Re-detached
3	0.6	nil	nil	Flat
4	0.6	0.5	nil	Flat

GROUP D Encirclement without drainage. Four patients, in all of whom the anterior chamber became shallow (Table V)

#### *Special remarks on Group D*

**Case 2** The retina was flat for 5 weeks, but massive vitreous retraction then occurred with total retinal detachment. Re-operation with a shoe under the encircling strap and intravitreal silicone fluid failed to achieve flattening.

**Case 4** Choroidal detachments were seen at the 5th postoperative day but had resolved by the 10th.

Early shallowing of the anterior chamber occurred in 28 cases and deepening in one of the total of 56 cases. Seventeen of the 28 followed local procedures and eleven followed encirclements. 52 per cent. of those treated by local procedures and 48 per cent. of the encirclements showed shallowing in the early postoperative period. Our findings of corneal thickening postoperatively were the same as those of Fiore and Newton (1970), *viz.* a temporary small increase in nearly all eyes, returning to normal within 2 weeks.

The amount of cryotherapy and the size of the plomb or strap did not seem to influence this shallowing. The present series is not large enough to show whether drainage or non-drainage of subretinal fluid influences the anterior chamber depth, but it was interesting to observe how rapidly the normal depth was fully recovered after major buckling procedures, with no drainage of subretinal fluid and with paracentesis.

#### **Discussion**

In a series of 160 cases of retinal detachment investigated by Sebastyen, Schepens, and Rosenthal (1962), postoperative raised intraocular pressure was seen to occur in fourteen. The authors noted that the following factors were related to this increase:

- (1) Changes in the filtration angle (such as peripheral anterior synechiae, narrow angle, and increased pigmentation of the trabecular meshwork)
- (2) Duration of the detachment
- (3) Multiple retinal operations
- (4) Surgical aphakia.

The series studied in this paper were fresh phakic cases, thus eliminating two of the factors mentioned. None had any previous or family history of glaucoma or obvious angle abnormality preoperatively.

The cases in which the anterior chamber became shallow tended to fall into two groups: (1) Those with a small amount of shallowing (0.1 — 0.6 mm.) which was usually present

on the first postoperative day and was unaccompanied by other changes except for increased corneal thickness. Preoperative anterior chamber depth was usually regained within 1 month.

(2) Those with a greater degree of shallowing (approximately 1.00 mm.), which was not necessarily present on the first day, but developed within the first week. The shallowing was accompanied by other changes: notably a change in intraocular pressure and choroidal detachments.

Permanent shallowing tended to occur in unsuccessfully treated cases. As can be seen from the Tables, there was a large proportion of the Group I type of shallowing in those treated by the local procedures. In contrast, there was a larger proportion of Group II response in the cases treated by encirclement.

The Group I response is a benign condition and possibly occurred because of temporary displacement of the ciliary body by the buckling procedure, causing relaxation of the lens iris diaphragm and allowing the lens to move forwards.

The Group II response is a serious condition with a poor prognosis for success. The eyes in this group were presumably poorly able to deal with the physiological readjustments subsequent to the detachment surgery. Possibly this is because the eyes are originally on the border-line of anterior segment ischaemia. Several of our patients showed ischaemic manifestations postoperatively. This group is being further examined at present.

### Summary

56 patients were examined before and after retinal detachment surgery. Special emphasis was placed on changes in anterior chamber depth, intraocular tension, and the type of surgery. 52 per cent. of the local buckling procedures and 48 per cent. of the encirclements were followed by shallowing of the anterior chamber in the early postoperative period. Permanent shallowing occurred more commonly after encirclements and carried a poor prognosis.

We should like to acknowledge the help of Messrs. L. G. Fison and J. H. Redmond Smith, and to thank them and the surgeons of the Western Ophthalmic Hospital for permission to report their cases.

### References

- FIGORE, J. V., and NEWTON, J. C. (1970) *Arch. Ophthalm. (Chicago)*, **84**, 284  
LOWE, R. F. (1966) *Amer. J. Ophthalm.*, **62**, 7  
SEBESTYEN, J., SCHEPENS, C. L., and ROSENTHAL, M. L. (1962) *Arch. Ophthalm. (Chicago)*, **67**, 736  
SMITH, T. R. (1967) *Amer. J. Ophthalm.*, **63**, 1807