

Report

Primate *DAX1*, *SRY*, and *SOX9*: Evolutionary Stratification of Sex-Determination Pathway

Megha Patel,¹ Karin S. Dorman,² Yao-Hua Zhang,³ Bing-Ling Huang,³ Arthur P. Arnold,¹ Janet S. Sinsheimer,^{2,4,5} Eric Vilain,^{3,4,6} and Edward R. B. McCabe^{3,6}

¹Department of Physiological Science, University of California Los Angeles; Departments of ²Biomathematics, ³Pediatrics, ⁴Human Genetics, and ⁵Biostatistics, University of California Los Angeles School of Medicine; and ⁶Mattel Children's Hospital at UCLA, Los Angeles

The molecular evolution of *DAX1*, *SRY*, and *SOX9*, genes involved in mammalian sex determination, was examined in six primate species. *DAX1* and *SRY* have been added to the X and Y chromosomes, respectively, during mammalian evolution, whereas *SOX9* remains autosomal. We determined the genomic sequences of *DAX1*, *SRY*, and *SOX9* in all six species, and calculated K_a , the number of nonsynonymous substitutions per nonsynonymous site, and compared this with the K_s , the number of synonymous substitutions per synonymous site. Phylogenetic trees were constructed by means of the *DAX1*, *SRY*, and *SOX9* coding sequences, and phylogenetic analysis was performed using maximum likelihood. Overall measures of gene and protein similarity were closer for *DAX1* and *SOX9*, but *DAX1* exhibited nonsynonymous amino acid substitutions at an accelerated frequency relative to synonymous changes, similar to *SRY* and significantly higher than *SOX9*. We conclude that, at the protein level, *DAX1* and *SRY* are under less selective pressure to remain conserved than *SOX9*, and, therefore, diverge more across species than does *SOX9*. These results are consistent with evolutionary stratification of the mammalian sex determination pathway, analogous to that for sex chromosomes.

DAX1 is a member of the nuclear hormone receptor superfamily (Zanaria et al. 1994). Mutations in this transcription factor result in developmental and functional abnormalities of the steroidogenic axis, as well as the clinical phenotypes of adrenal hypoplasia congenita (MIM 300200) and hypogonadotropic hypogonadism (Muscatelli et al. 1994; Zanaria et al. 1994; Guo et al. 1995; Habiby et al. 1996; McCabe 2000). Since *DAX1* maps to the dosage-sensitive sex reversal (DSS [MIM 300018]) critical region (Bardoni et al. 1994; Zanaria et al. 1994), and transgenic overexpression of *DAX1* in mice with a Y chromosome containing a weak *SRY* allele can result in XY sex reversal (Swain et al. 1998), *DAX1* is a candidate gene for DSS. However, experimental and clinical evidence suggests that the role of *DAX1* in sex determination is more complex. Targeted deletion of *DAX1* exon 2 in mice fails to give the expected sex-

reversed phenotype (Yu et al. 1998), and a female human who is genotypically homozygous for a *DAX1* mutation also does not exhibit the expected phenotype (Merke et al. 1999). It is possible, however, that *DAX1* acts as a part of a dosage-sensitive multiprotein complex, that time of expression is important, and that *DAX1* functions differently in human and mouse (Swain et al. 1998; Goodfellow and Camerino 1999). *DAX1* is a relatively recent addition to the X chromosome, since it is autosomal in marsupials (Pask et al. 1997) and maps to the stratum that was acquired by the eutherian mammal X chromosome ~80–130 million years ago (Lahn and Page 1999).

SRY encodes a protein containing a “high-mobility group” domain (HMG box), which enables it to bind DNA (Sinclair et al. 1990; Giese et al. 1994). Mutations that map to the HMG box disrupt the *SRY*-DNA complex and result in XY sex reversal (Pontiggia et al. 1994), and XX mice carrying the *Sry* transgene may develop as males (Koopman et al. 1991). Among eutherian mammals, the *SRY* sequence is rapidly evolving outside the HMG box (Tucker and Lundrigan 1993; Whitfield et al. 1993). *SRY* is a relatively recent addition to the sex-

Received September 11, 2000; accepted for publication November 13, 2000; electronically published December 7, 2000.

Address for correspondence and reprints: Dr. E. R. B. McCabe, Department of Pediatrics, UCLA School of Medicine, Los Angeles, CA 90095-1752. E-mail: emccabe@mednet.ucla.edu

© 2001 by The American Society of Human Genetics. All rights reserved. 0002-9297/2001/6801-0032\$02.00




Figure 1 Genomic sequencing strategy for *DAX1*. Overlapping PCR primers (sets A–G) were used to amplify primate *DAX1*; 5' primers (A.5–G.5) are shown above the gene and 3' primers (A.3–G.3) are shown below. The 5' end of *DAX1* was amplified by means of the conserved SF1-response element upstream of the start codon as a primer-anchoring site.

determination pathway, appearing for the first time in mammals (Graves and Foster 1994).

SOX9, located on human chromosome 17, has also been implicated in the sex-determination pathway (Foster et al. 1994; Wagner et al. 1994). *SOX9*, like *SRY*, is an HMG family member, and mutations in a single allele of *SOX9* are responsible for XY sex reversal and campomelic dysplasia (MIM 114290) (Foster et al. 1994; Wagner et al. 1994; Morais da Silva et al. 1996). *SOX9* duplications in humans (Huang et al. 1999) can result in XX males in the absence of *SRY*. *SOX9* is much older evolutionarily than *DAX1* or *SRY* (Morais da Silva et al. 1996; McBride et al. 1997), may be part of an ancient “core” sex-determination mechanism that predates the mammalian radiation (Koopman 1999) and has important roles in sex determination and bone development (Bi et al. 1999). These factors may explain the highly conserved nature of *SOX9*.

The purpose of our investigations was to examine the molecular evolution of *DAX1* in relation to *SRY* and to *SOX9*. Genomic DNA was obtained from cell lines for each species ATCC numbers CRL-1857: *Pan troglodytes* (chimpanzee); CRL-1850: *Pongo pygmaeus* (orangutan); TIB-201: *Hylobates* species (gibbon); CRL-1773: *Callithrix jacchus* (marmoset); and from ClonTech Laboratories: *Homo sapiens* (human) and *Macaca mulatta* (rhesus) (fig. 1). A PCR-based approach was used to sequence the primate homologue of each gene. We used or adapted previously published primers for *SRY* (Whitfield et al. 1993) and *SOX9* (Wagner et al. 1994). PCR primers used for *DAX1* were A5 (TTGAACTACC-GAGGTCATGGG) and A3 (CATGTTGTAGAGGAT-GCTGCC), B5 (GGGCTGCAGGAGCCGCGGGCC) and B3 (GCAGCGGTACAGGAGTGCCAC), C5 (GGGTGGGCAGAAAAGGGCTGC) and C3 (GGTGCTGCCCTGCTGCGGGTG), D5 (GGCCAGGGGGTAGAGAGGCGC) and D3 (GATCTTCTGCAGCATGCTGG), E5 (GAGACTGTAGAAGTCTCGGAG) and E3 (GTAGGCGTACTCCTTGGTACT, F5 (AGTACCAAGGAGTACGCCTAC and F3 (CCACTGGAGTCCCT-

GAATGTTCTT), and G5 (ACGGGACGTGCCGGCA-GTGC) and G3 (CACTTGTGTGGCCCACATGAC). Sequences were deposited into GenBank. PCR products were sequenced using ABI 375 (PE Biosystems), and sequence data were formatted and analyzed using the GCG software package, version 10.0.

The ratios of nonsynonymous to synonymous (K_a/K_s) were estimated with the use of codeml, from the phylogenetic analysis using maximum likelihood (PAML) package (Goldman and Yang 1994). We imposed the commonly accepted primate topology (Pilbeam 1984) and estimated kappa (κ) and omega (ω , i.e., K_a/K_s), with alpha (α) and rho (ρ) fixed at zero. A single ω was used for the entire topology, since a model with separate ω for each branch did not give a significantly better fit. Codon frequencies were determined by means of the 12-parameter $f_3 \times 4$ model, but parameter estimates were largely independent of the codon-frequency model used. The tree-shape comparisons among genes were performed by likelihood-ratio tests. We compared a restricted model, having a constant ratio between respective branches of the trees for the two genes but all other evolutionary parameters unrestricted, with a more general model having separate evolutionary parameters (including the branch lengths) for the two genes. The maximum likelihood of the restricted model (18 parameters) and the maximum likelihood of the general model (26 parameters) were calculated by use of the HKY nucleotide-substitution model (Hasegawa et al. 1985), implemented in a modification of Schadt et al. (1998) (fig. 2). Our results showed that at the nucleotide level, *SRY* was highly divergent (92.5% average pairwise identity) and




Figure 2 Phylogenetic trees for primate *SRY*, *DAX1*, and *SOX9*, constructed using the HKY nucleotide-substitution model (Hasegawa et al. 1985) and the commonly accepted topology. Branch lengths correlate to relative rates of evolution.




Figure 3 Amino acid alignment of primate DAX1. Amino acid identities are denoted by periods. Amino acid differences are noted.

SOX9 was very conserved (98.9% average pairwise identity) (table 1). When the overall nucleotide sequence of DAX1 was examined, it was also conserved (98.1% average pairwise identity), falling closer to SOX9 than to SRY.

We investigated the protein products of these genes at the amino acid level. We aligned the deduced DAX1 amino acid sequences for these primate species (fig. 3). The results were consistent with the nucleotide analysis. The average pairwise identities at the amino acid level were 85.8% for SRY, 96.6% for DAX1, and 99.8% for SOX9 (table 1). Inspection of the alignment suggests a clustering of amino acid changes between residues 180 and 240 of DAX1.

Proteins subjected to different functional selective pressures should accumulate amino acid differences in varying proportions relative to the silent mutations. Therefore, we calculated the frequency of nonsynonymous (K_a) and synonymous (K_s) nucleotide changes for our six primate species (Li et al. 1985) (table 2). When

K_a values for SRY, DAX1, and SOX9 were examined within individual species, it was clear that SRY had the highest values, with DAX1 intermediate and SOX9 lowest. K_s values were consistently lower for DAX1 than for SRY or for SOX9. K_a values, however, were consistently higher for DAX1 than for SOX9.

To examine the relative frequency of nonsynonymous nucleotide changes in comparison with the frequency of synonymous nucleotide changes, we calculated the K_a/K_s ratio for all six primate species (Goldman and Yang 1994). This ratio controls for different intrinsic rates of mutation focusing on the relative preference for amino acid changing substitutions. We found that DAX1 ($K_a/K_s = .43 \pm .08$) was not significantly different from SRY ($K_a/K_s = .62 \pm .11$) and was much higher than SOX9 ($K_a/K_s = .0061 \pm .0036$) (table 1).

Phylogenetic trees were constructed by means of the DAX1, SRY, and SOX9 coding sequences. PAML was performed (Goldman and Yang 1994) (fig. 2). *Homo sapiens* (human) and *P. troglodytes* (chimpanzee) were always most closely related, and *M. mulatta* (rhesus) and *C. jacchus* (marmoset) were also sister taxa. The relationship of *Hylobates* species and *P. pygmaeus* could not definitely be resolved for the two genes DAX1 and SOX9.

The shapes of the phylogenetic trees for DAX1, SRY, and SOX9, however, were distinct. We compared the tree shapes under the commonly accepted primate topology (Pilbeam 1984). Tree shape under a given topology was compared by testing whether the ratio of branch length i , for gene 1 (t_i), and branch length i , for gene 2 (s_i), is the same for all branches ($t_i/s_i = r$ for all branches i) (Goldman and Yang 1994). DAX1 and SRY did not differ ($\chi^2 = 8.10$, df 8, $P = .424$), whereas DAX1 differed from SOX9 ($\chi^2 = 16.68$, df 8, $P = .0336$), and SRY differed from SOX9 ($\chi^2 = 30.120$, df 8, $P = .0002$). Therefore, the SOX9 tree had a distinct

Table 1

Overall Measure of Sequence Divergence for Primate SRY, DAX1, and SOX9 Nucleotide Sequences

Parameter	SRY	DAX1	SOX9
Nucleotide identity ^a	92.5%	98.1%	98.9%
Amino acid identity ^a	85.8%	96.6%	99.8%
K_a/K_s ^b	.62	.43	.0061
K_a/K_s SE ^b	.11	.08	.0036
Phylogenetic tree branch length ^c	.71	.18	.11

^a Average pairwise comparisons.

^b K_a/K_s ratios estimated by means of PAML (Goldman and Yang 1994). Imposing the topologies in figure 1 gave similar results with alternative topology for DAX1.

^c The sum of branch lengths in the maximum-likelihood tree, as estimated by PAML, when the topology was imposed (Goldman and Yang 1994). Similar results were obtained with the use of an alternative topology.

Table 2 **K_a and K_s Values for Nonhuman Primate to Human *SRY*, *DAX1*, and *SOX9* Nucleotide Sequences**

Species and Measure	<i>SRY</i>	<i>DAX1</i>	<i>SOX9</i>
<i>P. troglodytes</i> :			
$K_a \times 10^2$ (95% CI)	1.44 (.35–2.53)	.47 (.05–.88)	.00 (.00–.17) ^a
$K_s \times 10^2$ (95% CI)	.82 (.00–2.46)	.30 (.00–.91)	.29 (.00–.87)
<i>P. pygmaeus</i> :			
$K_a \times 10^2$ (95% CI)	2.52 (1.05–3.98)	1.03 (.41–1.66)	.00 (.00–.17) ^a
$K_s \times 10^2$ (95% CI)	2.51 (.00–5.45)	1.83 (.33–3.33)	4.53 (2.15–6.91)
<i>Hylobates</i> :			
$K_a \times 10^2$ (95% CI)	3.56 (1.82–5.32)	1.60 (.82–2.37)	.08 (.00–.26)
$K_s \times 10^2$ (95% CI)	5.87 (1.37–10.38)	2.46 (.71–4.21)	5.98 (3.27–8.70)
<i>M. mulatta</i> :			
$K_a \times 10^2$ (95% CI)	4.63 (2.63–6.63)	1.12 (.47–1.77)	.00 (.00–.17) ^a
$K_s \times 10^2$ (95% CI)	13.22 (6.16–20.29)	2.78 (.91–4.65)	4.20 (1.91–6.49)
<i>C. jacchus</i> :			
$K_a \times 10^2$ (95% CI)	12.9 (9.44–16.43)	2.69 (1.68–3.71)	.17 (.00–.41)
$K_s \times 10^2$ (95% CI)	27.6 (16.52–38.59)	6.45 (3.53–9.34)	6.63 (3.74–9.52)

NOTE.— K_a and K_s values were estimated by means of the pairwise comparison method of Li et al. (1985).

^a Upper bound calculated assuming a maximum of two nonsynonymous substitutions.

tive shape, differing from the tree shapes for *DAX1* and *SRY*, which were similar to each other. These results were consistent with a discontinuity in the pressures driving the evolution of *SOX9* versus *SRY* and *DAX1*.

An evaluation of rates of primate *DAX1* evolution indicates that *DAX1* may be under selective pressure for its amino acid sequence to evolve at a rate faster than *SOX9*. The K_a/K_s ratio would increase when the number of nonsynonymous changes (K_a) are well tolerated or selected, relative to the number of synonymous substitutions (K_s). Using the maximum-likelihood method for estimating K_a/K_s ratios (Goldman and Yang 1994), we found that, for the six primates that we examined, *DAX1* has a K_a/K_s ratio that was statistically indistinguishable from *SRY* and was much higher than *SOX9* (table 1). Therefore, whereas *DAX1* is more highly conserved than *SRY* at the nucleotide level, the K_a/K_s ratio distinguished *DAX1* and *SOX9* as being under different evolutionary pressures. Analysis of the phylogenetic tree shapes for these three genes confirmed this distinction. Our data suggest that, although *DAX1* is a relatively conserved protein overall, it is, nevertheless, experiencing amino acid substitutions at a rate higher than is *SOX9*. We speculate that this selective pressure may be a consequence of the relatively recent evolutionary addition of *DAX1* to the sex-determination pathway (Pask et al. 1997; Lahn and Page 1999) and/or to a role in spermatogenesis (Yu et al. 1998), since male reproductive genes evolve rapidly (Wyckoff et al. 2000). The more highly conserved nature of *DAX1*, relative to *SRY*, may reflect additional constraints on *DAX1* stemming from its role in adrenal development.

DAX1 and *SRY* may be recent additions to an evolutionary ancient “core” sex-determination program

that involves *SOX9*. Our data on the molecular evolution of *SRY*, *DAX1*, and *SOX9* support this hypothesis for a stepwise evolution of the sex-determination pathway. Just as the human sex chromosomes may be evolving by punctuated sequential events (Lahn and Page 1997, 1999), our results are consistent with evolutionary stratification of the sex-determination pathway. We speculate that *SRY* and *DAX1* may be evolving dynamically as more recent additions to an evolutionarily stratified sex-determination pathway.

Acknowledgment

This work was supported, in part, by grants R01 HD22563, R01 HD39322, and P30 HD34610 from the National Institute of Child Health and Human Development.

Electronic-Database Information

Accession numbers and URLs for data in this article are as follows:

GenBank Overview, <http://www.ncbi.nlm.nih.gov/Genbank/GenbankOverview.html> (for *P. troglodytes* *SRY* [accession number X86380], *DAX1* [accession number AF322892], and *SOX9* [accession number AF322902]; *P. pygmaeus* *SRY* [accession number X86383], *DAX1* [accession number AF322893], and *SOX9* [accession number AF322898]; *Hylobates* *SRY* [accession number X86384], *DAX1* [accession number AF322894], and *SOX9* [accession number AF322897]; *M. Mulatta* *SRY* [accession number AF322901], *DAX1* [accession number AF322896], and *SOX9* [accession number AF322900]; and *C. jacchus* *SRY* [accession number X86386], *DAX1* [accession number AF322895], and *SOX9* [accession number AF322899])

Online Mendelian Inheritance in MAN (OMIM), <http://www.ncbi.nlm.nih.gov/Omim> (for adrenal hypoplasia congenita [MIM 300200], DSS [MIM 300018], SRY [MIM 480000], and campomelic dysplasia [MIM 114290])

References

- Bardoni B, Zanaria E, Guioli S, Florida G, Worley KC, Tonini G, Ferrante E, Chiumello G, McCabe ERB, Fraccaro M, Zuffardi O, Camerino G (1994) A dosage sensitive locus at chromosome Xp21 is involved in male to female sex reversal. *Nat Genet* 7:497–501
- Bi W, Deng JM, Zhang Z, Behringer RR, de Crombrughe B (1999) Sox9 is required for cartilage formation. *Nat Genet* 22:85–89
- Foster JW, Dominguez-Steglich MA, Guioli S, Kwok G, Weller PA, Stevanovic M, Weissenbach J, Mansour S, Young ID, Goodfellow PN, Brook JD, Schafer AJ (1994) Campomelic dysplasia and autosomal sex reversal caused by mutations in an SRY-related gene. *Nature* 372:525–530
- Giese K, Pagel J, Grosschedl R (1994) Distinct DNA-binding properties of the high mobility group domain of murine and human SRY sex-determining factors. *Proc Natl Acad Sci USA* 91:3368–3372
- Goldman N, Yang Z (1994) A codon-based model of nucleotide substitution for protein-coding DNA sequences. *Mol Biol Evol* 11:725–736
- Goodfellow PN, Camerino G (1999) DAX-1, an ‘antitestis’ gene. *Cell Mol Life Sci* 55:857–863
- Graves JAM, Foster JW (1994) Evolution of mammalian sex chromosomes and sex-determining genes. *Int Rev Cytol* 154:191–259
- Guo W, Mason JS, Stone CG, Jr, Morgan SA, Madu SI, Baldini A, Lindsay EA, Biesecker LG, Copeland KC, Horlick MNB, Pettigrew AL, Zanaria E, McCabe ERB (1995) Diagnosis of X-linked adrenal hypoplasia congenita by mutation analysis of the DAX1 gene. *JAMA* 274:324–330
- Habiby RL, Boepple P, Nachtigall L, Sluss PM, Crowley WF Jr, Jameson JL (1996) Adrenal hypoplasia congenita with hypogonadotropic hypogonadism: evidence that DAX-1 mutations lead to combined hypothalamic and pituitary defects in gonadotropin production. *J Clin Invest* 98:1055–1062
- Hasegawa M, Kishino H, Yano T (1985) Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. *J Mol Evol* 22:160–174
- Huang B, Wang S, Ning Y, Lamb AN, Bartley J (1999) Autosomal XX sex reversal caused by duplication of SOX9. *Am J Med Genet* 87:349–353
- Koopman P (1999) Sry and Sox9: mammalian testis determining genes. *Cell Mol Life Sci* 55:839–856
- Koopman P, Gubbay J, Vivian N, Goodfellow P, Lovell-Badge R (1991) Male development of chromosomally female mice transgenic for Sry. *Nature* 351:117–121
- Lahn BT, Page DC (1997) Functional coherence of the human Y chromosome. *Science* 278:675–679
- (1999) Four evolutionary strata on the human X chromosome. *Science* 286:964–967
- Li WH, Wu CI, Luo CC (1985) A new method for estimating synonymous and nonsynonymous rates of nucleotide substitution considering the relative likelihood of nucleotide and codon changes. *Mol Biol Evol* 2:150–174
- McBride D, Sang H, Clinton M (1997) Expression of Sry-related genes in the developing genital ridge/mesonephros of the chick embryo. *J Reprod Fertil* 109:59–63
- McCabe ERB (2000) Adrenal hypoplasias and aplasias. In: Scriver CR, Beaudet AL, Sly WS, Valle D, Childs B, Vogelstein B (eds) *The metabolic and molecular bases of inherited disease*, 8th ed. McGraw-Hill, New York
- Merke DP, Tajima T, Baron J, Cutler GB Jr (1999) Hypogonadotropic hypogonadism in a female caused by an X-linked recessive mutation in the DAX1 gene. *N Engl J Med* 340:1248–1252
- Morais da Silva S, Hacker A, Harley V, Goodfellow P, Swain A, Lovell-Badge R (1996) Sox9 expression during gonadal development implies a conserved role for the gene in testis differentiation in mammals and birds. *Nat Genet* 14:62–68
- Muscatelli F, Strom TM, Walker AP, Zanaria E, Recan D, Meindl A, Bardoni B, Guiolo S, Zehetner G, Rabl W, Schwarz HP, Kaplan J-C, Camerino G, Meitinger T, Monaco AP (1994) Mutations in the DAX-1 gene give rise to both X-linked adrenal hypoplasia congenita and hypogonadotropic hypogonadism. *Nature* 372:672–676
- Pask A, Toder R, Wilcox SA, Camerino G, Graves JAM (1997) The candidate sex reversing DAX-1 gene is autosomal in marsupials: implication for the evolution of sex determination in mammals. *Genomics* 41:422–426
- Pilbeam D (1984) The descent of hominoids and hominids. *Sci Am* 250:84–96
- Pontiggia A, Rimini R, Harley VR, Goodfellow PN, Lovell-Badge R, Bianchi ME (1994) Sex-reversing mutations affect the architecture of SRY-DNA complexes. *EMBO J* 13:6115–6124
- Schadt EE, Sinsheimer JS, Lange K (1998) Computational advances in maximum likelihood methods for molecular phylogeny. *Genome Res* 8:222–233
- Sinclair AH, Berta P, Palmer MS, Hawkin JR, Griffiths BL, Smith MJ, Foster JM, Frischauf AM, Lovell-Badge R, Goodfellow PN (1990) A gene from the human sex determining region encodes a protein with homology to a conserved DNA binding motif. *Nature* 346:240–244
- Swain A, Narvaez V, Burgoyne P, Camerino G, Lovell-Badge R (1998) Dax1 antagonizes Sry action in mammalian sex determination. *Nature* 391:761–767
- Tucker PK, Lundrigan BL (1993) Rapid evolution of the sex determining locus in Old World mice and rats. *Nature* 364:715–717
- Wagner T, Wirth J, Meyer J, Zabel B, Held M, Zimmer J, Pasantes J, Bricarelli FD, Keutel J, Hustert E, Wolf U, Tommerup N, Schempp W, Scherer G (1994) Autosomal sex reversal and campomelic dysplasia are caused by mutations in and around the SRY-related gene SOX9. *Cell* 79:1111–1120
- Whitfield LS, Lovell-Badge R, Goodfellow PN (1993) Rapid sequence evolution of the mammalian sex-determining gene SRY. *Nature* 364:713–715
- Wyckoff GJ, Wang W, Wu C-I (2000) Rapid evolution of male reproductive genes in the descent of man. *Nature* 403:304–309
- Yu RN, Ito M, Saunders TL, Camper SA, Jameson JL (1998)

Role of *Abcb* in gonadal development and gametogenesis.
Nat Genet 20:353–357
Zanaria E, Muscatelli F, Bardoni B, Strom TM, Guioli S, Guo
W, Lalli E, Moser C, Walker AP, McCabe ERB, Meitinger

T, Monaco AP, Sassone-Corsi P, Camerino G (1994) An
unusual member of the nuclear hormone receptor superfam-
ily responsible for X-linked adrenal hypoplasia congenita.
Nature 372:635–641