

The function and expansion of the Patched- and Hedgehog-related homologs in *C. elegans*

Olivier Zugasti,¹ Jeena Rajan,² and Patricia E. Kuwabara^{1,3}

¹University of Bristol, Department of Biochemistry, School of Medical Sciences, Bristol BS8 1TD, United Kingdom

The Hedgehog (Hh) signaling pathway promotes pattern formation and cell proliferation in *Drosophila* and vertebrates. Hh is a ligand that binds and represses the Patched (Ptc) receptor and thereby releases the latent activity of the multipass membrane protein Smoothed (Smo), which is essential for transducing the Hh signal. In *Caenorhabditis elegans*, the Hh signaling pathway has undergone considerable divergence. Surprisingly, obvious Smo and Hh homologs are absent whereas PTC, PTC-related (PTR), and a large family of nematode Hh-related (Hh-r) proteins are present. We find that the number of PTC-related and Hh-r proteins has expanded in *C. elegans*, and that this expansion occurred early in Nematoda. Moreover, the function of these proteins appears to be conserved in *Caenorhabditis briggsae*. Given our present understanding of the Hh signaling pathway, the absence of Hh and Smo raises many questions about the evolution and the function of the PTC, PTR, and Hh-r proteins in *C. elegans*. To gain insights into their roles, we performed a global survey of the phenotypes produced by RNA-mediated interference (RNAi). Our study reveals that these genes do not require Smo for activity and that they function in multiple aspects of *C. elegans* development, including molting, cytokinesis, growth, and pattern formation. Moreover, a subset of the PTC, PTR, and Hh-r proteins have the same RNAi phenotypes, indicating that they have the potential to participate in the same processes.

[Supplemental material is available online at www.genome.org. The following individuals kindly provided reagents, samples, or unpublished information as indicated in the paper: B. Grant, M. Labouesse, and G. Seydoux.]

The membrane protein Patched (Ptc) and its ligand, the morphogen Hedgehog (Hh), control numerous processes during embryonic development, ranging from segmentation in *Drosophila* to neural tube differentiation and axon guidance in vertebrates (Ingham and McMahon 2001; Wetmore 2003). Hh is also involved in regulating stem cell maintenance and proliferation (Taipale and Beachy 2001; Zhang and Kalderon 2001). In humans, dysfunction of Hh/Ptc signaling is associated with a variety of malignancies. Sporadic PTCH mutations can cause basal cell carcinomas and medulloblastomas (Johnson et al. 1996), and hereditary mutations are responsible for Gorlin's syndrome, which is associated with developmental abnormalities and a predisposition to develop cancer (Hahn et al. 1996).

Studies to date indicate that signal transduction mediated through Hh and Ptc requires Smoothed (Smo), a seven-pass membrane protein related to the Frizzled family of Wingless receptors (Ingham and McMahon 2001; Lum and Beachy 2004). In the absence of Hh, Ptc represses the intrinsic signaling activity of Smo. It has been proposed that Ptc regulates the transport or the distribution of an endogenous small molecule capable of influencing Smo activity (Chen et al. 2002; Taipale et al. 2002), although earlier studies indicated that Ptc repressed Smo activity via a direct protein-protein interaction (Stone et al. 1996; Murone et al. 1999). Inhibition of Smo is released when Hedgehog (Hh) binds to Ptc. In turn, the Hh signal is transduced and *Cubitus interruptus* transcriptionally activates Hh targets.

The genome of *Caenorhabditis elegans* encodes two PTC homologs, PTC-1 and PTC-3, and a pseudogene, PTC-2 (Kuwabara et al. 2000). In addition, 24 proteins with sequence and topological similarities to PTC have also been identified and named PTR (for Patched-related). The PTC and PTR proteins are distinguished by the presence of sterol sensing-domains (SSDs) (Kuwabara et al. 2000). A common feature shared by many SSD proteins is their participation in processes that variously involve the transport of lipids, sterols, or sterol-modified proteins (Kuwabara and Labouesse 2002). Despite the presence of PTC homologs, database searches of the complete *C. elegans* genome sequence have failed to identify obvious Hh or Smo homologs (The *C. elegans* Sequencing Consortium 1998), so the role of the PTC and PTR proteins in *C. elegans* is unclear. However, we previously demonstrated that PTC-1 is required in the *C. elegans* germline for cytokinesis and maintenance of autonomous germ cell development within the gonad syncytium (Kuwabara et al. 2000).

In *Drosophila* and vertebrates, Hh is a bipartite protein that follows a three-step maturation process to achieve full biological activity (Jeong and McMahon 2002). The C-terminal "Hog" domain autoproteolytically cleaves the Hh precursor to generate a secreted N-terminal "Hedge" signaling activity, which is modified at the C terminus by the addition of cholesterol and at the N terminus by palmitoylation (Pepinsky et al. 1998; Chamoun et al. 2001). The genome of *C. elegans* encodes >60 proteins that will be collectively referred to as Hedgehog-related (Hh-r) (Bürglin 1996; Aspöck et al. 1999), although they bear no direct sequence similarity with the "Hedge" of Hh (Aspöck et al. 1999). It has been proposed that the *C. elegans* Hh-r proteins are the evolutionary homologs of Hh (Aspöck et al. 1999), because some of the *C. elegans* Hh-r proteins are also bipartite, carrying a C-terminal Hog domain and an N-terminal "Hedge-related" sequence that

²Current address: The Wellcome Trust Sanger Institute, Hinxton, Cambridge, CB10 1SA, UK.

³Corresponding author.

Email p.kuwabara@bristol.ac.uk; fax: 44-117-928-8274.

Article and publication are at <http://www.genome.org/cgi/doi/10.1101/gr.3935405>.

Table 1. The *C. elegans ptc* and *ptr* genes promote molting

	N2			<i>rrf-3</i>			ESTs
	No. ^a	% Mlt	Stage ^b	No. ^a	% Mlt	Stage ^b	
control	204	0		185	0		
<i>ptc-1</i> ^c	375	0.8		200	1		11
<i>ptc-3</i> ^d	237	70.4	L2-L3	200	45	L1-L2	16
<i>ptr-1</i>	191	0		252	13.1	L4-A	0
<i>ptr-2</i> ^e	157	25	L4-A	188	32	L4-A	13
<i>ptr-3</i>	170	6	L4-A	225	8	L4-A	3
<i>ptr-4</i> ^d	235	94	L4-A	320	100	L3-L4	16
<i>ptr-5</i>	255	0		310	3.2	L4-A	12
<i>ptr-6</i>	276	0		270	11.1	L4-A	3
<i>ptr-7</i>	253	0		340	0		1
<i>ptr-8</i>	234	0		167	4.8	L4-A	9
<i>ptr-9</i>	287	0		266	0		0
<i>ptr-10</i>	236	2.5	L4-A	253	6.3	L4-A	2
<i>ptr-11</i>	237	0		265	0		5
<i>ptr-12</i>	240	0		198	4.5	L4-A	2
<i>ptr-13</i>	286	0		222	5.4	L4-A	0
<i>ptr-14</i>	258	0		360	6.4	L4-A	8
<i>ptr-15</i>	270	0		281	0		0
<i>ptr-16</i>	207	0		333	31.2	L2-L3	0
<i>ptr-17</i>	222	0		302	0		1
<i>ptr-18</i>	292	18	L4-A	262	91.6	L3-L4	5
<i>ptr-19</i>	239	0		282	0		0
<i>ptr-20</i>	245	4	L4-A	312	11.5	L4-A	4
<i>ptr-21</i>	249	0		198	9.6	L4-A	0
<i>ptr-22</i>	240	0		200	8	L3-L4	0
<i>ptr-23</i>	276	81.5	L4-A	330	94	L4-A	19
<i>ptr-24</i>	153	0		298	0		15

^aNo. represents results from at least three independent experiments.

^bThe earliest stage showing molting defects.

^cRNAi was performed on L1 stage animals to bypass the embryonic requirement for *ptc-1* activity (Kuwabara et al. 2000).

^d100% of *ptc-3(RNAi)* and *ptr-4(RNAi)* animals subsequently die.

^e90% of *ptr-2(RNAi)* animals display embryonic lethality (Emb); hence, only animals surviving this lethality were scored.

could have signaling potential (Aspöck et al. 1999). In addition, *C. elegans* encodes a number of proteins consisting only of the secreted Hedge-related domain and a single protein with only a Hog domain. The Hedge-related domains have been further divided into families based on sequence: *warthog* (*wrt*), *groundhog* (*grd*), *ground-like* (*grl*), and *quahog* (*qua*) (Bürglin 1996; Aspöck et al. 1999).

Given the importance of Hh signaling in other organisms, the absence of obvious Hh and Smo homologs in *C. elegans* has raised numerous questions about the roles played by the *C. elegans* PTC, PTR, and Hh-r proteins in development. To gain insights into the functions of these genes, we surveyed the phenotypes produced by RNA-mediated interference (RNAi) of the *ptc*, all 24 *ptr*, and 27 *hh-r* genes, and we show that these genes are functional and participate in multiple aspects of *C. elegans* development, despite the absence of Smo. We find that some of the *ptc*, *ptr*, and *hh-r* genes participate in the same processes, which share in common the need to transport proteins, lipids, or sterols. Although most of the *ptr* genes are individually capable of developing an RNAi phenotype; we demonstrate that some members share partial functional redundancy. We show that the expansion of the *ptr* family appears to be a feature of Nematoda as *Drosophila*, mouse, and human each have only a single *ptr* gene. Moreover, the activities of the *ptc* and *ptr* genes are conserved between *C. elegans* and *Caenorhabditis briggsae*. Finally, we provide evidence that some of the *ptr* genes participate in pro-

cesses requiring sterols and the trafficking of proteins, sterols, and lipids.

Results

The *ptc* and *ptr* genes promote molting

The phenotypes produced by inhibiting the activities of the *ptc-1*, *ptc-3*, and the 24 *ptr* genes were surveyed by RNAi using the wild-type N2 Bristol and *rrf-3*, a mutant hypersensitive to RNAi (Simmer et al. 2002, 2003). Table 1 reveals that RNAi inhibition of many of these genes produces a Mlt (molting-defective) phenotype. Mlt animals have problems in shedding old cuticle, which is the collagenous outer layer of the hypodermis (Fig. 1). Thus, animals retaining remnants of old cuticle, displaying circumferential constrictions, or dying from starvation because they are encapsulated in old cuticle are scored as Mlt.

When RNAi was repeated by using *rrf-3*, the number of *ptr* genes generating a Mlt phenotype more than doubled; we found that *ptc-1*, *ptc-3*, and 17 *ptr* genes all produced a Mlt phenotype (Table 1). For most genes, the use of *rrf-3* also increased the penetrance of the Mlt phenotype and made it possible to detect Mlt worms earlier in development (Table 1). However, *rrf-3* mutants are not Mlt (Table 1). It is interesting to note that the absence of *lrp-1/megalyn*, which encodes a receptor required for cholesterol endocytosis and is also capable of endocytosing Sonic Hh, also produces a similar RNAi Mlt phenotype (Yochem et al. 1999; McCarthy et al. 2002).

Redundancy in the activity of *ptr* genes

Large gene families can contain members that are functionally redundant (Prince and Pickett 2002). To determine whether this is true of the *ptr* family, we selected six genes—*ptr-1*, *-6*, *-10*, *-16*, *-20*, and *-21*—that individually develop a weak RNAi Mlt phenotype in *rrf-3* (Table 1), and we systematically analyzed the effect of simultaneously disrupting combinations of two or three of these genes by RNAi. Of the 35 possible combinations, the simultaneous RNAi disruption (combi-RNAi) of *ptr-1:ptr-16* or *ptr-1:ptr-6:ptr-10* had a striking synergistic effect; the severity and the percentage of animals displaying a Mlt phenotype was enhanced (Fig. 2). Thus, some members of the *ptr* gene family are partially redundant in function.




Figure 1. Molting defects (Mlt) associated with *ptr(RNAi)*. (A–C) Examples of Mlt defects (arrowheads) associated with *ptr-5*, *ptr-4*, and *ptr-21(RNAi)*, respectively. Scale bar, 50 μ m.




Figure 2. The *ptr* genes are partially redundant. All 35 double and triple combinations of *ptr-1*, *-6*, *-10*, *-16*, *-20*, and *-21* were subjected to RNAi in *rrf-3*; however, only the single disruptions and those combinations showing synergistic phenotypic enhancement are shown. Gray bar indicates Mlt animals; black bar, dead animals.

The *ptr* genes promote growth and cell patterning in hermaphrodites and males

The *ptr* genes not only are involved in molting but also are active in *C. elegans* growth and morphogenesis (Fig. 3). RNAi disruption of 15 *ptr* genes impairs cell growth and produces Sma (small) animals that are up to 40% shorter than same-stage N2 animals (Fig. 3A,E). Inhibition of four *ptr* genes also leads to vulval defects (Fig. 3B,E).

The genome-wide RNAi screens, performed so far in *C. elegans*, have focused on elucidating the RNAi phenotypes produced by hermaphrodites and not by males. To explore whether members of the *ptr* family have male-specific functions, we inhibited all of the genes listed in Table 1 by RNAi using the male producing strains, *him-8* or *him-8;rrf-3*. Inhibition of *ptr-1*, *-2*, *-4*, *-18*, *-20*, and *-23* by RNAi produced male molting defects (Supplement 1). Disruption of these genes also caused defects in the patterning of the spicules, fan, and sensory rays of the male tail; in the most severe cases, these structures were absent (Supplement 1; data not shown). Thus, at least six *ptr* genes promote male tail patterning.

ptr-2 is involved in somatic cytokinesis

ptr-2(RNAi) produces a phenotype that is markedly distinct from that of all other *ptr* genes. Disruption of *ptr-2* by RNAi causes embryos to arrest with multinucleated cells arising from a probable defect in somatic cytokinesis (data not shown). However, animals that escape from embryonic lethality (Emb) are also Mlt (Table 1). Similarly, *ptc-1*(RNAi) embryos arrest from defects occurring during germline cytokinesis (Kuwabara et al. 2000); however, when *ptc-1*(RNAi) is performed on post-embryonic animals, animals can also display molting defects (Table 1). It should be noted that *ptc-1*(RNAi) animals do not display defects in somatic cytokinesis.

Disruption of *ptr* gene activity impairs exocytosis and adult alae development

A re-examination by Nomarski DIC revealed that RNAi inhibition of 18 *ptr* genes caused the accumulation of abnormal fluid-filled vacuoles in the intestine and hypodermis (Fig. 3C,E). These

animals were also Unc because of their tendency to curl and to become immobile as adults, although they remained sensitive to touch. This immobile Unc phenotype strongly resembles that displayed by *sec-23* (yeast *SEC* homolog) and *lrp-1* mutants, which are known to be deficient in vesicle trafficking (Yochem et al. 1999; Roberts et al. 2003). A second, apparently unrelated Unc phenotype was also detected, whereby animals generate sinusoidal body waves but fail to progress forward or backward; this defect is probably caused by tears in the cuticle (data not shown). In addition, disruption of three *ptr* genes led to the formation of cuticular blisters (Bli) that might result from defects in the structure of the cuticle or the failure to secrete proteases that process collagen precursors (Fig. 3D,E; Thacker et al. 1995).

CHE-14, the *C. elegans* homolog of the SSD protein Disp, promotes the exocytosis of proteins from the hypodermis; animals lacking *che-14* display severe defects in the formation of alae (Michaux et al. 2000). The adult alae are secreted by seam cells and are composed of three raised cuticular ridges that form lateral stripes above the seam cell cords. Given the potential involvement of many *ptr* in trafficking, we asked whether the *ptr* genes could affect alae formation. We found that disruption of six *ptr* genes by RNAi led to the generation of four classes of defective alae: (1) multiple alae composed of four instead of three cuticular ridges, (2) discontinuous alae, (3) bifurcated alae, and (4) multiply branched alae (Fig. 4). We propose that similar to *che-14*, the defects in alae development resulting from *ptr*(RNAi) might be caused by an impairment in protein secretion (Michaux et al. 2000). Hence, it is possible that the subset of *ptr* genes affecting alae formation could have properties that are more similar to that of *che-14/Disp* than to *ptc*.




Figure 3. Disruption of *ptc* and *ptr* genes by RNAi leads to growth and developmental defects. (A) A *ptr-4*(RNAi) adult hermaphrodite (bottom) is 30% shorter than is a similarly staged wild-type hermaphrodite (top). (B) An everted vulva in a *ptr-11*(RNAi) animal. (C) *ptr-1*(RNAi) animals accumulate fluid-filled vacuoles in the hypodermis. (D) Blisters caused by *ptr-3*(RNAi). (E) Summary of phenotypes produced by RNAi, not including molting. All the *ptc* and *ptr* genes were examined; however, only those showing RNAi phenotypes are displayed. Scale bar, 50 μ m.




Figure 4. Elimination of *ptr* or *hh-r* gene activity by RNAi causes defects in adult alae formation. (A) Wild-type alae in N2. (B–D) Defective alae produced by RNAi of *ptr* or *hh-r* gene members. (B) Discontinuous alae (arrowheads). (C) Bifurcated alae. (D) Multiply branched alae. (C, D) Multiple alae composed of four instead of three cuticle ridges. (E) Summary of alae defects after RNAi of *ptr* or *hh-r* genes. All genes displayed in Tables 1 and 2 were examined; however, only those showing phenotypes are displayed. Scale bar, 10 μ m.

Inhibition of *ptr-4*, *ptr-18*, and *ptr-23* disrupts endocytosis

To explore the potential involvement of the *ptr* genes in trafficking, we asked whether disruption of the *ptr* genes inhibits endocytosis. To address this question, we used an assay developed by Grant and Hirsh (1999) that examines the ability of oocytes to endocytose a fluorescent yolk protein reporter, YP170::GFP. Normally, the yolk protein vitellogenin is secreted by the intestine and endocytosed by oocytes proximal to the spermatheca (Grant and Hirsh 1999). We found that the ability of oocytes to endocytose the YP170::GFP fusion protein is compromised in adult *ptr-4*(RNAi) animals; instead, ectopic vacuoles of yolk collect in the body cavity (Fig. 5). A milder form of this defect is present in *ptr-18*(RNAi) and *ptr-23*(RNAi) animals, which form ectopic vacuoles of yolk (data not shown).

The *hh*-related *ptc*, and *ptr* genes have similar RNAi phenotypes

In *Drosophila* and vertebrates, Ptc interacts directly with Hh to control cell patterning and proliferation (Incardona et al. 2000). To investigate whether the *C. elegans* Hh-r proteins have the potential to participate in similar activities as the PTC and PTR proteins, we performed a global survey of the RNAi phenotypes produced by a subset of 27 *hh-r* genes (Table 2). We observed that the disruption of these genes by RNAi produces phenotypes very similar to that seen with *ptc-3*(RNAi) and *ptr*(RNAi) (Table 2; Fig. 4), including defects in molting, growth, and vulval and male tail morphogenesis (Table 2, Supplement 2). In a few cases involving individual members of three paralogous gene pairs—*grd-3/grd-4*, *grd-5/grd-10*, and *grd-13/grd-14*—that share >70% identity at the nucleotide level, RNAi phenotypes are likely to arise from reducing the activity of both paralogs.

Expression patterns of the *ptc*, *ptr*, and *hh-r* genes

Expression profile analysis performed on selected *hh-r* genes revealed that they are expressed in a variety of tissues, including the hypodermis, seam cells, neurons, and neuron-associated cells (Aspöck et al. 1999). Because the *ptc*, *ptr* and *hh-r* genes share

similar RNAi phenotypes, we sought to determine if they were expressed in the same tissues by examining serial analysis of gene expression (SAGE) data obtained from the Genome BC *C. elegans* Gene Expression Consortium (<http://elegans.bcgsc.bc.ca/perl/sage>). Although there was some diversity in their expression profiles, most members of the *ptc*, *ptr*, and *hh-r* genes are expressed in the same cells (Supplement 3). In particular, the transcripts of *ptc-3*, eight *ptr*, and 10 of 27 *hh-r* genes under examination are enriched in hypodermal cells, the site where cuticle components are synthesized and secreted between molts (Supplement 3) (Johnstone 2000). We were unable to find additional expression data in the *C. elegans* fusion library database (<http://nematode.bsgsc.ca>) or the Nematode Expression Pattern Database (NEXTDB) (<http://nematode.lab.nig.ac.jp>). Taken together, the similarities between the RNAi phenotypes and the SAGE profiles allow for the possibility that the PTC, PTR, and Hh-r proteins act in the same pathways.

Expansion of the *ptr* gene family in *C. elegans*

The SSD family of proteins control multiple aspects of cholesterol biosynthesis and homeostasis and also participate in processes involving the transport of proteins and lipids (Kuwabara and Labouesse 2002). We previously showed that *C. elegans* encodes a seventh family of SSD proteins, consisting of the 24 *ptr* genes (Kuwabara et al. 2000). Given that complete genome sequences for *Drosophila*, mouse, and human were not available at the time of our initial analysis (Adams et al. 2000; Lander et al. 2001; Waterston et al. 2002), we performed BLAST analysis to identify all SSD-containing proteins encoded by these genomes, and aligned the sequences obtained by using Clustal W (Fig. 6) (see Experimental Procedures). Similar to the results obtained for *C. elegans*, we found that the SSD proteins cluster into seven families, although the genomes of fly, mouse, and human each encode only a single PTR homolog. These results indicate that




Figure 5. *ptr-4*(RNAi) disrupts endocytosis. (A–C) A YP170::GFP reporter is expressed in the intestine and endocytosed by the proximal-most oocyte (arrowheads) in wild-type hermaphrodites, as visualized by GFP fluorescence in *C. elegans*. (D–F) *ptr-4*(RNAi) hermaphrodites inefficiently localize YP170::GFP to oocytes (arrowheads). Panels A, B, D, and E are Nomarski DIC micrographs; panels C, F show YP170::GFP fluorescence. Scale bar, 50 μ m.

Table 2. RNAi phenotypes produced by a subset of *hh-r* genes

RNAi	N2			<i>rrf-3</i>			Growth defect	Fluid filled	Unc. Immobile	Blister	Everted vulva	Male tail defect
	n ^a	%Mlt	Stage ^b	n ^a	%Mlt	Stage ^b						
<i>wrt-1</i>	240	0		144	6.9	L4-A	■	■	■	■	■	■
<i>wrt-2</i>	183	0		174	4	L4-A	■	■	■	■	■	■
<i>wrt-3</i>	265	0		192	0		■	■	■	■	■	■
<i>wrt-4</i> ^c	159	7.5	L4-A	354	9.3	L3-L4	■	■	■	■	■	■
<i>wrt-5</i>	278	0		170	7.6	L4-A	■	■	■	■	■	■
<i>wrt-6</i>	248	0		264	12.1	L3-L4	■	■	■	■	■	■
<i>wrt-7</i> ^c	280	0		187	0		■	■	■	■	■	■
<i>wrt-8</i>	180	0		164	0		■	■	■	■	■	■
<i>wrt-9</i>	231	0		172	0		■	■	■	■	■	■
<i>wrt-10</i>	262	0		144	0		■	■	■	■	■	■
<i>grd-1</i>	242	0		201	11.4	L4-A	■	■	■	■	■	■
<i>grd-2</i>	187	0		189	3.7	L4-A	■	■	■	■	■	■
<i>grd-3</i>	288	0		280	10.7	L4-A	■	■	■	■	■	■
<i>grd-4</i>	251	0		303	8.2	L4-A	■	■	■	■	■	■
<i>grd-5</i>	308	0		308	5.8	L4-A	■	■	■	■	■	■
<i>grd-6</i>	274	0		154	0		■	■	■	■	■	■
<i>grd-7</i>	268	0		298	8.7	L4-A	■	■	■	■	■	■
<i>grd-8</i>	334	0		298	9.7	L4-A	■	■	■	■	■	■
<i>grd-9</i>	168	3.6	L4-A	312	4.8	L3-L4	■	■	■	■	■	■
<i>grd-10</i>	257	0		192	11.4	L3-L4	■	■	■	■	■	■
<i>grd-11</i>	222	0		207	0		■	■	■	■	■	■
<i>grd-12</i>	204	0		226	0		■	■	■	■	■	■
<i>grd-13</i>	175	0		215	0		■	■	■	■	■	■
<i>grd-14</i>	286	0		302	0		■	■	■	■	■	■
<i>grd-15</i>	232	0		209	0		■	■	■	■	■	■
<i>qua-1</i>	150	83.3	L3-L4	208	100	L1-L2	■	■	■	■	■	■
<i>hog-1</i>	212	0		144	6.9	L4-A	■	■	■	■	■	■

^an represents results from at least three independent experiments.

^bThe earliest stage showing molting defects.

^cAnimals also have a low penetrant multivulva phenotype.

the *ptr* genes are conserved across phyla, although the family has expanded in *C. elegans* (Fig. 6).

ptr gene expansion is a feature of Nematoda

The finding that the *ptr* genes are active during multiple aspects of *C. elegans* development and that their functions are not strictly redundant led us to ask whether they are present in other nematodes. We first assessed the orthology between the SSD proteins of *C. elegans* and *C. briggsae*, a rhabditid with a complete genome sequence that diverged from a common ancestor of *C. elegans* between 60 and 100 millions years ago (Stein et al. 2003). This was achieved by assigning each full-length *C. elegans* SSD protein a *C. briggsae* ortholog partner by reciprocal best BLAST (Stein et al. 2003). Second, we compared the syntenic conservation of genes flanking either side of each SSD encoding gene in *C. elegans* and *C. briggsae* (Kuwabara and Shah 1994). According to these criteria, all 31 *C. elegans* SSD proteins have a direct *C. briggsae* ortholog (Table 3; data not shown). Moreover, 23 of the 31 orthologous proteins share >80% amino acid identity (Table 3; Stein et al. 2003).

To ask whether PTC and PTR proteins are conserved in non-rhabditid branches of Nematoda, we searched the *Brugia malayi* genome sequence repository (<http://www.tigr.org/tdb/parasites/>); *B. malayi* is a filarial parasite and a member of the Spirurida order (Blaxter et al. 1998). Reciprocal best BLAST searches using predicted full-length *C. elegans* PTC and PTR sequences show that *B. malayi* encodes proteins with strong homology to *C. elegans* PTC-1, PTC-3, and at least 16 distinct PTR proteins (Supplement 4). We also searched for sequences encoding PTC and PTR proteins from the 3334 ESTs available for the nematode *Meloidogyne arenaria*, a plant parasite and member of the Tylenchida order (<http://www.nematode.net>) (Blaxter et al. 1998). Reciprocal best

BLAST analysis led to the identification of one PTC protein and at least two *M. arenaria* proteins sharing strong homology to PTR-2 and PTR-4 (data not shown). Thus, the presence of multiple *ptr* genes in nonrhabditid clades indicates that the expansion of *ptr* genes is a feature of Nematoda. However, assignments of orthology remain tentative because there is insufficient sequence data available to analyze syntenic associations, except in the case of *C. briggsae*.

Conservation of function between *C. elegans* and *C. briggsae* *ptr* and *ptr* genes

To provide evidence that the functions of the *ptr* and the *ptr* genes are conserved between *C. elegans* and *C. briggsae*, we disrupted the activities of the *C. briggsae* orthologs of *ptr-1*, *ptr-4*, *ptr-18*, and *ptr-23* by RNAi. Although the RNAi response in *C. briggsae* appeared to be less robust than in *C. elegans*, similar RNAi phenotypes were observed. Disruption of the *C. briggsae* *ptr-1* ortholog led to the appearance of multinucleate germ cells arising from a probable defect in germline cytokinesis, and molting defects were observed in *Cb-ptr-4(RNAi)*, *Cb-ptr-18(RNAi)*, and *Cb-ptr-23(RNAi)* animals (Kuwabara et al. 2000) (Supplement 5; data not shown). These results indicate that gene activity appears to be conserved between orthologous gene pairs.




Figure 6. Sterol sensing domain (SSD) proteins and their phylogeny. Full-length SSD proteins can be clustered into seven major families by Clustal W: HMG-CoAr, SCP, 7-DHCR, Disp, NPC, PTC, and PTR (Thompson et al. 1997, Kuwabara et al. 2000). Sequence prefix denotes species: *C. elegans* (Ce), *D. melanogaster* (Dm), *M. musculus* (Mm), and *H. sapiens* (Hs). GenBank accession numbers are available in the Methods section.

Table 3. Orthologous pairing of *C. elegans* and *C. briggsae* SSD proteins

Gene	Sequence name		Length (amino acids)		% Id	E-value
	<i>C. elegans</i>	<i>C. briggsae</i>	<i>C. elegans</i>	<i>C. briggsae</i>		
<i>ptc-1</i>	ZK675.1	CBG03294	1405	1402	89	0
<i>ptc-3</i>	Y110A2AL.8	CBG19582	1388	1380	86	0
<i>ptr-1</i>	C24B5.3	CBG19221	956	951	89	0
<i>ptr-2</i>	C32E8.8	CBG12004	933	933	93	0
<i>ptr-3</i>	C41D7.2	CBG06961	914	926	87	0
<i>ptr-4</i>	C45B2.7	CBG14821	960	771	70	0
<i>ptr-5</i>	C53C11.3	CBG15941	961	946	88	0
<i>ptr-6</i>	C54A12.1	CBG02358	1015	974	90	0
<i>ptr-7</i>	F31F6.5	CBG07607	955	915	86	0
<i>ptr-8</i>	F44F4.4	CBG20645	890	888	91	0
<i>ptr-9</i>	F54G8.5	CBG06803	413	410	85	4.0E - 191
<i>ptr-10</i>	F55F8.1	CBG14920	870	896	84	0
<i>ptr-11</i>	F56C11.2	CBG05104	820	819	87	0
<i>ptr-12</i>	K07A3.2	CBG04166	958	898	73	0
<i>ptr-13</i>	K07C10.1	CBG02953	690	689	84	3.8E - 319
<i>ptr-14</i>	R09H10.4	CBG04434	877	879	92	0
<i>ptr-15</i>	T07H8.6	CBG19049	840	840	89	0
<i>ptr-16</i>	T21H3.2	CBG01094	881	807	80	0
<i>ptr-17</i>	Y18D10A.7	CBG07932	817	922	70	0
<i>ptr-18</i>	Y38F1A.3	CBG02766	858	798	83	0
<i>ptr-19</i>	Y39A1B.2	CBG18310	1003	1004	94	0
<i>ptr-20</i>	Y53F4B.28	CBG03163	860	860	91	0
<i>ptr-21</i>	Y65B4BR.3	CBG05116	909	908	89	0
<i>ptr-22</i>	Y80D3A.7	CBG20322	859	833	76	0
<i>ptr-23</i>	ZK270.1	CBG19753	983	982	95	0
<i>ptr-24</i>	F46G10.5	CBG07780	765	830	68	1.9E - 319
<i>ncr-1</i>	F02E8.6	CBG14473	1383	1382	88	0
<i>ncr-2</i>	F09G8.4	CBG16653	1274	1222	65	0
<i>che-14</i>	F56H1.1	CBG12874	917	915	91	0
<i>ptd-2</i>	F07C3.1	CBG19582	936	871	64	0
<i>scp-1</i>	D213.8	CBG20225	1087	1035	77	0

Discussion

Homologs of the Hh signaling pathway in *C. elegans*

The Hh pathway has undergone extensive divergence in *C. elegans*, so we performed an RNAi survey to understand how the >60 Hh-r, 24 PTC-related proteins, and two PTC proteins function in the absence of Smo and Hh. Because the PTC and PTR proteins share overlapping functions, they will be discussed as a unit, although Clustal W analysis indicates that they can be subdivided into distinct SSD classes. In particular, *ptc-1* and *ptr-2* promote cytokinesis, and the *ptc-1*, *-3*, and most *ptr* genes are involved in molting (Kuwabara et al. 2000).

Our global survey of the RNAi phenotypes of the *C. elegans* *ptc*, *ptr*, and *hh-r* genes reveals for the first time that they share in common an involvement in cell growth, patterning, and molting, a process that is dependent on the availability of sterols. Molting plays an essential role in *C. elegans* growth and proliferation. This process involves the formation and degradation of the protective outer cuticle and depends on the periodic synthesis and secretion of proteins, such as collagens and proteolytic enzymes (Johnstone 2000). Below, we discuss how these processes provide clues to the roles played by the *ptc*, *ptr*, and *hh-r* genes in *C. elegans*. It should be noted that RNAi does not always result in a complete elimination of gene activity. Hence, it is possible that the *ptc*, *ptr*, and *hh-r* genes might have additional functions that will only be uncovered after null mutants corresponding to these genes are obtained.

Molting: The sterol connection

C. elegans is a cholesterol auxotroph that obtains sterols from exogenous sources. Cholesterol starvation or inhibition of *lrp-1*, which encodes a receptor with similarity to megalin and is essential for cholesterol endocytosis, leads to growth and molting defects (Yochem et al. 1999). Although *Drosophila* and *C. elegans* require sterols for viability, their sterol content is so low that it has been suggested that its primary role might be in signaling and not in membrane architecture (Kurzychalia and Ward 2003). In *Drosophila*, molting is controlled by the steroidal hormone 20-hydroxyecdysone (Riddiford 1993; Thummel 1996). In *C. elegans*, a sterol-derived hormone regulates dauer larvae formation and molting (Matyash et al. 2004). It was recently shown that the SSD of SCAP can bind directly to cholesterol (Radhakrishnan et al. 2004). Given that members of the SSD protein family collectively promote the transport of proteins, sterols and sterol-modified proteins, such as Hh, this raises the possibility that the PTC and PTR proteins play a similar role in molting (Kuwabara and Labouesse 2002). SCAP is also able to form a homotetramer; if PTC and PTR have the ability to form multimers, this could help to explain the partial functional redundancy of the PTR proteins.

Molting and protein transport

Protein transport and secretion play central roles in molting. Mutations in genes involved in protein secretion and transport, such as *rme-8*, which is required for receptor-mediated endocytosis, and *sec-23*, which is involved in ER-to-Golgi transport and encodes a component of COPII (coat protein complex II)-coated vesicles, impair molting (Yochem et al. 1999; Zhang and Kalderon 2001; Roberts et al. 2003). Multiple lines of evidence indicate that the molting defects caused by *ptc(RNAi)* and *ptr(RNAi)* could be caused by defects in protein, lipid, or sterol transport. First, the topology of the PTC and PTR proteins is similar to that of transporters, specifically those belonging to the resistance/nodulation/cell division (RND) family of prototypic bacterial multidrug efflux pumps (Paulsen et al. 1996; Kuwabara et al. 2000). Second, *ptc-1* and *ptr-2* are defective in cytokinesis because they fail to form or maintain a membranous cleavage furrow (Kuwabara et al. 2000; Strickland and Burgess 2004). Third, *ptr(RNAi)* disrupts alae formation, which is probably dependent on protein exocytosis, as demonstrated for *che-14* (Michaux et al. 2000). Finally, disruption by RNAi of a subset of *ptr* genes perturbs the endocytosis of the yolk protein vitellogenin.

The Hh-r proteins and their relationships with PTC and PTR proteins

The *hh-r*, *ptc*, and *ptr* genes not only promote molting but also affect organismal growth and post-embryonic patterning. Why does depletion of *hh-r* genes by RNAi produce a phenotype similar to that of *ptc* and *ptr(RNAi)*? Here we discuss hypotheses that are open to future investigation. First, given that many of the Hh-r proteins are likely to be secreted, there is a possibility that they could promote cell-to-cell signaling through an interaction with PTC or PTR proteins (Bürglin 1996; Aspöck et al. 1999). If such a receptor/ligand relationship were to exist, which at this time is purely speculative, then genetic arguments postulate that the Hh-r proteins are likely to activate and not inhibit the PTC and PTR proteins because they share similar phenotypes. By contrast, in all other organisms studied so far, Hh binding inhibits Ptc activity and promotes signal transduction via Smo (Ingham and McMahon 2001; Lum and Beachy 2004). Whether the PTC, PTR, and Hh-r proteins could then activate a signaling cascade is also unknown. However, if this were the case, it would be necessary to speculate that PTC and PTR could directly mediate signal transduction or function through a membrane protein other than Smo.

Second, in *Drosophila* and vertebrates, Hh proteins undergo a maturation process involving autoproteolysis, closely followed by the addition of cholesterol and lipid (Porter et al. 1996b). Similarly, it has been shown that the *C. elegans* Hh-r protein, WRT-1, undergoes autoproteolysis and has the potential to be modified by lipid and/or cholesterol (Porter et al. 1996a,b; Aspöck et al. 1999). Thus, it is possible that the PTC and the PTR proteins transport the lipid or cholesterol, which is necessary to facilitate these modifications. Third, the PTC and PTR proteins could be involved in the transport or endocytosis of Hh-r proteins. Fourth, the alae defects associated with RNAi of some *ptr* indicate that they could be mediating exocytosis as does CHE-14. Similar to Disp and Hh, the PTR proteins might be needed to exocytose the Hh-r proteins away from the cell from where they were synthesized (Michaux et al. 2000). In *Drosophila*, *Disp* and *Hh* mutants also share similar mutant phenotypes (Burke et al. 1999).

Finally, it should be stressed that there is no evidence directly linking the functions of the PTC and PTR proteins with those of the Hh-r proteins. It is equally possible that the activities of the PTC and PTR proteins are completely independent of any interaction with the Hh-r proteins. What it is clear, however, is that the Hh-r, PTC, and PTR proteins have functions that are independent of Smo, which might eventually be detected in other organisms. These activities could represent ancestral activities, such as the involvement of PTC-1 and PTR-2 in cytokinesis (Kuwabara et al. 2000; Skop et al. 2004).

Conclusions

The Hh signaling pathway, as deduced from studies in *Drosophila* and vertebrates, has undergone extensive evolutionary divergence in *C. elegans*. A number of key members of the pathway are absent, such as Hh and Smo, whereas other members, such as the Ptc, Ptc-related, and Hh-r proteins, are present and have also undergone an expansion in number. Moreover, the *C. elegans* TRA-1 ortholog of the zinc-finger transcription factor Ci/Gli, which is the terminal regulator of the Hh pathway, is used instead to control nematode sex determination (Hodgkin 1983; Mathies et al. 2004).

It remains unclear whether *C. elegans* ever had a complete Hh signaling pathway, or whether it was lost at some point in nematode evolution. An acceptable resolution of this question will depend, in part, on the outcome of an open debate regarding the robustness of the ecdysozoa clade assignment, which includes the molting animals *Drosophila* and *C. elegans* (Aguinaldo et al. 1997; Wolf et al. 2004). If nematodes, such as *C. elegans*, are indeed members of ecdysozoa, then there is a strong likelihood that most members of the Hh signaling pathway were present in both *Drosophila* and nematodes, but were subsequently lost in nematodes.

Methods

BLAST analysis and protein phylogeny

Sequences similarity searches were performed by using BLAST (Altschul et al. 1997). Cluster analysis was performed by using Clustal X, a Windows interface for the Clustal W multiple sequence alignment program (<ftp://ftp-igbmc.u-strasbg.fr/pub/ClustalX/>) (Thompson et al. 1997). Phylogenetic trees were constructed by using TreeView (<http://taxonomy.zoology.gla.ac.uk/rod/treeview.html>) (Page 1996). A total of 1000 bootstraps were run to assess the significance of branching order. *C. elegans* and *C. briggsae* sequences are available in Wormbase (<http://www.wormbase.org/>). Other nematode sequences are available from <http://www.tigr.org/tdb/> (The Institute for Genome Research [TIGR]). GenBank accession numbers for non-nematode sequences are as follows: DsHMGCR (P14773), MmHMGCR (XP_127496), HsHMGCR (AAH33692), DmSCAP (NP_788277), MmSCAP (NP_001001144), HsSCAP (Q12770), Mm7DHCR (AAH06854), Hs7DHCR (AAD02816), DmDisp (NP_524734), MmDispA (NP_081142), MmDispB (NP_733481), HsDispA (NP_116279), HsDispB (NP_277045), DmNPC (NP_609357), MmNPC-1 (NP_997125), HsNPC1L1 (AAR97886), HsNPC-1 (NP_000262), DmPTC-1 (AAA28696), MmPTC-1 (NP_032983), MmPTC-2 (NP_032984), HsPTC-1 (NP_000255), HsPTC-2 (NP_003729), DmPTR (NP_610209), MmPTR (XP_109751), and HsPTR (XP_370541).

C. elegans methods

General methods for the handling and maintenance of *C. elegans* are as previously described (Brenner 1974). The wild-type reference strain is N2 Bristol. The following mutants and transgenic lines were used: *rrf-3* (*pk1426*) (Simmer et al. 2002), *him-8* (*e1489*), and *him-5;vit-2::GFP* (Grant and Hirsh 1999). The *him-8* (*e1489*); *rrf-3* (*pk1426*) double mutant was generated by using conventional genetic techniques.

RNAi constructs and application

PCR fragments were generated by using primers (sequences available on request from investigators of this study) capable of amplifying ~0.8 kb of exon-rich DNA sequence, and were cloned into the L4440 RNAi feeding vector (Timmons et al. 2001). Although the *ptc* and *ptr* genes encode protein homologs, in most cases it is unlikely that RNAi is eliciting cross-reactivity or secondary transitive effects because the regions affected by RNAi share <65% identity at the nucleotide level. The only possible exceptions include, *ptr-3/ptr-12*, which share a short (101/120-nucleotide) region of shared identity upstream of the target sequence and the known paralogs, *grd-5/grd-10*, *grd-3/grd-4*, and *grd-13/grd-14*, which share >70% identity at the nucleotide level. All RNAi feeding experiments were performed essentially as described by Timmons et al. (2001).

Acknowledgments

We thank Barth Grant for generously providing the *him-5;vit-2::GFP* reporter strain, Michel Labouesse and Geraldine Seydoux for sharing unpublished results, and Thomas Bürglin for discussion. We also thank Jonathan Hodgkin, Karen Yook, and members of the Kuwabara laboratory for discussion and comments on the manuscript. SAGE data were produced at the Michael Smith Genome Sciences Centre with funding from Genome Canada. Some nematode strains used in this work were provided by the *Caenorhabditis* Genetics Center, which is funded by the NIH National Center for Research Resources (NCRR). P.E.K. is a recipient of a Senior Non-Clinical Fellowship funded by the Medical Research Council.

References

- Adams, M.D., Celnicker, S.E., Holt, R.A., Evans, C.A., Gocayne, J.D., Amanatides, P.G., Scherer, S.E., Li, P.W., Hoskins, R.A., Galle, R.F., et al. 2000. The genome sequence of *Drosophila melanogaster*. *Science* **287**: 2185–2195.
- Aguinaldo, A.M., Turbeville, J.M., Linford, L.S., Rivera, M.C., Garey, J.R., Raff, R.A., and Lake, J.A. 1997. Evidence for a clade of nematodes, arthropods and other moulting animals. *Nature* **387**: 489–493.
- Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D.J. 1997. Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Res.* **25**: 3389–3402.
- Aspöck, G., Kagoshima, H., Niklaus, G., and Bürglin, T.R. 1999. *Caenorhabditis elegans* has scores of hedgehog-related genes: Sequence and expression analysis. *Genome Res.* **9**: 909–923.
- Blaxter, M.L., De Lay, P., Garey, J.R., Lin, L.X., Schedelman, P., Vierstraete, A., Vanfleteren, J.R., Mackley, L.Y., Dorris, M., Frisse, L.M., et al. 1998. A molecular evolutionary framework for the phylum Nematoda. *Nature* **392**: 71–75.
- Brenner, S. 1974. The genetics of *Caenorhabditis elegans*. *Genetics* **77**: 71–94.
- Bürglin, T.R. 1996. Warthog and groundhog: Novel families related to hedgehog. *Curr. Biol.* **6**: 1047–1050.
- Burke, R., Nellen, D., Bellotto, M., Hafen, E., Senti, K.A., Dickson, B.J., and Basler, K. 1999. Dispatched, a novel sterol-sensing domain protein dedicated to the release of cholesterol-modified hedgehog from signaling cells. *Cell* **99**: 803–815.
- The *C. elegans* Sequencing Consortium. 1998. Genome sequence of the nematode *C. elegans*: A platform for investigating biology. *Science* **282**: 2012–2018.
- Chamoun, Z., Mann, R.K., Nellen, D., von Kessler, D.P., Bellotto, M., Beachy, P.A., and Basler, K. 2001. Skinny hedgehog, an acyltransferase required for palmitoylation and activity of the hedgehog signal. *Science* **293**: 2080–2084.
- Chen, J.K., Taipale, J., Young, K.E., Maiti, T., and Beachy, P.A. 2002. Small molecule modulation of Smoothed activity. *Proc. Natl. Acad. Sci.* **99**: 14071–14076.
- Grant, B. and Hirsh, D. 1999. Receptor-mediated endocytosis in the *Caenorhabditis elegans* oocyte. *Mol. Biol. Cell* **10**: 4311–4326.
- Hahn, H., Wicking, C., Zaphropoulos, P.G., Gailani, M.R., Shanley, S., Chidambaram, A., Vorechovsky, I., Holmberg, E., Uden, A.B., Gillies, S., et al. 1996. Mutations of the human homolog of *Drosophila patched* in the nevoid basal cell carcinoma syndrome. *Cell* **85**: 841–851.
- Hodgkin, J. 1983. Two types of sex determination in a nematode. *Nature* **304**: 267–268.
- Incardona, J.P., Lee, J.H., Robertson, C.P., Enga, K., Kapur, R.P., and Roelink, H. 2000. Receptor-mediated endocytosis of soluble and membrane-tethered Sonic hedgehog by Patched-1. *Proc. Natl. Acad. Sci.* **97**: 12044–12049.
- Ingham, P.W. and McMahon, A.P. 2001. Hedgehog signaling in animal development: Paradigms and principles. *Genes & Dev.* **15**: 3059–3087.
- Jeong, J. and McMahon, A.P. 2002. Cholesterol modification of Hedgehog family proteins. *J. Clin. Invest.* **110**: 591–596.
- Johnson, R.L., Rothman, A.L., Xie, J., Goodrich, L.V., Bare, J.W., Bonifas, J.M., Quinn, A.G., Myers, R.M., Cox, D.R., Epstein, E.H., et al. 1996. Human homolog of *patched*, a candidate gene for the basal cell nevus syndrome. *Science* **272**: 1668–1671.
- Johnstone, I.L. 2000. Cuticle collagen genes: Expression in *Caenorhabditis elegans*. *Trends Genet* **16**: 21–27.
- Kurzchalia, T.V. and Ward, S. 2003. Why do worms need cholesterol? *Nat. Cell Biol.* **5**: 684–688.
- Kuwabara, P.E. and Labouesse, M. 2002. The sterol-sensing domain: Multiple families, a unique role? *Trends. Genet.* **18**: 193–201.
- Kuwabara, P.E. and Shah, S. 1994. Cloning by synteny: Identifying *C. briggsae* homologues of *C. elegans* genes. *Nucleic Acids Res.* **22**: 4414–4418.
- Kuwabara, P.E., Lee, M.H., Schedl, T., and Jefferis, G.S. 2000. A *C. elegans patched* gene, *ptc-1*, functions in germ-line cytokinesis. *Genes & Dev.* **14**: 1933–1944.
- Lander, E.S., Linton, L.M., Birren, B., Nussbaum, C., Zody, M.C., Baldwin, J., Devon, K., Dewar, K., FitzHugh, W., Funke, R., et al. 2001. Initial sequencing and analysis of the human genome. *Nature* **409**: 860–921.
- Lum, L. and Beachy, P.A. 2004. The Hedgehog response network: Sensors, switches, and routers. *Science* **304**: 1755–1759.
- Mathies, L.D., Schwarzstein, M., Morphy, K.M., Belloch, R., Spence, A.M., and Kimble, J. 2004. TRA-1/GLI controls development of somatic gonadal precursors in *C. elegans*. *Development* **131**: 4333–4343.
- Matyash, V., Entchev, E.V., Mende, F., Wilsch-Brauninger, M., Thiele, C., Schmidt, A.W., Knolker, H.J., Ward, S., and Kurzchalia, T.V. 2004. Sterol-derived hormone(s) controls entry into diapause in *Caenorhabditis elegans* by consecutive activation of DAF-12 and DAF-16. *PLoS Biol.* **2**: e280.
- McCarthy, R.A., Barth, J.L., Chintalapudi, M.R., Knaak, C., and Argraves, W.S. 2002. Megalin functions as an endocytic sonic hedgehog receptor. *J. Biol. Chem.* **277**: 25660–25667.
- Michaux, G., Gansmuller, A., Hindelang, C., and Labouesse, M. 2000. CHE-14, a protein with a sterol-sensing domain, is required for apical sorting in *C. elegans* ectodermal epithelial cells. *Curr. Biol.* **10**: 1098–1107.
- Murone, M., Rosenthal, A., and de Sauvage, F.J. 1999. Sonic hedgehog signaling by the patched-smoothed receptor complex. *Curr. Biol.* **9**: 76–84.
- Page, R.D. 1996. TreeView: An application to display phylogenetic trees on personal computers. *Comput. Appl. Biosci.* **12**: 357–358.
- Paulsen, I.T., Brown, M.H., and Skurray, R.A. 1996. Proton-dependent multidrug efflux systems. *Microbiol. Rev.* **60**: 575–608.
- Pepinsky, R.B., Zeng, C., Wen, D., Rayhorn, P., Baker, D.P., Williams, K.P., Bixler, S.A., Ambrose, C.M., Garber, E.A., Miatkowski, K., et al. 1998. Identification of a palmitic acid-modified form of human Sonic hedgehog. *J. Biol. Chem.* **273**: 14037–14045.
- Porter, J.A., Ekker, S.C., Park W.J., von Kessler, D.P., Young, K.E., Chen, C.H., Ma, Y., Woods, A.S., Cotter, R.J., Konnin, E.V., et al. 1996a. Hedgehog patterning activity: Role of a lipophilic modification mediated by the carboxy-terminal autoprocessing domain. *Cell* **86**: 21–34.
- Porter, J.A., Young, K.E., and Beachy, P.A. 1996b. Cholesterol modification of hedgehog signaling proteins in animal development. *Science* **274**: 255–259.
- Prince, V.E. and Pickett, F.B. 2002. Splitting pairs: The diverging fates of duplicated genes. *Nat. Rev. Genet.* **3**: 827–837.
- Radhakrishnan, A., Sun, L.P., Kwon, H.J., Brown, M.S., and Goldstein, J.L. 2004. Direct binding of cholesterol to the purified membrane region of SCAP: Mechanism for a sterol-sensing domain. *Mol. Cell* **15**: 259–268.
- Riddiford, L.M. 1993. Hormone receptors and the regulation of insect metamorphosis. *Receptor* **3**: 203–209.
- Roberts, B., Clucas, C., and Johnstone, I.L. 2003. Loss of SEC-23 in *Caenorhabditis elegans* causes defects in oogenesis, morphogenesis, and extracellular matrix secretion. *Mol. Biol. Cell* **14**: 4414–4426.
- Simmer, F., Tijsterman, M., Parrish, S., Koushika, S.P., Nonet, M.L., Fire, A., Ahringer, J., and Plasterk, R.H. 2002. Loss of the putative RNA-directed RNA polymerase RRF-3 makes *C. elegans* hypersensitive to RNAi. *Curr. Biol.* **12**: 1317–1319.
- Simmer, F., Moorman, C., van der Linden, A.M., Kuijck, E., van den Bergh, P.V., Kamath, R.S., Fraser, A.G., Ahringer, J., and Plasterk, R.H. 2003. Genome-wide RNAi of *C. elegans* using the hypersensitive *rrf-3* strain reveals novel gene functions. *PLoS Biol.* **1**: E12.
- Skop, A.R., Liu, H., Yates III, J., Meyer, B.J., and Heald, R. 2004. Dissection of the mammalian midbody proteome reveals conserved cytokinesis mechanisms. *Science* **305**: 61–66.
- Stein, L.D., Bao, Z., Blasari, D., Blumenthal, T., Brent, M.R., Chen, N., Chinwalla, A., Clarke, L., Clee, C., Coghlan, A., et al. 2003. The genome sequence of *Caenorhabditis briggsae*: A platform for comparative genomics. *PLoS Biol.* **1**: E45.
- Stone, D.M., Hynes, M., Armanini, M., Swanson, T.A., Gu, Q., Johnson, R.L., Scott, M.P., Pennica, D., Goddard, A., Phillips, H., et al. 1996. The tumour-suppressor gene *patched* encodes a candidate receptor for Sonic hedgehog. *Nature* **384**: 129–134.

- Strickland, L.I. and Burgess, D.R. 2004. Pathways for membrane trafficking during cytokinesis. *Trends Cell Biol.* **14**: 115–118.
- Taipale, J. and Beachy, P.A. 2001. The Hedgehog and Wnt signalling pathways in cancer. *Nature* **411**: 349–354.
- Taipale, J., Cooper, M.K., Maiti, T., and Beachy, P.A. 2002. Patched acts catalytically to suppress the activity of Smoothed. *Nature* **418**: 892–897.
- Thacker, C., Peters, K., Srayko, M., and Rose, A.M. 1995. The *bli-4* locus of *Caenorhabditis elegans* encodes structurally distinct *kex2*/subtilisin-like endoproteases essential for early development and adult morphology. *Genes & Dev.* **9**: 956–971.
- Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F., and Higgins, D.G. 1997. The CLUSTAL_X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* **25**: 4876–4882.
- Thummel, C.S. 1996. Flies on steroids: *Drosophila* metamorphosis and the mechanisms of steroid hormone action. *Trends. Genet.* **12**: 306–310.
- Timmons, L., Court, D.L., and Fire, A. 2001. Ingestion of bacterially expressed dsRNAs can produce specific and potent genetic interference in *Caenorhabditis elegans*. *Gene* **263**: 103–112.
- Waterston, R.H., Lindblad-Toh, K., Birney, E., Rogers, J., Abril, J.F., Agarwal, P., Agarwala, R., Ainscough, R., Alexandersson, M., An, P., et al. 2002. Initial sequencing and comparative analysis of the mouse genome. *Nature* **420**: 520–562.
- Wetmore, C. 2003. Sonic hedgehog in normal and neoplastic proliferation: Insight gained from human tumors and animal models. *Curr. Opin. Genet. Dev.* **13**: 34–42.
- Wolf, Y.I., Rogozin, I.B., and Koonin, E.V. 2004. Coelomata and not Ecdysozoa: Evidence from genome-wide phylogenetic analysis. *Genome Res.* **14**: 29–36.
- Yochem, J., Tuck, S., Greenwald, I., and Han, M. 1999. A gp330/megalin-related protein is required in the major epidermis of *Caenorhabditis elegans* for completion of molting. *Development* **126**: 597–606.
- Zhang, Y. and Kalderon, D. 2001. Hedgehog acts as a somatic stem cell factor in the *Drosophila* ovary. *Nature* **410**: 599–604.

Web site references

- <ftp://ftp-igbmc.u-strasbg.fr/pub/ClustalX/>; ClustalX.
<http://elegans.bcgsc.bc.ca>; *C. elegans* expression project.
<http://elegans.bcgsc.bc.ca/perl/sage>; *C. elegans* SAGE project.
<http://nematode.lab.nig.ac.jp>; Nematode Expression Pattern Database (NEXTDB).
<http://taxonomy.zoology.gla.ac.uk/rod/treeview.html>; TreeView.
<http://www.nematode.net>; Genome Sequencing Center.
<http://www.tigr.org/tdb/>; TIGR, the Institute for Genome Research.
<http://www.wormbase.org>; WormBase.

Received March 15, 2005; accepted in revised form June 27, 2005.