

Regulation of receptor protein-tyrosine phosphatase α by oxidative stress

Christophe Blanchetot, Leon G.J. Tertoolen and Jeroen den Hertog¹

Hubrecht Laboratory, Netherlands Institute for Developmental Biology, Uppsalalaan 8, NL-3584 CT Utrecht, The Netherlands

¹Corresponding author
e-mail: hertog@niob.knaw.nl

The presence of two protein-tyrosine phosphatase (PTP) domains is a striking feature in most transmembrane receptor PTPs (RPTPs). The function of the generally inactive membrane-distal PTP domain (RPTP-D2) is unknown. Here we report that an intramolecular interaction between the spacer region (Sp) and the C-terminus in RPTP α prohibited intermolecular interactions. Interestingly, stress factors such as H₂O₂, UV and heat shock induced reversible, free radical-dependent, intermolecular interactions between RPTP α and RPTP α -SpD2, suggesting an inducible switch in conformation and binding. The catalytic site cysteine of RPTP α -SpD2, Cys723, was required for the H₂O₂ effect on RPTP α . H₂O₂ induced a rapid, reversible, Cys723-dependent conformational change *in vivo*, as detected by fluorescence resonance energy transfer, with cyan fluorescent protein (CFP) and yellow fluorescent protein (YFP) flanking RPTP α -SpD2 in a single chimeric protein. Importantly, H₂O₂ treatment stabilized RPTP α dimers, resulting in inactivation. We propose a model in which oxidative stress induces a conformational change in RPTP α -D2, leading to stabilization of RPTP α dimers, and thus to inhibition of RPTP α activity.

Keywords: dimerization/FRET/oxidative stress/regulation/RPTP

Introduction

Protein-tyrosine phosphorylation is of major importance for cell proliferation, differentiation, migration and transformation within higher eukaryotic organisms. The regulation of tyrosine phosphorylation levels is mediated by the antagonistic activities of protein-tyrosine kinases (PTKs) and protein-tyrosine phosphatases (PTPs) (Hunter, 1995). A lot of work has been directed at elucidating the regulation of PTKs. In contrast, little is known about the mechanism of regulation of the PTPs (Neel and Tonks, 1997; den Hertog, 1999). The receptor PTPs (RPTPs) form a large subfamily of PTPs, next to the cytosolic ones. Genetic studies in *Drosophila* and *Caenorhabditis elegans* indicate a fundamental role for RPTPs in axon path finding and fibroblast growth factor (FGF) signaling (Desai *et al.*, 1996; Krueger *et al.*, 1996; Kokel *et al.*, 1998; Garrity *et al.*, 1999). Mouse knockout

studies have provided insight into the function of several RPTPs in vertebrates (Van Vactor *et al.*, 1998; den Hertog, 1999) and demonstrate that the RPTPs have important and specific functions. However, little is known about the regulation of their PTP activity.

RPTPs contain variable extracellular domains. It is tempting to speculate that RPTPs are regulated by extracellular ligands, like the RPTKs. To date, ligands have been found for only a few RPTPs, and only RPTP β/ζ was reported to be regulated by its ligand, pleiotrophin (Meng *et al.*, 2000). Therefore, other means of regulation may exist for RPTPs. Interestingly, RPTP α was recently found to be regulated during mitosis by GRB2 binding and serine phosphorylation (Zheng and Shalloway, 2001).

RPTPs contain a single transmembrane domain and in most cases two conserved intracellular PTP domains (D1 and D2) separated by a so-called spacer region (Krueger *et al.*, 1990). Interestingly, the membrane-proximal PTP domain (RPTP-D1) contains most if not all the activity (Wang and Pallen, 1991). Several reports suggest that the RPTP-D1 activity of RPTP α and CD45 is negatively regulated by dimerization (Desai *et al.*, 1993; Jiang *et al.*, 1999). The crystal structure of RPTP α -D1 shows a direct reciprocal interaction of a helix–loop–helix ‘wedge’ structure in the juxtamembrane region with the catalytic site of the opposing monomer (Bilwes *et al.*, 1996). Moreover, RPTP α and CD45 are inhibited by forced dimerization, which is dependent on the wedge (Majeti *et al.*, 1998; Jiang *et al.*, 1999). The topology of the two RPTP-D1s in the dimer is crucial for inhibition of activity, since active dimers were identified as well (Jiang *et al.*, 1999). Finally, RPTP α dimerizes constitutively in cells, as detected using cross-linkers and fluorescence resonance energy transfer (FRET), and multiple domains are involved in dimerization, including the transmembrane domain (Jiang *et al.*, 2000; Tertoolen *et al.*, 2001). RPTP dimerization may not be a general phenomenon, since dimers were not found in the crystal structure of LAR-D1D2 and RPTP μ -D1 (Hoffmann *et al.*, 1997; Nam *et al.*, 1999).

The membrane-distal PTP domains, RPTP-D2s, have striking features. They are largely conserved in sequence and are inactive (or much less active than RPTP-D1s). Furthermore, RPTP-D2s have a highly conserved three-dimensional structure (Nam *et al.*, 1999; A.M.Bilwes and J.P.Noel, personal communication). Mutations in only two residues, which are otherwise highly conserved in active RPTP-D1s, are responsible for the low activity in RPTP-D2s (Lim *et al.*, 1998; Buist *et al.*, 1999; Nam *et al.*, 1999). The conservation of inactive RPTP-D2s raises the question of their biological function. Several studies indicate that RPTP-D1s and RPTP-D2s interact, either intramolecularly or intermolecularly. However, the function of the RPTP-D1–D2 interaction is still unclear.




Fig. 1. The spacer and the C-terminal region of RPTPα together block binding to RPTPα. (A) Schematic representation of the constructs used here. HA, HA tag; ED, extracellular domain; D1, RPTP-D1; spacer, region between RPTP-D1 and RPTP-D2; D2, RPTP-D2; Ct, C-terminus; Myc, Myc tag. (B–D) Co-immunoprecipitation experiments. 293 cells were transiently co-transfected with the constructs indicated. HA-tagged RPTPα was immunoprecipitated using anti-HA antibodies (12CA5), resolved on SDS–PAGE gels and blotted. The gels were probed with anti-Myc antibody (9E10; top panels) and anti-HA (12CA5; middle panels). Expression of Myc-tagged RPTPα-D2s (Mt-RPTPα-D2s) in the lysates was monitored in the bottom panels. (B) Co-transfection of Mt-RPTPα-D2 alone (–), with full-length HA-tagged RPTPα (FL) or mutant RPTPα lacking the second domain (ΔD2). (C) Co-transfection of HA-RPTPα-ΔD2 with various Mt-RPTPα-D2 constructs [depicted in (A)]. (D) Co-transfection of full-length HA-RPTPα with Mt-RPTPα-D2, -SpD2 or SpD2Y772F.

RPTPδ-D2 was found to directly inhibit RPTPσ-D1 activity concomitant with binding to the juxtamembrane region of RPTPσ-D1 (Wallace *et al.*, 1998). Inter- and intramolecular interactions between CD45-D1 and CD45-D2 may negatively regulate CD45-D1–CD45-D1 homodimerization, leading to activation of CD45-D1 (Felberg and Johnson, 1998). We recently showed binding of RPTPα to various RPTP-D2s, involving multiple binding sites, suggesting that cross-talk between RPTPs may be a broad mechanism of regulation. The involvement of the wedge in RPTP-D1–RPTP-D2 interactions suggested a possible role for D2s in the regulation of RPTP-D1–D1 dimerization (Wallace *et al.*, 1998; Blanchetot and den Hertog, 2000). The juxtamembrane region of RPTPμ was shown to interact intramolecularly with both PTP domains of RPTPμ, thus regulating RPTPμ-D1 activity (Feiken *et al.*, 2000). Finally, the crystal structure of the complete cytoplasmic domain of LAR provided some structural evidence for D1–D2 interactions. LAR-D1 sits on top of LAR-D2 with extensive contacts through the spacer region containing a four-residue-long ‘linker’ (Nam *et al.*, 1999). Despite a number of reports on interactions between RPTP-D1s and RPTP-D2s, whether and how these interactions are regulated is still largely unknown.

In the present study, we show that the spacer region (the region between D1 and D2; Sp) and the C-terminal region

of RPTPα interacted intramolecularly with each other to form a ‘closed’ conformation, which was unable to interact intermolecularly with RPTPα. Binding of RPTPα-SpD2 to RPTPα was inducibly restored by stress factors like H₂O₂, UV and heat shock. In addition, the catalytic site cysteine in RPTPα-D2 was required for H₂O₂-induced binding. Using FRET, we demonstrate that H₂O₂ evoked a conformational change in RPTPα-SpD2, which was rapid and reversible. *In vitro*, H₂O₂ induced a conformational change as well, suggesting a direct effect. The change in conformation was concomitant with stabilization of RPTPα dimers and complete inactivation of RPTPα activity. In conclusion, we provide evidence that intra- and intermolecular interactions of RPTPα are regulated by extracellular stimuli, including oxidative stress, and suggest a mechanism for regulation of RPTP complex formation and function.

Results

RPTPα interacts with RPTPα-D2

We have previously reported that RPTPα binds to multiple RPTP-D2s, including RPTPα-D2, LAR-D2, RPTPσ-D2, RPTPδ-D2 and RPTPμ-D2 (Blanchetot and den Hertog, 2000). An interesting feature of the interaction is the difference in affinity of RPTP-D2s for either full-length HA-tagged RPTPα (HA-RPTPα) or HA-RPTPα-ΔD2, a




Fig. 2. Binding of RPTP α -SpD2 to RPTP α is induced by oxidative stress and is reversible. **(A)** 293 cells were co-transfected with HA-RPTP α or HA-RPTP α - Δ D2 and Mt-RPTP α -SpD2. After stimulation with (final concentration) 10% newborn calf serum (NCS), pervanadate (VO₄; 1 mM VO₄ + 1 mM H₂O₂), 1 mM H₂O₂, 50 ng/ml TPA, UV (200 J/m²) or heat shock (HS; 42°C) for the time indicated, HA-RPTP α was immunoprecipitated. After separation on SDS-PAGE and blotting of the gel, the blot was probed with anti-Myc (top panel) and anti-HA (middle panel) antibodies. Equal expression of the Myc-tagged RPTP-D2s was monitored in the lower panel. **(B)** 293 cells were transiently transfected with HA-RPTP α and Mt-RPTP α -SpD2, pre-treated (+) or not (-) with 10 mM NAC for 16 h and subjected to 100 or 200 J/m² UV radiation, and left to recover for 10 min. Immunoprecipitation and immunoblotting were as in (A). **(C)** Transiently transfected 293 cells were treated with H₂O₂ (1 mM) for the time indicated. Alternatively, after 10 min of stimulation with H₂O₂, the medium was replaced with fresh pre-warmed medium for the time indicated (Recovery). Note that media change by itself did not affect H₂O₂-induced binding (last lane). Immunoprecipitation and immunoblotting were as in (A).

construct lacking part of the spacer region and the second domain (Figure 1A). Indeed, as depicted in Figure 1B, less myc-tagged RPTP α -D2 (Mt-RPTP α -D2) bound to full-length HA-RPTP α than to HA-RPTP α - Δ D2, while similar amounts of HA-RPTP α and HA-RPTP α - Δ D2 were expressed. This result suggests that the intrinsic RPTP α -D2 in full-length RPTP α had a negative effect on the binding of other RPTP-D2s, which may be due to occupation of the binding site by the intrinsic RPTP α -D2, or to intrinsic RPTP α -D2 holding RPTP α in a 'closed' conformation, thereby restricting binding to other RPTP-D2s.

Intramolecular interaction of the spacer region with the C-terminal helix of RPTP α -SpD2

In order to understand better the binding observed between RPTP α -D1 and RPTP α -D2, we used a panel of deletion mutants. The longest form of RPTP α -D2 (RPTP α -SpD2) encompassed a large part of the spacer region, Sp, the region between D1 and D2 (Figure 1A). To our surprise, this construct did not co-immunoprecipitate with HA-RPTP α (Figure 1C). Deletion of the C-terminus of RPTP α -SpD2 restored binding to HA-RPTP α (Figure 1C) despite the fact that deletion of the C-terminus in RPTP α -D2, without the spacer region, led to a decrease in binding (compare the first and second lane, Figure 1C) (Blanchetot and den Hertog, 2000). These results indicate

that the presence of both the spacer and the C-terminal region prohibits binding to RPTP α . A single mutation exactly at the interface between the spacer region and the C-terminal helix, Tyr772 to Phe, partially restored binding to HA-RPTP α , thus partially reproducing the effect of a deletion (Figure 1D). Presumably, Tyr772 stabilized the 'closed' conformation. RPTP α -SpD2 was not phosphorylated on tyrosine (data not shown), indicating that the effects of mutation of Tyr772 were not due to lack of tyrosine phosphorylation. Moreover, mutation of Tyr789, an *in vivo* phosphorylation site (den Hertog *et al.*, 1994), did not affect binding between RPTP α -SpD2 and RPTP α (data not shown). In conclusion, our results suggest that RPTP α -D2 exists in a 'closed' conformation. Alteration of the intramolecular interaction between the spacer and the C-terminal region by deletion or subtle mutation switches RPTP α -SpD2 to an 'open' conformation, allowing binding to RPTP α in an intermolecular fashion.

H₂O₂, UV and heat shock induced binding of RPTP α -SpD2 to RPTP α

We next investigated whether RPTP α -SpD2 could be induced to bind to RPTP α in response to a panel of stimuli. Oxidative stress like H₂O₂, UV, as well as heat shock strongly induced binding of RPTP α -SpD2 to RPTP α (Figure 2A). Similarly, RPTP α -SpD2 bound to RPTP α - Δ D2 after H₂O₂, UV or heat shock (Figure 2A and data not




Fig. 3. H₂O₂-induced binding of RPTP α -SpD2 to RPTP α is dependent on catalytic site Cys723. (A) 293 cells were co-transfected with HA-RPTP α and wild-type Mt-RPTP α -SpD2 (wt) or mutant Mt-RPTP α -SpD2 with Cys723 mutated to Ser (C723S). The cells were treated with H₂O₂ (1 mM, 15 min) (+) or left untreated (-). HA-RPTP α was immunoprecipitated, separated on SDS-PAGE gels and blotted. The blot was probed with anti-Myc (top panel) and anti-HA (middle panel) antibodies. Expression of the Mt-RPTP α -D2s in the lysate was monitored in the lower panel. (B) Mt-RPTP α -SpD2 was co-transfected in 293 cells with full-length wild-type HA-RPTP α (wt) or mutants, as indicated. The cells were stimulated with H₂O₂ (1 mM, 15 min). Immunoprecipitation and immunoblotting for (B–D) were carried out as in (A). (C) Mt-RPTP α -SpD2 was co-transfected with full-length HA-RPTP α (wt), HA-RPTP α -C433S/C723S or PSG5 (-) and stimulated with H₂O₂ (1 mM, 15 min). (D) 293 cells were transiently co-transfected with HA-RPTP α and various Mt-RPTP α -SpD2 mutants with Cys to Ser mutations, as indicated, and treated with H₂O₂ (1 mM, 5 min).

shown). Other stimuli, like serum, tumor necrosis factor- α , lysophosphatidic acid (LPA), epidermal growth factor or 12-*O*-tetradecanoylphorbol 13-acetate (TPA), did not induce binding (Figure 2A and data not shown). Pretreatment of transfected cells with *N*-acetyl-cysteine (NAC), a free radical scavenger, blocked UV-induced RPTP α -SpD2 binding to RPTP α (Figure 2B), indicating that the effect of UV is mediated by the production of free radicals. H₂O₂-induced binding was strong and rapid, reaching a maximum within 1 min of stimulation (Figure 2C), suggesting a direct effect of oxidative stress on RPTP α . As long as H₂O₂ was present in the medium, binding was observed (Figure 2C). Removal of H₂O₂ by replacing the medium led to a reduction in binding. After 120 min, no binding was detected anymore, showing that it was reversible, although dissociation was much slower than association. Note that H₂O₂-induced binding was not affected by the medium replacement itself (Figure 2C, last lane). In conclusion, we show that intermolecular interactions between RPTP α and RPTP α -SpD2 were rapidly induced by oxidative stress factors like H₂O₂ and UV, required the production of free radicals and were reversible.

Involvement of the catalytic site cysteine of RPTP α -SpD2 in H₂O₂-induced binding

Since oxidative stress induced binding of RPTP α -SpD2 to RPTP α , and since it has long been proposed that free radicals act on PTPs by oxidation of their catalytic site cysteine (Knebel *et al.*, 1996; Denu and Dixon, 1998;

Denu and Tanner, 1998; Lee *et al.*, 1998), we mutated Cys723 in RPTP α -SpD2. Interestingly, mutation of Cys723 to Ser in RPTP α -SpD2 completely abolished H₂O₂-induced binding to RPTP α (Figure 3A). The same result was obtained when Cys723 was mutated to Ala (data not shown). Mutation of Cys723 in full-length RPTP α also abolished H₂O₂-induced binding to RPTP α -SpD2 (Figure 3B), presumably because full-length RPTP α -C723S was locked in a 'closed' conformation. Mutation of another residue in the second catalytic site (R729K) or mutation of the first catalytic site cysteine (C433S) of RPTP α had no or hardly any effect on H₂O₂-induced binding to RPTP α -SpD2 (Figure 3B). These results suggest that Cys723 is specifically and directly involved in sensing free radicals, leading to the conformational change in RPTP α -SpD2 necessary to allow binding to RPTP α . Along the same lines, H₂O₂ increased RPTP α -D2 (without spacer) binding to RPTP α , but not to RPTP α -C433/723S (Figure 3C), suggesting that opening of full-length RPTP α is dependent on Cys723. However, Cys723 is not necessary for direct binding since a mutant RPTP α -D2-C723S (without the spacer region) was still able to bind to RPTP α (data not shown). In conclusion, we show that Cys723, the catalytic site cysteine of RPTP α -D2, plays a direct role in the H₂O₂-induced intermolecular interaction between RPTP α and RPTP α -SpD2.

Free radical-induced oxidation of proteins may lead to the formation of disulfide bonds. It is noteworthy that H₂O₂ did not induce non-specific covalent intermolecular bonds between RPTP α -SpD2 and RPTP α since mono-

meric proteins were found after SDS-PAGE separation under non-reducing conditions (data not shown). We investigated whether other cysteines in the vicinity of Cys723S were involved. Two cysteines are located close to the reactive Cys723, Cys634 and Cys734, both at ~ 9 Å. Cys634 is highly conserved in all PTP domains and Cys734 is divergent between PTP domains. Mutation of the conserved Cys634 to Ser had no effect on the H₂O₂-induced binding between RPTP α and RPTP α -SpD2. However, mutation of Cys734 to serine induced constitutive binding independently of H₂O₂ (Figure 3D). These results suggest that Cys734 may be involved in the oxidative stress response, possibly by forming a disulfide bridge with Cys723 leading to a structural distortion of RPTP α -SpD2, which may form the basis for the conformational change.

Detection of conformational changes using FRET

FRET between two green fluorescent protein (GFP) derivatives has become widely used to detect protein-protein interactions (Cubitt *et al.*, 1995; Bastiaens and Pepperkok, 2000) and conformational changes of peptides (Miyawaki *et al.*, 1997; Nagai *et al.*, 2000). We used FRET to test the hypothesis that H₂O₂ induced a conformational change in RPTP α -SpD2. The N- and C-termini of RPTP α -SpD2 are very close together, according to the crystal structure (A.M.Bilwes and J.P.Noel, personal communication), similar to LAR-SpD2 (Nam *et al.*, 1999). CFP and YFP were fused to the N- and C-terminus of RPTP α -SpD2, respectively (Figure 4A). The emission spectra of single transfected cells were determined after excitation at 430 nm, a wavelength that only excited CFP. As expected from the structural data, FRET was detected between CFP and YFP in single-cell measurements, revealing the close proximity of the two fluorophores and consequently of the interaction between the spacer region and the C-terminus (Figure 4C). Interestingly, addition of H₂O₂ dramatically reduced FRET, as detected by a reduction in the YFP emission and a concomitant increase in CFP emission (reduction in energy transfer from CFP to YFP leads to enhanced direct CFP emission), indicating that a conformational change was indeed induced in RPTP α -SpD2 (Figure 4C and D). H₂O₂-induced changes in FRET were rapid, reaching maximal effects after 1–2 min, similar to the binding experiments (Figure 4D). Interestingly, the catalytic site cysteine mutant (C723S) was insensitive to H₂O₂ in that no changes in FRET and thus in conformation were observed in response to H₂O₂ (Figure 4E and F), consistent with the binding experiments (Figure 3). The change in FRET was specific since H₂O₂ had no effect on the fluorescence properties of CFP and YFP themselves (Figure 4B). Continuous dual wavelength excitation measurement of FRET indicated that the conformational changes were very rapid, reaching half-maximal levels in ~ 30 s (Figure 4G). Similar to the interaction between RPTP α and RPTP α -SpD2, the conformational change was reversible and with much slower kinetics (Figure 4G and H). As a control, we fused CFP or YFP to the C- and N-termini of RPTP α -SpD2, generating CC and YY, respectively (Figure 4A). Following co-transfection of CC and YY, we did not detect any FRET (Figure 4I),

demonstrating that FRET, observed in CFP-RPTP α -SpD2-YFP, was due to an intramolecular interaction.

The very rapid and total change in conformation induced by H₂O₂ suggested a direct effect of H₂O₂ on RPTP α -SpD2. To test this hypothesis, transiently transfected cells were lysed and the cell lysates were then tested for FRET in a spectrophotometer. Under these conditions, FRET was still observed (Figure 5A). Stimulation of cells with H₂O₂ prior to lysis (*in vivo*) led to loss of FRET in the lysate. Addition of H₂O₂ directly to the lysate (*in vitro*) also led to a large decrease in FRET (Figure 5B), suggesting that the effect of H₂O₂ was direct. Taken together, using FRET *in vivo* and *in vitro*, we show that oxidative stress directly provoked reversible changes in the conformation of RPTP α -SpD2 in a catalytic site cysteine-dependent fashion.

H₂O₂-induced dimerization of full-length RPTP α

The formation of RPTP α dimers has been reported using chemical cross-linkers on the one hand and FRET on the other (Jiang *et al.*, 2000; Tertoolen *et al.*, 2001). However, co-immunoprecipitation experiments using differently tagged RPTP α were always unsuccessful. For instance, we were unable to reproducibly co-immunoprecipitate myc-tagged RPTP α (Mt-RPTP α) with HA-tagged full-length RPTP α , suggesting that RPTP α dimers were loosely associated and dissociate during the immunoprecipitation procedure. Since H₂O₂ affected intermolecular interactions between RPTP α , we tested co-immunoprecipitation following H₂O₂ treatment. Stimulation with H₂O₂ led to a large increase in the amount of Mt-RPTP α that co-immunoprecipitated with HA-RPTP α (Figure 6A), suggesting stabilization of dimers by intermolecular interactions. UV had the same effect as H₂O₂ on RPTP α complex formation, which was blocked by pre-incubation with NAC (data not shown). Mutation of the second domain catalytic site cysteine severely reduced H₂O₂-induced complex formation (Figure 6A). In addition, co-immunoprecipitation of full-length RPTP α was reversible after the removal of H₂O₂ (data not shown). Cys433 in RPTP α -D1 appeared to be involved in H₂O₂-induced dimerization as well, since mutation of both catalytic cysteines led to complete loss of H₂O₂-induced binding (Figure 6A). Mutation of Cys433 by itself had only minor effects on the H₂O₂-induced complex formation (data not shown). In conclusion, we show that H₂O₂ led to more stable complex formation of full-length RPTP α in a D2 catalytic site cysteine-dependent fashion.

We used the non-cell-permeable cross-linker bis[sulfosuccinimidyl]suberate (BS³) to investigate whether H₂O₂ induced a large increase in dimer formation. Figure 6B shows that H₂O₂ did not affect the extent of BS³-mediated cross-linking of RPTP α , suggesting that H₂O₂ does not induce *de novo* dimerization, but rather leads to stabilization of existing dimers. It is noteworthy that no higher order protein complexes were detected after H₂O₂ treatment and BS³-mediated cross-linking, indicating that H₂O₂ did not induce non-specific aggregation of RPTP α .

In order to obtain insight into the mechanism underlying H₂O₂-induced stabilization of RPTP α dimers, we investigated binding of RPTP α -SpD2 to RPTP α -D1 and RPTP α -D2. Following H₂O₂ treatment, RPTP α -SpD2 bound to both RPTP α -D1 and RPTP α -D2 (Figure 6C).




Fig. 4. H_2O_2 -induced conformational change in RPTP α -D2. (A) Schematic representation of the constructs used here: CFP-RPTP α -SpD2-YFP2.1 (WT), CFP-RPTP α -SpD2-C723S-YFP2.1 (C723S), CFP-RPTP α -SpD2-CFP (CC) and YFP-RPTP α -SpD2-YFP (YY). Sp, spacer region; Ct, C-terminus. (B) The spectral properties of a single 293 cell, transfected with CFP alone (excitation at 430 nm, emission maximum at 480 nm), or YFP2.1 alone (excitation at 490 nm, emission maximum at 525 nm), before (rest) and after a wash with fresh medium (wash) and during incubation in medium containing 1 mM H_2O_2 for the periods of time indicated. (C) Emission spectra of a single transfected 293 human embryonic kidney cell (excitation 430 nm), expressing wild-type CFP-RPTP α -SpD2-YFP2.1 (WT). H_2O_2 treatment (1 mM) as indicated. (D) Time course of the ratio of the emission intensities at 525 and 480 nm from (C). (E) Emission spectra as in (C) of a single cell expressing mutant CFP-RPTP α -SpD2-C723S-YFP2.1 (C723S). (F) Time course of the ratio of the emission intensities at 525 and 480 nm from (E). (G) Real-time FRET analysis was performed on a single cell expressing CFP-RPTP α -SpD2-YFP2.1 (WT) by continuous dual excitation wavelength measurements [excitation at 430 and 490 nm, emission at 535(30) nm; for details see Tertoolen *et al.*, 2001]. Treatment with 1 mM H_2O_2 and subsequent recovery were followed. The half-time of decay was calculated ($\tau = 30$ s). (H) A single cell expressing CFP-RPTP α -SpD2-YFP2.1 (WT) was treated with 1 mM H_2O_2 for 10 min, the medium was replaced by regular medium and emission spectra were obtained at regular intervals. The emission ratio at 525 and 480 nm is plotted against time. (I) Emission spectra of a single cell expressing both CC and YY (excitation at 430 nm, red trace; and to control for YFP expression, excitation at 490 nm, blue trace). All single-cell measurements were repeated at least three times. Representative experiments are depicted here.




Fig. 5. The effect of H₂O₂ on RPTP α -D2 is direct. (A) 293 cells were transfected with wild-type CFP-RPTP α -SpD2-YFP2.1 and lysed in cell lysis buffer. Emission spectra were determined for these lysates (excitation 430 nm). The cells had been left untreated (rest) or were stimulated with 1 mM H₂O₂ for 5 min prior to lysis (*in vivo*). (B) As in (A), emission spectra were determined for cell lysates. The cell lysates were left untreated (rest) or treated with 1 mM H₂O₂ (*in vitro*). The experiments were repeated at least three times and representative measurements are depicted here.

This result was surprising, since previously we had not been able to detect direct D2–D2 interactions in the yeast two-hybrid system or using GST–RPTP α -D2 binding assays. Apparently, the individual RPTP α -D2 domains did interact, which may be due to the fact that only small epitope tags were fused to RPTP α -SpD2, instead of larger GAL4 or GST proteins. Interestingly, co-transfection of HA-RPTP α -SpD2 with Mt-RPTP α -D1 and Mt-RPTP α -D2 demonstrated that PTP α -D2 bound preferentially to Mt-RPTP α -D2 (Figure 6C, compare top panel with bottom panel). Taken together, we demonstrate here that H₂O₂ treatment induced stabilization of RPTP α dimers, which appears to be mediated by D2–D2 interactions.

Inactivation of RPTP α activity *in vivo* by H₂O₂ is partially dependent on Cys723

Since H₂O₂ induced the formation of stable RPTP α complexes, we investigated the effect of H₂O₂ on RPTP α activity. To this end, HA-RPTP α and mutants were immunoprecipitated and assayed in a PTP assay, using pNPP as a substrate. In order to reduce experimental variation and to get significant PTP activity above background level, several dishes of (stimulated) transfected cells were collected and pooled. Addition of H₂O₂ *in vivo* led to reproducible complete inactivation of RPTP α activity (Figure 7). Surprisingly, H₂O₂ only partially reduced the activity of mutant RPTP α -C723S and RPTP α - Δ D2 (40–60%) (Figure 7). Yet, addition of

H₂O₂ (1 mM) to immunoprecipitated RPTP α -C723S completely inhibited its activity, like wild-type RPTP α (Figure 7). Wild-type RPTP α and RPTP α -C723S are equally sensitive to H₂O₂ treatment *in vitro*, in that different H₂O₂ concentrations inhibit wild-type and mutant RPTP α to the same extent (e.g. 0.1 mM H₂O₂ led to a 70% reduction in PTP activity of both wild-type and C723S; data not shown). These results correlate with the observation that wild-type RPTP α formed much more stable dimers than RPTP α -C723S, and suggest that H₂O₂-induced RPTP α dimers are inactive. In conclusion, we show that the effect of H₂O₂ on the catalytic activity of RPTP α -D1 is dependent on the catalytic site cysteine of RPTP α -D2, which is likely to be due to the H₂O₂-induced conformational change in RPTP α -D2.

Discussion

More and more evidence shows that RPTP-D2s interact with RPTPs. In some cases, RPTP-D2s have been shown to regulate RPTP activity. However, whether and how the interactions are regulated is not known. Here, we propose a model for the regulation of the interaction between RPTP α -D2 and RPTP α in which intramolecular interactions between the spacer and the C-terminal region of RPTP α -SpD2 lock RPTP α -SpD2 in a ‘closed’ conformation, prohibiting intermolecular binding to RPTP α . We show that oxidative stress induced a conformational change, which was accompanied by more stable RPTP α dimers and inactivation of RPTP α activity. Taken together, our results suggest a mechanism by which intra- and intermolecular interactions of RPTP α are regulated by oxidative stress.

The change in conformation was directly detected *in vivo* and *in vitro* using FRET between two GFP derivatives fused N- and C-terminally to RPTP α -SpD2. FRET between GFP derivatives is a valuable tool to study *in vivo* protein–protein interactions. Recently, FRET has been used to detect conformational changes in peptides as a result of calcium binding (Miyawaki *et al.*, 1997) or PKA phosphorylation (Nagai *et al.*, 2000). Here, we used FRET to detect a conformational change in a complete protein domain, and we followed the dynamics of this conformational change in living cells in real time (Figure 4). Analysis of FRET between GFP derivatives, fused to an entire protein domain, is a new approach to study the dynamics of conformational changes in proteins.

Extracellular stimuli, such as H₂O₂, UV and heat shock, induced a conformational change followed by intermolecular interactions between RPTP α *in vivo*. The catalytic site cysteine of RPTP α -SpD2 was necessary for H₂O₂-induced binding to RPTP α . The involvement of the catalytic site cysteine of RPTP α -D2 is very interesting. PTP catalytic site cysteines have been shown to be preferential targets of free radicals, and their oxidation a potential reversible mode to directly regulate PTP activity (Knebel *et al.*, 1996; Denu and Tanner, 1998; Fauman *et al.*, 1998). H₂O₂ induced a reversible disulfide bridge between the catalytic site cysteine and a nearby cysteine in LMW-PTP, Cdc25A and KAP (Caselli *et al.*, 1998; Fauman *et al.*, 1998; Song *et al.*, 2001). Importantly, the formation of a disulfide bridge induced detectable structural changes outside the catalytic pocket of LMW-PTP




Fig. 6. Oxidative stress stabilized RPTP α dimerization. (A) 293 cells were co-transfected with HA-RPTP α full length [wild-type (WT) or with the catalytic site cysteine single mutant (C723S) or double mutant (C433/C723S)] and Myc-tagged full-length RPTP α (Mt-RPTP α). Cells were treated with H₂O₂ (1 mM, 10 min) (+) or left untreated (-). After anti-HA immunoprecipitation and immunoblotting, the blot was probed with anti-Myc (top panel) and anti-HA (middle panel) antibodies. Equal expression of the Mt-RPTP α in the lysate was monitored in the lower panel. (B) 293 cells were transfected with HA-RPTP α , treated with H₂O₂ (1 mM, 10 min) (+) or left untreated. Membrane proteins were cross-linked using the non-cell-permeable cross-linker BS³. An immunoblot of whole-cell lysates is shown with the position of RPTP α monomers and dimers indicated. (C) HA-RPTP α -SpD2 was co-transfected with Mt-RPTP α -D1, Mt-RPTP α -D2 or both. H₂O₂ treatment (10 min, 1 mM), immunoblotting of anti-HA tag immunoprecipitates and lysates were performed as indicated.

(Caselli *et al.*, 1998). A similar intramolecular disulfide bridge was suggested for several other PTPs, including SHP-1 (Pregel and Storer, 1997; Denu and Tanner, 1998). However, disulfide bridges have been excluded for PTP-1B (Lee *et al.*, 1998) and VHR PTPs (Denu and Tanner, 1998), which preferentially form sulfenic acid after oxidative stress. Our results suggest that next to the catalytic site Cys723, Cys734 (at ~9 Å from Cys723) may be involved in the conformational change induced by oxidative stress, possibly through the formation of a disulfide bridge. Interestingly, in the H₂O₂-sensing transcription factor OxyR, two cysteines, 17 Å apart, were found to form a disulfide bridge after oxidative stress, resulting in significant structural changes (Choi *et al.*, 2001), suggesting that the distance of ~9 Å between Cys723 and Cys734 may not *per se* prohibit disulfide bridge formation. Alternatively, Cys734 may be involved otherwise in the H₂O₂-induced conformational change. Furthermore, we cannot exclude the possibility that mere oxidation of Cys723 to sulfenic acid triggers the conformational change. More work will be required to show definitively the involvement of a disulfide bridge or of other modifications, like sulfenic acid formation in RPTP α -D2.

Strikingly, all the effects of oxidative stress on RPTP α shown here are dependent on the catalytic site cysteine of RPTP α -D2. These effects include: (i) conformational changes in RPTP α -D2; (ii) binding of RPTP α -D2 to RPTP α ; (iii) formation of a stable RPTP α dimer complex; and (iv) complete inactivation of RPTP α . This suggests that all these effects may be linked, which is further supported by the kinetics of these effects (rapid induction within 1 min and slow recovery in ~2 h).

We propose a multistep regulatory mechanism for RPTP α as a result of oxidative stress (Figure 8). Before




Fig. 7. Oxidative stress-induced change in RPTP α activity. HA-RPTP α wild type (WT), HA-RPTP α -C723S (C723S) or HA-RPTP α -ΔD2 (ΔD2), lacking D2, were expressed in 293 cells. Cells were stimulated with H₂O₂ (1 mM, 5 min) (vivo), HA-RPTP α proteins were immunoprecipitated and PTP activity was assayed using pNPP as a substrate. Alternatively, H₂O₂ (1 mM) was added after immunoprecipitation of RPTP α from unstimulated cells (vitro). The activity was corrected for RPTP α expression.

stimulation, an equilibrium exists between monomeric and dimeric RPTP α . RPTP α dimerizes constitutively and extensively in living cells (Jiang *et al.*, 2000; Tertoolen *et al.*, 2001), suggesting that the dimeric state may be preferred. Whether dimeric RPTP α is active in the pre-stimulation situation depends on the rotational coupling in the dimers (Jiang *et al.*, 1999). Oxidative stress leads to rapid oxidation of catalytic sites in PTPs, partially inactivating PTP activity (Denu and Tanner, 1998; Lee *et al.*, 1998). RPTP α is completely inactivated following H₂O₂ treatment, while inactivation of RPTP α -C723S (and RPTP α -ΔD2) is partial, suggesting that additional events are responsible for complete inactivation of wild-type




Fig. 8. A model for regulation of RPTP α by oxidative stress. Under normal conditions, RPTP α monomers and dimers are in equilibrium (pre-stimulation). Whether the dimers are active or inactive depends on their rotational coupling. Oxidative stress (e.g. H₂O₂) leads to rapid inactivation by oxidation of the catalytic site cysteine in D1 (green to red). Oxidative stress also induces a conformational change in D2, which may drive the formation of stabilized dimers, perhaps through D2–D2 interactions. The stabilized dimers are completely inactive. Mutant RPTP α -C723S lacks a catalytic site cysteine in D1, does not form stabilized dimers and is not inactivated completely following oxidative stress. After reduction, RPTP α slowly returns to the pre-stimulation state following a conformational change in D2. See text for details.

RPTP α . Oxidative stress rapidly induces a conformational change in D2, but not D2-C723S. This conformational change induces intermolecular interactions of RPTP α -D2, with D2–D2 interactions being preferred. We have previously demonstrated that many domains in RPTP α are involved in RPTP α dimerization, including the transmembrane domain (Jiang *et al.*, 2000; Tertoolen *et al.*, 2001). The oxidative stress-induced D2–D2 interactions may drive stabilization of the dimers, facilitating additional interactions in the dimer, including for instance D1–wedge interactions (Bilwes *et al.*, 1996), which may be responsible for complete inactivation of RPTP α . Reduction of the catalytic site cysteine in D1 in the highly reducing intracellular milieu is fast (Lee *et al.*, 1998). The kinetics of the conformational change and SpD2 binding indicate that reversion to the pre-stimulation state is slow, suggesting that RPTP α activity is only slowly returning to pre-stimulation levels too. Our model suggests that oxidative stress leads to complete inactivation of RPTP α activity, allowing efficient propagation of phosphotyrosine-dependent signaling.

Activation of multiple RPTKs leads to local H₂O₂ production, suggesting that RPTK activation itself may lead to complete and/or sustained RPTP α inactivation. We were unable to detect any effect of growth factors or other known stimuli that produce free radicals on RPTP α -SpD2 binding capacity or conformational changes. This is most likely due to the local and low production of H₂O₂ in response to growth factors. RPTP α -SpD2 is cytoplasmic and may, therefore, not be localized correctly to perceive ligand-induced H₂O₂ production. UV is more potent and induces massive tyrosine phosphorylation, which is thought to be mediated by free radicals, leading to inactivation of PTPs (Lee *et al.*, 1998; Gross *et al.*, 1999). H₂O₂ has been known for a long time for its

mitogenic effect, which may be mediated by its inhibitory effect on PTPs (Knebel *et al.*, 1996; Bae *et al.*, 1997). The fact that H₂O₂ regulates the interaction between RPTP-D2s and RPTPs provides an interesting new insight into the mode of action of H₂O₂, as well as into the function of the conserved RPTP-D2s.

RPTPs interact with (multiple) RPTP-D2s, and inter- and intramolecular interactions may regulate RPTP activity. The presence of an extended N-terminal region forming a helix interacting extensively with the C-terminal helix, as observed in RPTP α -D2, is a conserved structural feature found in most PTP domains crystallized to date, including RPTP-D1s and RPTP-D2s, and to a lesser extent cytosolic PTPs. Competition in binding between PTP domains and spacer regions or C-terminal regions has been described for CD45 and RPTP μ , in which the presence of an extended N-terminal region prohibited intermolecular interactions in favor of intramolecular interactions (Feiken *et al.*, 2000; Hayami-Noumi *et al.*, 2000). Interestingly, these interactions also affected the activity of the PTPs (Felberg and Johnson, 1998; Feiken *et al.*, 2000; Aricescu *et al.*, 2001). It will be very interesting to see whether these interactions in RPTP μ and CD45 are regulated by any stimulus as well.

Most RPTP-D2s contain a catalytic site cysteine, suggesting that the model for oxidative stress-induced regulation of RPTP dimerization and activity we propose here (Figure 8) may be applicable to other RPTPs. However, the crystal structure of LAR suggests that D1–D1 interactions are prohibited due to steric hindrance by LAR-D2, therefore opposing the dimerization model suggested by the crystal structure of RPTP α -D1 (Bilwes *et al.*, 1996; Nam *et al.*, 1999). Regulated changes in the conformation of D2 and in the interaction between D1 and D2 may unify both crystal structures in that a specific stimulus affecting the conformation of D2 may allow for D1–D1 interactions to occur in LAR. Confirmation that RPTP topology can change will require the crystallization of other full cytoplasmic regions of RPTPs, before and after stimulation. In this respect, it would be very interesting to see the structural arrangement of the two PTP domains of RPTP α , and the extent of the conformational changes induced by H₂O₂.

We have previously suggested cross-talk between RPTPs, involving RPTP-D1–D2 interactions (Blanchetot and den Hertog, 2000). The existence of stimulus-dependent binding between domains of RPTPs raises the possibility of regulated cross-talk, in which one or two (distinct) stimuli induce specific interactions between RPTPs. Such reshuffling of RPTP complexes may lead to specific (de)phosphorylation of (other) substrates, thus triggering a new set of downstream events. Knowledge of regulated interactions between (distinct) RPTP-D1s and RPTP-D2s, as well as between full-length RPTPs, may help to understand the mechanism of regulation and cooperation. In conclusion, we show that oxidative stress affected RPTP α 's conformation, interactions and activity, which led us to propose a model for the regulation of RPTP α . In the future, it will be interesting to extend this model to other RPTPs, which may lead to the identification of distinct stimuli that regulate RPTP complex formation and function.

Materials and methods

Constructs

PSG5-13-HA-RPTP α full length, deletions and point mutations, and PCS+MT (myc-tagged vector) RPTP α -D2 (amino acids 537–793) were described previously (Blanchetot and den Hertog, 2000). RPTP α -SpD2 (amino acids 504–793) was cloned in PCS2+MT by PCR. PCS2+MT RPTP α -SpD2 Δ C-t was made in the same way as RPTP α -D2 Δ C-t, by cloning a *NcoI*–*EcoRI* fragment in PCS2+MT, deleting residues 774–792 and the endogenous stop codon, which led to replacement of the RPTP α sequence by some vector sequence, leaving the overall size of the protein virtually identical. Point mutations were made by site-directed mutagenesis, and verified by sequencing.

Cell cultures and transfections

293 human embryonic kidney (HEK) cells were routinely grown in DF medium supplemented with 7.5% fetal calf serum. Cells were transfected using the standard calcium phosphate method. Briefly, 10-cm dishes were transfected with a total of 20 μ g of DNA. The next day, the medium was refreshed, and left another 16 h before harvesting.

Immunoprecipitation and immunoblotting

Subconfluent (stimulated) cells were washed twice with ice-cold phosphate-buffered saline, and lysed in cell lysis buffer CLB (50 mM HEPES pH 7.4, 150 mM NaCl, 1 mM MgCl₂, 10% glycerol, 1% TX-100 and protease inhibitors, including benzamide, aprotinin and leupeptin) for 20 min on ice, harvested, and centrifuged at 14 000 *g* for 15 min. Heat shock was carried out by placing the dish in a 42°C pre-warmed waterbath, and UV treatment by placing an open dish in a Stratagene (Stratagene). Cross-linking was performed using BS³ (Pierce) as described (Tertoolen *et al.*, 2001). Lysates were added to 12CA5 antibodies that had been pre-coupled to protein A–Sepharose beads. After 2 h of incubation, the beads were carefully washed four times with HNTG buffer (20 mM HEPES pH 7.4, 150 mM NaCl, 0.1% TX-100, 10% glycerol), mixed with 2 \times Laemmli buffer and loaded on a 7.5% SDS–PAGE gel. The proteins were transferred to PVDF membrane using a semidry transfer system. After Coomassie Blue staining, the membrane was blocked for 1 h with 5% milk in TBS–Tween (50 mM Tris pH 8.0, 150 mM NaCl, 0.05% Tween-20) at room temperature, incubated with the first antibody [anti-HA tag, 12CA5; or anti-Myc tag, 9E10 (Santa Cruz)] for 1.5 h, washed four times with TBS–Tween, incubated with horseradish peroxidase-conjugated anti-mouse antibody (Transduction Laboratories) for 1 h, washed four times with TBS–Tween, and developed using enhanced chemiluminescence.

PTP activity assays

After immunoprecipitation (see above) supplemented with two washes with PTP buffer (0.1 mM succinic acid pH 6.0, 150 mM NaCl, 1 mM EDTA), the beads were incubated with 200 μ l of PTP buffer containing 10 mM pNPP and left at 30°C. After sufficient time (15–60 min), aliquots (80 μ l) were taken and the reaction was stopped by addition of NaOH (100 μ l, 2 M) and the OD_{415 nm} measured. Several aliquots were taken at different times of incubation. The phosphatase activity was corrected for the amount of RPTP α present in the phosphatase assay. Aliquots of the phosphatase assay were taken and loaded on an SDS–PAGE gel. The gel was blotted and RPTP α was quantified using the Attophos detection system (Amersham) and a FluorImager (Molecular Dynamics).

FRET analysis

A Leitz orthoplan upright microscope (Leitz GMBH, Wetzlar, Germany) was used, equipped with an epi-illumination fluorescence detection system and a temperature-controlled specimen holder at 33°C. Measurements were made in a buffer containing 145 mM NaCl, 5 mM KCl, 2 mM MgCl₂, 2 mM CaCl₂, 10 mM HEPES, 10 mM glucose and 0.5% bovine serum albumin. As an excitation source, a SPEX Fluorolog (SPEX Industries, Edison, NJ) with two excitation monochromators was used. Excitation of CFP and YFP was performed at 430(8) and 490(8) nm, respectively. For the spectral analysis of CFP and the FRET measurements, a filter cube fitted with a 455 nm dichroic mirror was used. The YFP spectra were measured with a dichroic mirror >510 nm. Spectral data were recorded with an integration time of 0.5 s/nm, slit 8 nm. Real-time dual excitation measurements were made by alternating excitation at 430 and 490 nm and analysis of YFP emission at 535(30) nm, essentially as described (Tertoolen *et al.*, 2001). All measurements were made with a Leitz 50 \times NA 1.0 water immersion objective. The spectra from cell lysates were collected at room temperature using a Perkin Elmer LS50B

(UK) and a quartz ultra micro cell (120 μ l; Hellma, Rijswijk, The Netherlands), excitation 430(10) nm.

293 cells were cultured on glass coverslips and transfected with the appropriate construct. The constructs used for the FRET measurements were CFP–RPTP α -SpD2–YFP2.1, mutant CFP–RPTP α -SpD2–C723S–YFP2.1, CFP–RPTP α -SpD2–CFP and YFP–RPTP α -SpD2–YFP. CFP, YFP2.1 and RPTP α -SpD2 (residues Ser504–Lys793) were amplified by PCR, introducing convenient restriction sites, and cloned into pCS2+. The resulting constructs were checked by sequencing. YFP2.1 contains two extra point mutations (Val68 to Leu and Glu69 to Lys) that render YFP less sensitive to pH changes (Miyawaki *et al.*, 1999). Importantly, the original YFP is highly sensitive to H₂O₂ treatment, which induced decreases in YFP emission of up to 50% (C.B., L.G.J.T. and J.d.H., unpublished), while YFP2.1 is practically insensitive to H₂O₂ (Figure 4B).

Acknowledgements

We thank John Overvoorde for his technical assistance, and Astrid van der Sar and Jaap van Hellemond for helpful discussions. These investigations were (in part) supported by the Research Council for Earth and Life sciences (ALW) with financial aid from the Netherlands Organization for Scientific Research (NWO).

References

- Aricescu, A.R., Fulga, T.A., Cismasiu, V., Goody, R.S. and Szedlaczek, S.E. (2001) Intramolecular interactions in protein tyrosine phosphatase RPTP μ : kinetic evidence. *Biochem. Biophys. Res. Commun.*, **280**, 319–327.
- Bae, Y.S., Kang, S.W., Seo, M.S., Baines, I.C., Tekle, E., Chock, P.B. and Rhee, S.G. (1997) Epidermal growth factor (EGF)-induced generation of hydrogen peroxide. Role in EGF receptor-mediated tyrosine phosphorylation. *J. Biol. Chem.*, **272**, 217–221.
- Bastiaens, P.I. and Pepperkok, R. (2000) Observing proteins in their natural habitat: the living cell. *Trends Biochem. Sci.*, **25**, 631–637.
- Bilwes, A.M., den Hertog, J., Hunter, T. and Noel, J.P. (1996) Structural basis for inhibition of receptor protein-tyrosine phosphatase- α by dimerization. *Nature*, **382**, 555–559.
- Blanchetot, C. and den Hertog, J. (2000) Multiple interactions between receptor protein-tyrosine phosphatase (RPTP) α and membrane-distal protein-tyrosine phosphatase domains of various RPTPs. *J. Biol. Chem.*, **275**, 12446–12452.
- Buist, A., Zhang, Y.L., Keng, Y.F., Wu, L., Zhang, Z.Y. and den Hertog, J. (1999) Restoration of potent protein-tyrosine phosphatase activity into the membrane-distal domain of receptor protein-tyrosine phosphatase α . *Biochemistry*, **38**, 914–922.
- Caselli, A., Marzocchini, R., Camici, G., Manao, G., Moneti, G., Pieraccini, G. and Ramponi, G. (1998) The inactivation mechanism of low molecular weight phosphotyrosine-protein phosphatase by H₂O₂. *J. Biol. Chem.*, **273**, 32554–32560.
- Choi, H., Kim, S., Mukhopadhyay, P., Cho, S., Woo, J., Storz, G. and Ryu, S. (2001) Structural basis of the redox switch in the OxyR transcription factor. *Cell*, **105**, 103–113.
- Cubitt, A.B., Heim, R., Adams, S.R., Boyd, A.E., Gross, L.A. and Tsien, R.Y. (1995) Understanding, improving and using green fluorescent proteins. *Trends Biochem. Sci.*, **20**, 448–455.
- den Hertog, J. (1999) Protein-tyrosine phosphatases in development. *Mech. Dev.*, **85**, 3–14.
- den Hertog, J., Tracy, S. and Hunter, T. (1994) Phosphorylation of receptor protein-tyrosine phosphatase α on Tyr789, a binding site for the SH3-SH2-SH3 adaptor protein GRB-2 *in vivo*. *EMBO J.*, **13**, 3020–3032.
- Denu, J.M. and Dixon, J.E. (1998) Protein tyrosine phosphatases: mechanisms of catalysis and regulation. *Curr. Opin. Chem. Biol.*, **2**, 633–641.
- Denu, J.M. and Tanner, K.G. (1998) Specific and reversible inactivation of protein tyrosine phosphatases by hydrogen peroxide: evidence for a sulfenic acid intermediate and implications for redox regulation. *Biochemistry*, **37**, 5633–5642.
- Desai, D.M., Sap, J., Schlessinger, J. and Weiss, A. (1993) Ligand-mediated negative regulation of a chimeric transmembrane receptor tyrosine phosphatase. *Cell*, **73**, 541–554.
- Desai, C.J., Gindhart, J.G.J., Goldstein, L.S. and Zinn, K. (1996) Receptor tyrosine phosphatases are required for motor axon guidance in the *Drosophila* embryo. *Cell*, **84**, 599–609.
- Fauman, E.B., Cogswell, J.P., Lovejoy, B., Rocque, W.J., Holmes, W.,

- Montana, V.G., Piwnica-Worms, H., Rink, M.J. and Saper, M.A. (1998) Crystal structure of the catalytic domain of the human cell cycle control phosphatase, Cdc25A. *Cell*, **93**, 617–625.
- Feiken, E., van Etten, I., Gebbink, M.F., Moolenaar, W.H. and Zondag, G.C. (2000) Intramolecular interactions between the juxtamembrane domain and phosphatase domains of receptor protein-tyrosine phosphatase RPTP μ . Regulation of catalytic activity. *J. Biol. Chem.*, **275**, 15350–15356.
- Felberg, J. and Johnson, P. (1998) Characterization of recombinant CD45 cytoplasmic domain proteins. Evidence for intramolecular and intermolecular interactions. *J. Biol. Chem.*, **273**, 17839–17845.
- Garrity, P.A., Lee, C.H., Salecker, I., Robertson, H.C., Desai, C.J., Zinn, K. and Zipursky, S.L. (1999) Retinal axon target selection in *Drosophila* is regulated by a receptor protein tyrosine phosphatase. *Neuron*, **22**, 707–717.
- Gross, S., Knebel, A., Tenev, T., Neining, A., Gaestel, M., Herrlich, P. and Bohmer, F.D. (1999) Inactivation of protein-tyrosine phosphatases as mechanism of UV-induced signal transduction. *J. Biol. Chem.*, **274**, 26378–26386.
- Hayami-Noumi, K., Tsuchiya, T., Moriyama, Y. and Noumi, T. (2000) Intra- and intermolecular interactions of the catalytic domains of human CD45 protein tyrosine phosphatase. *FEBS Lett.*, **468**, 68–72.
- Hoffmann, K.M., Tonks, N.K. and Barford, D. (1997) The crystal structure of domain 1 of receptor protein-tyrosine phosphatase μ . *J. Biol. Chem.*, **272**, 27505–27508.
- Hunter, T. (1995) Protein kinases and phosphatases: the yin and yang of protein phosphorylation and signaling. *Cell*, **80**, 225–236.
- Jiang, G., den Hertog, J., Su, J., Noel, J., Sap, J. and Hunter, T. (1999) Dimerization inhibits the activity of receptor-like protein-tyrosine phosphatase- α . *Nature*, **401**, 606–610.
- Jiang, G., den Hertog, J. and Hunter, T. (2000) Receptor-like protein tyrosine phosphatase α homodimerizes on the cell surface. *Mol. Cell Biol.*, **20**, 5917–5929.
- Knebel, A., Rahmsdorf, H.J., Ullrich, A. and Herrlich, P. (1996) Dephosphorylation of receptor tyrosine kinases as target of regulation by radiation, oxidants or alkylating agents. *EMBO J.*, **15**, 5314–5325.
- Kokel, M., Borland, C.Z., DeLong, L., Horvitz, H.R. and Stern, M.J. (1998) *clr-1* encodes a receptor tyrosine phosphatase that negatively regulates an FGF receptor signaling pathway in *Caenorhabditis elegans*. *Genes Dev.*, **12**, 1425–1437.
- Krueger, N.X., Streuli, M. and Saito, H. (1990) Structural diversity and evolution of human receptor-like protein tyrosine phosphatases. *EMBO J.*, **9**, 3241–3252.
- Krueger, N.X., Van Vactor, D., Wan, H.I., Gelbart, W.M., Goodman, C.S. and Saito, H. (1996) The transmembrane tyrosine phosphatase DLAR controls motor axon guidance in *Drosophila*. *Cell*, **84**, 611–622.
- Lee, S.R., Kwon, K.S., Kim, S.R. and Rhee, S.G. (1998) Reversible inactivation of protein-tyrosine phosphatase 1B in A431 cells stimulated with epidermal growth factor. *J. Biol. Chem.*, **273**, 15366–15372.
- Lim, K.L., Kolatkar, P.R., Ng, K.P., Ng, C.H. and Pallen, C.J. (1998) Interconversion of the kinetic identities of the tandem catalytic domains of receptor-like protein-tyrosine phosphatase PTP α by two point mutations is synergistic and substrate-dependent. *J. Biol. Chem.*, **273**, 28986–28993.
- Majeti, R., Bilwes, A.M., Noel, J.P., Hunter, T. and Weiss, A. (1998) Dimerization-induced inhibition of receptor protein tyrosine phosphatase function through an inhibitory wedge. *Science*, **279**, 88–91.
- Meng, K., Rodriguez-Pena, A., Dimitrov, T., Chen, W., Yamin, M., Noda, M. and Deuel, T.F. (2000) Pleiotrophin signals increased tyrosine phosphorylation of β β -catenin through inactivation of the intrinsic catalytic activity of the receptor-type protein tyrosine phosphatase β/ζ . *Proc. Natl Acad. Sci. USA*, **97**, 2603–2608.
- Miyawaki, A., Llopis, J., Heim, R., McCaffery, J.M., Adams, J.A., Ikura, M. and Tsien, R.Y. (1997) Fluorescent indicators for Ca²⁺ based on green fluorescent proteins and calmodulin. *Nature*, **388**, 882–887.
- Miyawaki, A., Griesbeck, O., Hein, R. and Tsien, R.Y. (1999) Dynamic and quantitative Ca²⁺ measurements using improved cameleons. *Proc. Natl Acad. Sci. USA*, **96**, 2135–2140.
- Nagai, Y., Miyazaki, M., Aoki, R., Zama, T., Inouye, S., Hirose, K., Iino, M. and Hagiwara, M. (2000) A fluorescent indicator for visualizing cAMP-induced phosphorylation *in vivo*. *Nature Biotechnol.*, **18**, 313–316.
- Nam, H.J., Poy, F., Krueger, N.X., Saito, H. and Frederick, C.A. (1999) Crystal structure of the tandem phosphatase domains of RPTP LAR. *Cell*, **97**, 449–457.
- Neel, B.G. and Tonks, N.K. (1997) Protein tyrosine phosphatases in signal transduction. *Curr. Opin. Cell Biol.*, **9**, 193–204.
- Pregel, M.J. and Storer, A.C. (1997) Active site titration of the tyrosine phosphatases SHP-1 and PTP1B using aromatic disulfides. Reaction with the essential cysteine residue in the active site. *J. Biol. Chem.*, **272**, 23552–23558.
- Song, H., Hanlon, N., Brown, N.R., Noble, M.E., Johnson, L.N. and Barford, D. (2001) Phosphoprotein–protein interactions revealed by the crystal structure of kinase-associated phosphatase in complex with phosphoCDK2. *Mol. Cell*, **7**, 615–626.
- Tertoolen, L.G., Blanchetot, C., Jiang, G., Overvoorde, J., Gadella, T.W.J., Hunter, T. and den Hertog, J. (2001) Dimerization of receptor protein-tyrosine phosphatase α in living cells. *BMC Cell Biol.*, **2**, 8.
- Van Vactor, D., O'Reilly, A.M. and Neel, B.G. (1998) Genetic analysis of protein tyrosine phosphatases. *Curr. Opin. Genet. Dev.*, **8**, 112–126.
- Wallace, M.J., Fladd, C., Batt, J. and Rotin, D. (1998) The second catalytic domain of protein tyrosine phosphatase δ (PTP δ) binds to and inhibits the first catalytic domain of PTP σ . *Mol. Cell Biol.*, **18**, 2608–2616.
- Wang, Y. and Pallen, C.J. (1991) The receptor-like protein tyrosine phosphatase HPTP α has two active catalytic domains with distinct substrate specificities. *EMBO J.*, **10**, 3231–3237.
- Zheng, X.M. and Shalloway, D. (2001) Two mechanisms activate PTP α during mitosis. *EMBO J.*, **20**, 6037–6049.

Received August 9, 2001; revised December 12, 2001;
accepted December 21, 2001