

## Integrated Recombinant Protein Expression and Purification Platform Based on *Ralstonia eutropha*

Gavin C. Barnard,<sup>1</sup> Jesse D. McCool,<sup>1</sup> David W. Wood,<sup>2</sup> and Tillman U. Gerngross<sup>1\*</sup>

*Thayer School of Engineering, Department of Biological Sciences, and Department of Chemistry, Dartmouth College, Hanover, New Hampshire, 03755,<sup>1</sup> and Department of Chemical Engineering, Princeton University, Princeton, New Jersey 08544<sup>2</sup>*

Received 24 February 2005/Accepted 13 May 2005

**Protein purification of recombinant proteins constitutes a significant cost of biomanufacturing and various efforts have been directed at developing more efficient purification methods. We describe a protein purification scheme wherein *Ralstonia eutropha* is used to produce its own “affinity matrix,” thereby eliminating the need for external chromatographic purification steps. This approach is based on the specific interaction of phasin proteins with granules of the intracellular polymer polyhydroxybutyrate (PHB). By creating in-frame fusions of phasins and green fluorescent protein (GFP) as a model protein, we demonstrated that GFP can be efficiently sequestered to the surface of PHB granules. In a second step, we generated a phasin-intein-GFP fusion, wherein the self-cleaving intein can be activated by the addition of thiols. This construct allowed for the controlled binding and release of essentially pure GFP in a single separation step. Finally, pure, active  $\beta$ -galactosidase was obtained in a single step using the above described method.**

We have previously reported the development of a novel high cell density protein expression platform based on the gram-negative bacterium *Ralstonia eutropha* (22, 23). This system has been developed to overcome some of the shortcomings associated with recombinant protein expression in *Escherichia coli* (e.g., poor fermentation performance at high cell density, and inclusion body formation). Expression of organophosphohydrolase, an enzyme originally isolated from *Pseudomonas diminuta* (20) and prone to inclusion body formation in *Escherichia coli* (4, 28, 29), was demonstrated at high levels. Titters of active, soluble organophosphohydrolyase, in excess of 10 g/liter were obtained in high cell density fermentation (3), representing at least a 100-fold increase over those previously reported in *E. coli*.

While the successful expression of a recombinant protein is a necessary requirement, recovery and purification still remain a significant cost in recombinant protein production. We thus sought to integrate the existing *R. eutropha* protein expression platform with a protein purification strategy to simplify the expression and purification of recombinant proteins. This specific approach uses the natural ability of *R. eutropha* to produce a polymer known as polyhydroxybutyrate (PHB), which accumulates as insoluble granules within the cell. PHB is a member of the polyhydroxyalkanoate class of polymers, synthesized by many bacteria, as carbon storage compounds (2, 15, 16, 26, 30, 31, 32). Polyhydroxyalkanoates have received attention as biodegradable polymers and can be obtained by fermentation processes utilizing cheap, abundant renewable carbon sources (2, 24). Polyhydroxyalkanoates have been produced industrially by ZENECA Bioproducts (26) and Monsanto (10).

PHB synthesis in *R. eutropha* has been the model system for studying polyhydroxyalkanoate biosynthesis in bacteria (10, 15, 16, 18, 26). The biogenesis of polyhydroxyalkanoate granules involves two distinct proteins, the polyhydroxyalkanoate synthase (PhaC) and phasins (PhaP). Phasins are low-molecular-weight proteins whose role in polyhydroxyalkanoate formation is not well understood (10, 11, 15, 16, 18, 24, 26, 30, 31, 32). Phasins accumulate during PHB synthesis, bind to PHB granules, and promote further PHB synthesis (32). It has been shown that *phaP* mutants form only one large PHB granule and that up-regulating the *phaP* gene increases the number of PHB granules while reducing their size (15, 26). Phasins accumulate at high levels in cells that are synthesizing PHB, and as much as 5% of total cellular protein can be PhaP (16). Phasins have high affinity for PHB granules and are the predominant protein present on the granule surface (24, 26).

In this study, we exploit the specific affinity between PhaP and the PHB granules for the purpose of purifying a recombinant model protein (GFP). In essence, this yields an affinity-based purification scheme wherein the cell synthesizes its own chromatography matrix and PhaP is used as the affinity tag to sequester a protein of interest to the PHB granule surface. The recombinant protein can then be recovered by cell disruption followed by a centrifugation step that separates the insoluble high-density polymer from other soluble cellular components. In an improved version, the protein of interest is linked to PhaP through an intein allowing its release by thiol induced cleavage (see Fig. 3).

Inteins are self-splicing proteins that occur as in-frame insertions in specific host proteins throughout nature, and have been adapted for use in recombinant protein expression and purification schemes (9, 27). Intein cleavage can be mediated by pH changes or the addition of thiols. The *Mxe* GyrA intein is a 198-amino-acid polypeptide, with N-terminal cleavage activity in the presence of thiols (25). By incorporating this intein into a PhaP-intein-GFP fusion we were able to show (i) the

\* Corresponding author. Mailing address: Thayer School of Engineering, Department of Biological Sciences, and Department of Chemistry, Dartmouth College, 8000 Cummings Hall, Hanover, NH 03755. Phone: (603) 646-3161. Fax: (603) 646-2277. E-mail: tillman.gerngross@dartmouth.edu.

TABLE 1. Strains and plasmids used in this study

Strain or plasmid	Description <sup>a</sup>	Reference or source
<i>R. eutropha</i> strains		
NCIMB 40124	Wild type; gentamicin resistant	National Collections of Industrial, Food & Marine Bacteria, Aberdeen, Scotland
G	pG introduced into <i>R. eutropha</i> wt	This study
PG	pPG introduced into <i>R. eutropha</i> wt	This study
PIG	pPIG introduced into <i>R. eutropha</i> wt	This study
GP	pGP introduced into <i>R. eutropha</i> wt	This study
GIP	pGIP introduced into <i>R. eutropha</i> wt	This study
PIL	pPIL introduced into <i>R. eutropha</i> wt	This study
<i>E. coli</i> strains		
TOP10	Host strain for plasmids derived from pCR2.1-TOPO	Invitrogen
S-17	Host strain for plasmids derived from pKNOCK-Cm	8, 17, 21
Plasmids		
pG	<i>phaPp::gfp</i> transcriptional fusion introduced into pKNOCK-Cm; plasmid used to create strain <i>R. eutropha</i> G	This study
pPG	<i>phaP</i> ORF:: <i>gfp</i> translational fusion introduced into pKNOCK-Cm; plasmid used to create strain <i>R. eutropha</i> PG	This study
pPIG	<i>phaP</i> ORF:: <i>Mxe</i> GyrA intein:: <i>gfp</i> translational fusion in pKNOCK-Cm; plasmid used to create strain <i>R. eutropha</i> PIG	This study
pGP	<i>phaPp::gfp::phaP</i> transcriptional fusion in pKNOCK-Cm; plasmid used to create strain <i>R. eutropha</i> GP	This study
pGIP	<i>phaPp::gfp::Mxe</i> GyrA intein:: <i>phaP</i> transcriptional fusion in pKNOCK-Cm; plasmid used to create strain <i>R. eutropha</i> GIP	This study
pPIL	<i>phaP</i> ORF:: <i>Mxe</i> GyrA intein:: <i>lacZ</i> translational fusion in pKNOCK-Cm; plasmid used to create strain <i>R. eutropha</i> PIL	This study
pKNOCK-Cm	Suicide vector used for introducing plasmids into <i>R. eutropha</i> chromosome	1
pGY1a+	Vector containing <i>phaPp::gfpmut2</i> translational fusion	30
pTWIN1	Commercially available expression vector containing <i>Ssp</i> DnaB intein and <i>Mxe</i> GyrA intein	New England Biolabs
pUCPPCm	Plasmid vector containing <i>phaP</i> promoter	22
pGB27	pKNOCK containing the <i>phaP</i> promoter	This study
pGB73	<i>Mxe</i> GyrA intein introduced into pCR2.1-TOPO	This study
pGB76	<i>phaP</i> ORF introduced into pCR2.1-TOPO	This study
pGB80	<i>phaPp::gfp::phaP</i> transcriptional fusion in pKNOCK-Cm	This study
pGB82	<i>phaPp::gfp::NEB</i> <i>Mxe</i> GyrA intein:: <i>phaP</i> transcriptional fusion in pKNOCK-Cm	This study
pGB85	<i>phaP::gfp</i> transcriptional fusion in pCR4-TOPO	This study
pGB91	<i>phaP::gfp</i> transcriptional fusion in pKNOCK-Cm	This study
pGB93	<i>phaP::NEB</i> <i>Mxe</i> GyrA intein:: <i>gfp</i> transcriptional fusion in pKNOCK-Cm	This study
pGB96	<i>Peptide linker::phaP</i> transcriptional fusion in pCR2.1-TOPO	This study
pGB97	<i>phaP::peptide linker</i> transcriptional fusion in pCR2.1-TOPO	This study
pGB470	<i>lacZ</i> ORF fragment cloned introduced into pCR4Blunt-TOPO	This study

<sup>a</sup> wt, wild type.

expression of a PhaP-intein-GFP fusion protein, (ii) its sequestration to PHB granules, and (iii) the subsequent release of GFP from the PHB granule by treating the cell debris with dithiothreitol.

#### MATERIALS AND METHODS

**Strains, plasmids and oligonucleotides.** Strains and plasmids that were used in this study are listed in Table 1. Oligonucleotides used in this study are listed in Table 2. Standard procedures were used for the preparation and manipulation of DNA and for PCR. All PCR products were subcloned into pCR2.1-TOPO (Invitrogen), pCR4Blunt-TOPO (Invitrogen) or pCRII-Blunt-TOPO (Invitrogen) and sequence verified at the Molecular Biology Core Facility at Dartmouth College. Methods for introducing plasmids into the *R. eutropha* chromosome have previously been described (22, 23). In brief, all pKNOCK-Cm derived plasmids are introduced into *E. coli* S17 (17, 21) before being incorporated into the *R. eutropha* chromosome by simple biparental mating.

**Growth media, antibiotics and cultivation conditions.** *E. coli* strains were grown in Luria-Bertani (LB) medium (13). *R. eutropha* strains were grown in one of the following media depending on the application: LB medium or Lee me-

dium (20 g/liter glucose, 3 g/liter Na<sub>2</sub>HPO<sub>4</sub> · 7H<sub>2</sub>O, 1 g/liter KH<sub>2</sub>PO<sub>4</sub>, 2 g/liter NH<sub>4</sub>Cl, 0.2 g/liter MgSO<sub>4</sub> · 7H<sub>2</sub>O, 1 ml/liter Corn Steep Liquor [Sigma], 2.4 ml/liter trace element solution [22]). Antibiotics were added to the growth medium to the following concentrations depending on the application: chloramphenicol (50 µg/ml), kanamycin (50 µg/ml), and gentamicin (10 µg/ml). *R. eutropha* and *E. coli* strains were cultivated at 30°C and 37°C, respectively.

**Fluorescence microscopy.** To prepare cells for fluorescence microscopy, cells were transferred from LB agar plates into 200 µl of buffer (phosphate-buffered saline) and resuspended thoroughly; 10 µl of this cell suspension were transferred to a single well in a 15-well slide pretreated with 1% poly-L-lysine. Microscopy was carried out using a Leica epifluorescence light microscope. An ORCA-ER charge-coupled device camera (Hamamatsu) and OPENLAB software (Improvision) were used for all image acquisition and processing.

**Sucrose gradient fractionation.** Strains were cultivated in 50 ml of Lee medium to an approximate optical density at 600 nm of 10. The cultures were centrifuged and the cells resuspended in 2 ml of buffer B1 (20 mM Tris, 500 mM NaCl, 1 mM EDTA, pH 8.5). Cells were sonicated in a Fisher Scientific Sonic Dismembrator 550 in ten pulsed cycles (2 seconds ON, 0.5 second off, 30 second duration, 5 min cooling on ice between cycles); 1 ml of the lysate was loaded onto a sucrose density gradient. The sucrose density gradient consists of nine layered 1 ml fractions of buffer B1 containing 0 to 2 M sucrose (0.25 M increments). The






FIG. 1. Representative GFP localization. Fluorescence microscopy and sucrose density gradient fractionation of cell lysates. a) Wild-type *R. eutropha*. b) *R. eutropha* G (expressing GFP). c) *R. eutropha* PG (expressing PhaP-GFP). d) *R. eutropha* PIG (expressing PhaP-intein-GFP). Cell lysates were generated by sonication and added to the top of a sucrose density gradient. Fluorescence of individual sucrose density gradient fractions is expressed in relative fluorescence units divided by 100 and shown in green. Corresponding PHB concentrations are expressed in mg/liter and shown in tan.

SacI/AscI fragment and replaced with the 0.6-kb SacI/AscI fragment of pGB97 to yield plasmids pPG and pPIG, respectively.

**Construction of plasmid pPIL (*phaP*::peptide linker::intein::lacZ translational fusion in pKNOCK-Cm).** The *lacZ* ORF was PCR amplified from *E. coli* S17 genomic DNA using oligonucleotides oGB216 and oGB217. The 3.0-kb fragment was subcloned into pCRII-Blunt-TOPO to generate pGB470. The *lacZ* ORF from pGB470 was cloned into pPIG as an XbaI/XhoI fragment, yielding plasmid pPIL.

***R. eutropha* strain generation.** *Ralstonia eutropha* recombinant strains were generated according to methods previously described (3, 22, 23). *R. eutropha* G, was generated using plasmid pG, which carries a transcriptional fusion between the *phaP* promoter and the *gfpmut2* ORF (6) (*phaPp::gfp*). Plasmid pG is a suicide plasmid and is integrated at the *phaP* promoter locus of the *R. eutropha* chromosome. Since integration occurs within the promoter region, the wild-type *phaP* gene remains intact. *R. eutropha* PG and *R. eutropha* PIG were generated using plasmids pPG and pPIG, respectively. Plasmid pPG contains an in-frame translational fusion between the *phaP* ORF and *gfpmut2* ORF (*phaP::gfp*). Plasmid pPIG is isogenic to pPG, with the exception of the in-frame insertion of the *Mxe* GyrA intein between the two open reading frames (*phaP::Mxe* GyrA intein::gfp). Plasmids pPG and pPIG do not contain the *phaP* promoter and the *phaP* ORF serves as the homologous recombination locus. Therefore in *R. eutropha* PG and *R. eutropha* PIG, the wild-type *phaP* gene has been replaced by a translational fusion encoding *phaP::gfp* and *phaP::intein::gfp*, respectively.

## RESULTS

Both fluorescence microscopy and sucrose density gradient fractionation of cell lysates were used to examine localization of GFP in *R. eutropha* strains. The fluorescence microscopy images in Fig. 1 show that the wild type exhibited no autofluorescence and that GFP is evenly distributed throughout the cell in *R. eutropha* G. However, Fig. 1 shows fluorescent foci throughout the cells in *R. eutropha* PG and *R. eutropha* PIG, presumably where GFP is localized on the surface of PHB granules.

Sucrose density gradient fractionation of cell lysates was performed to further examine GFP localization (see Materials and Methods). *R. eutropha* strains were cultivated in Lee medium, a phosphate limited growth medium that induces both PHB formation and transcription of genes under the control of the *phaP* promoter. Cells were recovered, washed, resuspended in buffer B1, and sonicated. Cell lysates were loaded onto a sucrose gradient (density from 1.02 g/ml to 1.29 g/ml) and equilibrated by centrifugation. PHB granules have a density of approximately 1.20 g/ml (19) and accumulate near the bottom of the sucrose density gradient. In contrast, soluble proteins accumulate in the low density fractions at the top of the sucrose density gradient. A fluorescence spectrophotometer was used to measure the fluorescence of each individual fraction of the sucrose gradient. *R. eutropha* G showed fluorescence predominantly in the top fractions, consistent with fluorescence micrographs that suggest that GFP is present as a soluble protein in the cytoplasm and not localized to PHB granules.

*R. eutropha* PG and *R. eutropha* PIG showed a strong fluorescent signal in fraction 7, which coincides with the fraction containing PHB. These results strongly suggest that in *R. eutropha* PG and *R. eutropha* PIG, the GFP is localized to the PHB granules.

A fluorescent signal also appeared in the upper fractions of the *R. eutropha* PG and *R. eutropha* PIG density gradients. We propose several possible explanations for the presence of soluble GFP in these strains: (i) excess PhaP-GFP and PhaP-intein-GFP cannot bind because the PHB granule binding capacity has been exhausted, (ii) affinity is reduced due to the

C-terminal fusion of GFP to the PhaP protein, or (iii) the minor fluorescence could represent a small amount of desorption that occurs during sonication and sucrose density gradient fractionation.

From the fluorescence microscopy and sucrose density fractionation, we concluded that PhaP-GFP and PhaP-intein-GFP fusions are localized in vivo to PHB granules. A cleavage experiment was designed to demonstrate the release of pure GFP from whole cell debris (see Materials and Methods). Briefly, *R. eutropha* strains were cultivated in Lee medium, harvested, resuspended in buffer B1 and sonicated. The lysate was centrifuged and the supernatant fraction, containing the soluble protein fraction, was discarded. The pellet was washed in buffer B1. To induce intein cleavage, the pellet was resuspended in buffer B2, and incubated overnight at 37°C. The mixture was then centrifuged and the pellet and supernatant fractions both retained. The pellet was again washed with buffer B1.




FIG. 2. Intein mediated cleavage of GFP and beta-galactosidase from whole cell debris. *R. eutropha* strains were lysed by sonication, the supernatant discarded and the insoluble pellet containing PHB granules retained. Intein mediated cleavage was activated by incubating the washed pellet overnight in buffer B2 at 37°C. After incubation, the pellet and supernatant fractions were isolated. (a) Fluorometry. Green bars show the fluorescence of the supernatant fractions. Tan colored bars denote the fluorescence of the resulting pellet fraction. (b) SDS-PAGE of fractions. Lanes 2 and 3: *R. eutropha* wild-type pellet and supernatant following dithiothreitol treatment, respectively. Lanes 4 and 5: *R. eutropha* G pellet and supernatant. Lanes 6 and 7: *R. eutropha* PG pellet and supernatant. Lanes 8 and 9: *R. eutropha* PIG pellet and supernatant. (c) SDS-PAGE. Lane 2: *R. eutropha* PIG supernatant.




FIG. 3. Flowchart of a recombinant protein purification scheme based on PhaP mediated PHB granule sequestration. *Ralstonia eutropha* recombinants expressing phasin-intein-GFP (PIG) are cultivated in a shake flask or a bioreactor. Cells are harvested by centrifugation, washed and resuspended in buffer B1 prior to cell disruption. The cell lysate is centrifuged, the supernatant fraction discarded and the insoluble fraction retained. After washing the insoluble pellet, the fraction is resuspended in buffer B2. The intein is thiol activated and GFP is released from the whole cell debris into the supernatant. The pure protein is recovered by centrifugation, discarding the pellet and retaining the supernatant fraction.

Figure 2A shows the fluorescence of the pellet and supernatant fractions. Neither the *R. eutropha* wild-type pellet nor the corresponding supernatant showed appreciable fluorescence. Similarly, the pellet and supernatant fractions of *R. eutropha* G showed no appreciable fluorescence as expected. As expected, *R. eutropha* PG showed strong fluorescence on the pellet with no appreciable fluorescence present in the supernatant. In contrast, *R. eutropha* PIG showed very strong fluorescence in the supernatant fraction, indicating that GFP had been released from the pellet into the supernatant fraction. Although the bulk of the total fluorescence was present in the supernatant, a minor amount of fluorescence remained on the PHB granule. To confirm the dithiothreitol-mediated release of GFP and the absence of pH effects, an identical experiment was performed in which the cell debris fraction generated from *R. eutropha* PIG was resuspended in buffer B1 (i.e., buffer B2 lacking dithiothreitol). No appreciable fluorescence was found in the supernatant fraction and the fluorescence of the cell debris remained unchanged (data not shown).

Figure 2B shows an SDS-PAGE gel for the pellet and supernatant fractions with the corresponding fluorescence data depicted in Fig. 2A. As expected, the whole cell debris for each strain contains numerous proteins. No protein is visible on the gel for the supernatant fractions of *R. eutropha* wild-type, *R. eutropha* G, and *R. eutropha* PG. The PhaP-intein-GFP fusion protein is expected to be 70 kDa in size. If intein-mediated cleavage occurs, a protein of 49 kDa, corresponding to an intein-GFP (IG) fusion, should be released. Figure 2B, lane 9, shows that IG was the only protein present in the supernatant fraction. This observation confirms that intein-mediated cleavage, activated by thiol addition, released GFP from the granule in the cell debris of *R. eutropha* PIG.

To further investigate the robustness of this method we attempted to purify a relatively large, multimeric, catalytically active protein ( $\beta$ -galactosidase,  $4 \times 116$  kDa). *R. eutropha* PIL was generated using plasmid pPIL (see Table 1). Plasmid pPIL is isogenic to plasmid pPIG with the exception that the *gfpmut2* ORF in pPIG has been replaced with the *lacZ* ORF in pPIL. Therefore, in *R. eutropha* PIL, the wild-type *phaP* gene has been replaced by a *phaP::Mxe GyrA intein::lacZ* translational fusion.

Intein mediated release of pure  $\beta$ -galactosidase, using cell extracts of *R. eutropha* PIL was demonstrated. Craven et al. (7) reported a specific activity of 40,000 U/nmol (340,000 U/mg, 28°C, pH 7.0) for purified  $\beta$ -galactosidase obtained by three purification steps (ammonium sulfate precipitation, size exclusion chromatography and DEAE chromatography). Colby and Hu (5) reported 19,000 U/nmol (160,000 U/mg, 30°C, pH 7.0) for purified  $\beta$ -galactosidase obtained by five purification steps (ammonium sulfate precipitation, electrophoresis, DEAE chromatography, size exclusion chromatography and crystallization). Following the above described method, a specific activity of 53,000 U/nmol was measured in the supernatant following dithiothreitol treatment of whole cell debris. Thus, a single purification step, without external chromatography; results in a protein fraction of high purity consistent with previously reported data for pure  $\beta$ -galactosidase and SDS-PAGE (see Fig. 2C).

We have also explored the effect of orientation by using

PhaP as the C-terminal fusion partner (i.e., creating plasmids pGIP and pLIP). Similar to results generated using PhaP as the N-terminal fusion partner, we have shown that soluble, active GFP and  $\beta$ -galactosidase can be recovered in a single purification step when PhaP was used as the C-terminal fusion partner (data not shown).

## DISCUSSION

In this study, we report the development of an integrated protein expression and purification approach that obviates the need for external chromatography. By replacing the wild-type *phaP* gene with a triple translational fusion (*phaP* ORF, *Mxe GyrA* intein and *gfpmut2*), we were able to show that the fusion protein can be localized to the PHB granule and separated from the remaining cytosolic protein fraction by centrifugation. In a subsequent step, we were able to release pure GFP by resuspending whole cell debris (insoluble fraction of cell lysate, containing PHB granules) in a buffer containing dithiothreitol (see Fig. 3). We also expressed, recovered and purified  $\beta$ -galactosidase to demonstrate the applicability of this method to purify catalytically active proteins. We showed that the enzyme is essentially pure and that the specific activity is comparable to previously reported literature values for pure  $\beta$ -galactosidase.

The single step purification eliminates the need for elaborate and costly protein purification schemes. Moreover, adapting the use of inteins eliminates the need for specific endopeptidases, which are routinely used to release recombinant protein from affinity matrixes. It is contemplated that this system could be adapted for use in other hosts that either produce PHB granules naturally or hosts that have been engineered to produce PHB.

By integrating high-level recombinant protein expression with a simple, protein purification step, this system has the potential to improve upon current technologies for the large-scale production of commodity enzymes, therapeutic proteins, vaccines, and peptides.

## ACKNOWLEDGMENTS

This work was supported by ARO grant DAAD19-00, DOJ grant 2000-DT-CX-K001(S-1), and NIST grant 60NANB1D0064.

We thank George O' Toole (Dartmouth Medical School) for the use of his fluorescence microscopy equipment.

## REFERENCES

- Alexeyev, M. F. 1999. The pKNOCK series of broad-host-range mobilizable suicide vectors for gene knockout and targeted DNA insertion into the chromosome of Gram-negative bacteria. *BioTechniques* **26**:824-826.
- Anderson, A. J., and E. A. Dawes. 1990. Occurrence, metabolism, metabolic role, and industrial uses of bacterial polyhydroxyalkanoates. *Microbiol. Rev.* **54**:450-472.
- Barnard, G. C., G. H. Henderson, S. Srinivasan, and T. U. Gerngross. 2004. High level recombinant protein expression in *Ralstonia eutropha* using T7 RNA polymerase based amplification. *Protein Express. Purif.* **38**:264-271.
- Cha, H. J., C. F. Wu, J. J. Valdes, G. Rao, and W. E. Bentley. 2000. Observations of green fluorescent protein as a fusion partner in genetically engineered *Escherichia coli*: monitoring protein expression and solubility. *Bio-technol.* **18**:565-574.
- Colby, C., Jr., and A. S. Hu. 1968. Purification and comparison of the  $\beta$ -galactosidase synthesized by *Escherichia coli* F-lac+ and *Proteus mirabilis* F-lac+. *Biochim. Biophys. Acta* **157**:167-177.
- Cormack, B. P., R. H. Valdivia, and S. Falkow. 1996. FACS-optimized mutants of the green fluorescent protein (GFP). *Gene* **173**:33-38.
- Craven, G. R., E. Steers, Jr., and C. B. Anfinsen. 1965. Purification, com-

- position, and molecular weight of the  $\beta$ -galactosidase of *Escherichia coli* K12. *J. Biol. Chem.* **240**:2468–2477.
8. de Lorenzo, V., M. Herrero, U. Jakubzik, and K. N. Timmis. 1990. Mini-Tn5 transposon derivatives for insertion mutagenesis, promoter probing, and chromosomal insertion of cloned DNA in gram-negative eubacteria. *J. Bacteriol.* **172**:6568–6572.
  9. Derbyshire, V., D. W. Wood, W. Wu, J. T. Dansereau, J. Z. Dalgaard, and M. Belfort. 1997. Genetic definition of a protein-splicing domain: functional min-inteins support structure predictions and a model for intein evolution. *Proc. Natl. Acad. Sci. USA* **94**:11466–11471.
  10. Hanley, S. Z., D. J. Pappin, D. Rahman, A. J. White, K. M. Elborough, and A. R. Slabas. 1999. Re-evaluation of the primary structure of *Ralstonia eutropha* phasin and implications for polyhydroxyalkanoic acid granule binding. *FEBS Lett.* **447**:99–105.
  11. Jurasek, L., and R. H. Marchessault. 2002. The role of phasins in the morphogenesis of poly(3-hydroxybutyrate) granules. *Biomacromolecules* **3**:256–261.
  12. Karr, D. B., J. K. Waters, and D. W. Emerich. 1983. Analysis of poly- $\beta$ -hydroxybutyrate in *Rhizobium japonicum* bacteroids by ion-exclusion high-pressure liquid chromatography and UV detections. *Appl. Environ. Microbiol.* **60**:3952–3958.
  13. Luria, S. E., J. N. Adams, and R. C. Ting. 1960. Transduction of lactose-utilizing ability among strains of *E. coli* and *S. dysenteriae* and the properties of the transducing phage particles. *Virology* **12**:348–390.
  14. New England Biolabs. 2003. IMPACT-TWIN instruction manual, version 1.2, 12/03. New England Biolabs, Beverly, Mass.
  15. Pieper-Fürst, U., M. H. Madkour, F. Mayer, and A. Steinbüchel. 1995. Identification of the region of a 14-kilodalton protein of *Rhodococcus ruber* that is responsible for the binding of this phasing to polyhydroxyalkanoic acid granules. *J. Bacteriol.* **177**:2513–2523.
  16. Pötter, M., M. H. Madkour, F. Mayer, and A. Steinbüchel. 2002. Regulation of phasin expression and polyhydroxyalkanoate (PHA) granule formation in *Ralstonia eutropha* H16. *Microbiology* **148**:2413–2426.
  17. Quandt, J., and M. F. Hynes. 1993. Versatile suicide vectors which allow direct selection for gene replacement in Gram-negative bacteria. *Gene* **127**:15–21.
  18. Rehm, B. H., and A. Steinbüchel. 1999. Biochemical and genetic analysis of PHA synthases and other proteins required for PHA synthesis. *Int. J. Biol. Macromol.* **25**:3–19.
  19. Resch, S., K. Gruber, G. Wanner, S. Slater, D. Dennis, and W. Lubitz. 1998. Aqueous release and purification of poly(beta-hydroxybutyrate) from *Escherichia coli*. *J. Biotechnol.* **65**:173–182.
  20. Serdar, C. M., and D. T. Gibson. 1985. Enzymatic-hydrolysis of organophosphates-cloning and expression of a parathion hydrolase gene from *Pseudomonas diminuta*. *Biotechnology* **3**:567–571.
  21. Simon, R., U. Priefer, and A. Pühler. 1983. A broad host-range mobilization system for in vivo genetic engineering: transposon mutagenesis in Gram-negative bacteria. *Biotechnology* **1**:784–791.
  22. Srinivasan, S., G. C. Barnard, and T. U. Gerngross. 2002. A novel high cell density protein expression system based on *Ralstonia eutropha*. *Appl. Environ. Microbiol.* **68**:5925–5932.
  23. Srinivasan, S., G. C. Barnard, and T. U. Gerngross. 2003. Production of recombinant proteins using multiple copy gene integration in high cell density fermentations of *Ralstonia eutropha*. *Biotechnol. Bioeng.* **84**:114–120.
  24. Steinbüchel, A., R. Wiczorek, and N. Krüger. 1996. PHA biosynthesis, its regulation and application of C1-utilizing microorganisms for polyester production, p. 237–244. In M. E. Lidstrom and F. R. Tabita (ed.), *Microbial growth on C<sub>1</sub> compounds: proceedings of the 7th International Symposium on Microbial Growth on C1 Compounds*, San Diego, Calif., 27 August–1 September 1995.
  25. Telenti, A., M. Southworth, F. Alcaide, S. Daugelat, W. R. Jacobs, Jr., and F. B. Perler. 1997. The *Mycobacterium xenopi* GyrA protein splicing element: characterization of a minimal intein. *J. Bacteriol.* **179**:6378–6382.
  26. Wiczorek, R., A. Pries, A. Steinbüchel, and F. Mayer. 1995. Analysis of a 24-kilodalton protein associated with the polyhydroxyalkanoic acid granules in *Alcaligenes eutrophus*. *J. Bacteriol.* **177**:2425–2435.
  27. Wood, D. W., W. Wu, G. Belfort, V. Derbyshire, and M. Belfort. 1999. A genetic system yields self-cleaving inteins for bioseparations. *Nat. Biotechnol.* **17**:889–892.
  28. Wu, C. F., H. J. Cha, G. Rao, J. J. Valdes, and W. E. Bentley. 2000. A green fluorescent protein fusion strategy for monitoring the expression, cellular location, and separation of biologically active organophosphorus hydrolase. *Appl. Microbiol. Biotechnol.* **54**:78–83.
  29. Wu, C. F., H. J. Cha, G. Rao, J. J. Valdes, and W. E. Bentley. 2001. Enhancement of organophosphorus hydrolase yield in *Escherichia coli* using multiple gene fusions. *Biotechnol. Bioeng.* **75**:100–103.
  30. York, G. M., B. H. Junker, J. Stubbe, and A. J. Sinskey. 2001. Accumulation of the PhaP phasin of *Ralstonia eutropha* is dependent on production of polyhydroxybutyrate in cells. *J. Bacteriol.* **183**:4217–4226.
  31. York, G. M., J. Stubbe, and A. J. Sinskey. 2001. New insight into the role of the PhaP phasing of *Ralstonia eutropha* in promoting synthesis of polyhydroxybutyrate. *J. Bacteriol.* **183**:2394–2397.
  32. York, G. M., J. Stubbe, and A. J. Sinskey. 2002. The *Ralstonia eutropha* PhaR protein couples synthesis of the PhaP phasing to the presence of polyhydroxybutyrate in cells and promotes polyhydroxybutyrate production. *J. Bacteriol.* **184**:59–66.