

Differentiation of *Paenibacillus larvae* subsp. *larvae*, the Cause of American Foulbrood of Honeybees, by Using PCR and Restriction Fragment Analysis of Genes Encoding 16S rRNA

Adriana M. Alippi,^{1*} Ana Claudia López,² and O. Mario Aguilar²

Centro de Investigaciones de Fitopatología, Facultad de Ciencias Agrarias y Forestales,¹ and Instituto de Bioquímica y Biología Molecular, Facultad de Ciencias Exactas,² Universidad Nacional de La Plata, 1900 La Plata, Argentina

Received 13 December 2001/Accepted 17 April 2002

A rapid procedure for the identification of *Paenibacillus larvae* subsp. *larvae*, the causal agent of American foulbrood (AFB) disease of honeybees (*Apis mellifera* L.), based on PCR and restriction fragment analysis of the 16S rRNA genes (rDNA) is described. Eighty-six bacterial strains belonging to 39 species of the genera *Paenibacillus*, *Bacillus*, *Brevibacillus*, and *Virgibacillus* were characterized. Amplified rDNA was digested with seven restriction endonucleases. The combined data from restriction analysis enabled us to distinguish 35 profiles. Cluster analysis revealed that *P. larvae* subsp. *larvae* and *Paenibacillus larvae* subsp. *pulvificiens* formed a group with about 90% similarity; however, the *P. larvae* subsp. *larvae* restriction fragment length polymorphism pattern produced by endonuclease *HaeIII* was found to be unique and distinguishable among other closely related bacteria. This pattern was associated with DNA extracted directly from honeybee brood samples showing positive AFB clinical signs that yielded the restriction profile characteristic of *P. larvae* subsp. *larvae*, while no amplification product was obtained from healthy larvae. The method described here is particularly useful because of the short time required to carry it out and because it allows the differentiation of *P. larvae* subsp. *larvae*-infected larvae from all other species found in apiarian sources.

American foulbrood (AFB) disease caused by the spore-forming bacterium *Paenibacillus larvae* subsp. *larvae* (15) (formerly *Bacillus larvae*) is a highly contagious, cosmopolitan disease of bacterial origin affecting the larval and pupal stages of honeybees (*Apis mellifera* L.). Infected individuals turn brown and then black, and the resultant mass becomes a hard scale of material deposited on the side of the cell. AFB is one of the few bee diseases capable of killing a colony, and it presents unique problems for prevention and control because the spores can remain viable for long periods and survive under adverse environmental conditions (17, 17a, 17b). The disease spreads when spores are carried on drifting bees, hive parts, beekeepers' clothes, and contaminated pollen or honey.

Govan et al. (13) and Dobbelaere et al. (8) reported the use of PCR for rapid identification of *P. larvae* subsp. *larvae* by using primers derived from gene regions encoding 16S rRNA (rDNA). Specific primers designed by Govan et al. (13) produced a single amplicon, whereas those designed by Dobbelaere et al. (8) produced four amplicons. The results of their analysis of a limited number of species from apiarian sources did not allow them to differentiate *P. larvae* subsp. *larvae* from *Paenibacillus larvae* subsp. *pulvificiens*, the cause of powdery scale disease (15, 16), because both subspecies showed the same pattern. Dobbelaere et al. (8) concluded that the high degree of similarity between 16S rRNA genes of the two sub-

species, about 99.44%, does not permit the design of specific primers for either of the two subspecies.

In addition, several *Paenibacillus* species and species of the genera *Bacillus*, *Brevibacillus*, and *Virgibacillus* were consistently reported as being isolated from apiarian sources (2, 9, 10, 11, 12). The complex microbial community of sporeformers includes *Paenibacillus alvei*, *Brevibacillus laterosporus*, and *Paenibacillus apiarius*, which are considered secondary bacterial invaders of larvae infected with European foulbrood (EFB), and also *P. larvae* subsp. *pulvificiens* and *Bacillus coagulans*, which cause diseases of minor economical impact (1, 5, 16, 19, 20). Nevertheless, these bacteria can easily contaminate and overgrow plates of the slow-growing fastidious *P. larvae* subsp. *larvae*, making the correct diagnosis of AFB difficult unless selective media are used (1, 2).

The aim of this study was to assess the feasibility of using restriction fragment length polymorphism analysis (RFLP) of PCR-amplified 16S rDNAs for the differentiation of *P. larvae* subsp. *larvae* from other *Paenibacillus* organisms and from other spore-forming bacteria from apiarian sources and to assess its applicability to the direct and rapid diagnosis of AFB.

Strains and media. Eighty-six bacterial strains from diverse origins used in this study are listed in Table 1. For the isolation of *P. larvae* subsp. *larvae* strains from brood combs affected by AFB and from honey samples, previously described techniques were employed (2, 3). *Brevibacillus laterosporus* BLA 168 was isolated from honeybee larvae exhibiting symptoms of EFB, and Argentinian strains of *P. alvei*, *Bacillus cereus*, *Bacillus mycoides*, and *Bacillus megaterium* were recovered from honey as reported before (2).

* Corresponding author. Mailing address: Centro de Investigaciones de Fitopatología, Facultad de Ciencias Agrarias y Forestales, Universidad Nacional de La Plata, calles 60 y 118, c.c. 31, 1900 La Plata, Argentina. Phone: 54 221 4236758, ext. 423. Fax: 54 221 425-2346. E-mail: alippi@biol.unlp.edu.ar.

TABLE 1. Restriction patterns of PCR-amplified 16S rDNA genes among *Paenibacillus*, *Bacillus*, *Brevibacillus*, and *Virgibacillus* species and origins of the strains used in this study

| Species, strain designation, and geographical origin ^f | Pattern obtained with restriction enzyme: | | | | | | |
|--|---|--------------|----------------|------------------|--------------|--------------|--------------|
| | <i>Alu</i> I | <i>Msp</i> I | <i>Hae</i> III | <i>Hinf</i> I | <i>Cfo</i> I | <i>Rsa</i> I | <i>Taq</i> I |
| <i>Paenibacillus larvae</i> subsp. <i>larvae</i> | | | | | | | |
| ATCC 2574, United States | B | B | A | NRS ^m | A | A | A |
| PL113, PL225, Argentina ^a | B | B | A | NRS | A | A | A |
| PL295, PL296, United States ^a | B | B | A | NRS | A | A | A |
| PL201, PL203, Italy ^a | B | B | A | NRS | A | A | A |
| PL212, PL213, Canada ^a | B | B | A | NRS | A | A | A |
| PL228, PL230, France ^a | B | B | A | NRS | A | A | A |
| PL252, PL254, Spain ^b | B | B | A | NRS | A | A | A |
| PL284, PL286, Uruguay ^c | B | B | A | NRS | A | A | A |
| PL289, PL290, Japan ^d | B | B | A | NRS | A | A | A |
| PL56, PL57, Sweden ^e | B | B | A | NRS | A | A | A |
| PL29, PL31, New Zealand ^f | B | B | A | NRS | A | A | A |
| PL90, PL91, Germany ^g | B | B | A | NRS | A | A | A |
| PL68, PL70, Poland ^h | B | B | A | NRS | A | A | A |
| PL75 (CCM4483), PL76 (CCM4485), Czech Republic | B | B | A | NRS | A | A | A |
| PL301, PL302, UK ⁱ | B | B | A | NRS | A | A | A |
| PL100, Tunisia ^a | | | | | | | |
| PL304, PL305, Belgium ^j | | | | | | | |
| <i>Paenibacillus larvae</i> subsp. <i>pulvifaciens</i> | | | | | | | |
| CCM 38 (CCM), Czech Republic | B | B | B | B | A | A | A |
| NRRL B-3688, NRRL B-3670 | B | B | B | B | A | A | A |
| NRRL B-14154, NRRL B-14152 | B | B | B | B | A | A | A |
| ATCC 13537 | B | B | B | B | A | A | A |
| SAG 4689-3, SAG 4689-6, United States ^b | B | B | B | B | A | A | A |
| <i>Paenibacillus lentimorbus</i> NRRL B-2522 | F | D | NRS | F | A | A | A |
| <i>Paenibacillus macquariensis</i> NRRL B-14306 | B | D | D | D | A | C | A |
| <i>Paenibacillus glucanolyticus</i> NRRL B-14679 | B | E | G | E | A | E | D |
| <i>Paenibacillus peoriae</i> NRRL B-14750 | B | E | H | G | A | A | A |
| <i>Paenibacillus curdlanolyticus</i> NRRL B-23243 | J | E | NRS | K | E | D | B |
| <i>Paenibacillus kobensis</i> NRRL B-23299 | D | J | I | B | G | B | E |
| <i>Paenibacillus dendritiformis</i> NRRL B-666 | N | E | E | C | A | B | C |
| <i>Paenibacillus lautus</i> NRRL NRS-1000 | B | E | G | D | A | A | A |
| <i>Paenibacillus validus</i> NRRL NRS-1347 | N | E | E | C | A | B | C |
| <i>Paenibacillus alginolyticus</i> NRRL NRS-1351 | H | D | J | J | D | C | B |
| <i>Paenibacillus chondroitinus</i> NRRL NRS-1356 | B | D | D | H | A | B | A |
| <i>Paenibacillus illinoisensis</i> | B | D | H | D | A | A | A |
| <i>Paenibacillus alvei</i> | | | | | | | |
| NRRL B-383 | I | E | NRS | H | A | A | A |
| m437a, m361, Argentina ^a | I | E | NRS | H | A | A | A |
| <i>Paenibacillus amylolyticus</i> NRRL B142 | E | E | F | C | A | A | C |
| <i>Paenibacillus apiarius</i> ATCC 29575 | B | E | NRS | | A | C | A |
| <i>Paenibacillus macerans</i> NRRL NRS-924 | A | H | C | B | B | D | B |
| <i>Paenibacillus pabuli</i> NRRL B-510 | B | F | H | D | A | F | A |
| <i>Paenibacillus polymyxa</i> NRRL B 510 | A | F | D | B | A | D | B |
| <i>Paenibacillus azotofixans</i> NRRL B-142 | G | F | H | G | A | A | A |
| <i>Paenibacillus chibensis</i> ATCC 11377 | B | E | L | G | A | A | A |
| <i>Paenibacillus thiaminolyticus</i> | A | D | C | C | B | D | B |
| <i>Paenibacillus popilliae</i> ATCC 14706 | B | D | NRS | C | C | A | A |
| <i>Paenibacillus borealis</i> KK19, Finland ^k | B | D | D | D | A | C | A |
| <i>Bacillus azotoformans</i> NRRL B-14310 | A | I | O | L | B | D | B |
| <i>Bacillus circulans</i> ATCC 4515 | C | C | C | C | A | B | B |
| <i>Bacillus cereus</i> | | | | | | | |
| ATCC 11778 | D | E | E | C | A | B | C |
| m432, m436, Argentina ^a | D | E | E | C | A | B | C |
| <i>Bacillus coagulans</i> ATCC 35670 | A | D | C | B | B | D | B |
| <i>Bacillus licheniformis</i> NRRL B-1001 | D | D | C | B | A | D | B |
| <i>Bacillus megaterium</i> | | | | | | | |
| NRRL B-939 | A | E | C | C | A | B | C |
| m412, m440a, Argentina ^a | A | E | C | C | A | B | C |
| <i>Bacillus mycoides</i> | | | | | | | |
| ATCC 10206 | K | E | K | C | A | B | C |
| m425, m440b, Argentina ^a | K | E | K | C | A | B | C |
| <i>Bacillus thuringiensis</i> ATCC 10792 | D | E | E | C | A | B | C |
| <i>Bacillus pumilus</i> ATCC 7061 | A | D | C | B | B | D | B |

Continued on following page

TABLE 1—Continued

| Species, strain designation, and geographical origin ^a | Pattern obtained with restriction enzyme: | | | | | | |
|--|---|-------------|---------------|--------------|-------------|-------------|-------------|
| | <i>AluI</i> | <i>MspI</i> | <i>HaeIII</i> | <i>HinfI</i> | <i>CfoI</i> | <i>RsaI</i> | <i>TaqI</i> |
| <i>Bacillus subtilis</i> ATCC 10783 | A | D | C | B | B | D | B |
| <i>Bacillus sphaericus</i> ATCC 245 | A | E | C | C | A | D | C |
| <i>Bacillus firmus</i> ATCC 8247 | G | F | D | C | A | A | C |
| <i>Virgibacillus pantothenicus</i> ATCC 14567 | L | E | M | B | F | G | C |
| <i>Brevibacillus laterosporus</i> CCT 31 (CCT) | M | G | N | C | A | B | B |
| BLA 168, Argentina ^a | M | G | N | C | A | B | B |

^a A. M. Alippi, Centro de Investigaciones de Fitopatología, Facultad de Ciencias Agrarias y Forestales, UNLP, La Plata, Argentina.

^b P. Avalos, Laboratorio y Estación Cuarentenaria Lo Aguirre, SAG, Chile.

^c C. Piccini, Laboratorio de Microbiología, Instituto de Investigaciones Biológicas Clemente Estable, Montevideo, Uruguay.

^d A. Kataoka, Research Institute for Animal Science in Biochemistry and Toxicology, Kanawaga, Japan.

^e I. Fries, Bee Division, Swedish University of Agricultural Sciences, Uppsala, Sweden.

^f M. R. Goodwin, Apicultural Research Unit, Ruakura Research Center, Hamilton, New Zealand.

^g U. Rdest, Biozentrum der Universität Würzburg, Lehrstuhl Mikrobiologie, Würzburg, Germany.

^h M. Jelinski, Instytut Weterynarii, Swarzedz, Poland.

ⁱ B. Dancer, School of Pure and Applied Biology, University of Wales, Cardiff, United Kingdom.

^j W. Dobbelaere, Veterinary and Agrochemical Research Centre, Brussels, Belgium.

^k S. Elo, Department of Biosciences, Division of General Microbiology, University of Helsinki, Finland.

^l ATCC, American Type Culture Collection, Rockville, Md.; CCM, Czech Collection of Microorganisms, Brno, Czech Republic; CCT, Coleção de Culturas Tropical, Fundação André Tosello, Campinas, SP, Brazil; NRRL, Northern Regional Research Laboratory, Peoria, Ill.

^m NRS, no recognition site.

P. larvae subsp. *larvae*, *P. larvae* subsp. *pulvifaciens*, *P. amylolyticus*, *P. lautus*, *P. illinoisensis*, and *P. chibensis* strains were grown on MYPGP agar (6) at 37°C for 48 h; the other *Paenibacillus* species were grown on MYPGP agar at 30°C for 24 to 48 h, except *P. macquariensis*, which was grown at 22°C, and *P. dendritiformis*, which was grown on Luria-Bertani agar at 37°C. *Bacillus*, *Brevibacillus*, and *Virgibacillus* species were grown on tryptic soy agar at 30°C for 24 h, with the exception of *B. coagulans*, which was incubated at 37°C. Purity was confirmed by colony morphology and microscopic examination of bacterial smears.

DNA preparation. Bacterial cells for DNA extraction were grown at the appropriate temperature and medium under aerobic conditions for 24 to 48 h according to the species used. For bacterial DNA preparation, a rapid procedure using whole cells from plates was employed (3). After centrifugation to remove bacterial debris and resin, the supernatant was used as the DNA template.

PCR amplification and RFLP analysis of 16S rDNA. Primers U1 and U2 described by Ash et al. were used for PCR amplification of 16S rRNA genes from *Bacillus*, *Paenibacillus*, *Brevibacillus*, and *Virgibacillus* species (4). These primers were derived from conserved regions and capable of amplifying about 1.1 kb of 16S rDNA from *Bacillus* species and closely related genera. The PCR mixtures, which contained 1.5 µl of deoxynucleotide mixture (2 mM each), 1.25 µl of a mixture of both primers (10 mM each), 1.5 µl of Promega (Buenos Aires, Argentina) reaction buffer, 1.0 µl of MgCl₂ (25 µM), 5 µl of supernatant DNA, and sterile deionized water to bring the final volume to 25 µl, were pretreated at 94°C before 1 U of *Taq* polymerase (Promega Corp.) was added. PCR amplification was carried out according to the protocol of Ash et al. (4). PCR products were examined by using agarose (0.8%) gel electrophoresis and visualized by using ethidium bromide and UV light.

After amplification, subsamples of about 5 µl were incubated overnight with endonucleases *RsaI*, *HaeIII*, *MspI*, *AluI*, *HinfI*, *TaqI*, and *CfoI* (Promega) according to the manufacturer's specifications. RFLP analysis was performed by electrophoresis in a 2% agarose gel at 80 V for 2.30 h.

We found that species belonging to the genera *Paenibacillus*, *Bacillus*, *Brevibacillus*, and *Virgibacillus* consistently yielded a PCR amplification product of about 1,100 bp. In our analysis of the 16S rRNA gene, we assayed the variation at seven restriction sites that were thought to provide an RFLP pattern diagnostic of *P. larvae* subsp. *larvae*. It was found that analysis of 86 strains from different sources and classified as belonging to 39 species allowed us to place them in 35 composite profiles following digestion with *RsaI*, *HaeIII*, *MspI*, *AluI*, *HinfI*, *TaqI*, and *CfoI* by using the "combined gels" option of Gelcompare. The program FreeTree (14) was used for the construction of a phylogenetic tree (Fig. 1B) and for jackknife analysis by using a binary matrix based on RFLP characters (Nei-Li distances; neighbor-joining tree-construction method; 1,000 resampled data sets).

The result of the analysis shown in Fig. 1 revealed that *P. larvae* subsp. *larvae* and *P. larvae* subsp. *pulvifaciens* formed a group with about 90% similarity. However, the *HaeIII* restriction pattern of *P. larvae* subsp. *larvae* was found to be unique and allowed us to distinguish it from other closely related bacteria. Indeed, none of eight *P. larvae* subsp. *pulvifaciens* strains we examined showed the two *HaeIII* fragments of about 300 and 470 bp, respectively, which were characteristic of *P. larvae* subsp. *larvae* strains ($n = 32$) (Fig. 2A). Further evidence that supported subspecies differentiation was obtained, extending this analysis to 365 isolates of *P. larvae* subsp. *larvae* from different origins that showed the same *HaeIII* restriction pattern (data not shown); in addition, *HinfI* restriction analysis, unlike with *P. larvae* subsp. *pulvifaciens*, revealed no recognition site in *P. larvae* subsp. *larvae* strains (Table 1). On the




FIG. 1. (A) Combined restriction patterns of PCR-amplified 16S rDNA from representative species of the genera *Paenibacillus*, *Bacillus*, *Brevibacillus*, and *Virgibacillus* obtained by using the endonucleases *AluI*, *CfoI*, *HaeIII*, *HinfI*, *MspI*, *RsaI*, and *TaqI*. (B) Phylogenetic tree constructed on the basis of RFLP data by the neighbor-joining method using FreeTree software. Jackknife values are indicated at the branching points (1,000 replicates).

other hand, profiles obtained with *AluI*, *CfoI*, *RsaI*, and *TaqI* were found to be identical in both subspecies. In addition, the *MspI* restriction patterns shown by *P. larvae* subsp. *larvae* and subsp. *pulvificiens* were identical, whereas they differed from that of other species we examined (Fig. 2B). This relatedness between *P. larvae* subsp. *larvae* and *P. larvae* subsp. *pulvificiens* is in agreement with previous evidence obtained by using a polyphasic approach (15). Differences between pairs of restriction patterns, as found here, could be simply explained in terms of gain or loss of only one or two restriction sites, which indicates indeed that these two subspecies are genetically closely related.

In a few cases, pairs or groups (e.g., *B. cereus* and *B. thuringiensis*; *B. subtilis*, *B. coagulans*, and *B. pumilus*; and *P. borealis* and *P. macquariensis*) were not differentiated by the set of endonucleases that we used (Table 1; Fig. 1A). The use of other endonucleases or DNA sequencing may provide a basis for their differentiation.

We conclude that the 16S rRNA gene is polymorphic among

the aerobic spore-forming bacterial species predominant in apiarian sources. However, intraspecies polymorphism was not detected among the 32 *P. larvae* subsp. *larvae* strains obtained from diverse geographic regions. More interesting, we found that restriction pattern analysis revealed a distinct genotype for *P. larvae* subsp. *larvae* which may be useful for its identification, since the use of the endonucleases *MspI*, *HinfI*, and *HaeIII* would result in the recognition of *P. larvae* subsp. *larvae* among apiarian bacteria.

Direct detection in honeybee larva samples. Current procedures to detect AFB disease are based on direct field inspection of the hives and on the use of selective bacterial growth media combined with PCR methods (1, 2, 3, 13). Overall, they possess some limitations, since occasionally clinical symptoms are ambiguous and several days are required to reach a conclusive diagnosis. Therefore, in order to assess whether the 16S rDNA-RFLP analysis might be useful to reveal and confirm *P. larvae* subsp. *larvae* infection in hives, we carried out the following assays. Larvae exhibiting clinical symptoms of AFB




FIG. 2. (A) Gel electrophoresis of a PCR-amplified 16S rDNA fragment of 1,100 bp digested with *Hae*III. Similar restriction patterns were grouped and assigned the same letter (A to N). Lanes: W, molecular weight marker (100-bp ladder; Biodynamics); 1, *P. larvae* subsp. *larvae* ATCC 2574; 2, *P. larvae* subsp. *larvae* PI 113; 3, *P. larvae* subsp. *pulvifaciens* CCM 38; 4, *P. larvae* subsp. *pulvifaciens* SAG; 5, *Bacillus circulans* ATCC 4515; 6, *Bacillus firmus* ATCC 8247; 7, *Paenibacillus azotofixans* NRRL B-14372; 8, *Brevibacillus laterosporus* CCT 31; 9, *Bacillus subtilis* ATCC 10783; 10, *Paenibacillus macerans* ATCC 8244; 11, *P. alvei* NRRL B-383; and 12, *Bacillus thuringiensis* ATCC 10792. (C) Agarose gel showing PCR-RFLP results from healthy and diseased larvae. Lanes: M, molecular weight marker (100-bp ladder; Biodynamics); 1, PCR of AFB-diseased larvae using primers U1/U2; 2, PCR of healthy larvae; 3, PCR-amplified 16S rDNA fragment of 1,100 bp from *P. larvae* subsp. *larvae* ATCC 2574 digested with *Hae*III; 4, PCR-amplified 16S rDNA fragment from AFB-infected larvae digested with *Hae*III.

were removed from the cells by using a toothpick and thoroughly mixed with 1 ml of sterile distilled water (two larval remains or scales per tube). One hundred microliters of this mixture was diluted in 900 μ l of sterile distilled water, vortex mixed, and centrifuged at $3,200 \times g$ for 5 min. Fifty microliters of the supernatant was heated at 95°C for 15 min (8) and centrifuged at $3,200 \times g$ for 5 min. Subsamples of the supernatant were used as DNA templates in the PCR amplification as described above. Similar treatment was applied to healthy larvae 2, 5, and 19 days old and also to larval remains infected with chalkbrood caused by the fungus *Ascosphaera apis* (18) and EFB caused by the bacterium *Melissococcus plutonius* (formerly *Melissococcus pluton*) (5, 7), which were assessed as controls. AFB, EFB, and chalkbrood were confirmed by using standard microscopic and microbiological techniques (1, 5, 7, 18, 19). The results in Fig. 2C reveal that DNA extracted from larva samples associated with AFB symptoms consistently amplified the 1,100-bp fragment which, after incubation with endonuclease *Hae*III, gave an RFLP pattern identical to that found to be characteristic of *P. larvae* subsp. *larvae*. No amplification was detected with extracts from healthy larvae, EFB-diseased larvae, or chalkbrood mummies (dried dead larvae affected by chalkbrood disease). Furthermore, by using larval samples carrying mixed spore-forming bacterial populations that had been described by Alippi (1), a unique *Hae*III restriction pattern identical to that of *P. larvae* subsp. *larvae* was observed (data not shown). We assume that the high level of *P. larvae* subsp. *larvae* spores present in larva samples may indicate that DNA from *P. larvae* subsp. *larvae* outcompetes those from other bacteria as a template in the PCR.

Finally, DNA fingerprint analysis using the primers BOX, REP, and ERIC revealed four different genotypes within the *P. larvae* subsp. *larvae* collection we examined, which were demonstrated to be genetically diverse even though the 16S rDNA-RFLP pattern was identical (3; Alippi et al., unpublished data).

Our study provides a method that appears to be helpful in distinguishing *P. larvae* subsp. *larvae* from other *Paenibacillus* organisms, particularly those that are closely related, such as *P. larvae* subsp. *pulvifaciens*, and also from the spore-forming species which are commonly found in samples from apiarian environments. Since this procedure allows the identification of *P. larvae* subsp. *larvae* obtained either from culture or from

(B) Gel electrophoresis of a PCR-amplified 16S rDNA fragment of 1,100 bp digested with *Msp*I. Similar restriction patterns were grouped and assigned the same letter (A to G). Lanes: W, molecular weight marker (100-bp ladder; Biodynamics); 1, *P. larvae* subsp. *larvae* ATCC 2574; 2, *P. larvae* subsp. *larvae* PI 113; 3, *P. larvae* subsp. *pulvifaciens* CCM 38; 4, *P. larvae* subsp. *pulvifaciens* SAG; 5, *Bacillus circulans* ATCC 4515; 6, *Bacillus firmus* ATCC 8247; 7, *Paenibacillus azotofixans* NRRL B-14372; 8, *Brevibacillus laterosporus* CCT 31; 9, *Bacillus subtilis* ATCC 10783; 10, *Paenibacillus macerans* ATCC 8244; 11, *P. alvei* NRRL B-383; and 12, *Bacillus thuringiensis* ATCC 10792. (C) Agarose gel showing PCR-RFLP results from healthy and diseased larvae. Lanes: M, molecular weight marker (100-bp ladder; Biodynamics); 1, PCR of AFB-diseased larvae using primers U1/U2; 2, PCR of healthy larvae; 3, PCR-amplified 16S rDNA fragment of 1,100 bp from *P. larvae* subsp. *larvae* ATCC 2574 digested with *Hae*III; 4, PCR-amplified 16S rDNA fragment from AFB-infected larvae digested with *Hae*III.

larvae, we believe it can be applied to the reliable and rapid diagnosis of AFB (in about 4 h), in contrast to classical microbiological methods, which require at least 2 days.

This research was supported by a grant from ANPCyT, Argentina (BID 1201/OC-AR PICT 08-03857), and from the IFS, Sweden (A.M.A.). A.M.A. and O.M.A. are Career Investigators of CIC and CONICET, respectively, and A.C.L. is a recipient of a scholarship from ANPCyT, Argentina.

We thank the curator of the NRRL Collection, N. Nakamura, for supplying bacterial cultures, those (listed in Table 1) who provided AFB samples or bacterial isolates, and F. J. Reynaldi for collecting larval samples.

REFERENCES

- Alippi, A. M. 1991. A comparison of laboratory techniques for the detection of significant bacteria of the honey bee, *Apis mellifera*, in Argentina. *J. Apic. Res.* **30**:75–80.
- Alippi, A. M. 1995. Detection of *Bacillus larvae* spores in Argentinian honeys by using a semi-selective medium. *Microbiologia* **11**:343–350.
- Alippi, A. M., and M. Aguilar. 1998. Characterization of isolates of *Paenibacillus larvae* subsp. *larvae* from diverse geographical origin by the polymerase chain reaction and BOX primers. *J. Invertebr. Pathol.* **72**:21–27.
- Ash, C., F. G. Priest, and M. D. Collins. 1993. Molecular identification of rRNA group 3 bacilli (Ash, Farrow, Wallbanks and Collins) using a PCR probe test. Proposal for the creation of a new genus *Paenibacillus*. *Antonie Leeuwenhoek J. Microbiol.* **64**:253–260.
- Bailey, L. 1983. *Melissococcus pluton*, the cause of European foulbrood of honeybees (*Apis* spp.). *J. Appl. Bacteriol.* **55**:65–69.
- Dingman, D. W., and D. P. Stahly. 1983. Medium promoting sporulation of *Bacillus larvae* and metabolism of medium components. *Appl. Environ. Microbiol.* **46**:860–869.
- Djordjevic, S. P., K. Noone, L. Smith, and M. Hornitzky. 1998. Development of a hemi-nested PCR assay for the specific detection of *Melissococcus pluton*. *J. Apic. Res.* **37**:165–174.
- Dobbelaere, W., D. C. De Graaf, J. E. Peeters, and F. J. Jacobs. 2001. Development of a fast and reliable method for American foulbrood disease (*Paenibacillus larvae* subsp. *larvae*) using a 16S rRNA gene based PCR. *Apidologie* **32**:363–370.
- Gilliam, M. 1978. Bacteria belonging to the genus *Bacillus* isolated from selected organs of queen honey bees, *A. mellifera*. *J. Invertebr. Pathol.* **31**:389–391.
- Gilliam, M. 1979. Microbiology of pollen and bee bread: the genus *Bacillus*. *Apidologie* **10**:269–274.
- Gilliam, M., and D. B. Prest. 1987. Microbiology of feces of the larval honey bee, *Apis mellifera*. *J. Invertebr. Pathol.* **49**:70–75.
- Gilliam, M., and D. K. Valentine. 1976. Bacteria isolated from the intestinal contents of foraging worker honey bees, *Apis mellifera*: the genus *Bacillus*. *J. Invertebr. Pathol.* **28**:275–276.
- Govan, V. A., M. H. Allsopp, and S. Davidson. 1999. A PCR detection method for rapid identification of *Paenibacillus larvae*. *Appl. Environ. Microbiol.* **65**:2243–2245.
- Hampel, V., A. Pavlicek, and J. Flegr. 2001. Construction and bootstrap analysis of DNA fingerprinting-based phylogenetic trees with the freeware program FreeTree; application to trichomonad parasites. *Int. J. Syst. Evol. Microbiol.* **51**:731–735.
- Heyndrickx, M., K. Vandemeulebroecke, B. Hoste, P. Janssen, K. Kersters, P. De Vos, N. A. Logan, N. Ali, and R. C. W. Kerkeley. 1996. Reclassification of *Paenibacillus* (formerly *Bacillus*) *pulvifaciens* (Nakamura 1984) Ash et al., 1994, a later subjective synonym of *Paenibacillus* (formerly *Bacillus*) *larvae* (White 1906) Ash et al. 1994, as a subspecies of *P. larvae*, with emended descriptions of *P. larvae* as *P. larvae* subsp. *larvae* and *P. larvae* subsp. *pulvifaciens*. *Int. J. Syst. Bacteriol.* **46**:270–279.
- Katznelson, H. 1950. *Bacillus pulvifaciens* (n. sp.), an organism associated with powdery scale of honey-bee larvae. *J. Bacteriol.* **59**:153–155.
- Matheson, A., and M. Reid. 1992. Strategies for the prevention and control of American foulbrood. Part I. *Am. Bee J.* **132**:399–402.
- Matheson, A., and M. Reid. 1992. Strategies for the prevention and control of American foulbrood. Part II. *Am. Bee J.* **133**:471–475.
- Matheson, A., and M. Reid. 1992. Strategies for the prevention and control of American foulbrood. Part III. *Am. Bee J.* **143**:534–547.
- Rose, J. B., M. Christensen, and W. T. Wilson. 1984. *Ascospaera* species inciting chalkbrood in North America and a taxonomic key. *Mycotaxon* **19**:41–55.
- Shimanuki, H., and D. A. Knox. 1991. Diagnosis of honey bee diseases. Agricultural handbook no. AH-690. U.S. Department of Agriculture, Washington, D.C.
- Vandenberg, J. D., and H. Shimanuki. 1990. Isolation and characterization of *Bacillus coagulans* associated with half-moon disorder of honey bees. *Apidologie* **21**:233–241.