

# Subpicosecond resonance Raman spectra of the early intermediates in the photocycle of bacteriorhodopsin


R. van den Berg, Du-Jeon-Jang,\* Herbert C. Bitting, and M. A. El-Sayed

Department of Chemistry and Biochemistry, University of California, Los Angeles, Los Angeles, California 90024-1569 USA

**ABSTRACT** The resonance Raman spectra are presented for the species formed during the photocycle of bacteriorhodopsin (bR) on a timescale of 800–900 fs. In the ethylenic stretch region two intermediates were found with frequencies of 1,510 and 1,518  $\text{cm}^{-1}$ , corresponding to species with optical absorption maxima at 660 and 625 nm, respectively. This leads to the assignment of the 1,518  $\text{cm}^{-1}$  band to the  $J_{625}$  intermediate. In the fingerprint region, the appearance of a vibration at 1,195  $\text{cm}^{-1}$  strongly suggests that the isomerization indeed has taken place in a time less than the pulsewidth of our laser. This supports the previous proposals made on the basis of the optical spectra. The spectra are compared with those observed in tens of picoseconds up to nanoseconds.

## INTRODUCTION

Upon absorption of light, the membrane protein bacteriorhodopsin (bR) pumps protons across the cell membrane of *Halobacterium halobium*, thus producing an electrochemical gradient that drives the bacteria's metabolic processes (Stoeckenius and Bogomolni, 1982). Excitation of the bR chromophore, retinal, initiates a photocycle which is completed within 10 ms at room temperature. The kinetics of this cycle, which consists of intermediates with lifetimes varying from femtoseconds to milliseconds, have been extensively studied with optical transient absorption techniques (Lozier et al., 1975; Kaufmann et al., 1976). The different intermediates are labeled by a letter and a number, which is their approximate absorption maximum:



The retinal is connected to a lysine residue of the protein backbone through a protonated Schiff base (PSB) linkage. This PSB deprotonates during the  $L_{550} \rightarrow M_{412}$  step, leading to the proton pumping process, and reprotonates in the following step.

The primary photochemical event in the photocycle has been a subject of considerable interest. Early picosecond optical studies suggested the formation of a red-shifted primary photoproduct  $K_{610}$  in 11 ps (Applebury et al., 1978). In later studies a number of experiments showed a redshifted precursor of  $K_{610}$  which forms in ~500 fs and relaxes to  $K_{610}$  on a 3 ps timescale (Sharkov et al., 1985; Nuss et al., 1985; Polland et al., 1986; Mathies et al.,

1988; Dobler et al., 1988). This precursor was labeled  $J_{625}$ , although there was some uncertainty about its absorption maximum.

Although transient absorption spectra are obtainable in the short-time domain, they are generally very broad and structureless, and give little or no information on configurational or conformational changes. Time-resolved resonance Raman spectroscopy of picosecond transients was demonstrated in 1980 by Turner et al. for the carboxyhemoglobin photointermediate and in 1981 by Hsieh et al. for bR. The vibrational spectra are greatly simplified when the laser is tuned to the electronic absorption wavelength of the retinal, because only retinal vibrations which mimic a molecular distortion in the excited state are enhanced. Using microbeam and flow techniques with picosecond excitation and optical multichannel analyzer (OMA) detection, it has proven possible to detect the resonance Raman spectra of picosecond transients (for a review see Turner and El-Sayed, 1985).

Bacteriorhodopsin has been the subject of a great number of resonance Raman studies. It has been shown that upon light absorption an *all-trans* to *13-cis* isomerization occurs in the retinal. This isomerization has already taken place in  $K_{610}$ , i.e., before 10–40 ps (Hsieh et al., 1981; Stern and Mathies, 1985; Atkinson et al., 1989). The resonance Raman spectrum of  $K_{610}$  was further thoroughly studied up to 25 ns (Smith et al., 1983; Hsieh et al., 1983; Stern and Mathies, 1985), but essentially similar results were obtained. Whether isomerization has already taken place in the  $J_{625}$  intermediate has been an open question. However, from a time-resolved pump/probe study of bR reconstituted with a sterically hindered retinal, 9,12-phenyl-retinal, using 3-ps pulses Polland et

Dr. Jang's present address is Korea Standards Research Institute, Daedeok Science Town, Taejeon, Choongnam 301-340, Republic of Korea.

al. (1984) concluded that formation of the  $K_{610}$  intermediate in bR is accompanied by an *all-trans* to *13-cis* isomerization. The pulsewidth used might allow for a similar conclusion on  $J_{625}$ .

In this paper we present the first resonance Raman spectrum which can exclusively be attributed to  $J_{625}$ . Spectra of the different vibrational regions were recorded. It is shown by a study of the "fingerprint region", which contains the vibrations sensitive to the isomeric configuration of the retinal, that isomerization occurs in a time interval equal to or shorter than the photolyzing laser pulse of 800–900 fs. In the ethylenic stretch region, apart from a peak at  $1,518\text{ cm}^{-1}$  corresponding to the  $J_{625}$  intermediate, another species is found at this timescale showing a shoulder at  $\sim 1,510\text{ cm}^{-1}$  corresponding to a  $\lambda_{\text{max}}$  of 660 nm. The results are discussed and compared with spectra reported in the literature on a timescale varying from femto- to nanoseconds.

## MATERIALS AND METHODS

### Sample preparation

The purple membrane was purified from the ET1-001 strain of *Halobacterium halobium* according to the method of Oesterhelt and Stoekenius (1974) and Becher and Cassim (1975). The purified purple membrane was suspended in doubly deionized water at a concentration of  $\sim 60\ \mu\text{M}$  as measured by the retinal absorption at 570 nm.

### Experimental set-up

The experimental arrangement for obtaining the subpicosecond resonance Raman spectra is similar to the one first reported by Terner et al. (1979a and b) and Hsieh et al. (1983), the only difference being that another laser system was used with a shorter pulsewidth.

In the present work a cw Nd:YAG laser (model 3800, Spectra-Physics Inc., Mountain View, CA), mode-locked at 82 MHz, produces a train of  $\sim 80$ -ps pulses at an average power of 14 W. The beam was directed into a Spectra Physics fiber-grating pulse compressor (model SP 3695), which shortens the 1,064-nm pulses to  $< 6$  ps. An SP 3225 frequency doubler is directly mounted in the compressor, producing  $\sim 1.2$  W at 532 nm. The green output, which is further stabilized to  $\sim 800$  mW by an acousto-optic stabilizer (model SP 3275), is used to synchronously pump a Spectra Physics 375B dye-laser, with Rhodamine 6G in ethylene glycol, fitted with an SP 344S cavity dumper. A two-plate birefringent is used as a wavelength tuning device. The bandwidth of the dye-laser is  $\sim 16$ – $18\text{ cm}^{-1}$ . At a dumping rate of 800 kHz, the average output power of the dye-laser system at the excitation wavelength of 587 nm was 25 mW (31 nJ/pulse). This wavelength was selected to simultaneously increase the photolysis quantum yield and to make efficient use of the resonance Raman enhancement. Furthermore it should be noted that the red fluorescence of bR obscures the resonance Raman signals when much longer wavelengths are used. The power used is sufficiently small to minimize stimulated processes. The temporal width of the excitation pulse was  $\sim 800$ – $900$  fs, as measured by autocorrelation (model 5-14A, Inrad Inc., Northvale, NJ) assuming a  $\text{sech}^2$  pulse shape.

The beam was directed via a microscope objective (Zeiss Neofluar 40 $\times$ ) onto a vertically flowing free jet of bR suspension, and was thus

tightly focused to a  $\sim 10$ - $\mu\text{m}$  spot. The sample was kept light-adapted by a lamp and was circulated from the sample vial through a syringe needle with a 110- $\mu\text{m}$  diameter by a peristaltic pump (Masterflex, Cole-Parmer Instrument Co., Chicago, IL). The measured flow rate was 25 m/s which, combined with the 10- $\mu\text{m}$  spot size, gave a sample residence time of 0.4  $\mu\text{s}$ . Because the time between the laser pulses is 1.25  $\mu\text{s}$ , each pulse should have encountered a fresh volume of unphotolyzed light-adapted bR.

The scattered light was collected at  $90^\circ$  and focused onto the slit of a model 1870 0.5-m spectrograph (Spex Industries Inc., Edison, NJ), and the Rayleigh scattering was blocked by appropriate Corning cut-off filters (2-61 and 2-62 for the ethylenic stretching region; 2-62 and 2-63 for the fingerprint region). The resolution of the monochromator was determined to be  $3\text{ cm}^{-1}$ . The detection system consisted of a diode array detector, thermoelectrically cooled to  $-20^\circ\text{C}$  (model 1455, EG&G Princeton Applied Research Corp., Princeton, NJ) and an OMA (model 1460) with a 1462 detector controller and 1462/99 14-bit A/D controller (EG&G Princeton Applied Research Corp.).

Wavelengths were calibrated with the lines of a neon discharge lamp. Vibrational frequencies are estimated to be accurate to  $\pm 3\text{ cm}^{-1}$  in the original spectra and to  $\pm 5\text{ cm}^{-1}$  in the computer-generated "difference" spectra (see below).

### Measurement procedure

In a single laser resonance Raman experiment on bR, each laser pulse serves both as the photolysis and Raman excitation light source. A spectrum of the unphotolyzed bR $_{570}$  sample is obtained at a low average laser power. The excitation conditions can be described in terms of a photolability parameter  $F = I_0 \cdot t$  (Mathies et al., 1976; Atkinson et al., 1985), in which  $t$  is the residence time of the sample in the laser beam and

$$I_0 = \phi \cdot \sigma_{\text{bR}}(\lambda) \cdot I(\lambda, r), \quad (1)$$

in which  $\sigma_{\text{bR}}(\lambda)$  is the absorption cross-section of bR $_{570}$  at the excitation wavelength,  $\phi$  is the photochemical quantum yield for reaction, and  $I(\lambda, r)$  is the laser intensity. The latter can be calculated from the peak power of the laser and the diameter of the focused laser spot in the jet.  $F$  was varied between 0.15 and 1.9 by attenuating the average laser power with neutral density filters. At low  $F$  values ( $F < 0.2$ ) a spectrum similar to that of bR $_{570}$  is obtained, whereas higher  $F$  values ( $F > 1$ ) are needed to produce significant amounts of photoproducts.

The "difference" spectra were obtained by subtracting an arbitrarily scaled low-power spectrum from a high-power spectrum. As was noted by Hsieh et al. (1983) caution should be exercised when information is extracted from these "difference" spectra. They are to be used strictly as a guide to interpreting the actual spectral changes.

## RESULTS AND DISCUSSION

### Resonance Raman spectrum of bR $_{570}$

To identify the resonance Raman spectra of the photocycle intermediates of bR, an unambiguous, well-characterized spectrum of bR $_{570}$  should be obtained first. Very low laser powers (corresponding to  $F$  values of  $\sim 0.1$ – $0.2$ ) were used to record such a spectrum, which is shown in Figs. 1 and 2 (upper spectra) for two different spectral regions: 1,450–1,600  $\text{cm}^{-1}$  and 900–1,400  $\text{cm}^{-1}$ , respectively. The









FIGURE 1 Resonance Raman spectra of bR in the ethylenic stretch region around  $1530\text{ cm}^{-1}$ . The upper spectrum was obtained at low laser power ( $F = 0.15$ ) and is the result of unphotolyzed, ground-state bR<sub>570</sub> only. The middle spectrum was recorded at high laser power ( $F = 1.9$ ) and represents the contributions of both bR<sub>570</sub> and J<sub>625</sub>. A clear broadening and shift are observed, showing J<sub>625</sub> has a lower ethylenic stretching frequency than bR<sub>570</sub>. The lower part shows the difference spectrum of the high and low laser power spectra. The  $\nu_{C-C}$  of the J<sub>625</sub> intermediate is observed around  $1518\text{ cm}^{-1}$ , in accordance with a linear relation between the absorption maximum of the intermediate and this ethylenic stretching frequency. The occurrence of a shoulder  $\sim 1510\text{ cm}^{-1}$  may indicate the formation of a more red-shifted intermediate with an absorption maximum at  $660\text{ nm}$ .

FIGURE 2 Resonance Raman spectra of the fingerprint region of bR. The upper spectrum represents the contribution of unphotolyzed, ground-state bR<sub>570</sub> only, because it was obtained at very low laser powers. The middle spectrum clearly shows the effects of the initiation of the photocycle at high laser powers and represents a spectrum of bR<sub>570</sub> and J<sub>625</sub>. The retinal *all-trans* to *13-cis* isomerization has taken place as can be seen from the filling in of the valley between the  $1,174$  and  $1,203\text{ cm}^{-1}$  peaks. Furthermore, an increase in the region between  $960$  and  $1,000\text{ cm}^{-1}$  can be distinguished. The lower part shows the difference spectrum of the high and low laser power spectra of Fig. 2. The isomerization of the retinal is evident from the peak at  $1,195\text{ cm}^{-1}$ . The  $1,164\text{ cm}^{-1}$  vibration, which is unique to the J<sub>625</sub> spectrum, shows that the isomeric configuration at these early times ( $800\text{--}900\text{ fs}$ ) is clearly different from that of the K<sub>610</sub> intermediate.

latter contains the C—C stretches and C—C—H in-plane bends between 1,100 and 1,300  $\text{cm}^{-1}$ , which characterize the isomeric configuration of the retinal (Smith et al., 1987a), and the hydrogen-out-of-plane (HOOP) and C—CH<sub>3</sub> rocking vibrations, between 900 and 1,100  $\text{cm}^{-1}$ , which are sensitive to conformational distortions along the C<sub>7</sub>—C<sub>15</sub> hydrocarbon chain (Smith et al., 1987a). The region around 1,500  $\text{cm}^{-1}$  has been assigned to a C=C stretching mode. Its frequency has been shown to correlate linearly with the electronic absorption maximum of the chromophore (Rimai et al., 1973; Aton et al., 1977).

The bR<sub>570</sub> spectrum, shown in Figs. 1 and 2 (upper panels) is in agreement with spectra reported previously (Terner et al., 1979a and b; Stern and Mathies, 1985; Atkinson et al., 1985, 1989). Moreover, a complete vibrational assignment has been given by Smith et al. (1987a), based on an extensive set of <sup>13</sup>C- and <sup>2</sup>H-labeled derivatives. Many of the vibrational features in the bR<sub>570</sub> spectrum are characteristic of an *all-trans* structure, although each frequency is found to be  $\sim 10 \text{ cm}^{-1}$  higher than the corresponding frequency in the *all-trans* retinal protonated Schiff base. This was explained as a result of increased  $\pi$ -electron delocalization of the retinal in the protein.

### Resonance Raman spectrum in the ethylenic stretch region

The spectrum obtained in the region around 1,500  $\text{cm}^{-1}$  at a high value of  $F$  is shown in Fig. 1 (*middle*). Photolysis is evident from the clear broadening of the 1,526  $\text{cm}^{-1}$  band compared to the 1,530  $\text{cm}^{-1}$  band in the spectrum of unphotolyzed bR<sub>570</sub>. Fig. 1 (*bottom*) shows the intense residual band around 1,518  $\text{cm}^{-1}$  obtained by subtracting the upper spectrum of Fig. 1 from the middle.

It is known that a linear relation exists between the absorption maximum of a bR photocycle intermediate and the frequency of its ethylenic stretch mode (Rimai et al., 1973; Aton et al., 1977). In Fig. 3 this relation is shown for the different bR intermediates. It should be noted, that some uncertainty exists regarding the  $\nu_{\text{C}=\text{C}}$  of both O<sub>640</sub> (Terner et al., 1979b; Smith et al., 1983) and K<sub>610</sub> (Terner et al., 1979a; Smith et al., 1983; Stern and Mathies, 1985). We have decided to take average values, and plot error bars according to the minimum and maximum values. From this figure the C=C stretch frequency of 1,518  $\text{cm}^{-1}$  as found from the difference spectrum of Fig. 1 (*bottom*) would correspond to an intermediate with an absorption maximum of 625 nm and might therefore be attributed to J<sub>625</sub>. Furthermore, it should be noted that the 1,518  $\text{cm}^{-1}$  band is asymmetric on the low-frequency side with a shoulder  $\sim 1,510 \text{ cm}^{-1}$ . This frequency corresponds to an intermediate with an absorption maximum at 660 nm and might indicate that




FIGURE 3 Plot of the dependence between the frequency of the ethylenic stretch mode ( $\nu_{\text{C}=\text{C}}$ ) and the absorption maximum ( $\lambda_{\text{max}}$ ) of the different bR photocycle intermediates. The intermediates that have been plotted are: M<sub>412</sub> (1,570  $\text{cm}^{-1}$ ), L<sub>550</sub> (1,539  $\text{cm}^{-1}$ ), bR<sub>570</sub> (1,531  $\text{cm}^{-1}$ ), K<sub>610</sub> (1,523  $\text{cm}^{-1}$ ) and O<sub>640</sub> (1,514  $\text{cm}^{-1}$ ). The C=C stretch frequencies of 1,510 and 1,518  $\text{cm}^{-1}$  would correspond to absorption maxima of 660 and 625 nm, respectively. They have been put in the figure with different symbols. A linear relation is obtained.

during the 800–900-fs photolysis pulse either another intermediate is formed, which has an even more red-shifted absorption spectrum than J<sub>625</sub>, or that an intermediate is formed corresponding to J<sub>625</sub> from another type of bR.

A large number of studies have been devoted to the earliest intermediates in the bR photocycle. Applebury et al. (1978) observed a red-shifted (625 nm) precursor of K<sub>610</sub> within the 6-ps time width of the excitation pulse, and Polland et al. (1986) observed an absorption maximum of J<sub>625</sub> around 615 nm. Further investigations clearly established the formation ( $\sim 0.5$  ps) and decay times ( $\sim 3$  ps) of J<sub>625</sub> and agreed that its absorption is red-shifted compared to that of K<sub>610</sub>. However, no spectra were shown (Nuss et al., 1985; Sharkov et al., 1985; Dobler et al., 1988). Very recently, Mathies et al. (1988) carried out a careful time-resolved transient absorption/emission study using 60-fs pump and 6-fs probe pulses, which shows an increased absorption around 640–660 nm on a timescale of 200–900 fs.

The absorption maximum which has been attributed to J<sub>625</sub> seems to be shifted to the red when it is probed at shorter timescales. At times varying between 1 and 3 ps the absorption maximum seems to be around 625 nm, the value commonly assumed for the J intermediate. The C=C stretch frequency of 1,518  $\text{cm}^{-1}$  would then correspond to J<sub>625</sub>, which would be in complete accord with the linear relation between  $\lambda_{\text{max}}$  and  $\nu_{\text{C}=\text{C}}$  (Fig. 3) (Rimai et al., 1973; Aton et al., 1977). At earlier times, however, as shown by the differential transmittance spectra of Mathies et al. (1988) there seems to be a redshift: at short times

(~220 fs) there is an absorption around ~660 nm, whereas at longer times (~990 fs), the transient absorption has shifted to ~630–640 nm. It is therefore possible that the 1,510  $\text{cm}^{-1}$  band is due to an absorption of an intermediate before  $J_{625}$ , which absorbs around 660 nm and is present on the 200–800 fs timescale. It could also be due to the corresponding  $J_{625}$  intermediate in a parallel cycle, that has an absorption in  $K_{610}$  not very different from that of the first cycle. The Raman spectrum of  $L_{550}$  indeed shows two types of L which are now believed to result from parallel cycles (Diller and Stockburger, 1988). The presence of two types of bR was first suggested from the observation of two risetimes of the  $M_{412}$  intermediate (Hanamoto et al., 1984).

### Resonance Raman spectrum in the fingerprint region

A “high-power” spectrum in the fingerprint region is shown in the middle part of Fig. 2. A substantial amount of photochemistry has been induced at this higher power, producing a significant concentration of  $J_{625}$ . This can be seen from the increased scattering in the region between 960 and 1,000  $\text{cm}^{-1}$ . These frequencies are in the HOOP region and are particularly sensitive to conformational distortions along the polyene chain (Smith et al., 1987a). A similar observation was made by Atkinson (Atkinson et al., 1985; Brack and Atkinson, 1989) on a timescale of ~10 ps and was interpreted as arising from a twisting motion of the conjugated polyene chain. The observed scattering enhancement of these C—H vibrations would occur when the  $\sigma$  and  $\pi^*$  electron systems become intermixed upon twisting of the polyene chain (Hsieh et al., 1981). The broadening to the low-frequency side might suggest that scattering occurs from hot unrelaxed vibrational levels, as was observed in the C=C stretching region (Brack and Atkinson, 1990). Furthermore, as was observed before (Hsieh et al., 1981), slight differences in the unrelaxed protein environment around the twisted retinal might also induce further broadening in this region.

A variety of experiments have argued that the initial photochemistry in bR involves an *all-trans* to *13-cis* isomerization of the retinal. Such an isomerization has already taken place in the  $J_{625}$  intermediate as shown by the filling in of the valley between the 1,174 and the 1,203  $\text{cm}^{-1}$  peaks, i.e., the appearance of a new band around 1,195  $\text{cm}^{-1}$ . It is this vibrational region which is sensitive to the isomeric state of the retinal chromophore. In the difference spectrum of Fig. 2 (*bottom*) this new band is clearly resolved and so is another band at a lower frequency of 1,164  $\text{cm}^{-1}$ . The latter confirms the observation appearing in their 8-ps resonance Raman spectrum. Stern and Mathies (1985) observed this same band on a

7-ps timescale at 1,163  $\text{cm}^{-1}$ , but attributed it to differences in the excitation wavelength. This seems to be unlikely, because it has already been shown that the  $bR_{570}$  spectrum undergoes only minor changes in relative intensities, when excited at four different wavelengths: 676 nm (Braiman and Mathies, 1982), 568 nm (Terner et al., 1979a), 514 nm (Stockburger et al., 1979) and 457 nm (Marcus and Lewis, 1978). A similar observation was made in the resonance Raman spectra of the retinal protonated Schiff base when excited at 413 nm (Braiman and Mathies, 1980) and at 647 nm (Aton et al., 1977). Furthermore, Atkinson et al. (1989) clearly showed that when exciting around 590 nm, the 1,163–1,164- $\text{cm}^{-1}$  band is not part of the  $K_{610}$  spectrum. However, both the 8-ps spectrum of Atkinson (Atkinson et al., 1989) and the 7-ps spectrum of Stern and Mathies (1985) contain a considerable contribution from  $K_{610}$ , because the decay time of  $J_{625}$  is 3 ps. This contribution could not be resolved. The spectrum presented here is the first that can be exclusively attributed to  $J_{625}$ , because the 800–900-fs pulsewidth does not allow for appreciable  $K_{610}$  formation. Furthermore it has been shown that resonance Raman scattering from the electronic excited state of  $bR_{570}$  is unlikely, because of its low concentration (due to the extremely short lifetime of ~200 fs) and its small absorption coefficient at the laser wavelength (Polland et al., 1986).

The occurrence of the 1,195  $\text{cm}^{-1}$  band is indicative of isomerization at the  $C_{13}=C_{14}$  position, i.e., from *all-trans* to *13-cis* retinal. This band was also observed in spectra obtained on a 40-ps (Hsieh et al., 1981, 1983), a 200-ps (Stern and Mathies, 1985), and a 20–30-ns timescale (Hsieh et al., 1983; Smith et al., 1983; Stern and Mathies, 1985). It is also observed for the  $K_{610}$  intermediate which Braiman and Mathies (1982) have shown to have a *13-cis* configuration. The 1,164- $\text{cm}^{-1}$  vibration on the other hand was not observed in any of the spectra mentioned above, and therefore does show that the retinal isomerization at these very early times (<1 ps) is of a different nature than at much longer times. It is, however, very difficult to determine the exact configuration at  $C_{13}=C_{14}$ , because there are many factors which influence the features appearing in the fingerprint region.  $bR_{570}$ , which contains an *all-trans* retinal, has a band at ~1,171  $\text{cm}^{-1}$ . If one assumes that upon isomerization its frequency changes with time, one observes the following. In our spectrum we observe a transient at 1,164  $\pm$  3  $\text{cm}^{-1}$  ( $\lambda_{\text{exc}} = 587$  nm). This same band might be found at 1,163  $\text{cm}^{-1}$  in 7 ps ( $\lambda_{\text{exc}} = 576$  nm) (Stern and Mathies, 1985), at 1,170  $\text{cm}^{-1}$  at times >10 ps ( $\lambda_{\text{exc}} = 590$  nm) (Brack and Atkinson, 1989), at 1,178  $\text{cm}^{-1}$  in 40 ps ( $\lambda_{\text{exc}} = 587$  nm) (Hsieh et al., 1981, 1983), and at 1,185  $\text{cm}^{-1}$  in 0.2 ns ( $\lambda_{\text{exc}} = 514$  nm) (Stern and Mathies, 1985) or longer: 20–30 ns ( $\lambda_{\text{exc}} = 587$  nm) (Hsieh et al., 1983), ( $\lambda_{\text{exc}} =$

514 nm) (Stern and Mathies, 1985) and 100 ns ( $\lambda_{\text{exc}} = 552.3$  nm) (Hsieh et al., 1981). Realizing that *13-cis* retinal in dark-adapted bR has a strong band at  $1,183\text{ cm}^{-1}$  (Terner et al., 1979c; Smith et al., 1987b) one is tempted to suggest that the  $1,164\text{-cm}^{-1}$  vibration in *all-trans* retinal in bR changes into the  $1,185\text{ cm}^{-1}$  vibration during the isomerization process. If so, then the isomerization starts in  $<500$  fs (Dobler et al., 1988; Mathies et al., 1988) and is completed in  $<200$  ps. In that case, the fact that no corresponding changes in the optical absorption of the retinal have been reported on this timescale might suggest that this bond is very sensitive to the protein environment, and that the time dependence of its frequency simply reflects the fast relaxation rates of the amino acid residues within the active site. Its frequency in these early intermediates should therefore be sensitive to genetic substitution of the residues known to be present within the active site (Mogi et al., 1989).

This work was supported by the Department of Energy (Office of Basic Energy Sciences) under grant DE-FG03-88ER13828. R. van den Berg gratefully acknowledges Richard F. Savage for the loan of many optical components, Steve F. Vitorino for a thorough training on the femtosecond laser systems, the SHELL Research Laboratory, Amsterdam (Koninklijke/Shell-Laboratorium, Amsterdam) for financial support and the Netherlands Organization for Scientific Research (Nederlandse organisatie voor Wetenschappelijk Onderzoek) for a research fellowship.

## REFERENCES

- Applebury, M. L., K. S. Peters, and P. M. Rentzepis. 1978. Primary intermediates in the photochemical cycle of bacteriorhodopsin. *Biophys. J.* 23:375-382.
- Atkinson, G. H., T. L. Brack, I. Grieger, G. Rumbles, D. Blanchard, and L. M. Siemanowski. 1985. Picosecond time-resolved resonance Raman spectroscopy of bacteriorhodopsin intermediates. In *Time-resolved Vibrational Spectroscopy*. G. H. Atkinson, editor. Gordon and Breach, New York. 55-82.
- Atkinson, G. H., T. L. Brack, D. Blanchard, and G. Rumbles. 1989. Picosecond time-resolved resonance Raman spectroscopy of the initial *trans* to *cis* isomerization in the bacteriorhodopsin photocycle. *Chem. Phys.* 131:1-15.
- Aton, B., A. G. Doukas, R. H. Callender, B. Becher, and T. G. Ebrey. 1977. Resonance Raman studies of the purple membrane. *Biochemistry*. 16:2995-2999.
- Becher, B. M., and J. Y. Cassim. 1975. Improved isolation procedures for the purple membrane of *Halobacterium halobium*. *Prep. Biochem.* 5:161-178.
- Brack, T. L., and G. H. Atkinson. 1989. Picosecond time-resolved resonance Raman spectrum of the  $K_{590}$  intermediate in the room temperature bacteriorhodopsin photocycle. *J. Mol. Spectr.* 214:289-303.
- Brack, T. L. and G. H. Atkinson. 1990. Vibrationally excited retinal in the bacteriorhodopsin photocycle: picosecond time-resolved anti-Stokes resonance Raman scattering. *J. Phys. Chem.* In press.
- Braiman, M., and R. A. Mathies. 1980. Resonance Raman evidence for the *all-trans* to *13-cis* isomerization in the proton pumping cycle of bacteriorhodopsin. *Biochemistry*. 19:5421-5428.
- Braiman, M., and R. A. Mathies. 1982. Resonance Raman spectra of bacteriorhodopsin's primary photoproduct: evidence for a distorted *13-cis* retinal chromophore. *Proc. Natl. Acad. Sci. USA.* 79:403-407.
- Diller, R., and M. Stockburger. 1988. Kinetic resonance Raman studies reveal different conformational states of bacteriorhodopsin. *Biochemistry*. 27:7641-7651.
- Dobler, J., W. Zinth, W. Kaiser, and D. Oesterhelt. 1988. Excited-state reaction dynamics of bacteriorhodopsin studied by femtosecond spectroscopy. *Chem. Phys. Lett.* 144:215-220.
- Hanamoto, J. H., P. Dupuis, and M. A. El-Sayed. 1984. On the protein (tyrosine)-chromophore (protonated Schiff base) coupling in bacteriorhodopsin. *Proc. Natl. Acad. Sci. USA.* 81:7083-7087.
- Hsieh, C.-L., M. Nagumo, M. Nicol, and M. A. El-Sayed. 1981. Picosecond and nanosecond resonance Raman studies of bacteriorhodopsin. Do configurational changes of retinal occur in picoseconds? *J. Phys. Chem.* 85:2714-2717.
- Hsieh, C.-L., M. A. El-Sayed, M. Nicol, M. Nagumo, and J. H. Lee. 1983. Time-resolved resonance Raman spectroscopy of the bacteriorhodopsin photocycle on the picosecond and nanosecond time-scales. *Photochem. Photobiol.* 38:83-94.
- Kaufmann, K. J., P. M. Rentzepis, W. Stoeckenius, and A. Lewis. 1976. Primary photochemical processes in bacteriorhodopsin. *Biochem. Biophys. Res. Commun.* 68:1109-1115.
- Lozier, R., R. A. Bogomolni, and W. Stoeckenius. 1975. Bacteriorhodopsin: a light-driven proton pump in *Halobacterium halobium*. *Biophys. J.* 15:955-962.
- Marcus, M. A., and A. Lewis. 1978. Resonance Raman spectroscopy of the retinylidene chromophore in bacteriorhodopsin ( $bR_{570}$ ),  $bR_{560}$ ,  $M_{412}$ , and other intermediates: structural conclusions based on kinetics, analogues, models and isotopically labeled membranes. *Biochemistry*. 17:4722-4735.
- Mathies, R., A. R. Oseroff, and L. Stryer. 1976. Rapid-flow resonance Raman spectroscopy of photolabile molecules: rhodopsin and isorhodopsin. *Proc. Natl. Acad. Sci. USA.* 73:1-5.
- Mathies, R. A., C. H. Brito Cruz, W. T. Pollard, and C. V. Shank. 1988. Direct observation of the femtosecond excited-state *cis-trans* isomerization in bacteriorhodopsin. *Science (Wash. DC)*. 240:777-779.
- Mogi, T., T. Marti, and H. G. Khorana. 1989. Structure function studies on bacteriorhodopsin. IX. Substitutions of tryptophan residues affect protein retinal interactions in bacteriorhodopsin. *J. Biol. Chem.* 264:14197-14201.
- Nuss, M. C., W. Zinth, W. Kaiser, E. Kölling, and D. Oesterhelt. 1985. Femtosecond spectroscopy of the first events of the photochemical cycle in bacteriorhodopsin. *Chem. Phys. Lett.* 117:1-6.
- Oesterhelt, D., and W. Stoeckenius. 1974. Isolation of the cell membrane of *Halobacterium halobium* and its fractionation into red and purple membrane. *Methods Enzymol.* 31:667-678.
- Pollard, H. J., M. A. Franz, W. Zinth, W. Kaiser, E. Kölling, and D. Oesterhelt. 1984. Optical picosecond studies of bacteriorhodopsin containing a sterically fixed retinal. *Biochim. Biophys. Acta.* 767:635-639.
- Pollard, H. J., M. A. Franz, W. Zinth, W. Kaiser, E. Kölling, and D. Oesterhelt. 1986. Early picosecond events in the photocycle of bacteriorhodopsin. *Biophys. J.* 49:651-662.
- Rimai, L., M. E. Heyde, and D. Gill. 1973. Vibrational spectra of some carotenoids and related linear polyenes. A Raman spectroscopic study. *J. Am. Chem. Soc.* 95:4493-4501.

- Sharkov, A. V., A. V. Pakulev, S. V. Chekalin, and Y. A. Matveetz. 1985. Primary events in bacteriorhodopsin probed by subpicosecond spectroscopy. *Biochim. Biophys. Acta.* 808:94–102.
- Smith, S. O., M. Braiman, and R. A. Mathies. 1983. Time-resolved resonance Raman spectroscopy of the K<sub>610</sub> and O<sub>640</sub> photointermediates of bacteriorhodopsin. In *Time-resolved vibrational spectroscopy*. G. H. Atkinson, editor. Academic Press, Inc., New York. 219–230.
- Smith, S. O., M. S. Braiman, A. B. Meyers, J. A. Pardoen, J. M. L. Courtin, C. Winkel, J. Lugtenburg, and R. A. Mathies. 1987a. Vibrational analysis of the *all-trans* retinal chromophore in light-adapted bacteriorhodopsin. *J. Am. Chem. Soc.* 109:3108–3125.
- Smith, S. O., J. Pardoen, J. Lugtenburg, and R. A. Mathies. 1987b. Vibrational analysis of the *13-cis* retinal chromophore in dark-adapted bacteriorhodopsin. *J. Phys. Chem.* 91:804–819.
- Stern, D., and R. A. Mathies. 1985. Picosecond and nanosecond resonance Raman evidence for structural relaxation in bacteriorhodopsin's primary photoproduct. In *Time-resolved Vibrational Spectroscopy*. A. Laubereau and M. Stockburger, editors. Springer-Verlag, Berlin. 250–254.
- Stockburger, M., W. Klusmann, H. Gattermann, G. Massig, and R. Peters. 1979. Photochemical cycle of bacteriorhodopsin studied by resonance Raman spectroscopy. *Biochemistry.* 18:4886–4900.
- Stoeckenius, W., and R. A. Bogomolni. 1982. Bacteriorhodopsin and related pigments of *Halobacteria*. *Annu. Rev. Biochem.* 52:587–619.
- Terner, J., and M. A. El-Sayed. 1985. Time-resolved resonance Raman spectroscopy of photobiological and photochemical transients. *Acc. Chem. Res.* 18:331–338.
- Terner, J., C.-L. Hsieh, A. R. Burns, and M. A. El-Sayed. 1979a. Time-resolved resonance Raman spectroscopy of intermediates of bR: the bK<sub>590</sub> intermediate. *Proc. Natl. Acad. Sci. USA.* 76:3046–3050.
- Terner, J., C.-L. Hsieh, A. R. Burns, and M. A. El-Sayed. 1979b. Time-resolved resonance Raman characterization of the bO<sub>640</sub> intermediate of bacteriorhodopsin. Reprotonation of the Schiff base. *Biochemistry.* 18:3629–3634.
- Terner, J., C.-L. Hsieh, and M. A. El-Sayed. 1979c. Time-resolved resonance Raman characterization of the L<sub>550</sub> intermediate and the two dark-adapted bR<sub>560</sub> forms of bacteriorhodopsin. *Biophys. J.* 26:527–541.
- Terner, J., T. G. Spiro, M. Nagumo, M. F. Nicol, and M. A. El-Sayed. 1980. Resonance Raman spectroscopy in the picosecond timescale: the carboxyhemoglobin photointermediate. *J. Am. Chem. Soc.* 102:3238–3239.