- 12 Starfield B. Primary care: balancing health needs, services and technology. Oxford: Oxford University Press, 1998.
- 13 Haslam D. Schools and hospitals for education and health. BMJ 2003; 326:234-5.
- 14 Irvine D. The quiet revolution. William Pickles lecture 1975. J R Coll Gen Pract 1975:25:399-407.
- 15 Seddon MF, Marshall MN, Campbell SM, Roland M. Systematic review of studies of quality of clinical care in general practice in the United Kingdom, Australia and New Zealand. Qual Health Care 2001;10:152-8.
- 16 Baker R, Roland M. General practice: continuous quality improvement since 1948. Br J Gen Pract 2002;52(suppl):S2-3.
- 17 Campbell SM, Roland M, Middleton E, Reeves D. Improvements in the quality of clinical care in English general practice 1998-2003. BMJ (in press).
- 18 Sheaff R, Sibbald B, Campbell S, Roland M, Marshall MN, Pickard S, et al. Soft governance and attitudes to clinical quality in English general practice. I Health Serv Res Polico 2004;9:132-8.
- practice. J Health Serv Res Policy 2004;9:132-8.

 19 Health and Social Care Information Centre. Quality and outcomes framework information you by the Services (not faccessed 21 Sep. 2005).
- framework information. www.ic.nhs.uk/services/qof (accessed 21 Sep 2005).

 20 Commission for Healthcare Improvement. Local health services patient survey 2003. http://www.healthcarecommission.org.uk/assetRoot/04/00/46/21/04004621.pdf (accessed 2 Nov 2005).

 21 Bower P, Sheaff RS, Sibbald B, Campbell S, Roland M, Marshall MN, et al.
- 21 Bower P, Sheaff RS, Sibbald B, Campbell S, Roland M, Marshall MN, et al. Setting standards based on patients' views on access and continuity: secondary analysis of data from the general practice assessment survey. BMJ 2003:236:258-60.

- 22 Grol R, Wensing M, Mainz J, Jung H, Ferreira P, Hearnshaw H, et al. Patients in Europe evaluate general practice care: an international composition. B. L. Com. Parts 2000;58:89-7
- parison. Br J Gen Pract 2000;50:882-7. 23 Watt G. The inverse care law today. Lancet 2002;360:252-4.
- 24 Freeman G, Hjortdahl P. What future for continuity of care in general practice? BMJ 1997;314:1870-3.
- 25 Manious A, Baker R, Love M, Pereira Gray DJ, Gill JM. Continuity of care and trust in one's physician: evidence from primary care in the US and UK. Fam Med 2001;33:22-7.
- 26 Dovey SM, Meyers DS, Phillips RL Jr, Green LA, Fryer GE, Galliher JM, et al. A preliminary taxonomy of medical errors in family practice. *Qual Saf Health Care* 2002:11:233–8.
- 27 Sheaff R, Schofield J, Mannion R, Dowling B, Marshall M, McNally R. Organisational factors and performance: a review of the literature. London: National Coordinating Centre for Service Delivery and Organisation, 2003. www.sdol.shtm.ac.uk/pdf/studyinghealthcare_sheaff_report.pdf (accessed 2 Nov 2005).
- 28 Marshall M, Noble J, Davies H, Walshe K, Waterman H, Sheaff R, et al. Producing information about general practice services that makes sense to patients and the public; final project report. Manchester: National Primary Care Research and Development Centre. 2005.
- Care Research and Development Centre, 2005.
 29 Department of Health. Building on the best; choice, responsiveness and equity in the NHS. London: DoH, 2003.
- Robinson J, Luft H. Competition, regulation and hospital costs, 1982 to 1986. JAMA 1988;260:2676-81.

The NHS revolution: health care in the market place

What do patients and the public want from primary care?

Angela Coulter

The government hopes that getting patients' views on their priorities for primary care will ensure support for its plans. It is likely to find patients care more about quality of care than structural or financial reform

The UK government has stated it wants the public to help shape the future of the health service. In the run-up to the planned publication of a white paper on care outside hospitals, Patricia Hewitt, secretary of state for health in England, is leading a big public engagement exercise to "genuinely involve patients, public and staff in designing family health and social care to meet the challenges of the 21st century." The secretary of state's commitment to engaging directly with the public is commendable if it is a genuine attempt to listen and learn, but she should also take account of the extensive body of research evidence on what patients and the public want. Patients have diverse needs and expectations leading to different, and sometimes conflicting, views on priorities,2 but it is possible to discern themes. What does the evidence show?

Structure of primary care

A distinction can be made between what patients want as individual healthcare users and what they hope for as citizens or taxpayers (box 1). In general, patients care more about the quality of their everyday interactions with health professionals than about how the service is organised. Furthermore, although there is scope for improvement in primary care, changes that seem to undermine the founding principles of the NHS are likely to be strongly resisted.

Interpersonal care

Patients want primary care professionals who are good communicators and have sound, up to date clinical knowledge and skills. They also want professionals who



National Citizens summit organised to get patients' views on health care

are interested and sympathetic, involve them in decisions, give them sufficient time and attention, and provide advice on health promotion and self care. A systematic review of the literature on patients' priorities for general practice care, which examined 19 studies published between 1966 and 1995, found that the most important factor was "humaneness," which ranked highest in 86% of studies that included this aspect. This was followed by "competence/accuracy" (64%), "patients' involvement in decisions" (63%), and "time for care" (60%).

Most patients who consult their general practitioner have specific expectations—for example, they want an explanation of their symptoms, treatment, or investigation. Many have their own ideas about what is wrong and what may have caused it, but they do not This article is part of a series examining the government's planned market reforms to healthcare provision

Picker Institute Europe, Oxford OX1 1RX Angela Coulter chief executive

angela.coulter@ pickereurope.ac.uk

BMJ 2005;331:1199-200

Box 1: Healthcare aspirations of patients and citizens

Patients

Fast access to reliable health advice
Effective treatment delivered by trusted professionals
Participation in decisions and respect for preferences
Clear, comprehensible information and support for self care
Attention to physical and environmental needs
Emotional support, empathy, and respect
Involvement of, and support for, family and carers

Citizens

Affordable treatment and care, free at the point of use Safety and quality

Health protection and disease prevention

Continuity of care and smooth transitions

Accessible local services and national centres of excellence

Universal coverage; geographical and social equity

Responsiveness, flexibility, and choice

Participation in service developments

Transparency, accountability, and opportunity to influence policy decisions

always articulate these.⁵ Failure to engage with the patient's agenda can lead to misunderstandings, dissatisfaction, and poor outcomes.⁶ ⁷ The good news is that most British patients report positive experiences in primary care. Box 2 gives highlights from the latest national patient survey carried out in all primary care trusts in England, which obtained responses from 116 939 patients (a response rate of 47%).⁸

The results of this survey show the strengths of our primary care system—good clinician-patient relationships, free access, and continuity of care—but they also suggest it is failing to meet patients' expectations in relation to providing information, involvement in treatment decisions, access to records, preventive advice, and support for self care. Clinical targets for care of chronic diseases are mostly met, according to the quality and outcomes framework review,⁹ but we urgently need to improve support for self care and self management, for which the current general medical services contract does not provide incentives.⁹

The relatively paternalistic nature of the UK system is also reflected in studies comparing patients' experiences in different countries. The UK scores

Box 2: Key findings from the 2005 primary care patient survey⁸

Positive experiences

92% said they were treated with dignity and respect by the doctor 82% said the doctor listened carefully to them

74% said they definitely had enough time with the doctor to discuss their problem

76% said they had complete confidence and trust in their doctor 85% had complete confidence and trust in other primary care staff

Negative experiences

41% would have liked more say in decisions about medicines 39% of those prescribed new drugs wanted more information about side effects

70% of patients referred to a specialist were not given copies of referral letters 57% of patients who had phoned the practice had had difficulty contacting the practice

19% of smokers who wanted help to quit smoking had not been offered it

high on doctor-patient communication, continuity, and affordability but low on information and choice.¹¹ ¹²

Access

Patients want easier and more flexible access to certain primary care services, including free choice of practitioners, more physiotherapy and complementary therapies, and longer clinic opening hours.¹³ ¹⁴ Most patients are willing to be seen and treated by nurses and other primary care practitioners as well as doctors, suggesting scope to vary skill mix and encourage team working as a means of improving access.¹⁵

Alternative ways of accessing health advice are becoming more popular. Internet use is growing; a recent UK survey of people aged 45 and over found that 30% of respondents had searched for health information on the web. Not surprisingly, use of the internet was more prevalent among middle aged people and those in higher socioeconomic groups, but use of telephone helplines, such as NHS Direct, was popular among people from lower socioeconomic groups. Email access to general practitioners and other primary care staff is still relatively rare, however. In a recent international survey, only 13% of those in the UK said they could communicate with their doctor by email, compared with 22% in New Zealand, and 20% in the United States. 11

Choice and continuity

The government wants to promote greater choice in the primary care system. Choice of provider is popular among those waiting for elective treatment, ¹⁶ and most patients value the opportunity to choose their general practitioner. ¹² There is almost certainly scope to extend the choices available, but the heaviest users of primary care (older people and those with chronic conditions) place particular value on continuity of care from professionals they know. ¹⁷ Better community support for carers and for people with long term mental health problems are also perennial themes in patient surveys. ¹⁸ These groups require well coordinated local services rather than a diversity of competing providers.

Younger patients, commuters, and those with urgent needs are more willing to trade continuity for faster access to alternative primary care services. ¹⁹ Nevertheless, there is little evidence of a desire to dismantle the registered list system, despite the apparent popularity of walk-in centres. Some patients may want freer access to hospital based specialists, but tampering with the referral system risks undermining the important coordination role provided by general practitioners and other primary care staff. Greater provider choice will not be worth having if it undermines the foundations of a system that works reasonably well at present.

Shared decision making

Patients want the opportunity to make choices in relation to their health care, but choosing a provider is only one of the choices they can make. Knowing about the various treatment options available and having a say in these is more important to most patients than having a

choice of where to be treated. Failures in communication of information about illness and treatment are the most frequent source of patient dissatisfaction.²⁰

The traditional model of decision making assumed that doctors and patients shared the same goals, that only the doctor was sufficiently informed and experienced to decide what should be done, and that patient involvement should be confined to giving or withholding consent to treatment. However, this paternalistic approach now seems seriously outdated. Many, if not most, patients now expect to be given information about their condition and the treatment options, and they want clinicians to take account of their preferences.¹⁹ Some expect to go further: to be actively engaged in the decision making process or even to take the decision themselves. Evidence is growing that engaging patients in treatment decisions and in managing their healthcare can lead to more appropriate and cost effective use of health services and better health outcomes.21

Equity and participation

Patients and citizens do seem to want more personalised and flexible services, but this does not mean they want to have to shop around for all their primary care needs. Geographical equity and needs based allocation are fundamental NHS values which politicians ignore at their peril. The British public remains strongly wedded to the idea that services everywhere should be high quality and equally available to all.22 Nevertheless, the notion that healthcare providers should be required to account for their performance also attracts strong support, as does the need for independent regulation at arm's length from government, coupled with open reporting on performance.22

Including patients in planning developments to the service has had some positive results, but direct involvement is likely to remain a minority activity.²³ Most people are not members of organised patients' groups, and only a minority want to sit on policy committees. Many of the current NHS initiatives that go under the banner of patient and public involvement are little more than window dressing. Including a few token patients on committees is relatively easy, but it does little to tackle the heart of the problem. The really important changes need to occur at the level of individual interactions between patients and health professionals. Improving the quality of these and achieving a shift from paternalism to a partnership approach has the potential to increase the effectiveness of health care.

Conclusion

What most patients and citizens want is the security of knowing that health services will be there when they need them, that their views and preferences will be taken account of by health professionals, that they will be given the help they need to help themselves, that they can access reliable information about their condition and the treatment options, and that they won't have to worry about the financial consequences of being ill. They also want to be sure that these benefits are equitably distributed and that public resources are

Summary points

The UK government wants its reforms of primary care to reflect public priority

Research shows the patients want fast access to good care

They want to have a say in their care and be helped to help

Free and equitable care is also important

being used efficiently for the good of all. Social solidarity and trust will continue to be the essential underpinnings of a sustainable health system.

Contributors and sources: AC has spent the past 20 years researching patients' experience of health care. She directs Picker Institute Europe, a charity which works with patients, professionals, and policy makers to promote understanding of the patient's perspective at all levels of healthcare policy and practice. This article draws on studies and reviews carried out by the Picker Institute

Competing interests: None declared.

- Department of Health. Hewitt asks the public to help shape care outside hospitals. Press release: 23 Jun 2005. www.dh.gov.uk (search for 2005/0217). Jung HP, Baerveldt C, Olesen F, Grol R, Wensing M. Patient characteris-
- tics as predictors of primary health care preferences: a systematic literature analysis. *Health Expect* 2003;6:160-81.
- Little P. Everitt H. Williamson I. Warner G. Moore M. Gould C. et al. Preferences of patients for patient centred approach to consultation in primary care: observational study. *BMJ* 2001;322:468-72.

 Wensing M, Jung HP, Mainz J, Olesen F, Grol R. A systematic review of the
- literature on patient priorities for general practice care. 1. Description of the research domain. *Soc Sci Med* 1998;47:1573-88.
- Cockburn J, Pit S. Prescribing behaviour in clinical practice: patients' expectations and doctors' perceptions of patients' expectations—a questionnaire study. *BMJ* 1997;315:520-3.

 McKinley RK, Middleton JF. What do patients want from doctors?
- Content analysis of written patient agendas for the consultation. Br J Gen Pract 1999;49:796-800.
- Stevenson FA, Cox K, Britten N, Dundar Y. A systematic review of the research on communication between patients and health care professionals about medicines: the consequences for concordance. Health Expect 2004:7:235-45.
- Healthcare Commission. *Primary care trust survey of patients* 2005. London: Healthcare Commission, 2005.
- Health and Social Care Information Centre. National quality and outcomes framework statistics for England 2004/05. London: Health and Social Care Information Centre, Government Statistical Service, 2005.
- $10\,$ Ellins J, Coulter A. How engaged are people in their health care? London: Health Foundation, 2005.
- 11 Schoen C, Osborn R, Huynh PT, Doty M, Davis K, Zapert K, et al. Primary care and health system performance: adults' experiences in five countries. Health Affairs 2004;W4:487.
- 12 Coulter A, Jenkinson C. European patients' views on the responsiveness of health systems and healthcare providers. Eur J Public Health 2005;15:355-60.
- 13 Cabinet Office. Open all hours? Results from the people's panel. London: Cabinet Office, 2000.

 14 YouGov. Daily Telegraph survey on GPs. www.yougov.com/archives/pdf/
- OMI050101083_Lpdf (accessed 4 Nov 2005).

 15 Kinnersley P, Anderson E, Parry K, Clement J, Archard L, Turton P, et al.
- Randomised controlled trial of nurse practitioner versus general practitioner care for patients requesting "same day" consultations in primary care. BMJ 2000;320:1043-8.

 16 Coulter A, Henderson L, Le Maistre N. Hospital choices: patients' experience
- of the London patient choice project. Oxford: Picker Institute Europe, 2004. Mainous AG, Baker R, Love MM, Pereira Gray D, Gill JM. Continuity of care and trust in one's physician: evidence from primary care in the United States and the United Kingdom. Fam Med 2001;33:22-7.
- 18 Picker Institute. Is the NHS getting better or worse? Oxford: Picker Institute Europe, 2005.
- 19 Coulter A, Magee H. The European patient of the future. Maidenhead: Open University Press, 2003
- 20 Coulter A, Cleary PD. Patients' experiences with hospital care in five
- 20 Coulter A, Cleary Fr. Fatterins experiences with hospital care in live countries. Health Affairs 2001;20:244-52.
 21 Coulter A. The automomous patient. London: Nuffield Trust, 2002.
 22 Magee H, Davis L-J, Coulter A. Public views on healthcare performance indicators and patient choice. JR Soc Med 2003;96:338-42.
- 23 Crawford MJ, Rutter D, Manley C, Weaver T, Bhui K, Fulop N, et al. Systematic review of involving patients in the planning and development of health care. *BMJ* 2002;325:1263-5.