

Novel 4-Chlorophenol Degradation Gene Cluster and Degradation Route via Hydroxyquinol in *Arthrobacter chlorophenolicus* A6

Karolina Nordin,^{1,2} Maria Unell,³ and Janet K. Jansson^{2,3*}

Department of Biochemistry and Biophysics, Stockholm University, 106 91 Stockholm, Sweden¹; Section for Natural Sciences, Södertörn University College, 141 89 Huddinge, Sweden²; and Department of Microbiology, Swedish University of Agricultural Sciences, Box 7025, 750 07 Uppsala, Sweden³

Received 8 December 2004/Accepted 13 June 2005

Arthrobacter chlorophenolicus A6, a previously described 4-chlorophenol-degrading strain, was found to degrade 4-chlorophenol via hydroxyquinol, which is a novel route for aerobic microbial degradation of this compound. In addition, 10 open reading frames exhibiting sequence similarity to genes encoding enzymes involved in chlorophenol degradation were cloned and designated part of a chlorophenol degradation gene cluster (*cph* genes). Several of the open reading frames appeared to encode enzymes with similar functions; these open reading frames included two genes, *cphA-I* and *cphA-II*, which were shown to encode functional hydroxyquinol 1,2-dioxygenases. Disruption of the *cphA-I* gene yielded a mutant that exhibited negligible growth on 4-chlorophenol, thereby linking the *cph* gene cluster to functional catabolism of 4-chlorophenol in *A. chlorophenolicus* A6. The presence of a resolvase pseudogene in the *cph* gene cluster together with analyses of the G+C content and codon bias of flanking genes suggested that horizontal gene transfer was involved in assembly of the gene cluster during evolution of the ability of the strain to grow on 4-chlorophenol.

Arthrobacter chlorophenolicus A6 is a gram-positive actinobacterium that was previously isolated from a soil slurry enriched with increasing concentrations of 4-chlorophenol (4-CP) (28). This bacterium can degrade unusually high concentrations of 4-CP (up to 2.7 mM) and other *p*-substituted phenols, such as 4-nitrophenol and 4-bromophenol (28). In addition, *A. chlorophenolicus* A6 can degrade 4-CP at low temperatures (5°C) and during temperature fluctuations between 28°C and 5°C (3). Strain A6 was previously tagged with marker genes encoding the green fluorescent protein (*gfp*) or firefly luciferase (*luc*) in order to specifically monitor its ability to survive in nonsterile soil as it degraded 4-CP (9). The tagged cells survived well and were metabolically active in soil contaminated with 4-CP (9, 12). A larger fraction of the cell population remained viable after incubation in soil at 5°C than after incubation in soil at 28°C (4).

There have been several reports of 4-CP degradation in other bacteria, but none of these bacteria have been shown to degrade concentrations of 4-CP as high as those degraded by *A. chlorophenolicus* A6. Successful mineralization of 4-CP by bacteria is usually accomplished by the oxidation of 4-CP to 4-chlorocatechol, followed by *ortho* cleavage of the aromatic ring. After ring cleavage, the chlorine is removed and the carbon skeleton is transformed into products that are assimilated into the central metabolism of the cell. By contrast, two strains belonging to the actinobacterium group, *Arthrobacter ureafaciens* CPR706 (5) and *Nocardioideis* sp. strain NSP41 (8), were reported to transform 4-CP into hydroquinone instead of 4-chlorocatechol. However, the degradation pathways in these strains have not been described further.

The aim of the present study was to elucidate the biochemistry and genetics of 4-chlorophenol degradation in *A. chlorophenolicus* A6. Our results suggest that hydroxyquinol is an intermediate in the 4-CP degradation route used by *A. chlorophenolicus* A6, a possibility which has not been reported previously for aerobic degradation of monosubstituted chlorophenols by bacteria. In addition, we identified a 4-CP catabolic gene cluster in this strain and constructed a mutant in which one of the key catabolic genes was interrupted in order to address its functional significance. The results of this study increase our understanding of the diversity and evolution of catabolic pathways.

MATERIALS AND METHODS

Culture conditions. A minimal medium (GM medium) was used for growth of *A. chlorophenolicus* A6 on 4-CP as described previously (28). 4-CP was added to a concentration of 1.2 or 1.9 mM, the cultures were incubated at 28°C, and the cells were normally harvested at the mid-log phase, at which time the cultures had an optical density at 600 nm (OD₆₀₀) of 0.1. When higher cell densities were desired, yeast extract was added to GM medium to a final concentration of 0.1% (wt/vol), or 10% Luria-Bertani broth (LB) was used. *Escherichia coli* was grown in LB or on LB agar plates at 37°C. When appropriate, ampicillin was added to a concentration of 50 µg/ml, chloramphenicol was added to a concentration of 34 µg/ml, or 2,2'-dipyridyl was added to a concentration of 1 mM.

Chloride ion measurements. *A. chlorophenolicus* A6 cells were grown in GM medium with 1.9 mM 4-CP to the stationary phase and inoculated into fresh medium to an OD₆₀₀ of approximately 0.04. Then 4-CP was added at different concentrations up to an inhibitory level of 3.5 mM to three 100-ml replicate cultures for each concentration, and the cultures were incubated at 28°C with shaking. The 4-CP concentration was measured in cell-free supernatants at 279 nm with a spectrophotometer (Beckman Coulter, Fullerton, CA). The chloride ion concentration was measured with an ion-selective electrode (Orion Research Inc., Beverly, MA), and prior to every measurement the accuracy of the electrode was confirmed by comparison to a standard curve.

Gas chromatography. Samples (5 ml) were reduced with 2 mg dithionite and extracted with 1 volume of ethyl acetate, and the organic phase was collected and dried over disodium sulfate. Derivatization by silylation was performed as described by Apajalahti and Salkinoja-Salonen (1). The samples were dissolved in ethyl acetate and analyzed using a Varian (Walnut Creek, CA) model 3800 gas chromatograph with a model 8200 autosampler equipped with a flame ionization

* Corresponding author. Mailing address: Department of Microbiology, Swedish University of Agricultural Sciences, Box 7025, 750 07 Uppsala, Sweden. Phone: 46-18-673201. Fax: 46-18-673393. E-mail: Janet.Jansson@mikrob.slu.se.

TABLE 1. Oligonucleotide primers used in this study

Primer	Sequence (5' to 3')	Purpose	Position in accession no. AY131335 sequence
CphAIf	ATGACGACCCGTCAGTAGC	Expression of <i>cphA-I</i>	8415–8434
CphAIr	CTTCAGATCAGGATTAGGAGCAA	Expression of <i>cphA-I</i>	9326–9304
CphAIIIf	ATGTTCGGGAATCATCCATGCC	Expression of <i>cphA-II</i>	12394–12374
CphAIIr	TGCGAGGCGGGGGCGAGGA	Expression of <i>cphA-II</i>	11486–11505
P1	CTGAAACAACCTCATGCAAGCCCT	Anneal to conserved region of hydroxyquinol dioxygenase gene	8523–8545
P2	GGAGCGAGAACGATATCGAA	Anneal to conserved region of hydroxyquinol dioxygenase gene	9310–9291
P3	TGGAACGGAC	Random primer	5652–5661
P4	GCGATGGTCT	Anneal to clone c1	8707–8698
P5	AGTGCCTGCTGC	Random primer	2181–2190
P6	CCGAAGTGGT	Anneal to clone c2	5981–5972
P7	GATCGTCAGC	Random primer	–9–0
P8	GGCATCGTTA	Anneal to clone c3	2405–2396
P9	GTCCAGGGTGACCCTTATAT	Anneal to clone c1	9147–9166
P10 ^a	TGCACCACAAGATNTGYCA	Anneal to conserved region of maleylacetate reductase	12832–12814
P11	CGGTTTCAGGA	Anneal to clone c6	13555–13564
P12	ATCACGAGGG	Random primer	15017–15008
P13	TGCCCTACGA	Anneal to clone c7	14653–14662
P14	CAGCCGACGT	Random primer	15476–15485
CmxF	AGCATGTAGAGGGCAAAGG	Anneal to internal fragment of Tn 1409Cβ	NA ^b
Tnp2	AGAAGACGTGCTGGCGTACTTCGA	Anneal to internal fragment of Tn1409Cβ	NA
Tra1	GCTGATAGGCGAACGCAACTGGTC	Screen <i>cph</i> cluster for transposon insertion	315–338
Tra2	GTTCGGAACCTGAACCGCTAACTCG	Screen <i>cph</i> cluster for transposon insertion	7500–7477
Tra3	CACCGCGAGTTTGTGCGAATTAT	Screen <i>cph</i> cluster for transposon insertion	7042–7065
Tra4	TCTGGCACCTTACGGCCAATTAC	Screen <i>cph</i> cluster for transposon insertion	13842–13819

^a Degenerate primer.

^b NA, not applicable.

detector. A 30-m DB1-ht column (100% methyl; inside diameter, 0.25 mm; film thickness, 0.1 μm; J&W Scientific, Folsom, CA) was used. The temperature program was as follows: 1 min at 50°C, followed by an increase in the temperature at a rate of 20°C/min to 150°C, a slower increase at a rate of 5°C/min to 200°C, and then an increase at a rate of 20°C/min to 320°C, which was held for 5 min. For unambiguous identification of compounds, samples were run on a Hewlett-Packard model 6890 gas chromatograph equipped with a model 5973 mass spectrometer (GC-MS) and the Chemstation analysis program (Hewlett-Packard, Kista, Sweden).

HPLC. *A. chlorophenolicus* A6 cells, grown to the late log phase in GM medium with 1.2 mM 4-CP, were harvested by centrifugation, washed, and subsequently resuspended to an OD₆₀₀ of 0.1 in 300 ml GM medium supplemented with 0.15 mM 4-CP, 4-chlorocatechol, or hydroquinone. The cultures were incubated with shaking at 28°C, and at regular intervals samples were taken and used for high-pressure liquid chromatography (HPLC) and cell mass analyses. Prior to HPLC analysis 0.3 mg sodium dithionite was added to cell-free supernatants obtained after centrifugation. The concentrations of 4-CP and metabolites were determined with an Agilent 1100 high-pressure liquid chromatograph (Agilent Technologies, Waldbronn, Germany).

PCR, cloning, and sequencing. Unless otherwise noted, the standard PCR conditions were as follows: each deoxynucleoside triphosphate at a concentration of 50 μM, each primer at a concentration of 0.7 μM, 1.5 mM MgCl₂, 2.5 U *Taq* polymerase (Amersham Biosciences, Buckinghamshire, England), and 10 ng *A. chlorophenolicus* A6 genomic DNA in a 50-μl reaction mixture. The reactions were performed with a Biometra Uno II thermocycler (Whatman Biometra, Göttingen, Germany).

Primers that annealed to conserved regions of hydroxyquinol 1,2-dioxygenase genes were designed by aligning the sequence of *npdB*, encoding this enzyme in *Arthrobacter* sp. strain JS443 (L. L. Perry and G. J. Zylstra, unpublished data), to the DNA sequence of ORF2 (20) and locating conserved regions. *Pfu* DNA polymerase (Promega, Madison, WI) was used for amplification from *A. chlorophenolicus* A6 genomic DNA.

A PCR walking system (15) was used to clone sequences from *A. chlorophenolicus* A6 (Table 1), and the PCR products were screened for similarity to cloned regions by Southern blotting. One set of primers was designed to anneal to conserved regions of maleylacetate reductase genes (Table 1). For PCR with

these primers, the Expand system (Roche Diagnostics, Mannheim, Germany) was used.

Part of the *cph* cluster (clone c6) was obtained by creating a subgenomic library. Southern blotting of EcoRI-digested *A. chlorophenolicus* A6 genomic DNA revealed that a 2,000-bp fragment hybridized to the *cphF-II* open reading frame (ORF) (this study). Therefore, EcoRI fragments in the size range from 1,800 to 2,200 bp were cloned into pUC18. Positive clones that hybridized to *cphF-II* were identified by colony hybridization, and the inserts were sequenced.

To overexpress *cphA-I* and *cphA-II*, the ORFs were PCR amplified from *A. chlorophenolicus* A6 genomic DNA (Table 1) and cloned into the vector pCRT7/CT-TOPO (Invitrogen), which created C-terminal six-His-tagged fusions to the genes. The constructs were transformed into competent BL21(DE3)/pLysS cells (Invitrogen), and cultivation and induction were performed as specified by the manufacturers.

Preparation of cell extracts. *A. chlorophenolicus* A6 cells were harvested by centrifugation, suspended in 50 mM phosphate buffer (pH 7.0), and passed once through a French press (164 MPa). The lysate was centrifuged at 9,000 × *g* for 10 min, and the supernatant was recovered.

Crude extracts of *E. coli* expressing CphA-I or CphA-II were prepared in 2 ml of 50 mM phosphate buffer (pH 7.0) and were disrupted by three sonication treatments (1 min each) at 0°C with a Branson Sonifier 250 at an output level of 5.5 (18 W). Each lysate was centrifuged at 14,000 × *g* for 30 min, and the supernatant was saved and used as a crude cell extract.

Enzyme activity assays. Hydroxyquinol removal, 4-chlorocatechol removal, and catechol removal were monitored separately in 50 mM phosphate buffer (pH 7.0) with 5 to 141 μg crude cell extract and 0.2 mM substrate in a 1-ml (final volume) (*E. coli*) or 0.5-ml (final volume) (*A. chlorophenolicus* A6) mixture using a Shimadzu UV-2501 PC or Hitachi U-2000 dual-beam spectrophotometer, as previously described (24). In the hydroxyquinol assay a maleylacetate peak at 245 nm was also detected. One unit of activity was defined as the amount of enzyme required to catalyze the oxidation of 1 μmol hydroxyquinol in 1 min. Catechol 1,2-dioxygenase activity was detected by measuring the increase in absorbance at 260 nm for *cis,cis*-muconate, and 1 U of activity was defined as the amount of enzyme that produced 1 μmol *cis,cis*-muconate in 1 min.

Mutant construction. *A. chlorophenolicus* A6 was transformed with plasmid pKGT452Cβ, which contains an *Arthrobacter*-derived, randomly inserting trans-




FIG. 1. Release of chloride ions by *A. chlorophenolicus* A6 during growth on different concentrations of 4-CP. The theoretical stoichiometric release is indicated by a straight line. The symbols indicate the means for three replicates, and the error bars indicate the standard deviations of the means ($n = 3$).

poson conferring chloramphenicol resistance (10). Preparation of *A. chlorophenolicus* A6 cells and electroporation were performed as described previously (18). After recovery in 10% LB for 2 h, the cells were inoculated onto agar plates containing 10% LB and 10 $\mu\text{g/ml}$ chloramphenicol and incubated at 28°C for 3 days. Chloramphenicol-resistant clones (approximately 2,000 clones) were transferred to plates containing GM medium with 1.2 mM 4-CP as the sole carbon source and 5 $\mu\text{g/ml}$ chloramphenicol. The wild-type strain *A. chlorophenolicus* A6 did not exhibit any visible growth on medium containing chloramphenicol. Clones that grew poorly on 4-CP were subjected to further screening. Transposon insertions in the *cph* gene cluster were screened using PCR primers Tra1, Tra2, Tra3, and Tra4 (Table 1) targeted to internal regions of the gene cluster. Single chromosomal insertion of a transposon was detected as a size increase of 3,409 bp in the amplified fragment combined with Southern blotting using a 414-bp fragment of the transposon's chloramphenicol resistance gene as a probe.

Nucleotide sequence accession number. The nucleotide sequence described here has been deposited in the GenBank database under accession number AY131335.

RESULTS

Stoichiometric release of chloride ions from 4-CP. Chloride ions were released in stoichiometric amounts from 4-CP at increasing substrate concentrations up to 2.7 mM, which is the highest 4-CP concentration on which *A. chlorophenolicus* A6 can grow (28) (Fig. 1). These data corroborate previous results from studies with ^{14}C -labeled 4-CP (29) indicating that *A. chlorophenolicus* mineralizes 4-CP to carbon dioxide, chloride ions, and water.

Detection of potential 4-CP metabolites. Four potential metabolites (designated compounds I, II, III, and IV) were detected in cultures of *A. chlorophenolicus* A6 growing on 4-CP.




FIG. 2. Results of a representative experiment showing metabolite accumulation during growth of *A. chlorophenolicus* A6 on 2.1 mM 4-CP.

Compounds I, II, and III were identified as hydroquinone, 4-chlorocatechol, and hydroxyquinol, respectively, by gas chromatography with flame ionization detection and by GC-MS analysis (Table 2). Compound IV did not cochromatograph with any standard. GC-MS analysis revealed that the fragmentation pattern of compound IV was consistent with that of a chlorohydroxyquinol, and the spectrum was similar to the chlorohydroxyquinol spectrum described by Apajalahti and Salkioja-Salonen (1) (Table 2).

4-CP degradation by *A. chlorophenolicus* A6 and production of intermediates were studied during growth on 4-CP as a sole carbon source at a concentration of 2.1 mM. After 20 h of incubation, when the cultures were in the mid-log phase, 4-chlorocatechol was the metabolite that accumulated to the highest levels (Fig. 2). At earlier incubation times (5 to 6 h, cells in the lag phase) hydroquinone and 4-chlorocatechol were the only detectable metabolites, and their concentrations were approximately 7 μM and 12 μM , respectively (data not shown). After 26 h of incubation, no intermediates were detected in the medium. When succinate or yeast extract was present as an auxiliary substrate, 4-CP degradation was delayed until the substrate was depleted (data not shown).

In a separate set of experiments, potential metabolites of 4-CP degradation were exogenously added to 4-CP-induced cells, and their rates of removal in triplicate cultures were monitored. The cell densities were adjusted to an OD_{600} of 0.1 as this value was the cell density in the mid-log phase during growth on 4-CP as a sole carbon source. We found that there was not a significant difference ($P < 0.01$) between the degradation rates of hydroquinone and 4-CP and that both com-

TABLE 2. Fragmentation patterns of 4-CP metabolites as determined by GC-MS analysis

Compound	Retention time (min)	m/z of observed fragment ions	Identification
I	6.15	254 (M^+), 239 ($\text{M} - \text{CH}_3$), 73 [$\text{Si}(\text{CH}_3)_3$]	Silylated hydroquinone
II	6.71	290, 288 (M^+), 275, 273 ($\text{M} - \text{CH}_3$), 200, 185, 170, 73 [$\text{Si}(\text{CH}_3)_3$]	Silylated 4-chlorocatechol
III	7.76	342 (M^+), 327 ($\text{M} - \text{CH}_3$), 254, 239, 73 [$\text{Si}(\text{CH}_3)_3$]	Silylated hydroxyquinol
IV	9.11	378, 376 (M^+), 363, 361 ($\text{M} - \text{CH}_3$), 288, 275, 273, 237, 179, 73 [$\text{Si}(\text{CH}_3)_3$]	Silylated chlorohydroxyquinol




FIG. 3. Proposed 4-CP degradation pathway in *A. chlorophenolicus* A6. Compound 1, 4-CP; compound 2, benzoquinone; compound 3, hydroquinone; compound 4, 4-chlorocatechol; compound 5, 5-chlorohydroxyquinol; compound 6, 2-hydroxy-1,4-benzoquinone; compound 7, hydroxyquinol; compound 8, maleylacetate. The transformation from compound 5 to compound 6 is theoretically due to reductive dechlorination of compound 5. Compound 6 is presented in a different orientation with respect to compound 5 for ease of presentation of the rest of the pathway. The presence of maleylacetate is inferred from genetic and biochemical evidence. Brackets indicate hypothetical intermediates supported, but not confirmed, by our data. An alternative hydroquinone cleavage route to hydroxymuconic semialdehyde that is not supported by our data is indicated by a dashed arrow.

pounds were degraded at a rate of 179 $\mu\text{mol/g}$ cells/min. By contrast, 4-chlorocatechol removal was significantly slower (78 $\mu\text{mol/g}$ cells/min), and there was a sharp decline in the rate after 1.5 h of incubation. The only metabolite that was detected upon incubation with 4-chlorocatechol in repeated experiments was chlorohydroxyquinol (average concentration, 3.0 μM ; $n = 3$). No 4-chlorocatechol or catechol dioxygenase activity was detected in 4-CP-induced cell extracts that exhibited confirmed activity with hydroxyquinol, indicating that clas-

sical 4-chlorocatechol metabolism via the *ortho* cleavage pathway was not induced.

We also studied the impact of an inhibitor of ring cleavage dioxygenases, 2,2'-dipyridyl (6), on the growth of *A. chlorophenolicus* cells. Actively 4-CP-degrading cultures changed color within minutes after 2,2'-dipyridyl addition and eventually became deep red, which was indicative of hydroxyquinol accumulation (14). In four separate experiments, hydroxyquinol accumulated to levels that were approximately 12% of the 4-CP concentration at the time of 2,2'-dipyridyl addition.

Another indication of hydroxyquinol dioxygenase activity was the ability of cell extracts from 4-CP-grown cells to deplete hydroxyquinol from the medium with an activity of 96 mU/mg protein (standard deviation, 4.4 mU/mg protein; $n = 3$). This activity was apparently induced in the presence of 4-CP since no depletion of hydroxyquinol was measured in extracts of succinate-grown cells (detection limit, 3.7 mU/mg protein). Based on these results, we propose a 4-CP degradation pathway via ring cleavage of hydroxyquinol (Fig. 3).

Cloning of a *cph* gene cluster. Southern blot analysis indicated that *A. chlorophenolicus* A6 contains a gene similar to the *npdB* gene, which encodes a hydroxyquinol 1,2-dioxygenase in *Arthrobacter* sp. strain JS443 (Perry and Zylstra, unpublished). A similar, but not identical, sequence was cloned from *A. chlorophenolicus* A6 using PCR primers designed to anneal to conserved regions in hydroxyquinol 1,2-dioxygenase genes (fragment c1 in Fig. 4; Table 1). Flanking sequences (c2 to c5, c7, and c8 in Fig. 4) were cloned and sequenced using a combination of primer walking and PCR amplification with primers targeted to conserved sequences of predicted genes (Table 1 and Fig. 4). One region (c6) (Fig. 4) was identified by screening a subgenomic library. The combined sequences generated a 15,475-bp contiguous stretch of *A. chlorophenolicus* A6 DNA (Fig. 4). The best matches with existing sequences in databases for the ORFs are shown in Table 3. Based on the results of




FIG. 4. Proposed *cph* gene cluster, including plasmid clones and primers (Table 1). The position of each clone is indicated below the line representing the consensus sequence, and primers are indicated by arrowheads. The triangle indicates the site of insertion of Tn1409C β in mutant T99.

TABLE 3. Amino acid sequence comparisons

Gene product	Position in accession no. AY131335 sequence	Similar protein	Function (gene, organism)	% Identity ^a	Score (bits)	E value	Accession no.
CphF-I	795–1859 (complement)	PnpD	Maleylacetate reductase, <i>Ralstonia</i> sp. strain SJ98	45.8	302	1e-80	AAS87585
CphC-I	1965–3572 (complement)	NpcA	Oxygenase component of 4-nitrophenol monooxygenase, <i>Rhodococcus opacus</i>	72.0	825	0.0	BAD30042
CphS	4060–4950 (complement)	TfuA	Involved in trifolixotoxin production, <i>Rhizobium leguminosarum</i> bv. trifolii T24	12.4	58	2e-07	AAB17513
CphR	4954–7098 (complement)	AcoK	<i>trans</i> -Acting regulatory protein of <i>aco</i> operon, <i>Klebsiella pneumoniae</i> CG43	17.2	95	6e-18	AAC44880
CphB	7169–7996	NpcB	Reductase component of 4-nitrophenol monooxygenase, <i>Rhodococcus opacus</i>	37.1	141	2e-32	BAD30041
CphA-I	8415–9329	ORF2	Hydroxyquinol 1,2-dioxygenase, <i>Arthrobacter</i> strain BA-5-17	75.7	467	e-130	BAA82713
CphX	9419–10048	ProX	Unknown function in phthalate degradation, <i>Pseudomonas straminea</i> NGJ1	28.9	74	2e-12	BAB21454
CphA-II	11483–12394 (complement)	ORF2	Hydroxyquinol 1,2-dioxygenase, <i>Arthrobacter</i> strain BA-5-17	77.6	491	e-138	BAA82713
CphF-II	12465–13541 (complement)	PnpD	Maleylacetate reductase, <i>Ralstonia</i> sp. strain SJ98	44.4	286	4e-76	AAS87585
CphC-II	13545–15341 (complement)	PheA	Phenol 2-monooxygenase, <i>Pseudomonas</i> sp. strain EST1001	62.9	762	0.0	AAC64901

^a For calculation of percent identity, gaps were treated as a 21st amino acid, but terminal gaps were ignored.

these analyses, the gene organization shown in Fig. 4 is proposed, and the putative genes were designated *cph* (chlorophenol degradation).

Several of the ORFs in the *cph* cluster were predicted to encode proteins with similar functions. For example, there were two hydroxyquinol 1,2-dioxygenases (encoded by *cphA-I* and *cphA-II*), two putative monooxygenases (*cphC-I* and *cphC-II*), and two putative maleylacetate reductases (*cphF-I* and *cphF-II*). The *cphB* gene could encode an NADH:flavin adenine dinucleotide oxidoreductase to supply the monooxygenase encoded by *cphC-I* with reduced flavin adenine dinucleotide (Table 3).

When the derived amino acid sequences of CphA-I and CphA-II were aligned with the sequences of other intradiol-cleaving enzymes, we found that two histidyl and tyrosyl residues coordinating the ferric iron in protocatechuate 3,4-dioxygenase from *Pseudomonas putida* were also conserved in CphA-I and CphA-II (23). The amino acid sequence of CphA-II was 78.4% identical to that of CphA-I.

The *cphX* ORF showed some similarity to a gene with an unknown function found in a phthalate degradation gene cluster in *Pseudomonas straminea* (17). There were also some similarities to molybdate transport proteins. However, due to the low quality of the database hits we were not able to assign a putative function to this ORF.

There were several indications that some of the ORFs with duplicated functions in the *cph* cluster were derived via a horizontal gene transfer event. To begin with, the region from position 10599 to position 11158 between *cphX* and *cphA-II* (Fig. 4) is a nonfunctional, putative resolvase pseudogene that was calculated to be, on the DNA level, 65% identical to the resolvase gene in the Tn21 family of the phthalate degradation pathway genes in *Arthrobacter keyseri* (accession number AF331043) and also 65% identical to Tn1721 in *E. coli* (accession number X02590). In addition, analyses of G+C contents revealed that the ORFs upstream of the resolvase gene in Fig. 4 had lower total G+C contents (56 to 59.6%) than the *A. chlorophenolicus* genome as a whole (65.1%), whereas the

ORFs downstream of the resolvase gene had similar G+C contents (62.1 to 67.1%). Moreover, the third-position G+C content and the effective number of codons (30) of the ORFs upstream of the resolvase gene differed from the values for the downstream ORFs, which had values more similar to those obtained for *Arthrobacter globiformis*, the type species of the genus (13; data not shown). Taken together, these results suggest that the region of the *cph* gene cluster upstream of the resolvase gene was acquired via horizontal gene transfer from another organism (22).

Expression of *cphA-I* and *cphA-II* in *E. coli*. Recombinant *cphA-I* and *cphA-II* were overexpressed in *E. coli*, and hydroxyquinol removal was studied in crude extracts with measured activities of 16 and 3.0 U/mg protein, respectively. In addition, CphA-I and CphA-II exhibited some *ortho* cleavage activity with catechol (0.67 mU/mg protein for CphA-I and 0.15 mU/mg protein for CphA-II), which has also been detected for other bacterial hydroxyquinol 1,2-dioxygenases (2, 20). Neither CphA-I (1.6 U) nor CphA-II (0.3 U) exhibited activity with 4-chlorocatechol, hydroquinone, or resorcinol (with 6 min of incubation and a reaction setup like that used for hydroxyquinol). Sodium dodecyl sulfate-polyacrylamide gel electrophoresis analysis revealed that the molecular masses of recombinant CphA-I and CphA-II were 40 kDa and 42 kDa, respectively (data not shown). Both of these observed molecular masses are slightly larger than the values expected (theoretically 33.5 and 33.0 kDa, respectively, plus the 3.3-kDa His tag), which could have been due to posttranslational modification of the proteins and/or running anomalies in the gel.

Generation and analysis of a *cphA-I* knockout mutant. We used transposon mutagenesis to disrupt the *cphA-I* gene, which generated mutant strain *A. chlorophenolicus* T99 (Fig. 4). Sequencing of the transposon insertion site revealed a duplication of 8 bases (positions 8862 to 8869 in the *cph* gene cluster) at the point of insertion, which is consistent with properties of the Tn1409C β transposon (10). The mutant contained a single transposon insertion, as verified by Southern blotting (data not shown). In repeated experiments we confirmed that unlike the

wild-type strain, the T99 mutant was not able to grow on 1.2 mM 4-CP as a sole carbon source, as shown by no visible increase in cell density after 40 h of incubation. In addition, a red-orange color accumulated during incubation of the mutant with 4-CP, which indicated that there was hydroxyquinol accumulation. Therefore, 4-CP catabolic ability was disrupted by the transposon insertion in the *cphA-I* gene.

DISCUSSION

Our results indicate that *A. chlorophenolicus* A6 degrades 4-CP via hydroxyquinol. To begin with, this compound was found in culture medium during 4-CP degradation. In addition, hydroxyquinol was removed from cell extracts derived from 4-CP-grown cells but not from extracts of cells grown on succinate. Therefore, there is clearly induction of the ability to remove hydroxyquinol when 4-CP is the growth substrate compared to when an alternative substrate is used. Furthermore, our results suggest that hydroxyquinol is the ring cleavage substrate during 4-CP degradation by *A. chlorophenolicus* A6 (Fig. 3). This hypothesis is supported by our observation that hydroxyquinol accumulates upon iron chelation (i.e., conditions under which the ring cleavage enzyme is inhibited). However, we did not observe a 1:1 accumulation of hydroxyquinol from 4-chlorophenol, which could have been due to the chemical instability of this compound (32) or incomplete inhibition.

In support of the biochemical evidence, a gene cluster containing genes encoding two hydroxyquinol 1,2-dioxygenases, CphA-I and CphA-II, was cloned from the *A. chlorophenolicus* A6 genome. Also, we successfully inactivated the *cphA-I* gene by transposon mutagenesis, and mutant strain T99 could not utilize 4-CP as a carbon source. Therefore, CphA-I (a hydroxyquinol 1,2-dioxygenase according to BLAST homology search results and the activity when it was expressed in *E. coli*) is required for 4-CP transformation and mineralization in this organism. However, we cannot completely disregard the possibility that the lack of growth was due to polar effects on *cphX*, whose function is currently unknown. Although the *cphA-II* gene was intact in the mutant strain and produced a functional protein when it was expressed in *E. coli*, it was insufficient to enable growth of the T99 mutant strain on 4-CP, and thus its functional significance in vivo is currently not known. Possibly, the gene is not optimally transcribed, or it is subject to negative gene regulation under these conditions. Taken together, the combined chemical and genetic evidence indicates that 4-CP degradation in *A. chlorophenolicus* A6 proceeds via hydroxyquinol, which is subsequently cleaved. To our knowledge, this is the first time that such a pathway has been described for aerobic 4-CP degradation by a bacterium.

4-Chlorocatechol, chlorohydroxyquinol, and hydroquinone were also found to be metabolites in addition to hydroxyquinol during growth on 4-CP. The logical positions of these compounds in the proposed degradation route are shown in Fig. 3; chlorohydroxyquinol is in brackets as it has not been confirmed to be an intermediate. In accordance with previous reports, the oxidation and simultaneous removal of a chlorine group from the aromatic ring by a monooxygenase are likely to yield a quinone that can then be reduced to the corresponding quinol (31), and although this possibility was not addressed in the present work, the resulting quinone is shown in brackets in Fig.

3. Although *A. chlorophenolicus* A6 removed 4-chlorocatechol at a rate that was approximately one-half the rate of hydroquinone removal, the 4-chlorocatechol branch shown in Fig. 3 could theoretically still significantly contribute to 4-CP degradation in this strain.

The proposed transformation of 4-CP via hydroquinone to hydroxyquinol in *A. chlorophenolicus* A6 is similar to the transformation commonly found in the degradation of tri- to pentachlorophenols; however, the hydroquinone is chlorinated in these cases (24). In addition, transformation of nonchlorinated hydroquinone to hydroxyquinol has been suggested for degradation of 4-aminophenol (26). Another possibility is that hydroquinone, and not hydroxyquinol, is the ring cleavage substrate in *A. chlorophenolicus* A6, and this possibility is indicated by the dashed line in Fig. 3. Hydroxyquinol could then be transformed into hydroquinone prior to ring cleavage, as has been reported to occur in 4-nitrophenol degradation (7). However, the observed accumulation of hydroxyquinol in 2,2'-dipyridyl-inhibited cultures and the results of the *cphA-I* mutant studies discussed above support the possibility that there is ring cleavage directly from hydroxyquinol.

We also detected chlorohydroxyquinol in culture medium during 4-CP degradation. This compound could be formed by hydroxylation of 4-chlorocatechol and then dechlorinated to hydroxyquinol; the latter reaction is analogous to transformation of 5-chlorohydroxyquinol to hydroxyquinol by *Burkholderia cepacia* AC1100 during 2,4,5-trichlorophenoxyacetic acid degradation (32). Chlorohydroxyquinol would then be an intermediate between 4-chlorocatechol and hydroxyquinol in 4-CP degradation by *A. chlorophenolicus* A6 (Fig. 3), but this possibility remains to be resolved.

A. chlorophenolicus A6 can also degrade 4-nitrophenol, and although it was outside the scope of this study to elucidate the degradation pathway for this compound, 4-nitrophenol has been reported to be degraded via cleavage of both hydroquinone (25) and hydroxyquinol (14). Kitagawa et al. (14) recently cloned a 4-nitrophenol catabolic gene cluster containing a hydroxyquinol 1,2-dioxygenase with sequence similarity to CphA-I and CphA-II and a two-component monooxygenase similar to the CphC-I and CphB in this study. Possibly, the same enzyme system could be used for 4-CP degradation and 4-nitrophenol degradation in *A. chlorophenolicus* A6.

Two different types of monooxygenases are predicted to be present in *A. chlorophenolicus* A6. The CphC-I and CphB sequences were most similar to a two-component monooxygenase involved in 4-nitrophenol degradation (Table 3) (14). This type of monooxygenase, in which the large subunit is the oxygenase which receives electrons from the small subunit, an NADH:flavin adenine dinucleotide oxidoreductase, has also been found to be involved in degradation of 2,4,6-trichlorophenol (16, 27). The second putative monooxygenase in the *cph* gene cluster, CphC-II, was similar to a different family of monooxygenases, the flavin monooxygenases, which are not dependent on an oxidoreductase but instead transfer electrons from NAD(P)H to a flavin which is situated at the active site (11). Another difference from CphC-I is that the deduced CphC-II protein was most similar to a phenol monooxygenase acting on nonchlorinated substrates (21). In future studies, expression of the putative monooxygenases and studies of their

functions may shed light on the transformations prior to ring cleavage.

Three of the steps in the degradation pathway (addition of hydroxyl groups to 4-CP, cleavage of the aromatic ring, and reduction of maleylacetate) are represented by double sets of genes in *A. chlorophenolicus* A6 (Fig. 4). The presence of some of the genes appears to be due to lateral gene transfer since in the *cph* gene cluster there is a marked difference in G+C content between ORFs and the cluster contains a putative resolvase pseudogene, both of which are typical indications of a past horizontal gene transfer event (22). The ORFs apparently encoding isofunctional enzymes in *A. chlorophenolicus* A6 showed marked sequence differences. The two predicted monooxygenases appeared to be different types, as discussed above. Also, detailed phylogenetic analysis revealed that CphA-I was more closely related to NpdB, an isofunctional enzyme from *Arthrobacter* sp. strain JS443 (Perry and Zylstra, unpublished), than to CphA-II (not shown). Similar to the results of the present work, several cases of multiple copies of genes for catechol degradation have previously been reported (19).

In summary, in this study we identified both a chlorophenol degradation pathway and a 4-CP degradation gene cluster in *A. chlorophenolicus* A6. In future studies we plan to clarify the roles of the remaining genes in the *cph* gene cluster, such as *cphX*, and to determine how these genes are regulated in this bacterium.

ACKNOWLEDGMENTS

We thank Karl-Heinz Gartemann and Rudolf Eichenlaub at Universität Bielefeld, Germany, for the kind gift of the pKGT452CB vector; Ingvar Sundh and Elisabet Börjesson at the Swedish Agricultural University for help with the GC-MS analysis; Åsa Perez-Bercoff for technical assistance with the codon bias analysis; and Gerben J. Zylstra at Rutgers University for the kind gift of the *npdB* gene.

This work was supported by grants from the Swedish Council for Engineering Science and the Swedish Foundation for Environmental Research (MISTRA).

REFERENCES

- Apajalahti, J. H. A., and M. S. Salkinoja-Salonen. 1987. Complete dechlorination of tetrachlorohydroquinone by cell extracts of pentachlorophenol-induced *Rhodococcus chlorophenolicus*. *J. Bacteriol.* **169**:5125–5130.
- Armengaud, J., K. N. Timmis, and R.-M. Wittich. 1999. A functional 4-hydroxysalicylate/hydroxyquinol degradative pathway gene cluster is linked to the initial dibenzo-*p*-dioxin pathway genes in *Sphingomonas* sp. strain RW1. *J. Bacteriol.* **181**:3452–3461.
- Backman, A., and J. K. Jansson. 2004. Degradation of 4-chlorophenol at low temperature and during extreme temperature fluctuations by *Arthrobacter chlorophenolicus* A6. *Microb. Ecol.* **48**:246–253.
- Backman, A., N. Maraha, and J. K. Jansson. 2004. Impact of temperature on the physiological status of a potential bioremediation inoculant, *Arthrobacter chlorophenolicus* A6. *Appl. Environ. Microbiol.* **70**:2952–2958.
- Bac, H. S., J. M. Lee, and S.-T. Lee. 1996. Biodegradation of 4-chlorophenol via a hydroquinone pathway by *Arthrobacter ureafaciens* CPR706. *FEMS Microbiol. Lett.* **145**:125–129.
- Chapman, P. J., and D. J. Hopper. 1968. The bacterial metabolism of 2,4-xyleneol. *Biochem. J.* **110**:491–498.
- Chauhan, A., A. K. Chakraborti, and R. K. Jain. 2000. Plasmid-encoded degradation of *p*-nitrophenol and 4-nitrocatechol by *Arthrobacter protophormiae*. *Biochem. Biophys. Res. Commun.* **270**:733–740.
- Cho, Y.-G., J.-H. Yoon, Y.-H. Park, and S.-T. Lee. 1998. Simultaneous degradation of *p*-nitrophenol and phenol by a newly isolated *Nocardioideis* sp. *J. Gen. Appl. Microbiol.* **44**:303–309.
- Elväng, A. M., K. Westerberg, C. Jernberg, and J. K. Jansson. 2001. Use of green fluorescent protein and luciferase biomarkers to monitor survival and activity of *Arthrobacter chlorophenolicus* A6 cells during degradation of 4-chlorophenol in soil. *Environ. Microbiol.* **3**:32–42.
- Gartemann, K.-H., and R. Eichenlaub. 2001. Isolation and characterization of IS1409, an insertion element of 4-chlorobenzoate-degrading *Arthrobacter* sp. strain TM1, and development of a system for transposon mutagenesis. *J. Bacteriol.* **183**:3729–3736.
- Harayama, S., and M. Kok. 1992. Functional and evolutionary relationships among diverse oxygenases. *Annu. Rev. Microbiol.* **46**:565–601.
- Jernberg, C., and J. K. Jansson. 2002. Impact of 4-chlorophenol contamination and/or inoculation with the 4-chlorophenol-degrading strain, *Arthrobacter chlorophenolicus* A6L, on soil bacterial community structure. *FEMS Microbiol. Ecol.* **42**:387–397.
- Keddie, R. M., M. D. Collins, and D. Jones. 1986. Genus *Arthrobacter* Conn and Dimmick 1947, 300^{AL}, p. 1228–1301. In P. H. A. Sneath, N. S. Mair, M. E. Sharpe, and J. G. Holt (ed.), *Bergey's manual of systematic bacteriology*, vol. 2. Williams & Wilkins, Baltimore, Md.
- Kitagawa, W., N. Kimura, and Y. Kamagata. 2004. A novel *p*-nitrophenol degradation gene cluster from a gram-positive bacterium, *Rhodococcus opacus* SAO101. *J. Bacteriol.* **186**:4894–4902.
- Lin, X., D. W. Kelemen, E. S. Miller, and J. C. H. Shih. 1995. Nucleotide sequence and expression of *kerA*, the gene encoding a keratinolytic protease of *Bacillus licheniformis* PWD-1. *Appl. Environ. Microbiol.* **61**:1469–1474.
- Louie, T. M., C. M. Webster, and L. Xun. 2002. Genetic and biochemical characterization of a 2,4,6-trichlorophenol degradation pathway in *Ralstonia eutropha* JMP134. *J. Bacteriol.* **184**:3492–3500.
- Murayama, K., M. Miwa, N. Tsujii, T. Nagai, N. Tomita, T. Harada, H. Sobajima, and H. Sugisaki. 2001. Cloning, sequencing, and expression of the gene encoding 4-hydroxy-4-methyl-2-oxoglutarate aldolase from *Pseudomonas ochraceae* NGJ1. *Biosci. Biotechnol. Biochem.* **65**:2701–2709.
- Möller, A., and J. K. Jansson. 1998. Detection of firefly luciferase-tagged bacteria in environmental samples. *Methods Mol. Biol.* **102**:269–283.
- Murakami, S., A. Takashima, J. Takemoto, S. Takenaka, R. Shinke, and K. Aoki. 1999. Cloning and sequence analysis of two catechol-degrading gene clusters from the aniline-assimilating bacterium *Frateuria* species ANA-18. *Gene* **226**:189–198.
- Murakami, S., T. Okuno, E. Matsumura, S. Takenaka, R. Shinke, and K. Aoki. 1999. Cloning of a gene encoding hydroxyquinol 1,2-dioxygenase that catalyzes both intradiol and extradiol ring cleavage of catechol. *Biosci. Biotechnol. Biochem.* **63**:859–865.
- Nurk, A., L. Kasak, and M. Kivisaar. 1991. Sequence of the gene (*pheA*) encoding phenol monooxygenase from *Pseudomonas* sp. EST1001: expression in *Escherichia coli* and *Pseudomonas putida*. *Gene* **102**:13–18.
- Ochman, H., J. G. Lawrence, and E. A. Groisman. 2000. Lateral gene transfer and the nature of bacterial innovation. *Nature* **405**:299–304.
- Ohlendorf, D. H., J. D. Lipscomb, and P. C. Weber. 1988. Structure and assembly of protocatechuate 3,4-dioxygenase. *Nature* **336**:403–405.
- Padilla, L., V. Matus, P. Zenteno, and B. González. 2000. Degradation of 2,4,6-trichlorophenol via chlorohydroxyquinol in *Ralstonia eutropha* JMP134 and JMP222. *J. Basic Microbiol.* **4**:243–249.
- Prakash, D., A. Chauhan, and R. K. Jain. 1996. Plasmid-encoded degradation of *p*-nitrophenol by *Pseudomonas cepacia*. *Biochem. Biophys. Res. Commun.* **224**:375–381.
- Takenaka, S., S. Okugawa, M. Kadowaki, S. Murakami, and K. Aoki. 2003. The metabolic pathway of 4-aminophenol in *Burkholderia* sp. strain AK-5 differs from that of aniline and aniline with C-4 substituents. *Appl. Environ. Microbiol.* **69**:5410–5413.
- Takizawa, N., H. Yokoyama, K. Yanagihara, T. Hatta, and H. Kiyohara. 1995. A locus of *Pseudomonas pickettii* DTP0602, *had*, that encodes 2,4,6-trichlorophenol-4-dechlorinase with hydroxylase activity, and hydroxylation of various chlorophenols by the enzyme. *J. Ferment. Bioeng.* **80**:318–326.
- Westerberg, K., A. M. Elväng, E. Stackebrandt, and J. K. Jansson. 2000. *Arthrobacter chlorophenolicus* sp. nov., a new species capable of degrading high concentrations of 4-chlorophenol. *Int. J. Syst. Appl. Microbiol.* **50**:2083–2092.
- Westerberg, K., C. Jernberg, A. M. Elväng, and J. K. Jansson. 2001. *Arthrobacter chlorophenolicus* A6—the complete story of a 4-chlorophenol degrading bacterium, p. 133–140. In V. S. Magar, G. Johnson, S. K. Ong, and A. Leeson (ed.), *Bioremediation of energetics, phenolics, and polynuclear aromatic hydrocarbons*. Proceedings of the Sixth International In Situ and On-Site Bioremediation Symposium. Battelle Press, Columbus, Ohio.
- Wright, F. 1990. The 'effective number of codons' used in a gene. *Gene* **87**:23–29.
- Xun, L., and C. M. Webster. 2004. A monooxygenase catalyzes sequential dechlorinations of 2,4,6-trichlorophenol by oxidative and hydrolytic reactions. *J. Biol. Chem.* **279**:6696–6700.
- Zaborina, O., D. L. Daubaras, A. Zago, L. Xun, K. Saido, T. Klem, D. Nikolic, and A. M. Chakraborty. 1998. Novel pathway for conversion of chlorohydroxyquinol to maleylacetate in *Burkholderia cepacia* AC1100. *J. Bacteriol.* **180**:4667–4675.