

# PsaC Subunit of Photosystem I Is Oriented with Iron-Sulfur Cluster $F_B$ as the Immediate Electron Donor to Ferredoxin and Flavodoxin

Ilya R. Vassiliev,\* Yean-Sung Jung,# Fan Yang,§ and John H. Golbeck\*

\*Department of Biochemistry and Molecular Biology, S-310 Frear Laboratory, The Pennsylvania State University, University Park, Pennsylvania 16802; §Department of Chemistry, University of Nebraska, Lincoln, Nebraska 68588; and #Department of Biochemistry, University of Nebraska, Lincoln, Nebraska 68588

**ABSTRACT** The PsaC subunit of photosystem I (PS I) binds two [4Fe-4S] clusters,  $F_A$  and  $F_B$ , functioning as electron carriers between  $F_X$  and soluble ferredoxin. To resolve the issue whether  $F_A$  or  $F_B$  is proximal to  $F_X$ , we used single-turnover flashes to promote step-by-step electron transfer between electron carriers in control (both  $F_A$  and  $F_B$  present) and HgCl<sub>2</sub>-treated ( $F_B$ -less) PS I complexes from *Synechococcus* sp. PCC 6301 and analyzed the kinetics of P700<sup>+</sup> reduction by monitoring the absorbance changes at 832 nm in the presence of a fast electron donor (phenazine methosulfate (PMS)). In control PS I complexes exogenously added ferredoxin, or flavodoxin could be photoreduced on each flash, thus allowing P700<sup>+</sup> to be reduced from PMS. In  $F_B$ -less complexes, both in the presence and in the absence of ferredoxin or flavodoxin, P700<sup>+</sup> was reduced from PMS only on the first flash and was reduced from  $F_X^-$  on the following flashes, indicating lack of electron transfer to ferredoxin or flavodoxin. In the  $F_B$ -less complexes, a normal level of P700 photooxidation was detected accompanied by a high yield of charge recombination between P700<sup>+</sup> and  $F_A^-$  in the presence of a slow donor, 2,6-dichlorophenol-indophenol. This recombination remained the only pathway of  $F_A^-$  reoxidation in the presence of added ferredoxin, consistent with the lack of forward electron transfer.  $F_A^-$  could be reoxidized by methyl viologen in  $F_B$ -less PS I complexes, although at a concentration two orders of magnitude higher than is required in wild-type PS I complexes, thus implying the presence of a diffusion barrier. The inhibition of electron transfer to ferredoxin and flavodoxin was completely reversed after reconstituting the  $F_B$  cluster. Using rate versus distance estimates for electron transfer rates from  $F_X$  to ferredoxin for two possible orientations of PsaC, we conclude that the kinetic data are best compatible with PsaC being oriented with  $F_A$  as the cluster proximal to  $F_X$  and  $F_B$  as the distal cluster that donates electrons to ferredoxin.

## INTRODUCTION

PsaC is a photosystem I-bound, 8.9-kDa polypeptide that contains two [4Fe-4S] clusters termed  $F_A$  and  $F_B$  (Hayashida et al., 1987). Although its three-dimensional structure has not yet been solved, the main-chain folding pattern of PsaC is presumed to be similar to the small bacterial dicluster ferredoxins from *Peptococcus asaccharolyticus* (Adman et al., 1976) (formerly *P. aerogenes*) and *Clostridium acidi-urici* (Duee et al., 1994), which contain two  $\alpha$ -helices near the iron-sulfur clusters and two regions of two-stranded antiparallel  $\beta$ -sheet. In common with these proteins, PsaC is likely to possess a pseudo- $C_2$  symmetry axis that is oriented perpendicular to a distance vector connecting the two iron-sulfur clusters,  $F_A$  and  $F_B$ . The amino acid sequence of 2[4Fe-4S] ferredoxins usually contains two Cxx-CxxCxxxCP iron-sulfur binding motifs (Dunn and Gray, 1988;

Oh-oka et al., 1988) in which the first three cysteines in one motif cooperate with the fourth cysteine in the other motif to bind one cubane cluster. The location of  $F_A$  and  $F_B$  relative to the cysteine ligands was first deduced by in vitro mutagenesis studies (Zhao et al., 1992) and has been confirmed using in vivo mutagenesis (Yu et al., 1997; Jung et al., 1997; Mannan et al., 1996).  $F_A$  with principal  $g$ -values of 1.86, 1.94, and 2.05 and a midpoint potential of  $-540$  mV is identified as the cluster ligated by cysteines 21, 48, 51, and 54.  $F_B$  with principal  $g$ -values of 1.89, 1.92, and 2.07 and a midpoint potential of  $-590$  mV is identified as the cluster ligated by cysteines 11, 14, 17, and 58 (for review see Brettel, 1997).

An electron paramagnetic resonance (EPR) study of membrane-oriented photosystem I (PS I) complexes (Guigliarelli et al., 1993) provided the first indication that the  $F_A$ - $F_B$  axis was tilted from the membrane plane, implying that electron transfer from  $F_X$  to ferredoxin occurs sequentially through the two iron-sulfur clusters. X-ray crystallographic studies showed that the center-to-center distances between  $F_X$  and the two PsaC-bound [4Fe-4S] clusters are 15 Å and 22 Å, respectively; the distance between  $F_A$  and  $F_B$  is  $12 \text{ Å} \pm 0.5 \text{ Å}$ ; and the distance vector connecting  $F_A$  and  $F_B$  is tilted  $54^\circ (\pm 5^\circ)$  from the membrane normal (Krauss et al., 1993; 1996). The full  $g$ -tensor orientation of  $F_A$  and  $F_B$  in single crystals of PS I further fixes the orientation of PsaC along a rotation axis that passes through  $F_A$  and  $F_B$  (Kamlowski et al., 1997a; 1997b). However, because of its pseudo- $C_2$  symmetric axis, there remains a twofold ambiguity in the orientation of PsaC, leaving the issue of whether  $F_A$  or  $F_B$  is proximal to  $F_X$  unresolved.

Received for publication 20 October 1997 and in final form 6 January 1998.

Address reprint requests to Dr. John H. Golbeck, Department of Biochemistry and Molecular Biology, S-310 Frear Laboratory, The Pennsylvania State University, University Park, PA 16802. Tel.: 814-865-1163; Fax: 814-863-7024; E-mail: jhg5@psu.edu.

Dr. Vassiliev is currently on leave from the Department of Biophysics, Faculty of Biology, M. V. Lomonosov Moscow State University, Moscow 119899, Russia.

Dr. Jung's present address is Department of Molecular Biology and Biochemistry, University of California, Irvine, California 92697

© 1998 by the Biophysical Society

0006-3495/98/04/2029/07 \$2.00

There are thus two possibilities for arranging  $F_A$  and  $F_B$  in the sequence of electron transfer from  $F_X$  to ferredoxin or flavodoxin. 1) The sequence  $F_X \rightarrow F_B \rightarrow F_A$  was invoked from the preferential  $F_B$  photoreduction in the presence of chemically reduced  $F_A$  (Heathcote et al., 1978) and the lack of  $F_A$  photoreduction upon  $F_B$  destruction by diazonium benzene sulfonate (Malkin, 1984). This arrangement also complies with the fact that  $F_A$  is more electropositive than  $F_B$ . Additional evidence for this arrangement follows from studies of rebinding of a mutant PsaC with specific amino acid substitutions around the two iron-sulfur cluster binding sites. Based on the premise of electrostatic interaction between D9 in PsaC (which is in close proximity of  $F_B$ ) and an arginine of one of the external loops of either PsaA or PsaB, Biggins et al. (1995; Rodday et al., 1996) suggested the most favorable orientation of PsaC has  $F_B$  as the proximal cluster to  $F_X$ . The lack of efficient reconstruction of a PS I core with a loop-deleted PsaC (Naver et al., 1996) also favors the  $F_X \rightarrow F_B \rightarrow F_A$  sequence. 2) The sequence  $F_X \rightarrow F_A \rightarrow F_B$  was invoked from the EPR data on efficient photoreduction of  $F_A$  in the presence of chemically reduced  $F_B$  (Cammack et al., 1979; Nugent et al., 1981) and negligible  $F_B$  photoreduction in the presence of chemically reduced  $F_A$  (Bearden and Malkin, 1976). Studies on Hg-treated PS I complexes provide the strongest arguments for  $tk_2F_A$  as the  $F_X$ -proximal cluster. Sakurai et al. (1991) reported that  $F_B$  was more easily extracted by  $HgCl_2$  treatment, whereas  $F_A$  was left almost intact, consistent with lower steric hindrance of  $F_B$ . The steady-state rates of electron transfer from plastocyanin to  $NADP^+$  or to ferredoxin in spinach PS I (He and Malkin, 1994) and from cytochrome  $c_6$  to  $NADP^+$  or to flavodoxin in *Synechococcus* sp. PCC 6301 PS I (Jung et al., 1995) were inhibited by  $\sim 70\%$  upon  $HgCl_2$ -treatment. In the latter study,  $NADP^+$  reduction was completely restored upon rebuilding of  $F_B$  cluster. This implies that  $F_B$  functions as the terminal electron acceptor bound to PS I complex. However, these results are not unambiguous because a lower quantum efficiency of electron transfer might not be apparent in a steady-state measurement and therefore cannot be excluded as an alternative interpretation of low  $NADP^+$  reduction rates seen in  $HgCl_2$ -treated samples.

In this study, we used single-turnover saturating flashes at room temperature to promote a step-by-step electron transfer to the terminal electron acceptor in control (containing both  $F_A$  and  $F_B$ ) and  $F_B$ -less PS I complexes in the absence and presence of electron acceptors. By analyzing the kinetics of  $P700^+$  reduction kinetics from external donors (phenazine methosulfate (PMS) and 2,6-dichlorophenol-indophenol (DCPIP)), we find that  $F_B$  is required for forward electron transfer to ferredoxin or flavodoxin. This result addresses the structural twofold uncertainty in the orientation of PsaC on the PS I complex by supporting the following sequence of electron transfer:  $F_X \rightarrow F_A \rightarrow F_B$ .

## MATERIALS AND METHODS

Trimeric PS I complexes from *Synechococcus* sp. PCC 6301 were isolated using Triton X-100 and sucrose gradient ultracentrifugation (Golbeck,

1995). Preparation of  $F_B$ -less TX-PS I complexes by treatment with  $HgCl_2$  and reinsertion of the  $F_B$  iron-sulfur cluster using  $FeCl_3$ ,  $Na_2S$ , and  $\beta$ -mercaptoethanol were performed as described previously (Jung et al., 1995). Recombinant ferredoxin from *Synechocystis* sp. PCC 7002 was overproduced in *Escherichia coli* strain BL21 cells harboring *petF* in the expression plasmid pSE280 (Mühlenhoff et al., 1996). Recombinant flavodoxin from *Synechocystis* sp. PCC 7002 was overproduced in *E. coli* strain BL21 cells harboring *isiB* in the expression plasmid pSE280 (Mühlenhoff et al., 1996). Ferredoxin and flavodoxin were purified as described elsewhere (Bottin and Lagoutte, 1992).


PS I complexes were suspended to a Chl *a* concentration of  $50 \mu g/ml$  in a 10-mm  $\times$  4-mm quartz cuvette. Kinetics of absorbance changes at 832 nm ( $\Delta A_{832}$ ) were measured with a spectrophotometer described previously (Vassiliev et al., 1997), except that the detection beam (power, 30 mW;  $\lambda$ , 832 nm) was provided by a PMT-25 laser diode assembly (Power Technology Inc., Little Rock, AR). Single turnover flashes were provided by a frequency-doubled ( $\lambda$ , 532 nm), Q-switched (FWHM, 10 ns) Nd-YAG laser model DCR-11 (Spectra-Physics, Mountain View, CA) at a flash energy of 10 mJ. Multiple flash excitation at 15-ms intervals was provided by a Xenon flash lamp (FWHM, 10  $\mu s$ , flash energy  $\sim 5$  mJ) Model 6100E-72 (Photochemical Research Associates, London, ON, Canada). The multiexponential fits of  $\Delta A_{832}$  kinetics were performed by the Marquardt algorithm in Igor Pro v. 3.03 (Wavemetrics, Lake Oswego, OR) on a Power Macintosh 7100/88 computer. In multiple flash experiments (Fig. 1), the four kinetics were analyzed by a global two-exponential fit to an equation  $A(t) = A_1 \times \exp(-t/\tau_1) + (A_0 - A_1 - A_3) \times \exp(-t/\tau_2) + A_3$ , in which  $\tau_1$  and  $\tau_2$  are the global lifetimes of the components,  $A_1$  is the free running amplitude of the first component,  $A_3$  is the free running baseline amplitude, and  $A_0$  is the global sum of all amplitudes; the amplitude of the second component  $A_2$  is represented by  $A_0 - A_1 - A_3$ .

Treatment of cyanobacterial PS I complexes with  $HgCl_2$  under the conditions specified in Jung et al. (1995) resulted in 90% destruction of the  $F_B$  iron-sulfur cluster and in the retention of 80% of the  $F_A$  iron-sulfur cluster when assayed by low temperature EPR spectroscopy (data not shown). To obtain kinetic confirmation for the removal of a single iron-sulfur cluster, we used  $\Delta A_{832}$  measurements and a multiple flash excitation protocol (Sauer et al., 1978) that enables quantitation of the number of photoactive electron acceptors at room temperature in PS I. This approach is based on a selected combination of two experimental conditions. The first condition is the use of a fast electron donor to  $P700^+$  (reduced PMS) provided at appropriate concentrations so that it reduces  $P700^+$  faster than the back-reaction of  $(F_A/F_B)^-$  but slower than the back-reaction of  $F_X^-$ . Given that the lifetimes of the back-reaction from  $(F_A/F_B)^-$  are  $\sim 10$  and 80 ms (1:4 amplitude ratio) and the lifetimes of the  $F_X^-$  back-reaction (in the presence of  $F_A^-$  and  $F_B^-$ ) are 450  $\mu s$  and 1.5 ms (5:1 amplitude ratio) (Vassiliev et al., 1997), the 6-ms electron donation from 50  $\mu M$  PMS provides a favorable forward electron transfer time. The second condition is the use of excitation flashes fired at time intervals (15 ms) that are shorter than the lifetime of  $P700^+$  ( $F_A/F_B$ ) $^-$  but longer than the forward transfer of electron donation from the exogenous donor. When these conditions are satisfied, the first two flashes lead to sequential reduction of  $F_A$  and  $F_B$  that remain reduced during the time interval before the third excitation flash. Hence, after the first two flashes,  $P700^+$  is reduced primarily from PMS, whereas on subsequent flashes  $P700^+$  is reduced primarily from  $F_X^-$ . The presence of the 6-ms component can therefore be used as an indicator of active forward electron transfer from  $F_X$  to one (or both) of the terminal clusters. If a soluble electron acceptor is present in the media, the efficiency of forward electron transfer from  $F_X^-$  will depend on the ability of  $F_A^-$  and  $F_B^-$  to be oxidized by this component.

## RESULTS

The global multiexponential fit of the kinetics of control PS I complexes (Fig. 1 A) yields two major components, which we denote as the  $F_X$ -component ( $\tau$ , 468  $\mu s$ ) and the PMS-component ( $\tau$ , 6.1 ms). Ideally, the multiple flash protocol

FIGURE 1 Kinetics of absorbance change at 832 nm ( $\Delta A_{832}$ ) of integral (left column), HgCl<sub>2</sub>-treated (middle column), and F<sub>B</sub>-reconstructed (right column) PS I complexes upon excitation with trains of four consecutive flashes at 15 ms intervals; flash numbers are indicated near the traces. The samples were suspended in 25 mM 2-[N-morpholino]ethanesulfonic acid buffer, pH 6.3, with 0.02% Triton X-100, 50  $\mu$ M PMS, and 2 mM sodium dithionite and incubated in the dark for 1 min before excitation. Three sets of four-flash trains separated by 1-min intervals were applied to each sample; then ferredoxin (2nd and 3rd rows) was added to the media at concentrations indicated in the figure, and four-flash trains were again applied at 1-min dark intervals. Flavodoxin (4th row) was added to a new sample instead of ferredoxin in identical experimental conditions. The kinetics in response to the flash trains were acquired as single traces, which were then averaged and cut into individual kinetics. The results of the global two-exponential fit are shown as solid lines.



calls for the appearance of the PMS-component only after the first two flashes and the appearance of the F<sub>X</sub>-component on all subsequent flashes. In practice, mixed kinetics occur in the experiment, which are due, in part, to the relatively close lifetime of the F<sub>X</sub><sup>-</sup> back-reaction and the forward electron donation time from PMS to P700<sup>+</sup>. The presence of the F<sub>X</sub>-component on the first two flashes is also a consequence of the unavoidable chemical reduction of a fraction of F<sub>A</sub> and/or F<sub>B</sub>, and the presence of the PMS-component on subsequent flashes indicates that PMS overrides the reduction of P700<sup>+</sup> from F<sub>X</sub><sup>-</sup> also in a small percentage of reaction centers. Finally, the small fraction of reaction centers with both clusters reduced is increased on the second flash due to photochemical reduction of the second PsaC-bound cluster in those centers where one of the clusters was chemically prereduced. This leads to an even higher contribution of the F<sub>X</sub>-component on the second flash. The net result is that the contribution of the PMS-component is 67% on the first flash and 46% on the second flash, whereas on the third and the fourth flashes it drops to 26 and 16%, respectively.

Consistent with the results of Sakurai et al. (1991), the kinetics of the HgCl<sub>2</sub>-treated PS I complex on the first flash differs dramatically from those on the second and all sub-

sequent flashes, which are nearly identical (Fig. 1 B). The contribution of the PMS-component is 64% on the first flash, whereas on the next three flashes it has lower values of 46, 45, and 46%, respectively. This indicates that unlike the control sample that has different contributions of the F<sub>X</sub><sup>-</sup> back-reaction on the second and subsequent flashes, the Hg-treated PS I complex has identical contributions of the F<sub>X</sub><sup>-</sup> back reaction on the second and subsequent flashes. Therefore, the F<sub>A</sub> cluster acts as an efficient electron acceptor from F<sub>X</sub> at room temperature as well as cryogenic temperatures (Jung et al., 1995). A higher contribution of the PMS-component in HgCl<sub>2</sub>-treated is due to a slight acceleration of the back-reaction between P700<sup>+</sup> and F<sub>A</sub><sup>-</sup>, which was uncovered in single-flash excitation experiments in the presence of a slow external donor to P700<sup>+</sup> (see below; Vassiliev et al., 1997).

The addition of 16  $\mu$ M ferredoxin affects the kinetic pattern of the  $\Delta A_{832}$  kinetics in the control but not in the HgCl<sub>2</sub>-treated PS I complexes (Fig. 1, D and E). An increase in ferredoxin concentration up to 48  $\mu$ M does not affect the kinetics in the HgCl<sub>2</sub>-treated complexes (Fig. 1, G and H). Addition of 21  $\mu$ M flavodoxin as an alternative electron acceptor also leads to a complete elimination of the flash-number-dependency in the control but not in the HgCl<sub>2</sub>-



treated PS I complexes (Fig. 1, *J* and *K*). A large increase in flavodoxin concentration up to 83  $\mu\text{M}$  has no additional effect on the  $\text{HgCl}_2$ -treated complexes (not shown). To show that the lack of effect of ferredoxin and flavodoxin on  $\text{HgCl}_2$ -treated complexes is due to the loss of the  $F_B$  cluster and not to damage to the ferredoxin/flavodoxin docking site, we reconstructed the  $F_B$  cluster in the  $\text{Hg}$ -treated PS I complexes and repeated the  $\Delta A_{832}$  kinetic measurements. As shown in Fig. 1 *C*, the normal flash number dependency is almost completely restored, and the addition of either ferredoxin (Fig. 1, *F* and *D*) or flavodoxin (Fig. 1, *L*) leads to the complete loss of the flash number dependency.

Additional evidence for the lack of electron transfer to ferredoxin and flavodoxin in  $\text{HgCl}_2$ -treated PS I complexes is provided by single flash experiments in the presence of reduced DCPIP, a slow electron donor to  $\text{P700}^+$ . Unlike our previous study (Vassiliev et al., 1997), we used aerobic rather than anaerobic conditions so as to provide oxygen as an electron trap when using methyl viologen as the immediate electron acceptor. Under aerobic conditions the terminal iron-sulfur cluster of PS I is reoxidized by oxygen present in solution, and the contribution of the component arising because of direct reduction of  $\text{P700}^+$  from DCPIP (4- to 10-s lifetime) is 35 to 45% (Fig. 2), which is about two times higher than under anaerobic conditions (Vassiliev et al., 1997). Addition of 30–100  $\mu\text{M}$  ferredoxin leads to a significant increase (up to 75%) of the contribution of the DCPIP-mediated component in the control PS I complex (Figs. 2 *A* and 3 *A*). In the  $\text{HgCl}_2$ -treated PS I complex, most of the back-reaction is derived from  $F_A^-$  with life times of 17 and 91 ms (47 and 27% amplitude, respectively) and with a negligible contribution of the slower DCPIP-component that can be resolved in most of these experiments only as a baseline (5 to 8% amplitude). The remainder of the  $\Delta A_{832}$  decay is brought about by components with lifetimes of 61  $\mu\text{s}$  (11%) and 1.9 ms (10%), which arise because of back-reactions of  $A_1^-$  and  $F_X^-$ . The contribution of the latter two components is lower than we found in the  $\text{HgCl}_2$ -treated sample in our previous work (Vassiliev et al., 1997) and indicates an even greater retention of  $F_A$  in these samples. Unlike the control PS I complex, addition of ferredoxin up to 430  $\mu\text{M}$  has no effect on the kinetics of the  $\text{HgCl}_2$ -treated PS I complex (Figs. 2 *B* and 3 *B*).

Methyl viologen is an efficient electron transport mediator between the acceptor side of PS I and molecular oxygen (Hiyama and Ke, 1971). As the photo-reduced terminal acceptor of PS I cannot participate in the recombination reaction because of its efficient oxidation by methyl viologen (Figs. 2 *A* and 3 *B*), greater than 90% of  $\text{P700}^+$  is reduced by DCPIP in the presence of 50  $\mu\text{M}$  methyl viologen. This concentration of methyl viologen has no effect on  $\text{HgCl}_2$ -treated complex, but an additional increase of its concentration to the millimolar range results in more than 80% of  $\text{P700}^+$  reduction occurring from DCPIP (Fig. 3 *B*; also see Fujii et al., 1990). Hence, unlike ferredoxin or flavodoxin, methyl viologen has the ability to accept electrons from  $F_A$ , but only at a concentration nearly two orders of magnitude greater than is required in control PS I complexes.




FIGURE 2 Kinetics of  $\Delta A_{832}$  decay in control (*A*) and  $\text{HgCl}_2$ -treated (*B*) PS I complexes in 25 mM Tris buffer, pH 8.3, with 0.02% Triton X-100, 10 mM sodium ascorbate, and 4  $\mu\text{M}$  DCPIP in the absence and in the presence of ferredoxin or methyl viologen upon laser flash excitation, average of 12 traces acquired at 150-s intervals. Multiexponential fits are shown as solid lines.

## DISCUSSION

Even though  $F_A$  and  $F_B$  were among the first bound electron transfer cofactors discovered in PS I, the issue of whether electron transfer proceeds from  $F_X \rightarrow F_A \rightarrow F_B$  or from  $F_X \rightarrow F_B \rightarrow F_A$  remained unclear. The reason for this uncertainty lies mostly in the identical optical signatures of  $F_A$  and  $F_B$ , which disallows the use of time-resolved optical spectroscopy to distinguish one acceptor from another at room temperature. Consequently, most of the functional data on the photoreduction of the iron-sulfur clusters has been provided by EPR measurements performed at cryogenic temperatures. However, even though  $F_A$  and  $F_B$  have distinguishable  $g$ -tensors, the time resolution of EPR is insufficient to perform the requisite kinetic measurements.

In attempting to resolve this issue, we examine our kinetic data in the context of previous work and in light of the two possible orientations of  $\text{PsaC}$ .

A possible orientation of  $\text{PsaC}$  would be that first  $F_B$  is proximal to  $F_X$ . In the absence of  $F_B$  the electron transfer between  $F_X$  and  $F_A$  would need to span a center-to-center distance of 22 Å. The distance between  $F_A$  and the iron-sulfur cluster on ferredoxin will likely be unchanged. Given




FIGURE 3 The dependency of the amplitude of the slow (4 to 10 s) component arising caused by direct reduction of  $P700^+$  from DCPIP on the concentration of ferredoxin (A) and methyl viologen (B) in control (circles) and  $HgCl_2$ -treated (squares) PS I complexes. The basic media consisted of 25 mM Tris buffer, pH 8.3, with 0.02% Triton X-100, 10 mM sodium ascorbate, and 4  $\mu$ M DCPIP. The amplitudes are normalized to the initial amplitude of the absorbance change derived from the multiexponential fit as shown in the inset.

that the photoreduction of  $F_A$  is highly efficient at both cryogenic (Jung et al., 1995) and room temperatures (this study), it is difficult in this model to rationalize the failure of ferredoxin and flavodoxin to become reduced. Although it might be argued that the binding site is destroyed by the  $Hg$ -treatment, it would need to be a reversible loss as steady-state rates of  $NADP^+$  photoreduction (Jung et al., 1995) and flash-induced ferredoxin and flavodoxin reduction (this study) are restored when  $F_B$  is rebuilt with  $\beta$ -mercaptoethanol, inorganic iron, and sulfide. Additionally, the binding site for ferredoxin and flavodoxin is likely to involve PsaD (and possibly PsaE) rather than PsaC (Chitnis et al., 1995). In principle, conformational changes in PsaC induced by the loss of  $F_B$  could be transmitted to PsaD (and perhaps PsaE). However, the absence of PsaD leads to large changes in the EPR spectrum of  $F_A$  and  $F_B$ , implying that the  $g$ -tensor is sensitive to protein conformation (Li et al., 1991; Chitnis et al., 1996). Therefore, conformational changes in PsaC should have resulted in significant changes in the EPR spectrum of  $F_A$ . Yet, with the exception of a slight upfield shift of the  $g_x$  resonance, the EPR spectrum of

$F_A$  is identical to that of the control (Jung et al., 1995). Any changes in the binding site for ferredoxin and flavodoxin also should be irrelevant to the rates of reduction of methyl viologen. Given that this is likely to be a diffusion-mediated process that does not require a docking site, it is difficult to rationalize the need for a two-order of magnitude increase in the concentration required for electron acceptance from  $F_A^-$ . This orientation of PsaC is also incompatible with distance versus rate considerations. Although no direct data on the rate of  $F_A$  photoreduction exist, analysis of various spectroscopic and electrometric data yields lifetime values of forward electron transfer from  $F_X^-$  ranging between 50 and 800 ns (Brettel, 1997). Assuming  $\lambda$  (a coefficient that depends on the intervening medium in propagating the wave function) of  $1.4 \text{ \AA}^{-1}$  for electron transfer in proteins and given an electron transfer rate constant of  $10^{13}$  at van der Waals contact (Moser et al., 1992), a 7- $\text{\AA}$  increase in distance leads to a prediction of an about 18,000-fold increase in the electron transfer time up to a value ranging from 900  $\mu$ s to 14 ms (Fig. 4 A for illustration). The  $P700^+$   $F_X^-$  recombination kinetics in the  $HgCl_2$ -treated PS I complex (with 100 mM dithionite) is mainly composed of two components with lifetimes of 270  $\mu$ s and 842  $\mu$ s at approximately 1:1 ratio (not shown). Such kinetics are consistent with those found in integral PS I complexes with pre-reduced  $F_A$  and  $F_B$  and in PsaC-devoid core preparations (Vassiliev et al., 1997). Therefore, if the  $F_B$  cluster is located between  $F_X$  and  $F_A$ , the large increase of the forward electron transfer rate from  $F_X^-$  should have led to a measurable increase in the contribution of the  $F_X^-$  back-reaction.

A second possible orientation of PsaC would be that  $F_A$  is proximal to  $F_X$ . EPR studies in urea-ferricyanide-treated




FIGURE 4 Schematic representation of possible electron transport pathways from  $F_X$  to exogenously added ferredoxin (Fd) and methyl viologen (MV) for two possible orientations of PsaC assuming that the cluster proximal to  $F_X$  is  $F_B$  (A) or  $F_A$  (B). The structural diagrams of PsaC are reprinted from Kamlowski, A., A. Van der Est, P. Fromme, N. Krauss, W. D. Schubert, O. Klukas, and D. Stehlik. 1997. The structural organization of the PsaC protein in Photosystem I from single crystal EPR and x-ray crystallographic studies, *Biochim. Biophys. Acta* 1319:185–198, with kind permission of Elsevier Science—NL, Sara Burgerhastraat 25, 1055, KV Amsterdam, The Netherlands.

PS I complexes that preferentially retain  $F_A$  rather than  $F_B$  (Golbeck and Warden, 1982), kinetic measurements in Hg-treated PS I complexes, which totally lack  $F_B$  (Sakurai et al., 1991), and the restoration of NADP<sup>+</sup> reduction after reconstitution of  $F_B$  in Hg-treated PS I complexes (Jung et al., 1995) provide the strongest arguments for  $F_A$  as the  $F_X$  proximal cluster. The one lingering issue is that a lower quantum efficiency of electron transfer caused by the irreversible loss of the ferredoxin and flavodoxin docking site cannot be excluded as an alternative interpretation of the results of He and Malkin (1994) and Jung et al. (1995). In this study, we eliminated this contingency by showing in a  $F_B$ -less PS I complex: 1) a normal level of photochemical activity of P700, based on the amplitudes of the photoinduced absorbance change at 832 nm; 2) an unimpaired functioning of  $F_A$  at room temperature based on a high yield of charge recombination on the tens-of-ms time scale between  $P700^+$  and  $F_A^-$ ; 3) lack of direct electron transfer from  $F_A^-$  to either ferredoxin or flavodoxin; and 4) a decreased efficiency of methyl viologen reduction in the absence of  $F_B$ , which implies a diffusion barrier but a normally functioning  $F_A$  cluster. In the absence of  $F_B$ , the electron transfer between  $F_X$  and  $F_A$  would remain at a center-to-center distance of 15 Å. This is compatible with the efficient reduction of  $F_A$  at both cryogenic (Jung et al., 1995) and room temperatures (this study). The distance between  $F_A$  and ferredoxin is difficult to judge given that the three-dimensional structure of Psac as well as the binding site and orientation of ferredoxin on the PS I complex are only known to approximation. Assuming for the purpose of argument that the distance vector between  $F_A$  and the [2Fe-2S] cluster in ferredoxin passes through  $F_B$ , an additional distance of 12 Å would need to be spanned without participation of a cofactor if  $F_B$  is absent (Fig. 4 B). Reduction of both ferredoxin and flavodoxin involves complex formation preceding electron transfer, which follows multicomponent kinetics (Hervas et al., 1992; Medina et al., 1992). Three phases with halftimes of 500 ns, 20 μs, and 100 μs have been attributed to reduction of ferredoxin on a single flash (Sétif and Bottin, 1994). The increased distance (without mediation by a redox-active cofactor) leads to the prediction of a  $2.1 \times 10^7$ -fold increase in the electron transfer time to ~10 s for the fastest phase of ferredoxin reduction. This is about two orders of magnitude slower than the  $P700^+$   $F_A^-$  back-reaction and would result in negligible quantum yield of ferredoxin reduction, which agrees with the experiment. The recovery of photoreduction of ferredoxin and flavodoxin on a single turnover flash (this study), as well as the recovery of steady-state NADP<sup>+</sup> photoreduction mediated by ferredoxin and flavodoxin (Jung et al., 1995), correlates with the restoration of a functional  $F_B$  cluster as an essential intermediate electron carrier. A small molecule such as methyl viologen with a mass of 186 is capable of accepting the electron from  $F_A^-$  but at a concentration of two orders of magnitude greater than is required in the control. We attribute this to steric hindrance in which a higher concentra-

tion of methyl viologen is required to overcome the diffusion barrier to the buried  $F_A$  cluster.

The finding that  $F_X$  fails to donate electrons to ferredoxin and flavodoxin when  $F_B$  is missing and that it succeeds in donating to ferredoxin and flavodoxin when  $F_B$  is restored is best compatible with an orientation of Psac with  $F_A$  as the  $F_X$  proximal iron-sulfur cluster. The implied electron transfer sequence is therefore  $F_X \rightarrow F_A \rightarrow F_B \rightarrow$  ferredoxin. Note that this orientation of Psac involves an electron transfer from  $F_A$  ( $E_m$ , -540 mV) to a more electronegative acceptor,  $F_B$  ( $E_m$ , -590 mV), which results in the presence of a positive Gibbs free energy step in PS I (given that the known midpoint potential values for the [4Fe-4S] clusters determined for PS I at cryogenic temperatures apply at room temperature and that reduced  $F_A^-$  does not influence the determination of midpoint potential of  $F_B$ ). It is noteworthy that ferredoxin reduction is not affected in the K52S/R(53)A mutant of Psac in which the preferential photoreduction of  $F_B$  is attributed to a more negative redox potential of  $F_A$  (Fischer et al., 1997). Recent studies indicate that small positive Gibbs free energy changes may be common in multicofactor enzymes; examples are found in two segments of mitochondrial respiratory chain (Ohnishi and Salerno, 1982), [NiFe] hydrogenase (Fontecilla-Camps, 1996), and the tetraheme cytochromes of bacterial reaction centers (Nitschke et al., 1993). The relevance, if any, of this uphill electron transfer step to PS I function is unknown. The important issue for an efficient electron transfer from the primary donor of PS I to ferredoxin is a net negative change in Gibbs free energy from  $A_1$  to ferredoxin, which still occurs in the  $A_1 \rightarrow F_X \rightarrow F_A \rightarrow F_B \rightarrow$  ferredoxin sequence.

We thank Art van der Est, Petra Fromme, Andreas Kamlowski, Norbert Krauss, Wolf-Dieter Schubert, and Dietmar Stehlik for helpful comments on the work. This work was funded by National Science Foundation Grants MCB-9696179 and MCB-972366.

## REFERENCES

- Adman, E. T., L. C. Sieker, and L. H. Jensen. 1976. The structure of a bacterial ferredoxin. *J. Biol. Chem.* 251:3801–3806.
- Bearden, A. J., and R. Malkin. 1976. Correlation of reaction-center chlorophyll (P-700) oxidation and bound iron-sulfur protein photoreduction in chloroplast photosystem I at low temperatures. *Biochim. Biophys. Acta.* 430:538–547.
- Biggins, J., S. Rodday, and L. Do. 1995. Interaction of the subunit Psac with its binding site on the PSI core heterodimer. In *Photosynthesis: from Light to Biosphere*, Vol. 2. P. Mathis, editor. Kluwer Academic Publishers, Dordrecht. 111–114.
- Bottin, H., and B. Lagoutte. 1992. Ferredoxin and flavodoxin from the cyanobacterium *Synechocystis* sp PCC 6803. *Biochim. Biophys. Acta.* 1101:48–56.
- Brettel, K. 1997. Electron transfer and arrangement of the redox cofactors in photosystem I. *Biochim. Biophys. Acta.* 1318:322–373
- Cammack, R., M. D. Ryan, and A. C. Stewart. 1979. The EPR spectrum of iron-sulfur center B in photosystem I of *Phormidium laminosum*. *FEBS Lett.* 107:422–426.
- Chitnis, P. R., Q. Xu, V. P. Chitnis, and R. Nechushtai. 1995. Function and organization of photosystem I polypeptides. *Photosynth. Res.* 44:23–40.
- Chitnis, V. P., Y. S. Jung, L. Albee, J. H. Golbeck, and P. R. Chitnis. 1996. Mutational analysis of photosystem I polypeptides: role of Psad and the



- lysyl 106 residue in the reductase activity of photosystem I. *J. Biol. Chem.* 271:11772–11780.
- Duee, E. D., E. Fanchon, J. Vicat, L. C. Sieker, J. Meyer, and J. M. Moulis. 1994. Refined crystal structure of the 2[4Fe-4S] ferredoxin from *Clostridium acid urici* at 1.84 Ångstrom resolution. *J. Mol. Biol.* 243: 683–695.
- Dunn, P. P. J., and J. C. Gray. 1988. Localization and nucleotide sequence of the gene for the 8 kDa subunit of photosystem I in pea and wheat chloroplast DNA. *Plant. Mol. Biol.* 11:311–319.
- Fischer, N., P. Sétif, and J. D. Rochaix. 1997. Targeted mutations in the *psaC* gene of *Chlamydomonas reinhardtii*: preferential reduction of  $F_B$  at low temperature is not accompanied by altered electron flow from photosystem I ferredoxin. *Biochemistry* 36:93–102.
- Fontecilla-Camps, J. C. 1996. The active site of Ni-Fe hydrogenases: model chemistry and crystallographic results. *J. Biol. Inorg. Chem.* 1:91–98.
- Fujii, T., E. Yokoyama, K. Inoue, and H. Sakurai. 1990. The sites of electron donation of photosystem I to methyl viologen. *Biochim. Biophys. Acta.* 1015:41–48.
- Golbeck, J. H. 1995. Resolution and reconstitution of photosystem I. In *CRC Handbook of Organic Photochemistry and Photobiology*. P. S. Song and W. M. Horspool, editors. CRC Press, Boca Raton. 1407–1419.
- Golbeck, J. H., and J. T. Warden 1982. Electron spin resonance studies of the bound iron-sulfur centers in photosystem I: photoreduction of center A occurs in the absence of center B. *Biochim. Biophys. Acta.* 681:77–84.
- Guigliarelli, B., J. Guillaussier, C. More, P. Sétif, H. Bottin, and P. Bertrand. 1993. Structural organization of the iron-sulfur centers in *Synechocystis* 6803 photosystem-I: EPR study of oriented thylakoid membranes and analysis of the magnetic interactions. *J. Biol. Chem.* 268:900–908.
- Hayashida, N., T. Matsubayashi, K. Shinozaki, M. Sugiura, K. Inoue, and T. Hiyama. 1987. The gene for the 9 kD polypeptide, a possible apoprotein for the iron-sulfur centers A and B of the photosystem I complex, in tobacco chloroplast DNA. *Curr. Genet.* 12:247–250.
- He, W. Z., and R. Malkin. 1994. Reconstitution of iron-sulfur center B of photosystem I damaged by mercuric chloride. *Photosynth. Res.* 41: 381–388.
- Heathcote, P., D. L. Williams-Smith, C. K. Sihra, and M. C. W. Evans. 1978. The role of the membrane-bound iron-sulfur centers A and B in the photosystem I reaction centre of spinach chloroplasts. *Biochim. Biophys. Acta.* 503:333–42.
- Hervas, M., J. Navarro, and G. Tollin. 1992. A laser flash spectroscopy study of the kinetics of electron transfer from spinach photosystem-I to spinach and algal ferredoxins. *Photochem. Photobiol.* 56:319–324.
- Hiyama, T., and B. Ke. 1971. P430: possible primary electron acceptor of photosystem I. *Arch. Biochem. Biophys.* 147:99–108.
- Jung, Y. S., I. R. Vassiliev, J. Yu, L. McIntosh, and J. H. Golbeck. 1997. Strains of *Synechocystis* sp. PCC. 6803 with altered PsaC .2. EPR and optical spectroscopic properties of  $F_A$  and  $F_B$  in aspartate, serine, and alanine replacements of cysteines 14 and 51. *J. Biol. Chem.* 272: 8040–8049.
- Jung, Y. S., L. Yu, and J. H. Golbeck. 1995. Reconstitution of iron-sulfur center  $F_A$  results in complete restoration of NADP(+) photoreduction in Hg-treated photosystem I complexes from *Synechococcus* sp PCC 6301. *Photosynth. Res.* 46:249–255.
- Kamlowski, A., A. Van der Est, P. Fromme, and D. Stehlik. 1997a. Low temperature EPR on photosystem I single crystals: orientation of the iron-sulfur centers  $F_A$  and  $F_B$ . *Biochim. Biophys. Acta.* 1319:185–198.
- Kamlowski, A., A. Van der Est, P. Fromme, N. Krauss, W. D. Schubert, O. Klukas, and D. Stehlik. 1997b. The structural organization of the PsaC protein in photosystem I from single crystal EPR and x-ray crystallographic studies. *Biochim. Biophys. Acta.* 1319:199–213.
- Krauss, N., W. Hinrichs, I. Witt, P. Fromme, W. Pritzkow, Z. Dauter, C. Betzel, K. S. Wilson, H. T. Witt, and W. Saenger. 1993. 3-Dimensional structure of System-I of photosynthesis at 6 Ångstrom resolution. *Nature.* 61:326–331.
- Krauss, N., W. D. Schubert, O. Klukas, P. Fromme, H. T. Witt, and W. Saenger. 1996. Photosystem I at 4 Ångstrom resolution represents the first structural model of a joint photosynthetic reaction centre and core antenna system. *Nat. Struct. Biol.* 3:965–973.
- Li, N., J. Zhao, P. Warren, J. Warden, D. Bryant, and J. Golbeck. 1991. PsaD is required for the stable binding of PsaC to the photosystem-I core protein of *Synechococcus* sp PCC 6301. *Biochemistry.* 30:7863–7872.
- Malkin, R. 1984. Diazonium modification of photosystem I: a specific effect on iron-sulfur center B. *Biochim. Biophys. Acta.* 764:63–69.
- Mannan, R. M., W. Z. He, S. U. Metzger, J. Whitmarsh, R. Malkin, and H. B. Pakrasi. 1996. Active photosynthesis in cyanobacterial mutants with directed modifications in the ligands for two iron-sulfur clusters in the PsaC protein of photosystem I. *EMBO J.* 15:1826–1833.
- Medina, M., M. Hervas, J. A. Navarro, M. A. De la Rosa, C. Gomez-Moreno, and G. Tollin. 1992. A laser flash absorption spectroscopy study of *Anabaena* sp PCC 7119 flavodoxin photoreduction by photosystem I particles from spinach. *FEBS Lett.* 313:239–242.
- Moser, C. C., J. M. Keske, K. Warncke, R. S. Farid, and P. L. Dutton. 1992. Nature of biological electron transfer. *Nature.* 355:796–802.
- Mühlenhoff, U., J. D. Zhao, and D. A. Bryant. 1996. Interaction between photosystem I and flavodoxin from the cyanobacterium *Synechococcus* sp PCC 7002 as revealed by chemical cross-linking. *Eur. J. Biochem.* 235:324–331.
- Naver, H., M. P. Scott, J. H. Golbeck, B. L. Møller, and H. V. Scheller. 1996. Reconstitution of barley photosystem I with modified PSI-C allows identification of domains interacting with PSI-D and PSI-A/B. *J. Biol. Chem.* 271:8996–9001.
- Nitschke, W., M. Jubaultbregler, and A. W. Rutherford. 1993. The reaction center associated tetraheme cytochrome subunit from *Chromatium vinosum* revisited: a reexamination of its EPR properties. *Biochemistry.* 32:8871–8879.
- Nugent, J. H. A., B. L. Møller, and M. C. W. Evans. 1981. Comparison of the EPR properties of photosystem I iron-sulfur centers A and B in spinach and barley. *Biochim. Biophys. Acta.* 634:249–255.
- Ohnishi, T., and J. C. Salerno. 1982. Iron-sulfur clusters in the mitochondrial electron-transport chain. In *Iron-Sulfur Proteins*, Vol. 4. T. G. Spiro, editor. Wiley Publishing Co., New York. 285–327
- Oh-oka, H., Y. Takahashi, K. Kuriyama, K. Saeki, and H. Matsubara. 1988. The protein responsible for center A/B in spinach photosystem I: isolation with iron-sulfur cluster(s) and complete sequence analysis. *J. Biochem. (Tokyo).* 103:962–968.
- Rodday, S. M., L. T. Do, V. Chynwat, H. A. Frank, and J. Biggins. 1996. Site-directed mutagenesis of the subunit PsaC establishes a surface-exposed domain interacting with the photosystem I core binding site. *Biochemistry.* 35:11832–11838.
- Sakurai, H., K. Inoue, T. Fujii, and P. Mathis. 1991. Effects of selective destruction of iron-sulfur center B on electron transfer and charge recombination in photosystem I. *Photosynth. Res.* 27:65–71.
- Sauer, K., P. Mathis, S. Acker, and J. A. Van Best. 1978. Electron acceptors associated with P-700 in Triton solubilized photosystem I particles from spinach chloroplasts. *Biochim. Biophys. Acta.* 503:120–134.
- Sétif, P. Q. Y., and H. Bottin. 1994. Laser flash absorption spectroscopy study of ferredoxin reduction by photosystem I in *Synechocystis* sp. PCC 6803: evidence for submicrosecond and microsecond kinetics. *Biochemistry.* 33:8495–8504.
- Vassiliev, I. R., Y. S. Jung, M. D. Mamedov, A. Y. Semenov, and J. H. Golbeck. 1997. Near-IR absorbance changes and electrogenic reactions in the microsecond-to-second time domain in photosystem I. *Biophys. J.* 72:301–315.
- Yu, J., I. R. Vassiliev, Y. S. Jung, J. H. Golbeck, and L. McIntosh. 1997. Strains of *Synechocystis* sp. PCC 6803 with altered PsaC .1: mutations incorporated in the cysteine ligands of the two [4Fe-4S] clusters  $F_A$  and  $F_B$  of photosystem I. *J. Biol. Chem.* 272:8032–8039.
- Zhao, J., N. Li, P. Warren, J. H. Golbeck, and D. A. Bryant. 1992. Site-directed conversion of a cysteine to aspartate leads to the assembly of a [3Fe-4S] cluster in PsaC of photosystem-I: the photoreduction of  $F_A$  is independent of  $F_B$ . *Biochemistry.* 31:5093–5099.