

# ATP-Independent Luminal Oscillations and Release of $\text{Ca}^{2+}$ and $\text{H}^+$ from Mast Cell Secretory Granules: Implications for Signal Transduction

Ivan Quesada,\* Wei-Chun Chin,<sup>†</sup> and Pedro Verdugo\*

\*Department of Bioengineering and Friday Harbor Laboratories, University of Washington, Seattle, Washington 98195; and <sup>†</sup>Department of Chemical Engineering, FAMU/FSU, Tallahassee, Florida 32310-6046

**ABSTRACT**  $\text{InsP}_3$  is an important link in the intracellular information network. Previous observations show that activation of  $\text{InsP}_3$ -receptor channels on the granular membrane can turn secretory granules into  $\text{Ca}^{2+}$  oscillators that deliver periodic trains of  $\text{Ca}^{2+}$  release to the cytosol (T. Nguyen, W. C. Chin, and P. Verdugo, 1998, *Nature*, 395:908–912; I. Quesada, W. C. Chin, J. Steed, P. Campos-Bedolla, and P. Verdugo, 2001, *Biophys. J.* 80:2133–2139). Here we show that  $\text{InsP}_3$  can also turn mast cell granules into proton oscillators.  $\text{InsP}_3$ -induced intraluminal  $[\text{H}^+]$  oscillations are ATP-independent, result from  $\text{H}^+/\text{K}^+$  exchange in the heparin matrix, and produce perigranular pH oscillations with the same frequency. These perigranular pH oscillations are in-phase with intraluminal  $[\text{H}^+]$  but out-of-phase with the corresponding perigranular  $[\text{Ca}^{2+}]$  oscillations. The low pH of the secretory compartment has critical implications in a broad range of intracellular processes. However, the association of proton release with  $\text{InsP}_3$ -induced  $\text{Ca}^{2+}$  signals, their similar periodic nature, and the sensitivity of important exocytic proteins to the joint action of  $\text{Ca}^{2+}$  and pH strongly suggests that granules might encode a combined  $\text{Ca}^{2+}/\text{H}^+$  intracellular signal. A  $\text{H}^+/\text{Ca}^{2+}$  signal could significantly increase the specificity of the information sent by the granule by transmitting two frequency encoded messages targeted exclusively to proteins like calmodulin, annexins, or syncollin that are crucial for exocytosis and require specific combinations of  $[\text{Ca}^{2+}]$  “and” pH for their action.

## INTRODUCTION

The dynamics of luminal pH in the secretory pathway is critical for the proper function of a broad range of cell processes including protein sorting, enzyme activation, and biogenic amines loading (Bell-Parikh et al., 2001). In secretory granules, changes in the preexocytotic luminal pH are thought to be necessary for the final steps of exocytosis (Williams and Webb, 2000; Barg et al., 2001; Han et al., 1999). The regulation of intraluminal pH ( $\text{pH}_G$ ) has been thought as a simple  $\text{H}^+$  influx/efflux equilibrium with pumps as  $\text{H}^+$  source and “leakage” to the cytosol as a  $\text{H}^+$  sink (Mitchell et al., 2001; Demaurex, 2002). However, the characteristic polyelectrolyte matrices present inside virtually all granules offer a novel alternative as intraluminal  $\text{H}^+$  sink/donors. The strong polyanionic properties of these polymer networks can function as efficient ion exchange resins controlling the bound/free turnover of intraluminal cations (Uvnas and Aborg, 1977, 1989; Verdugo, 1994; Nguyen et al., 1998; Quesada et al., 2001; Nanavati and Fernandez, 1993; Marszalek et al., 1997; Chin et al., 2002). Thus, the level of free ionized  $\text{Ca}^{2+}$ ,  $\text{K}^+$ , and  $\text{H}^+$  inside the granule results not only from the influx/efflux equilibrium of these ions in/from the granule but also from their complex

exchange with the secretory matrix. However, most of the work on the control of intraluminal cations, particularly  $\text{Ca}^{2+}$ , remains focused almost exclusively on the action of pumps and export channels (Mitchell et al., 2001; Demaurex, 2002). The key role of the secretory matrix as cation sink/donor in the granule has been highlighted by recent observations that reveal that the granule can function as an intracellular  $\text{Ca}^{2+}$  oscillator, and that  $\text{InsP}_3$ -induced intraluminal  $\text{Ca}^{2+}$  oscillations—and corresponding oscillatory release of  $\text{Ca}^{2+}$  to the cytosol—results from  $\text{Ca}^{2+}/\text{K}^+$  exchange in the matrix (Nguyen et al., 1998; Quesada et al., 2001). According to this new model, the frequency-encoded  $\text{Ca}^{2+}$  signaling system of the granule results from the interplay between the  $\text{Ca}^{2+}/\text{K}^+$  ion-exchange properties of the secretory matrix and two  $\text{Ca}^{2+}$ -sensitive channels located at close proximity on the membrane of secretory vesicles: an  $\text{ASK}_{\text{Ca}}$  channel that mediates  $\text{K}^+$  entry into the vesicular lumen, and an  $\text{InsP}_3$ -R channel that releases  $\text{Ca}^{2+}$  to the cytosol (see Fig. 1). Stimulation of the cell induces production of  $\text{InsP}_3$  leading to  $\text{InsP}_3$  binding to the  $\text{InsP}_3$ -R channel, release of  $\text{Ca}^{2+}$  from the granule, and decrease of intraluminal  $[\text{Ca}^{2+}]_{\text{IL}}$ . Cytosolic  $\text{Ca}^{2+}$  ( $[\text{Ca}^{2+}]_{\text{C}}$ ) increases around the granule, turning open nearby  $\text{ASK}_{\text{Ca}}$  channels and closing  $\text{InsP}_3$ -R channels.  $\text{K}^+$  imported into the vesicular lumen exchange for  $\text{Ca}^{2+}$  in the polyanionic matrix and together with the closure of the  $\text{InsP}_3$ -R channel results in an increase of  $[\text{Ca}^{2+}]_{\text{IL}}$  in the granule’s lumen. As diffusion and cytosolic  $\text{Ca}^{2+}$ -buffering restore the  $[\text{Ca}^{2+}]_{\text{C}}$  to lower levels, the  $\text{InsP}_3$ -R channel opens again, starting a new cycle that recurs for as long as the  $\text{InsP}_3$ -R remains activated (Nguyen et al., 1998; Quesada et al., 2001). In goblet cells, these  $[\text{Ca}^{2+}]_{\text{IL}}$  oscillations are accompanied by corresponding  $\text{pH}_G$  oscillations that can increase the gain of

Submitted March 15, 2003, and accepted for publication May 5, 2003.

Address reprint requests to Prof. P. Verdugo, Friday Harbor Laboratories, University of Washington, 620 University Rd., Friday Harbor, WA 98250. E-mail: verdugo@u.washington.edu.

**Abbreviations used:**  $\text{InsP}_3$ , Inositol-1,4,5-trisphosphate;  $\text{InsP}_3$ -R channel,  $\text{InsP}_3$ -receptor channel;  $\text{ASK}_{\text{Ca}}$  channel, apamin-sensitive  $\text{Ca}^{2+}$ -sensitive  $\text{K}^+$  channel;  $[\text{Ca}^{2+}]_{\text{IL}}$ , intraluminal  $\text{Ca}^{2+}$  concentration;  $[\text{Ca}^{2+}]_{\text{EL}}$ , extraluminal  $\text{Ca}^{2+}$  concentration;  $[\text{H}^+]_{\text{IL}}$ , intraluminal  $\text{H}^+$  concentration;  $[\text{H}^+]_{\text{EL}}$ , extraluminal  $\text{H}^+$  concentration.

© 2003 by the Biophysical Society

0006-3495/03/08/963/08 \$2.00




FIGURE 1 Dynamics of  $H^+$  and  $Ca^{2+}$  in secretory granules. This model involves two pools of intraluminal cations: a pool of free ionized cations in equilibrium with a pool of cations bound to the granule polyanionic matrix that functions as a  $H^+/K^+$  and  $Ca^{2+}/K^+$  ion exchange network (Verdugo, 1994; Marszalek et al., 1997). The model further assumes that vesicular  $Ca^{2+}$  uptake is driven by an undefined  $Ca^{2+}$ -ATPase (Mitchell et al., 2001), and that the activity of V-type  $H^+$ -ATPases is responsible for  $H^+$  transport to maintain a steady-state intraluminal pH (Demaurex, 2002). Oscillations of intraluminal  $H^+$  and  $Ca^{2+}$  result from the interaction of the granule polyanionic matrix and two  $Ca^{2+}$ -sensitive ion channels located in close proximity on the granular membrane: an  $ASK_{Ca}$  channel to mediate  $K^+$  flux into the granule, and an  $InsP_3$ -R channel to release  $Ca^{2+}$  to the cytosol (Nguyen et al., 1998; Quesada et al., 2001). See text for further details.

the  $Ca^{2+}/K^+$  exchange leading to increased  $Ca^{2+}$  unbinding and a rise in the flux of diffusion-driven  $Ca^{2+}$  release (Chin et al., 2002).

Although the profusion of intracellular  $Ca^{2+}$ -sensitive proteins explains the broad capacity of this cation to relay changes of functionality to intracellular sensor/effector molecules, it is insufficient to explain the specificity of the  $Ca^{2+}$  message. Here we show that  $InsP_3$  can turn the secretory vesicles of mast cells into a double ionic oscillator that broadcasts both  $Ca^{2+}$  and  $H^+$  signals, thereby constraining the granule's message exclusively to sensor/effector proteins that are sensitive to both  $Ca^{2+}$  and pH.

## MATERIALS AND METHODS

### Mast cell granule isolation and dye loading

Motion artifacts can be a critical problem when performing thin optical sections of secretory granules in intact cells. The advantages of the isolated mast cell granule preparation we used in these experiments are that, because of their large size, granules can be easily resolved by optical microscopy (Quesada et al., 2001; Monck et al., 1992), and they can be securely immobilized on poly-L-lysine coated glass. In these experiments, mast cells of beige mice ( $Bg^1/Bg^1$ ) (Jackson Laboratory, Bar Harbor, ME) were isolated by peritoneal lavage (Quesada et al., 2001; Marszalek et al., 1997). Granules were labeled as previously described (Nguyen et al., 1998; Quesada et al., 2001). Briefly, cells were washed twice in a  $Ca^{2+}$ -free Hanks' solution (pH = 7.2) and loaded for 30 min at 37°C with either 2  $\mu$ M of LysoSensor Green DND-189 (LS) ( $pK_a = 5.2$ ;  $\lambda_{exc} = 443$  nm,  $\lambda_{em} = 505$  nm) to monitor  $pH_G$  changes, or with 5  $\mu$ M of Calcium Orange-5N (CO-5N) ( $K_d = 20$   $\mu$ M;  $\lambda_{exc} = 545$  nm,  $\lambda_{em} = 580$  nm) (Molecular Probes, Eugene, OR) for 45 min at 37°C, to monitor  $[Ca^{2+}]_{IL}$  (see Fig. 2). To remove any excess dye these two pools of cells were then washed and resuspended in an intracellular buffer solution containing 140 mM  $K^+$  glutamate, 20 mM HEPES, 5 mM  $MgSO_4$ ,

2 mM ATP, and 100 nM  $Ca^{2+}$  buffered with ethylene glycol bis( $\beta$ -aminoethyl ether)- $N,N,N',N'$ -tetraacetic acid (EGTA), pH = 7.2. Secretory granules were extracted by sonication and separated by centrifugation at 10,000 rpm for 5 min. To detect extraluminal  $Ca^{2+}$  (Nguyen et al., 1998; Quesada et al., 2001; Belan et al., 1996), the granules were resuspended in intracellular buffer containing 10  $\mu$ g  $ml^{-1}$  of low-diffusivity nonpermeant dextran-conjugated Calcium Green-1 ( $K_d = 190$  nM;  $\lambda_{exc} = 506$  nm,  $\lambda_{em} = 531$  nm) or Calcium Crimson ( $K_d = 185$  nM;  $\lambda_{exc} = 570$  nm,  $\lambda_{em} = 610$  nm). Changes of extraluminal pH were reported by 10  $\mu$ g  $ml^{-1}$  of dextran-conjugated SNARF-1 (SN) ( $pK_a = 7.5$ ;  $\lambda_{exc} = 488$  nm,  $\lambda_{em} = 587$  nm) (Molecular Probes) diluted in intracellular buffer. Granule suspensions were then allowed to attach to poly-L-lysine-coated glass chambers for 5 min. The chambers were mounted and kept at 37°C on the thermoregulated stage of a Nikon inverted fluorescence microscope. Notice that our set-up allows detection of only one emission at a time. We can monitor two ions simultaneously if their fluorescent probes have similar spectral characteristics but are localized in different compartments. Our results report simultaneous measurements of fluctuations of intra- and extraluminal  $Ca^{2+}$ , or intra- and extraluminal  $H^+$ , or intraluminal  $H^+$  and extraluminal  $Ca^{2+}$ . In all these cases we used probes that segregate in these two compartments.

### $H^+/K^+$ ion exchange

To investigate the  $H^+/K^+$  ion exchange properties of the vesicular matrix, granules loaded with LS were equilibrated in ATP-free intracellular buffer containing heparin (100  $\mu$ g  $ml^{-1}$ ) and apamin (100 nM). Under these conditions, resting  $[Ca^{2+}]_{IL}$  remains stable ( $\sim 25$   $\mu$ M), suggesting that the  $InsP_3$ -R and the  $ASK_{Ca}$  channel were rendered inoperative (Nguyen et al., 1998; Quesada et al., 2001). To titrate the intraluminal  $[K^+]$ , the granules were exposed to the  $K^+$  ionophore valinomycin (20  $\mu$ M) while  $[K^+]$  in the intracellular buffer was increased from 0 to 140 mM. Ionic strength and osmolarity were kept constant by adjusting the concentration of monovalent organic cation  $NMG^+$ .

### Calibration of extraluminal pH

The pH/photon-count transfer function for SN emission was obtained by measuring the fluorescence in thin optical sections of solutions of SN similar to those used in experiments but in which the pH buffered with MES, HEPES, or Tris (20 mM) was progressively increased from 6 to 6.8, 7.2, 7.6, 8, and 9, yielding a  $pK_a$  of 7.4.

Although the uncertainty of the quantum yield of LS in the intraluminal milieu prevented us from conducting absolute measurement of pH inside the granule, oscillations of intraluminal pH were readily reported by relative variations of LS photon count emission.

### Optical sectioning

Granules were imaged with a Nikon Diaphot inverted fluorescence microscope using a 100 W mercury vapor epifluorescence source and a 100 $\times$ , 1.4 NA oil-immersion objective. Images were formed on the 336  $\times$  243 charge-coupled-device array of a thermoelectrically cooled, low dark noise (1.3 photoelectrons  $s^{-1}$  at  $-36^\circ C$ ) frame transfer digital camera with 16-bit resolution and  $10^5$  pixel  $s^{-1}$  maximum readout rate (Spectra Source Model 400, Westlake Village, CA). The camera was attached to the photoport of the microscope using a 20 $\times$  relay lens, yielding a final resolution of 10 pixels  $\mu m^{-1}$ . To avoid aliasing, we acquired three-line scans at a time, instead of the whole image, yielding a sampling rate of 3 scans  $s^{-1}$  with 300-ms exposure time and  $\sim 25$  samples/period of  $[Ca^{2+}]_{IL}$  or  $pH_G$  oscillation. Scans sampled an area of 0.3  $\mu m \times 24 \mu m$  containing one or more granules and were accumulated in a memory buffer forming 50- to 60-s long sequential scan stacks (inset in Fig. 2 B). Optical sections of  $\sim 0.2 \mu m$  for  $Ca^{2+}$  changes and extraluminal pH measurements and  $\sim 2 \mu m$  for  $pH_G$  were performed using a no-neighbors deconvolution algorithm




FIGURE 2  $\text{InsP}_3$ -induced  $\text{pH}_G$  oscillations. (A) Phase contrast and fluorescent images of intact mast cells (top panel) and isolated granules (bottom panel) loaded with the pH- and  $\text{Ca}^{2+}$ -sensitive fluorescent probes LS (green) and CO-5N (orange), respectively. The large size of beige mast cell granules allows unequivocal intraluminal and extraluminal fluorescence measurements (Quesada et al., 2001). Scale bars: 5  $\mu\text{m}$ . (B) Application of 3  $\mu\text{M}$   $\text{InsP}_3$  induced  $\text{pH}_G$  oscillations of  $\sim 0.1$ – $0.12$  Hz ( $n = 8$ ). Notice that LS fluorescence increases with  $[\text{H}^+]$ , i.e., decreasing with pH. Inset displays a line-scan from a deconvoluted image of an isolated secretory granule (Nguyen et al., 1998; Quesada et al., 2001), showing intraluminal fluorescence changes resulting from  $\text{pH}_G$  oscillations after exposure to  $\text{InsP}_3$ . Scale bar: 3  $\mu\text{m}$ . (C)  $\text{InsP}_3$ -induced  $\text{pH}_G$  oscillations were observed in isolated granules exposed to 0.5–1  $\mu\text{M}$  bafilomycin ( $n = 5$ ) or in the absence of ATP in the medium (not shown). (D) Removal of ATP (filled circles;  $n = 3$ ) or application of 500 nM bafilomycin (open circles;  $n = 5$ ) caused pH alkalization in isolated granules.

(Nguyen et al., 1998; Quesada et al., 2001; Monck et al., 1992). Validation of the optical sectioning method has been published elsewhere (Nguyen et al., 1998; Quesada et al., 2001). The time course of average fluorescence intensity in photoelectron-counts per pixel  $\text{s}^{-1}$  inside and outside the secretory granules was measured from the line scans. Free  $[\text{Ca}^{2+}]$  was calculated from the readouts of the line scans following published methods (Nguyen et al., 1998; Quesada et al., 2001; Kao, 1994).

## RESULTS

### $\text{InsP}_3$ induces intraluminal pH oscillations in secretory granules


We performed experiments in isolated mast cell granules using pH-sensitive and  $\text{Ca}^{2+}$ -sensitive fluorescent probes combined with digital sectioning (Nguyen et al., 1998; Quesada et al., 2001; Monck et al., 1992). Optical sections of isolated and in situ mast cell secretory granules loaded with the fluorescent probes LS and CO-5N are shown in Fig. 2 A. Isolated mast cell granules exposed to intracellular buffer containing 3  $\mu\text{M}$   $\text{InsP}_3$  exhibit periodic oscillations of  $\text{pH}_G$  with a frequency of  $\sim 0.12$  Hz (Fig. 2 B).  $\text{InsP}_3$ -induced  $\text{pH}_G$  oscillations were blocked by exposure to intracellular buffer containing heparin (100  $\mu\text{g ml}^{-1}$ ) (a blocker of  $\text{InsP}_3$ -R channels) or apamin (100 nM) (a blocker of  $\text{ASK}_{\text{Ca}}$  channels) (Nguyen et al., 1998; Quesada et al., 2001), or by replacement of  $\text{K}^+$  by  $\text{NMG}^+$  (not shown), suggesting that activation of the  $\text{InsP}_3$ -R and inflow of  $\text{K}^+$  into the granule are required for  $\text{pH}_G$  oscillations. Notwithstanding the important role of  $\text{H}^+$  pumps in the control of  $\text{pH}_G$  (Demaurex, 2002), removal of ATP from the intracellular solution or exposure of the granules to the  $\text{H}^+$  V-ATPase inhibitor bafilomycin (0.5  $\mu\text{M}$ ), failed to abolish the  $\text{pH}_G$  oscillations, implying that they must result from a mechanism other than  $\text{H}^+$ -pump activity (Fig. 2 C). However, in granules not treated with  $\text{InsP}_3$ , the removal of ATP or

exposure of isolated granules to bafilomycin (0.5  $\mu\text{M}$ ) resulted in intraluminal alkalization (Fig. 2 D). This outcome is consistent with previous reports that secretory granules have a resting  $\text{H}^+$  permeability leading to continuous efflux of  $\text{H}^+$  to the cytosol (Demaurex, 2002; Wu et al., 2001). Replacement of  $\text{K}^+$  glutamate by equimolar concentrations of KCl rendered similar results (not shown).

### $\text{H}^+/\text{K}^+$ exchange in the intraluminal matrix mediates $\text{pH}_G$ oscillations and oscillatory $\text{H}^+$ efflux from the granule

The experimental validation that  $\text{pH}_G$  oscillations can result from  $\text{H}^+/\text{K}^+$  exchange was conducted in situ, in isolated granules loaded with LS, and in vitro, by titration of  $\text{H}^+/\text{K}^+$  exchange in solutions of heparin. In valinomycin (20  $\mu\text{M}$ ) treated granules—in which both  $\text{InsP}_3$ -R and  $\text{ASK}_{\text{Ca}}$  channels were blocked by heparin (100  $\mu\text{g ml}^{-1}$ ) and apamin (100 nM), respectively—the increase of intraluminal  $\text{K}^+$  led to a concomitant acidification (Fig. 3). Heparin—the major constituent of the mast cell intraluminal matrix—had been shown to work as a histamine/ $\text{K}^+$  exchanger (Uvnas et al., 1989), and we found that it can function as a  $\text{H}^+/\text{K}^+$  exchanger as well. In dilute solutions of heparin (6 mg  $\text{ml}^{-1}$ ), increasing  $[\text{K}^+]$  decreased the pH (Fig. 3 B).

These observations suggest that  $\text{K}^+$  influx into the granule must drive both a  $\text{Ca}^{2+}/\text{K}^+$  exchange process—responsible for  $[\text{Ca}^{2+}]_{\text{IL}}$  oscillations (Nguyen et al., 1998; Quesada et al., 2001)—and a  $\text{H}^+/\text{K}^+$  exchange, that accounts for the periodic acidification of the granule during  $\text{pH}_G$  oscillations (Fig. 1). The corresponding periodic alkalization phases during  $\text{pH}_G$  oscillations probably result from the release of  $\text{Ca}^{2+}$  through  $\text{InsP}_3$ -R channels or from efflux of  $\text{H}^+$  from the granule. Since free  $\text{Ca}^{2+}$  and  $\text{H}^+$  are in equilibrium with




intracellular medium containing  $10 \mu\text{g ml}^{-1}$  of dextran-conjugated Calcium Green-1 and 100 nM apamin, to prevent K<sup>+</sup> influx. InsP<sub>3</sub>-induced release of Ca<sup>2+</sup> (open circles) was accompanied by a concomitant intraluminal alkalinization (filled circles;  $n = 4$ ). (D) Intraluminal and extraluminal pH was simultaneously measured by equilibrating LS-loaded granules in an intracellular solution containing  $10 \mu\text{g ml}^{-1}$  dextran-conjugated SN and 2 mM HEPES (pH = 7.2). SN fluorescence was captured at 587 nm. Addition of InsP<sub>3</sub> ( $3 \mu\text{M}$ ) led to extraluminal pH oscillations in the immediate vicinity of the granule (open circles, right axis). Notice that these extraluminal pH oscillations exhibit the same frequency ( $\sim 0.12$  Hz) and are in phase with pH<sub>G</sub> changes (filled circles) ( $n = 6$ ). In separate preparations in which granules were not loaded with LS, SN reported similar extraluminal pH oscillations in the perigranular space upon InsP<sub>3</sub> application (not shown).

their respective bound forms in the matrix, the release of Ca<sup>2+</sup> through InsP<sub>3</sub>-R channels and the concomitant decrease of  $[Ca^{2+}]_{IL}$  may displace bound Ca<sup>2+</sup> from the polyanionic network to restore the equilibrium with free Ca<sup>2+</sup>, leaving free negative sites which H<sup>+</sup> could occupy, causing alkalinization. A similar competition for binding sites—in this case, cytosolic binding sites—between Ca<sup>2+</sup> and H<sup>+</sup> has been suggested to explain the formation of a secondary H<sup>+</sup> signal in melanotrophs (Beatty et al., 1993). In fact, Fig. 3 C shows that InsP<sub>3</sub>-induced release of Ca<sup>2+</sup> from granules in the presence of apamine, which prevents K<sup>+</sup> influx, led to slight alkalinization. However, a more likely mechanism for intraluminal alkalinization is that the periodic increases of transmembrane pH gradient ( $\Delta\mu_{H^+}$ ) can result in higher efflux of H<sup>+</sup>, with periodic intraluminal alkalinization, and corresponding periodic acidification in the extraluminal side. This outcome is supported by our results and by several reports that have indicated that the major contributor to H<sup>+</sup> export from the granule is an endogenous H<sup>+</sup> permeability—or “leak”—driven by the transmembrane pH gradient ( $\Delta\mu_{H^+}$ ) (Demaurex, 2002; Wu et al., 2001; Schapiro and Grinstein, 2000; Farinas and Verkman, 1999). Although vesicular H<sup>+</sup> “leakage” has been thought to probably take place via voltage-gated H<sup>+</sup> channels (Demaurex, 2002; Schapiro and Grinstein, 2000), the specific mechanism of H<sup>+</sup> efflux from the granule remains controversial (Wu et al., 2001; Schapiro and Grinstein, 2000). To test if secretory vesicles can produce extraluminal oscillations of pH, we equilibrated granules in

an intracellular solution containing  $10 \mu\text{g ml}^{-1}$  of dextran-conjugated SN, a nonpermeant, low diffusivity fluorescent pH sensor. When these granules were exposed to InsP<sub>3</sub>, the pH in the immediate periphery of the granule started to oscillate at the same frequency ( $\sim 0.12$  Hz) and in phase with intraluminal pH oscillations (Fig. 3 D). Therefore, the intraluminal alkalinization we observed during pH<sub>G</sub> oscillations must result from H<sup>+</sup> efflux to the cytosol.

### Temporal relationship between intraluminal and extraluminal dynamics of Ca<sup>2+</sup> and H<sup>+</sup>

To investigate the relationship between Ca<sup>2+</sup> release from the granule and pH<sub>G</sub>, we equilibrated granules loaded with LS in an intracellular bathing solution (see Methods) containing  $10 \mu\text{g ml}^{-1}$  of Calcium Crimson, a dextran-conjugated Ca<sup>2+</sup> probe, to monitor  $[Ca^{2+}]_{EL}$ . The pH of the bathing solution was buffered at 7.2 by 40 mM of HEPES to prevent artifacts resulting from pH-dependent changes of quantum yield of Calcium Crimson. In agreement with previous results (Nguyen et al., 1998; Quesada et al., 2001), Fig. 4 shows that exposure of the granules to  $3 \mu\text{M}$  InsP<sub>3</sub> induced a train of  $[Ca^{2+}]_{IL}$  oscillations by triggering the release of Ca<sup>2+</sup> with the corresponding rise of  $[Ca^{2+}]_{EL}$  and decrease of  $[Ca^{2+}]_{IL}$ . Similarly, InsP<sub>3</sub> produced oscillations of  $[H^+]_{IL}$  of the same frequency but out of phase with the oscillations of  $[Ca^{2+}]_{EL}$  (Fig. 4 B), i.e., decreases of  $[H^+]_{IL}$  are accompanied by corresponding increases of  $[Ca^{2+}]_{EL}$  outside the granule. In isolated granules exposed to heparin



**FIGURE 4** Relationship between intraluminal and extraluminal  $\text{H}^+$  and  $\text{Ca}^{2+}$  oscillations. (A) The intraluminal and extraluminal changes of  $[\text{Ca}^{2+}]$  were monitored in granules loaded with CO-5N and equilibrated in an intracellular solution containing  $10 \mu\text{g ml}^{-1}$  Calcium Crimson. Application of  $3 \mu\text{M}$   $\text{InsP}_3$  provoked oscillations of  $[\text{Ca}^{2+}]_{\text{IL}}$  (filled circles, left axis) and  $[\text{Ca}^{2+}]_{\text{EL}}$  (open circles) of  $\sim 0.12$  Hz, which were  $\sim 180^\circ$  out of phase ( $n = 6$ ). Periodic release of  $\text{Ca}^{2+}$  from the granules results in a corresponding increase of  $[\text{Ca}^{2+}]$  outside the granule (Nguyen et al., 1998; Quesada et al., 2001). (B) Simultaneous monitoring of  $\text{pH}_G$  and  $[\text{Ca}^{2+}]_{\text{EL}}$  was performed in granules loaded with LS and CG as in Fig. 3 C.  $\text{InsP}_3$  provoked oscillations of  $[\text{H}^+]_{\text{IL}}$  (filled circles, left axis) with the same frequency but  $\sim 180^\circ$  out of phase with the  $[\text{Ca}^{2+}]_{\text{EL}}$  oscillations (open circles;  $n = 6$ ). The results in Fig. 3 D and Fig. 4, A and B, indicate that the release of  $\text{Ca}^{2+}$  and the efflux of  $\text{H}^+$  from the granule are  $180^\circ$  out of phase.

( $100 \mu\text{g ml}^{-1}$ ) and apamin ( $100 \text{ nM}$ ),  $[\text{H}^+]_{\text{IL}}$  was unaffected by raising the extraluminal  $[\text{Ca}^{2+}]$  to  $1 \text{ mM}$  (not shown), ruling out the potential involvement of  $\text{Ca}^{2+}/\text{H}^+$  exchangers on the granular membrane, in agreement with previous reports (Mitchell et al., 2001; Schapiro and Grinstein, 2000).

Notice that while the intraluminal and extraluminal oscillations of  $[\text{Ca}^{2+}]$  are phase-shifted (Fig. 4 A), the oscillations of  $[\text{H}^+]_{\text{IL}}$  and  $[\text{H}^+]_{\text{EL}}$  are in phase (Fig. 3 D). To explain this outcome we need to consider that, although the intraluminal  $[\text{Ca}^{2+}]$  and  $[\text{H}^+]$  oscillations are both coupled to  $\text{K}^+$  influx, the oscillations of  $[\text{Ca}^{2+}]_{\text{IL}}$  are modulated by the open/close dynamics of both the  $\text{InsP}_3\text{-R}$  and the  $\text{ASK}_{\text{Ca}}$  channels, while the oscillations of  $[\text{H}^+]_{\text{IL}}$  depend on the open/close dynamics of only the  $\text{ASK}_{\text{Ca}}$  channel and the leakage of this ion from the granule. In the case of  $\text{Ca}^{2+}$  (see model in Fig. 1), the  $\text{InsP}_3$ -mediated  $\text{Ca}^{2+}$  efflux results in a transient decrease in  $[\text{Ca}^{2+}]_{\text{IL}}$  and an increase of  $[\text{Ca}^{2+}]_{\text{EL}}$ . The rise of  $[\text{Ca}^{2+}]_{\text{EL}}$  in the vicinity of the granule both closes the  $\text{InsP}_3\text{-R}$  channel and turns on the  $\text{ASK}_{\text{Ca}}$  channel, activating the influx of  $\text{K}^+$  that results in  $\text{Ca}^{2+}/\text{K}^+$  exchange and rebound of  $[\text{Ca}^{2+}]_{\text{IL}}$ . As  $\text{Ca}^{2+}$  around the granule dissipates by diffusion and buffering, the  $\text{InsP}_3\text{-R}$  channel opens again and the cycle repeats for as long as the  $\text{InsP}_3$  remains bound to its receptor. In the case of  $\text{H}^+$  (see model in Fig. 1), the  $\text{H}^+/\text{K}^+$  exchange in the matrix that increase

$[\text{H}^+]_{\text{IL}}$  steps in when  $\text{ASK}_{\text{Ca}}$  channels open and influx of  $\text{K}^+$  takes place. Since  $\text{H}^+$  efflux is driven by its intraluminal concentration, the oscillations of  $[\text{H}^+]_{\text{IL}}$  changes. During the closed time of the  $\text{ASK}_{\text{Ca}}$  channel, the  $\text{H}^+/\text{K}^+$  exchange ceases but  $\text{H}^+$  still leaks out and  $[\text{H}^+]_{\text{IL}}$  decreases. An implication of these results is that the extraluminal  $[\text{Ca}^{2+}]$  and  $[\text{H}^+]_{\text{IL}}$  must be out of phase. In addition, the rate of  $\text{H}^+/\text{K}^+$  exchange from the heparin matrix must exceed the efflux of  $\text{H}^+$  leakage, otherwise efflux of  $\text{H}^+$  should result in increased  $[\text{H}^+]_{\text{EL}}$  but decreased  $[\text{H}^+]_{\text{IL}}$ . In the case of  $\text{Ca}^{2+}$ , the conductance of the  $\text{InsP}_3$  channel in the open conformation must be higher than the rate of  $\text{Ca}^{2+}$  unbinding from the matrix as  $[\text{Ca}^{2+}]_{\text{IL}}$  rebounds only when the  $\text{InsP}_3$  channel closes and the influx of  $\text{K}^+$  exchanges for a new batch of  $\text{Ca}^{2+}$  from the matrix. We can also infer that oscillations of  $[\text{Ca}^{2+}]_{\text{IL}}$  and  $[\text{H}^+]_{\text{IL}}$  are probably in phase because: 1)  $[\text{Ca}^{2+}]_{\text{IL}}$  and  $[\text{Ca}^{2+}]_{\text{EL}}$  oscillations are out of phase, 2)  $[\text{H}^+]_{\text{IL}}$  and  $[\text{H}^+]_{\text{EL}}$  are in phase, and 3)  $[\text{H}^+]_{\text{IL}}$  and  $[\text{Ca}^{2+}]_{\text{EL}}$  are out of phase (Fig. 4 A, Fig. 3 D, and Fig. 4 B, respectively).

## DISCUSSION

The polymer matrix found inside subcellular organelles—including the secretory granule—holds the answer to a highly significant set of questions in cell biology. From the polymer phase transition properties of the secretory matrix that allows the remarkable payload and efficient discharge of hormones and small molecules to the ion exchange properties of the intravesicular polymer networks, the granule offers one of the most elegant systems designed by evolution. The granule stores and releases material and signals its departure to the export machinery of the cell. Whereas the discovery of phase transitions of the granular matrix brought attention to storage and release in secretion (Verdugo, 1994; Marszalek et al., 1997), the study of the ion exchange properties of the matrix is shifting the focus to questions of signal transduction in secretory cells (Nguyen et al., 1998; Quesada et al., 2001). The  $\text{H}^+$  source/sink properties of the heparin matrix, and probably other secretory matrices, have a broad range of important implications, including pH regulation in subcellular organelles, phagosomal maturation, enzyme activation, protein packing, and sorting in the trans-Golgi network (Bell-Parikh et al., 2001; Reeves et al., 2002). However, the association of  $\text{H}^+$  release with the  $\text{InsP}_3$ -induced  $\text{Ca}^{2+}$  signal from the granule, their oscillatory nature, and the presence of exocytic proteins sensitive to the joint action of  $\text{Ca}^{2+}$  and pH strongly suggest that  $\text{Ca}^{2+}/\text{H}^+$  release from the secretory granule might encode a combined intracellular signal. According to our working model (Fig. 1), the activation of an extracellular receptor is relayed to the intracellular network by production of  $\text{InsP}_3$  (Berridge et al., 2000). The  $\text{InsP}_3$  signal is received by  $\text{InsP}_3\text{-R}$  channels of nearby secretory granules, turning them into double ion oscillators that respond with two spatially and temporally

constrained frequency-encoded signals of  $\text{Ca}^{2+}$  and  $\text{H}^+$ . These oscillations are independent of ATP-mediated active uptake of  $\text{Ca}^{2+}$  or  $\text{H}^+$ . Instead, they are brought about by the interaction of  $\text{InsP}_3$ -R and  $\text{ASK}_{\text{Ca}}$  channels of the granule (Nguyen et al., 1998; Quesada et al., 2001; Gerasimenko et al., 1996; Yoo, 2000; Thevenod, 2002), with opposite gating sensitivities to  $\text{Ca}^{2+}$ ; the  $\text{H}^+$  “leakage” properties of the granular membrane (Demaurex, 2002; Wu et al., 2001); and the unique  $\text{Ca}^{2+}/\text{K}^+$  and  $\text{H}^+/\text{K}^+$  ion exchange properties of the heparin granular matrix (Uvnas and Aborg, 1977, 1989; Verdugo, 1994; Nguyen et al., 1998; Quesada et al., 2001; Nanavati and Fernandez, 1993; Marszalek et al., 1997; Chin et al., 2002).

In the space domain, the release of  $\text{Ca}^{2+}$  and  $\text{H}^+$  affects an exceedingly small cytosolic volume that probably scales to intermolecular distances not much farther than the local Debye potential field present in the cleft between plasma and granular membranes before membrane fusion. With these boundary conditions, diffusional distances become irrelevant, and the local concentration of  $\text{Ca}^{2+}$  and  $\text{H}^+$  in the cleft could very well mirror the intravesicular concentration of these ions. Because of the buffering properties of the cytosol, these signals should be time and space limited, reaching strictly confined domains in the cleft and preventing undesired cross talk with other receptor proteins not involved in membrane fusion.

In the time domain, the observed 0.1 Hz frequency of oscillation of  $\text{Ca}^{2+}$  and pH signals allows scanning of a broad range of cytosolic  $[\text{Ca}^{2+}]$  and pH in 5-s periods. Diffusional delays are unlikely to occur because the sensor-effector proteins are already present in the cleft either in free form or anchored to the granule or plasma membranes (Sudhof, 1995), and the diffusion distance for  $\text{Ca}^{2+}$  and  $\text{H}^+$  to reach their targets across the cleft is extremely short. Thus, considering the typical  $\mu\text{s}$ -ms relaxation timescale of molecular conformational changes, effector proteins would have enough time to switch configuration (Subramaniam and Henderson, 2000; Rami and Udgaonkar, 2001). The pre-exocytic oscillations of  $\text{Ca}^{2+}$  and  $\text{H}^+$  in the narrow cleft existing between the two membranes exhibit broad overlapping. They scan a wide combination of concentrations of  $\text{Ca}^{2+}$  and  $\text{H}^+$  that could create multiple yet unique conditions, attuned to the specific optimal  $\text{Ca}^{2+}/\text{pH}$  dependency of the different exocytic proteins, perhaps triggering their individual fusogenic properties in a well programmed sequence.

Several proteins implicated in exocytosis including calmodulin, syncollin, or Rab3a exhibit high interdependent sensitivity to  $\text{Ca}^{2+}$  and pH (An et al., 2000; Kiss and Korn, 1999; Kajio et al., 2001; Hudmon et al., 1996; Kennedy et al., 1983). The interaction of calmodulin with different substrates requires not only changes of pH and  $[\text{Ca}^{2+}]$  but frequency-encoded signals of  $[\text{Ca}^{2+}]_{\text{C}}$  as well (De Koninck and Schulman, 1998). Protein kinase C is another protein involved in secretion that can also work as a decoder of

oscillatory signals (Oancea and Meyer, 1998). However, the family of annexins gives the most striking case of combined  $\text{Ca}^{2+}/\text{pH}$  dependence. These proteins are important mediators of exocytosis by means of their collective ability to fuse membranes in a  $\text{Ca}^{2+}$ -dependent manner (Caohuy and Pollard, 2001; Konig et al., 1998). Remarkably, recent studies have demonstrated that the fusogenic efficiency of these proteins exhibits a critical sensitivity to pH, requiring an acidic environment of lower pH than the one found in the bulk cytosol. Depending on each specific annexin, different acidic pH values are required with slight variations of the synergy between  $\text{Ca}^{2+}$  and  $\text{H}^+$  (Langen et al., 1998; Isas et al., 2000; Caohuy and Pollard, 2002). Since the requirements of these proteins for both ions are much higher than those found in the bulk cytosol, several groups have proposed that membrane fusion induced by annexins is possible because of local signals that generate confined areas of high concentration of both  $\text{Ca}^{2+}$  and  $\text{H}^+$  (Langen et al., 1998; Isas et al., 2000; Caohuy and Pollard, 2002).

The present results are in agreement with observations in intact cells. Several groups have seen preexocytotic granular pH changes in pancreatic  $\beta$ -cells, mast cells, and neurons, postulating an active role of pH in priming granules for release (Williams and Webb, 2000; Barg et al., 2001; Han et al., 1999; Renstrom et al., 2002). The idea of a  $\text{Ca}^{2+}/\text{H}^+$  signaling system is consistent with observations that both luminal  $\text{Ca}^{2+}$  efflux and the maintenance of granular  $\Delta\mu_{\text{H}^+}$  are needed for vacuole and granule fusion (Peters and Mayer, 1998; Ungermann et al., 1999; Peters et al., 2001; Scheenen et al., 1998; Mundorf et al., 2000; Yang et al., 2002). Although the mechanisms of acidification remain uncertain, the idea that pH changes may facilitate secretion by affecting exocytotic proteins, making them more fusogenic, has also been considered (Barg et al., 2001, 2002; Yang et al., 2002; Renstrom et al., 2002).

The search for how specificity is encoded in intracellular signal transduction remains one of the most interesting and challenging topics in cell biology. Instances of built-in conditional arguments are present in the intracellular web of information (Beatty et al., 1993; Berridge et al., 2000; Susini et al., 2000). However, the formalization of simple principles of information theory in this field remains virtually unexplored. Although both  $\text{Ca}^{2+}$  and  $\text{H}^+$  can readily induce conformational changes, switching on/off functional conformations in proteins or other polyions present in the cell, the broad effect of these cations can decrease their specificity. The assignment of their combination in signaling could represent a heuristic model of Boolean conditional signaling whereby the granule can target a specific group of sensor/effector proteins involved in implementing exocytosis.

We thank Drs. Peter B. Detwiler, Albert M. Gordon, Thomas R. Hinds, and Garrett M. Odell.



This work was supported by a grant from the National Science Foundation Biocomplexity Program to P.V. and Florida State University First-Year Assistant Professor and Center Development Award to W.C.C. I.Q. is a recipient of a Spanish Ministry of Education and Culture postdoctoral fellowship.

## REFERENCES

- An, S. J., N. J. Hansen, A. Hodel, R. Jahn, and J. M. Edwardson. 2000. Analysis of the association of syncollin with the membrane of the pancreatic zymogen granule. *J. Biol. Chem.* 275:11306–11311.
- Barg, S., L. Eliasson, E. Renstrom, and P. Rorsman. 2002. A subset of 50 secretory granules in close contact with L-type  $\text{Ca}^{2+}$  channels accounts for first-phase insulin secretion in mouse beta-cells. *Diabetes.* 51:S74–S82.
- Barg, S., P. Huang, L. Eliasson, D. J. Nelson, S. Obermuller, P. Rorsman, F. Thevenod, and E. Renstrom. 2001. Priming of insulin granules for exocytosis by granular  $\text{Cl}^-$  uptake and acidification. *J. Cell Sci.* 114: 2145–2154.
- Beatty, D. M., B. M. Chronwall, D. E. Howard, T. B. Wiegmann, and S. J. Morris. 1993. Calcium regulation of intracellular pH in pituitary intermediate lobe melanotopes. *Endocrinology.* 133:972–984.
- Belan, P. V., O. V. Gerasimenko, D. Berry, E. Saftenku, O. H. Petersen, and A. V. Tepikin. 1996. A new technique for assessing the microscopic distribution of cellular calcium exit sites. *Pflugers Arch.* 433:200–208.
- Bell-Parikh, L. C., B. A. Eipper, and R. E. Mains. 2001. Response of an integral granule membrane protein to changes in pH. *J. Biol. Chem.* 276:29854–29863.
- Berridge, M. J., P. Lipp, and M. D. Bootman. 2000. The versatility and universality of calcium signalling. *Nat. Rev. Mol. Cell Biol.* 1:11–21.
- Caohuy, H., and H. B. Pollard. 2001. Activation of annexin 7 by protein kinase C in vitro and in vivo. *J. Biol. Chem.* 276:12813–12821.
- Caohuy, H., and H. B. Pollard. 2002. Protein kinase C and guanosine triphosphate combine to potentiate calcium-dependent membrane fusion driven by annexin 7. *J. Biol. Chem.* 277:25217–25225.
- Chin, W. C., I. Quesada, T. Nguyen, and P. Verdugo. 2002. Oscillations of pH inside the secretory granule control the gain of  $\text{Ca}^{2+}$  release for signal transduction in goblet cell exocytosis. *Novartis Found. Symp.* 248: 132–149.
- De Koninck, P., and H. Schulman. 1998. Sensitivity of CaM kinase II to the frequency of  $\text{Ca}^{2+}$  oscillations. *Science.* 279:227–230.
- Demaurex, N. 2002. pH homeostasis of cellular organelles. *News Physiol. Sci.* 17:1–5.
- Farinas, J., and A. S. Verkman. 1999. Receptor-mediated targeting of fluorescent probes in living cells. *J. Biol. Chem.* 274:7603–7606.
- Gerasimenko, O. V., J. V. Gerasimenko, P. V. Belan, and O. H. Petersen. 1996. Inositol trisphosphate and cyclic ADP-ribose-mediated release of  $\text{Ca}^{2+}$  from single isolated pancreatic zymogen granules. *Cell.* 84:473–480.
- Han, W., D. Li, A. K. Stout, K. Takimoto, and E. S. Levitan. 1999.  $\text{Ca}^{2+}$ -induced deprotonation of peptide hormones inside secretory vesicles in preparation for release. *J. Neurosci.* 19:900–905.
- Hudmon, A., J. Aronowski, S. J. Kolb, and M. N. Waxham. 1996. Inactivation and self-association of  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase II during autophosphorylation. *J. Biol. Chem.* 271:8800–8808.
- Isas, J. M., J. P. Cartiailler, Y. Sokolov, D. R. Patel, R. Langen, H. Luecke, J. E. Hall, and H. T. Haigler. 2000. Annexins V and XII insert into bilayers at mildly acidic pH and form ion channels. *Biochemistry.* 39: 3015–3022.
- Kajio, H., S. Olszewski, P. J. Rosner, M. J. Donelan, K. F. Geoghegan, and C. J. Rhodes. 2001. A low-affinity  $\text{Ca}^{2+}$ -dependent association of calmodulin with the Rab3A effector domain inversely correlates with insulin exocytosis. *Diabetes.* 50:2029–2039.
- Kao, J. P. Y. 1994. Practical aspects of measuring  $[\text{Ca}^{2+}]$  with fluorescent probes. In *A Practical Guide to the Study of Calcium in Living Cells*. R. Nuccitelli, editor. Academic Press, San Diego. 155–181.
- Kennedy, M. B., T. McGuinness, and P. Greengard. 1983. A calcium/calmodulin-dependent protein kinase from mammalian brain that phosphorylates Synapsin I: partial purification and characterization. *J. Neurosci.* 3:818–831.
- Kiss, L., and S. J. Korn. 1999. Modulation of N-type  $\text{Ca}^{2+}$  channels by intracellular pH in chick sensory neurons. *J. Neurophysiol.* 81:1839–1847.
- Konig, J., J. Prenen, B. Nilius, and V. Gerke. 1998. The annexin II-p11 complex is involved in regulated exocytosis in bovine pulmonary artery endothelial cells. *J. Biol. Chem.* 273:19679–19684.
- Langen, R., J. M. Isas, W. L. Hubbell, and H. T. Haigler. 1998. A transmembrane form of annexin XII detected by site-directed spin labeling. *Proc. Natl. Acad. Sci. USA.* 95:14060–14065.
- Marszalek, P. E., B. Farrell, P. Verdugo, and J. M. Fernandez. 1997. Kinetics of release of serotonin from isolated secretory granules. II. Ion exchange determines the diffusivity of serotonin. *Biophys. J.* 73:1169–1183.
- Mitchell, K. J., P. Pinton, A. Varadi, C. Tacchetti, E. K. Ainscow, T. Pozzan, R. Rizzuto, and G. A. Rutter. 2001. Dense core secretory vesicles revealed as a dynamic  $\text{Ca}^{2+}$  store in neuroendocrine cells with a vesicle-associated membrane protein aequorin chimera. *J. Cell Biol.* 155:41–51.
- Monck, J. R., A. F. Oberhauser, T. J. Keating, and J. M. Fernandez. 1992. Thin-section ratiometric  $\text{Ca}^{2+}$  images obtained by optical sectioning of fura-2 loaded mast cells. *J. Cell Biol.* 116:745–759.
- Mundorf, M. L., K. P. Troyer, S. E. Hochstetler, J. A. Near, and R. M. Wightman. 2000. Vesicular  $\text{Ca}^{2+}$  participates in the catalysis of exocytosis. *J. Biol. Chem.* 275:9136–9142.
- Nanavati, C., and J. M. Fernandez. 1993. The secretory granule matrix: a fast-acting smart polymer. *Science.* 259:963–965.
- Nguyen, T., W. C. Chin, and P. Verdugo. 1998. Role of  $\text{Ca}^{2+}/\text{K}^+$  ion exchange in intracellular storage and release of  $\text{Ca}^{2+}$ . *Nature.* 395:908–912.
- Oancea, E., and T. Meyer. 1998. Protein kinase C as a molecular machine for decoding calcium and diacylglycerol signals. *Cell.* 95:307–318.
- Peters, C., M. J. Bayer, S. Buhler, J. S. Andersen, M. Mann, and A. Mayer. 2001. Trans-complex formation by proteolipid channels in the terminal phase of membrane fusion. *Nature.* 409:581–588.
- Peters, C., and A. Mayer. 1998.  $\text{Ca}^{2+}$ /calmodulin signals the completion of docking and triggers a late step of vacuole fusion. *Nature.* 396:575–580.
- Quesada, I., W. C. Chin, J. Steed, P. Campos-Bedolla, and P. Verdugo. 2001. Mouse mast cell secretory granules can function as intracellular ionic oscillators. *Biophys. J.* 80:2133–2139.
- Rami, B. R., and J. B. Udgaonkar. 2001. pH-jump-induced folding and unfolding studies of barstar: evidence for multiple folding and unfolding pathways. *Biochemistry.* 40:15267–15279.
- Reeves, E. P., H. Lu, H. L. Jacobs, C. G. Messina, S. Bolsover, G. Gabella, E. O. Potma, A. Warley, J. Roes, and A. W. Segal. 2002. Killing activity of neutrophils is mediated through activation of proteases by  $\text{K}^+$  flux. *Nature.* 416:291–297.
- Renstrom, E., R. Ivarsson, and S. B. Shears. 2002. Inositol 3,4,5,6-tetrakisphosphate inhibits insulin granule acidification and fusogenic potential. *J. Biol. Chem.* 277:26717–26720.
- Schapiro, F. B., and S. Grinstein. 2000. Determinants of the pH of the Golgi complex. *J. Biol. Chem.* 275:21025–21032.
- Scheenen, W. J., C. B. Wollheim, T. Pozzan, and C. Fasolato. 1998.  $\text{Ca}^{2+}$  depletion from granules inhibits exocytosis. A study with insulin-secreting cells. *J. Biol. Chem.* 273:19002–19008.
- Subramaniam, S., and R. Henderson. 2000. Crystallographic analysis of protein conformational changes in the bacteriorhodopsin photocycle. *Biochim. Biophys. Acta.* 1460:157–165.
- Sudhof, T. 1995. The synaptic vesicle cycle: a cascade of protein-protein interactions. *Nature.* 375:645–653.
- Susini, S., G. Van Haasteren, S. Li, M. Prentki, and W. Schlegel. 2000. Essentiality of intron control in the induction of *c-fos* by glucose and glucocorticoid peptides in INS-1 beta-cells. *FASEB J.* 14:128–136.

- Thevenod, F. 2002. Ion channels in secretory granules of the pancreas and their role in exocytosis and release of secretory proteins. *Am. J. Physiol. Cell Physiol.* 283:C651–C672.
- Ungermann, C., W. Wickner, and Z. Xu. 1999. Vacuole acidification is required for trans-SNARE pairing, LMA1 release, and homotypic fusion. *Proc. Natl. Acad. Sci. USA.* 96:11194–11199.
- Uvnas, B., and C. H. Aborg. 1977. On the cation exchanger properties of rat mast cell granules and their storage of histamine. *Acta Physiol. Scand.* 100:309–314.
- Uvnas, B., and C. H. Aborg. 1989. Role of ion exchange in release of biogenic amines. *News Physiol. Sci.* 4:68–71.
- Uvnas, B., C. H. Aborg, L. Lyssarides, and L. G. Danielsson. 1989. Intracellular ion exchange between cytoplasmic potassium and granule histamine, an integrated link in the histamine release machinery of mast cells. *Acta Physiol. Scand.* 136:309–320.
- Verdugo, P. 1994. Polymer gel phase transition in condensation-decondensation of secretory products. *Advances in Polymer Science.* 110:145–156.
- Williams, R. M., and W. W. Webb. 2000. Single granule pH cycling in antigen-induced mast cell secretion. *J. Cell Sci.* 113:3839–3850.
- Wu, M. M., M. Grabe, S. Adams, R. Y. Tsien, H. P. Moore, and T. E. Machen. 2001. Mechanisms of pH regulation in the regulated secretory pathway. *J. Biol. Chem.* 276:33027–33035.
- Yang, J., A. Hodel, and G. D. Holman. 2002. Insulin and isoproterenol have opposing roles in the maintenance of cytosol pH and optimal fusion of GLUT4 vesicles with the plasma membrane. *J. Biol. Chem.* 277:6559–6566.
- Yoo, S. H. 2000. Coupling of the IP<sub>3</sub> receptor/Ca<sup>2+</sup> channel with Ca<sup>2+</sup> storage proteins chromogranins A and B in secretory granules. *Trends Neurosci.* 23:424–428.