

ERRATA

Stx2 Subtyping of Shiga Toxin-Producing *Escherichia coli* Isolated from Cattle in France: Detection of a New Stx2 Subtype and Correlation with Additional Virulence Factors

Yolande Bertin, Karima Boukhors, Nathalie Pradel, Valerie Livrelli, and Christine Martin

Laboratoire de Microbiologie, Centre de Recherche INRA de Clermont-Ferrand-Theix, 63122 St-Genès Charniere, and Groupe de Recherche Pathogénie Bactérienne Intestinale, Faculté de Pharmacie, Université d'Auvergne, Clermont-Ferrand, France

Volume 39, no. 9, p. 3060–3065, 2001. Page 3065: References 11 to 29 should be numbered 10 to 28, respectively.

Multilocus Sequence Typing Scheme for *Enterococcus faecium*

Wieger L. Homan, David Tribe, Simone Poznanski, Mei Li, Geoff Hogg, Emile Spalburg, Jan D. A. van Embden, and Rob J. L. Willems

Research Laboratory for Infection Diseases, National Institute of Public Health and the Environment, Bilthoven, The Netherlands, and Department of Microbiology and Immunology and Microbiological Diagnostic Unit, University of Melbourne, Parkville, Victoria, Australia

Volume 40, no. 6, p. 1963–1971. Page 1964, column 2: The sequences for the primers gyd1 and purK2 should read as follows: for gyd1, “5'-CAA ACT GCT TAG CTC CAA TGG C-3',” and for purK2, “5'-TAC ATA AAT CCC GCC TGT TTY-3'.”