

Short report

HIV in semen: Still more to be learned

Pietro L Vernazza*

Address: Division of Infectious Diseases and Hospital Epidemiology, Department of Medicine, Cantonal Hospital, St. Gallen Switzerland

Email: Pietro L Vernazza* - pietro.vernazza@kssg.ch

* Corresponding author

Published: 03 December 2005

Received: 07 November 2005

AIDS Research and Therapy 2005, **2**:11 doi:10.1186/1742-6405-2-11

Accepted: 03 December 2005

This article is available from: <http://www.aidsrestherapy.com/content/2/1/11>

© 2005 Vernazza; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In 1983, during the earliest days of AIDS research, Deborah Anderson and her colleagues in Boston, Massachusetts hypothesized that AIDS was transmitted by virally-infected "Trojan horse leukocytes" in semen [1]. This prediction has been supported by numerous studies over the past two decades, although many questions remain concerning HIV infection of the male genital tract. In this issue of *AIDS Research and Therapy*, the Anderson group presents an important research tool to help address some of the critical unanswered questions in this area [2].

A series of salient studies have shaped current concepts on HIV-1 in semen. In 1984, Ho et al. described retroviral particles and infected cells in the semen of a homosexual man with AIDS [3]. Shortly thereafter Stewart et al. reported infection of four out of eight women following artificial insemination with semen from one seroconverting individual [4], leading to a mandatory semen quarantine requirement and HIV testing of semen donors in Assisted Reproduction clinics. At that time, the detection of HIV, which was still termed HTLV-III, was not routinely feasible from blood, let alone from semen. Since then, technological advances have enabled the detection and quantitation of HIV-1 RNA and proviral DNA and greatly improved our understanding of the dynamics of HIV-1 in semen and sexual transmission risks. HIV-infected white blood cells have been detected throughout the male genital tract, and in preejaculatory fluid and semen from HIV+men [5,8]; the weight of evidence suggests that sperm are not infectious [9], leading to the successful development of sperm wash procedures to reduce the risk of HIV transmission from HIV-infected men to uninfected partners through assisted reproduction techniques [10]. A combination of epidemiological and clinical research

studies have determined a relationship between HIV-1 RNA viral load in semen and the risk of sexual transmission. The most important factors associated with increased HIV viral loads in semen and risk of sexual transmission are: HIV-viremia and coinfections with other sexually transmitted pathogens [11,12]. HAART dramatically suppresses HIV-1 RNA viral loads in blood and semen, but HIV-1 proviral DNA can persist in semen WBCs for months after the initiation of HAART [13]. Data from other studies showing discordantly higher levels of HIV in semen than blood in some individuals support this finding. In addition, molecular sequencing studies indicate that the male genital tract is a compartment, like the central nervous system, in which HIV-1 replication and divergent evolution can occur under the influence of local factors [14]. Several clinically important questions remain: 1) Is HIV-1 primarily sexually transmitted by infected cells, cell-free virus or both? 2) What is the origin of cell-free and cell-associated HIV-1 in semen? 3) Are men on HAART with undetectable peripheral viral loads capable of sexually transmitting drug-resistant HIV-1?

Episomal HIV-1 c-DNA, a by-product of HIV-1 infection, is currently used in clinical trials as a marker of residual viral replication and potential evolution of drug resistance mutations in viral reservoir sites in individuals on HAART [15]. Such a marker would be useful for identifying sites of HIV-1 replication in the male genital tract, and for monitoring cryptic HIV-1 infection in the genital tract of men on antiretroviral therapy. The only reported study that measured episomal HIV-1 c-DNA in blood and semen of men before and after initiation of HAART failed to detect HIV episomal 2-LTR cDNA in semen [16]. The method used to recover HIV-infected cells from semen in

this study – separation of seminal WBC on Ficoll gradients – likely decreased the sensitivity of HIV episomal c-DNA detection because infected macrophages and a proportion of infected T-cells are lost through this approach. The paper by Xu et al. used a direct lysis technique optimizing recovery of DNA from HIV-infected cells in semen. Using this approach, combined with quantitative PCR and DNA sequencing, the investigators show that episomal 2-LTR cDNA is detectable in semen from a subset of men with other evidence of seminal HIV-1 infection. The marker was not detected in semen from 22 men at 1- and 6-months after peripheral viral suppression due to addition of indinavir to their ART regimen. This study is important because it provides a new tool for studying HIV infection of the male genital tract, and provides preliminary evidence that cryptic HIV-1 infection may not occur in the genital tract of men on HAART. Further studies will surely follow to confirm and extend these observations.

References

- Anderson DJ, Yunis EJ: "**Trojan Horse**" leukocytes in AIDS. *N Engl J Med* 1983, **309**:984-985.
- Xu C, Politch JA, Mayer KH, Anderson DJ: **Detection of Human Immunodeficiency Virus type-I episomal cDNA in semen**. *AIDS Res and Therapy* 2005:000-000.
- Ho DD, Schooley RT, Rota TR, Kaplan JC, Flynn T, Salahuddin SZ, Gonda, MA, Hirsch MS: **HTLV-III in the semen and blood of a healthy homosexual man**. *Science* 1984, **226**:451-453.
- Stewart GJ, Tyler JP, Cunningham AL, Barr JA, Driscoll GL, Gold J, Lamont BJ: **Transmission of human T-cell lymphotropic virus type III (HTLV-III) by artificial insemination by donor**. *Lancet* 1985, **2**:581-585.
- Van Voorhis BJ, Martinez A, Mayer K, Anderson DJ: **Detection of human immunodeficiency virus type I in semen from seropositive men using culture and polymerase chain reaction deoxyribonucleic acid amplification techniques**. *Fertil Steril* 1991, **55**:588-594.
- Pudney J, Anderson D: **Orchitis and human immunodeficiency virus type I infected cells in reproductive tissues from men with the acquired immune deficiency syndrome**. *Am J Pathol* 1991, **139**:149-160.
- Pudney J, Oneta M, Mayer K, Seage G, Anderson DJ: **Pre-ejaculatory fluid as potential vector for sexual transmission of HIV-1**. *Lancet* 1992, **340**:1470-1470.
- Xu C, Politch JA, Tucker L, Mayer KH, Seage GR, Anderson DJ: **Factors associated with increased levels of human immunodeficiency virus type I DNA in semen**. *J Infect Dis* 1997, **176**:941-947.
- Quayle AJ, Xu C, Mayer KH, Anderson DJ: **T lymphocytes and macrophages, but not motile spermatozoa, are a significant source of human immunodeficiency virus in semen**. *J Infect Dis* 1997, **176**:960-968.
- Semprini AE, Vucetich A, Hollander L: **Sperm washing, use of HAART and role of elective Caesarean section**. *Curr Opin Obstet Gynecol* 2004, **16**:465-470.
- Vernazza PL, Gilliam BL, Dyer JR, Fiscus SA, Eron JJ, Frank AC, Cohen MS: **Quantitation of HIV in semen: Correlation with antiviral treatment and immune status**. *AIDS* 1997, **11**:987-993.
- Cohen MS, Hoffman IF, Royce RA, Kazembe P, Dyer JR, Daly CC, Zimba D, Vernazza PL, Maida M, Fiscus SA, Eron JJ: **Reduction of concentration of HIV-1 in semen after treatment of urethritis: implications for prevention of sexual transmission of HIV-1**. *Lancet* 1997, **349**:1868-1873.
- Vernazza PL, Troiani L, Flepp MJ, Cone RW, Schock J, Roth F, Boggian K, Cohen MS, Fiscus SA, Eron JJ: **Potent antiretroviral treatment of HIV-infection results in suppression of the seminal shedding of HIV. The Swiss HIV Cohort Study**. *AIDS* 2000, **14**:117-121.
- Eron JJ, Vernazza PL, Johnston DM, Seillier-Moisewitsch F, Alcorn TM, Fiscus SA, Cohen MS: **Resistance to HIV-1 to antiretroviral agents in blood and seminal plasma: Implications for Transmission**. *AIDS* 1998, **12**:F189.
- Dornadula G, Nunnari G, Vanella M, Roman J, Babinchak T, DeSimone J, Stern J, Bruffman M, Zhang H, Pomerantz RJ: **Human immunodeficiency virus type I-infected persons with residual disease and virus reservoirs on suppressive highly active antiretroviral therapy can be stratified into relevant virologic and immunologic subgroups**. *J Infect Dis* 2001, **183**:1682-1687.
- Nunnari G, Otero M, Dornadula G, Vanella M, Zhang H, Frank I, Pomerantz RJ: **Residual HIV-1 disease in seminal cells of HIV-1-infected men on suppressive HAART: latency without ongoing cellular infections**. *AIDS* 2002, **16**:39-45.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

