

# The Protein Phosphatase *AtPP2CA* Negatively Regulates Abscisic Acid Signal Transduction in Arabidopsis, and Effects of *abh1* on *AtPP2CA* mRNA<sup>1[W]</sup>

Josef M. Kuhn<sup>2</sup>, Aurélien Boisson-Dernier<sup>2</sup>, Marie B. Dizon, Mohammad H. Maktabi, and Julian I. Schroeder\*

Division of Biological Sciences, Cell and Developmental Biology Section, and Center for Molecular Genetics, University of California, La Jolla, California 92093–0116

To identify new loci in abscisic acid (ABA) signaling, we screened a library of 35S::cDNA Arabidopsis (*Arabidopsis thaliana*)-expressing lines for ABA-insensitive mutants in seed germination assays. One of the identified mutants germinated on 2.5  $\mu\text{M}$  ABA, a concentration that completely inhibits wild-type seed germination. Backcrosses and F<sub>2</sub> analyses indicated that the mutant exhibits a dominant phenotype and that the ABA insensitivity was linked to a single T-DNA insertion containing a 35S::cDNA fusion. The inserted cDNA corresponds to a full-length cDNA of the *AtPP2CA* gene, encoding a protein phosphatase type 2C (PP2C). Northern-blot analyses demonstrated that the *AtPP2CA* transcript is indeed overexpressed in the mutant (named PP2CAox). Two independent homozygous T-DNA insertion lines, *pp2ca-1* and *pp2ca-2*, were recovered from the Arabidopsis Biological Resource Center and shown to lack full-length *AtPP2CA* expression. A detailed characterization of PP2CAox and the T-DNA disruption mutants demonstrated that, whereas ectopic expression of a 35S::*AtPP2CA* fusion caused ABA insensitivity in seed germination and ABA-induced stomatal closure responses, disruption mutants displayed the opposite phenotype, namely, strong ABA hypersensitivity. Thus our data demonstrate that the PP2CA protein phosphatase is a strong negative regulator of ABA signal transduction. Furthermore, it has been previously shown that the *AtPP2CA* transcript is down-regulated in the ABA-hypersensitive nuclear mRNA cap-binding protein mutant *abh1*. We show here that down-regulation of *AtPP2CA* in *abh1* is not due to impaired RNA splicing of *AtPP2CA* pre-mRNA. Moreover, expression of a 35S::*AtPP2CA* cDNA fusion in *abh1* partially suppresses *abh1* hypersensitivity, and the data further suggest that additional mechanisms contribute to ABA hypersensitivity of *abh1*.

The phytohormone abscisic acid (ABA), which regulates many agronomically important aspects of plant life, including seed development and dormancy, plays a critical role in plant stress responses such as drought, salinity, cold shock, wounding, and pathogen attack (Schroeder et al., 2001; Finkelstein et al., 2002; Hetherington and Woodward, 2003; Fan et al., 2004). These physiological responses to ABA are in large part due to changes in gene expression and a complex signal transduction network (Hoth et al., 2002; Seki et al., 2002; Leonhardt et al., 2004; Takahashi et al., 2004). Several transcription factors mediating ABA responses have been isolated (Giraudat et al., 1992; Finkelstein et al., 1998; Finkelstein and Lynch, 2000;

Uno et al., 2000; Himmelbach et al., 2002; Abe et al., 2003).

Whereas the biosynthesis of ABA is well understood (Seo and Koshiba, 2002), the mechanisms by which ABA regulates multiple plant responses is beginning to be revealed through genetic and physiological analyses in Arabidopsis (*Arabidopsis thaliana*). To date, genetic screens for ABA-hypersensitive mutants have indicated that processes including farnesylation (*era1*; Cutler et al., 1996; Pei et al., 1998), inositol 1,4,5-triphosphate (IP<sub>3</sub>) dephosphorylation (*fry1*; Xiong et al., 2001b), and RNA metabolism (*hyl1*; Lu and Fedoroff, 2000; *abh1*, Hugouvieux et al., 2001; *sad1*, Xiong et al., 2001a) are required to attenuate the ABA signal. However, surprisingly few non-transcription factor-encoding genes have been identified as recessive ABA-insensitive disruption mutants, namely, the G-protein  $\alpha$ -subunit *GPA1* (Wang et al., 2001), the *RCN1* protein phosphatase type 2A subunit (Kwak et al., 2002), the *OST1/SnRK2E* protein kinase (Mustilli et al., 2002; Yoshida et al., 2002), the *AtRBOHD/F* NADPH oxidases (Kwak et al., 2003), *ABI8* (Brocard-Gifford et al., 2004), *RPK1* (Osakabe et al., 2005), and *GCA2* (Himmelbach et al., 1998).

Many gene families in the Arabidopsis genome have large numbers of homologs relative to other sequenced genomes (Arabidopsis Genome Initiative, 2000). Therefore, the relatively low number of recessive

<sup>1</sup> This work was supported by National Institutes of Health (R01GM060396) and National Science Foundation (MCB0417118) grants (to J.I.S.), and by a European Molecular Biology Organization fellowship (to J.M.K.).

<sup>2</sup> These authors contributed equally to the paper.

\* Corresponding author; email julian@biomail.ucsd.edu; fax 858–534–7108.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors ([www.plantphysiol.org](http://www.plantphysiol.org)) is: Julian I. Schroeder ([julian@biomail.ucsd.edu](mailto:julian@biomail.ucsd.edu)).

<sup>[W]</sup> The online version of this article contains Web-only data.

Article, publication date, and citation information can be found at [www.plantphysiol.org/cgi/doi/10.1104/pp.105.070318](http://www.plantphysiol.org/cgi/doi/10.1104/pp.105.070318).

ABA-insensitive mutants is most likely due to redundancy in genes encoding ABA transducers, requiring analyses of double or multiple mutations in (partially) redundant genes (Kwak et al., 2003). In addition, the ABA signaling pathway is mediated by a network of events and interacts with many other signaling pathways including drought, salinity, cold, sugar, GA<sub>3</sub>, and ethylene (Finkelstein et al., 2002; Zhu, 2002; Himmelbach et al., 2003; Yamaguchi-Shinozaki and Shinozaki, 2005). Consequently, functional redundancy could also be explained by the fact that such complex networks have the ability to buffer a mutation's effects in a neighboring pathway (Cutler and McCourt, 2005).

To further enhance the chance of success in identifying new mutants in ABA signaling, we screened a library of 35S::cDNA Arabidopsis-expressing lines (LeClere and Bartel, 2001) for ABA-insensitive mutants in seed germination assays. In theory, the 35S::cDNA fusion can generate mutants due to (1) random insertional gene disruptions; (2) overexpression of the inserted full-length or truncated cDNA; or (3) silencing of the cDNA-corresponding endogenous gene (LeClere and Bartel, 2001). Consequently, this screen can identify both positive and negative regulators of a given signaling pathway. In this study, we report isolation of two strong ABA-insensitive mutants: a new insertional mutant of *ABI5*, coding for a basic Leu zipper transcription factor, a well-known positive regulator of ABA responses (Finkelstein and Lynch, 2000), as well as a constitutive overexpressor of *AtPP2CA* (named PP2CAox) encoding a protein phosphatase type 2C (PP2C).

Sixty-nine PP2Cs are encoded in the Arabidopsis genome (Kerk et al., 2002; Schweighofer et al., 2004), and a gene disruption phenotype has only been reported for one of these PP2Cs (Leonhardt et al., 2004; Saez et al., 2004). The protein phosphatase PP2CA belongs to group A of the Arabidopsis PP2C family, together with *ABI1*, *ABI2*, and *AtP2C-HA* (hereafter named *AtP2C-HAB1*; Schweighofer et al., 2004). Recently, reverse-genetics studies have demonstrated that *AtP2C-HAB1* is a negative regulator of ABA signaling (Leonhardt et al., 2004; Saez et al., 2004). Isolation and characterization of the dominant negative *abi1-1* and *abi2-1* PP2C mutants and their intragenic revertants also support a negative role for these two PP2Cs (Leung et al., 1994, 1997; Meyer et al., 1994; Rodriguez et al., 1998; Gosti et al., 1999; Merlot et al., 2001). However, in transgenic Arabidopsis plants overexpressing the *ABI1* gene, no altered ABA sensitivity was found in seed germination or in suppression of ABA-mediated gene induction (Wu et al., 2003). To date, no allele of *ABI1* and *ABI2* in which the corresponding protein is not produced has been reported.

*AtPP2CA* has been shown to block ABA-induced gene induction when transiently overexpressed in protoplasts (Sheen, 1998). However, stable *AtPP2CA* overexpression in planta and corresponding knock-out mutants have not been reported. In this study, the

isolation and detailed phenotypic characterization of PP2CAox and two insertional mutants, *pp2ca-1* and *pp2ca-2*, unequivocally demonstrate that PP2CA acts as a strong negative regulator of ABA signaling not only at the seed germination level but also in vegetative tissues. It has been previously shown that *AtPP2CA* transcripts are reduced in the ABA-hypersensitive mutant *abh1*, which encodes a nuclear mRNA cap-binding protein (Hugouvieux et al., 2001). We therefore also analyzed *AtPP2CA* mRNA splicing and the effects of the introduction of a 35S::*AtPP2CA* fusion in the *abh1* background to determine the role of *AtPP2CA* in mediating the ABA hypersensitivity of *abh1*.

## RESULTS

### Screening of a 35S::cDNA-Expressing Line Library for ABA-Insensitive Mutants Identifies an Overexpressor of *AtPP2CA*

Because studies suggest genetic and network redundancy in ABA signal transduction, a screen was pursued that can include dominant ABA-insensitive mutants. Roughly one million seeds were screened from approximately 60,000 activation-tagged lines for ABA-insensitive seed germination at 5  $\mu$ M ABA. Remarkably, after retesting putative mutants, no robust ABA-insensitive mutant line was isolated from the two activation-tagged populations that were tested twice independently. These findings may be attributed to the robustness of ABA signaling and the inherent limitations in the mutation rate of activation-tagged lines (Weigel et al., 2000).

To further enhance the frequency of obtaining dominant mutants in ABA signaling, we screened Arabidopsis lines expressing a library of random 35S::cDNAs (LeClere and Bartel, 2001) for ABA-insensitive mutants in seed germination assays. This library is composed of approximately 400,000 T2 seeds coming from 33,000 different T1 lines in the ecotype Columbia (Col-0) background, each expressing a random 35S::cDNA together with a Basta resistance gene inserted in the T-DNA of the 35SpBARN vector. From the original screen of 400,000 T2 seeds, 902 putative individual T2 seeds were able to germinate (radicle emergence plus expanded green cotyledons) 5 d after stratification on 2.5  $\mu$ M ABA, a concentration that completely inhibits wild-type seed germination. These young ABA-resistant seedlings were then transferred to soil and allowed to self pollinate. The progeny of each of the T2 plants were then tested again for ABA insensitivity in germination assays. From this secondary screen, the progeny of two T2 plants, named 54.7 and 393.1, exhibiting strong ABA resistance were selected and further characterized.

Southern-blot analyses revealed that both mutants contained tandem T-DNA insertions at one locus each (data not shown). Segregation analyses of both ABA and Basta resistance in the next generations as well as in F<sub>1</sub> and F<sub>2</sub> populations from backcrosses indicated

that both ABA insensitivities were linked to the corresponding T-DNAs. However, whereas 393.1 was heterozygous for the mutation and its ABA insensitivity was dominant, 54.7 was homozygous and its phenotype was due to a single recessive nuclear mutation.


Because the strong ABA insensitivity of the recessive mutant 54.7 was reminiscent of the loss-of-function phenotype of the classic positive regulators of ABA responses *ABI3*, *ABI4*, and *ABI5*, a PCR-based diagnosis was performed on 54.7 genomic DNA (see "Materials and Methods"). This analysis and sequencing showed that 54.7 has a tandem T-DNA insertion located in the first intron of *ABI5* approximately 500 bp before the start codon (data not shown). Thus, considering its strong ABA insensitivity and the location of the T-DNA insertion, 54.7 is most likely a new allele of *abi5*.

The cDNA contained within the T-DNA of the dominant 393.1 mutant was identified by PCR and perfectly matches the full-length cDNA of *AtPP2CA*, encoding a PP2C (Fig. 1A). The *AtPP2CA* gene belongs to group A of the Arabidopsis *PP2C* gene family (Schweighofer et al., 2004). A homozygous T3 line, PP2CAox (see below), was recovered from the progeny of the heterozygous 393.1 plant, and northern-blot analysis revealed that *AtPP2CA* was indeed overexpressed in PP2CAox compared to wild type (Fig. 1B). PP2CAox clearly exhibited ABA insensitivity in seed germination assays as it germinated almost completely on 2.5  $\mu\text{M}$  ABA, whereas wild-type seeds do not germinate at all (Fig. 2A).

To further test whether the 35S::*AtPP2CA* fusion caused the ABA-insensitive phenotype in PP2CAox, the full-length *AtPP2CA* cDNA was cloned back in the 35SpBARN binary vector (LeClere and Bartel, 2001; see "Materials and Methods") and transformed into wild-type Arabidopsis plants. Fourteen independent T1 35S::*AtPP2CA* lines were recovered as well as 10 T1 35SpBARN lines (empty-vector controls) and the germination rate of their T2 seeds was analyzed in the presence of 2.5  $\mu\text{M}$  ABA. Whereas none of the 10 T1 35SpBARN lines exhibited ABA insensitivity in their progeny compared to untransformed wild-type Arabidopsis (data not shown), all 14 35S::*AtPP2CA* lines displayed ABA resistance similar to the original heterozygous 393.1 mutant seeds (Fig. 2C). Thus, we conclude that ectopic expression of the 35S::*AtPP2CA* fusion is responsible for the ABA insensitivity observed for PP2CAox seeds.

#### Seeds from Two Independent T-DNA Insertion Mutants for the *AtPP2CA* Gene Are Strongly ABA Hypersensitive


To further elucidate the role of *AtPP2CA* in ABA signaling, we isolated an insertion mutant from the Salk Institute Genomic Analysis Laboratory (SIGnAL; Alonso et al., 2003) database corresponding to donor stock number SALK\_028132. Sequencing of the T-DNA flanking region indicated that the T-DNA is located at the end of exon II (position +864 [numbering refers to



**Figure 1.** The ABA-insensitive 393.1 mutant is an overexpressor of *AtPP2CA* encoding a PP2C. A, Schematic representation of the 393.1 mutant T-DNA containing the full-length cDNA of *AtPP2CA*. The *AtPP2CA* cDNA is inserted at the original *XhoI* and *NotI* restriction sites of the 35SpBARN vector. Arrows indicate the direction of transcription. 35S-F and NOS-R are the primers used for amplifying the cDNA. BARN-LB1 is the primer used for T-DNA left-border sequencing. Note that the *XhoI* restriction site no longer exists because it has been filled with T4 polymerase during the original cloning procedure (LeClere and Bartel, 2001). B, Northern-blot analysis of *AtPP2CA* in wild-type, *abh1*, *pp2ca-1*, and PP2CAox leaves. Hybridization signals with *ACT17* cDNA (*ACT7*) were used for standardization of RNA and the value obtained from wild-type leaves was set to 1.

the ATG start codon]; Fig. 3A). Plants homozygous for the T-DNA insertion recovered by PCR genotyping were renamed *pp2ca-1*. Northern-blot analysis failed to detect a full-length *AtPP2CA* transcript in the *pp2ca-1* mutant (Fig. 1B), although a shorter transcript could be detected (Fig. 1B, asterisk). In addition, reverse transcription (RT)-PCR analysis did not result in any product corresponding to a full-length cDNA in the T-DNA disruption allele *pp2ca-1* (Fig. 3B).

ABA germination assays were carried out with seeds from *pp2ca-1* in parallel with seeds from the PP2CAox




**Figure 2.** Constitutive expression of *AtPP2CA* results in ABA insensitivity, while disruption of *AtPP2CA* causes ABA hypersensitivity during seed germination. **A**, Comparison of germination rates of wild-type (Col-0; black circles), *pp2ca-1* (white triangles), *abh1* (black squares), and PP2CAox (white circles) seeds exposed to 0, 0.3, 0.5, 1, and 2.5  $\mu\text{M}$  ABA at 5 d. Data represent the mean  $\pm$  SEM of three independent experiments with 36 seeds per genotype and experiment. See Supplemental Figure 1A for similar data with *pp2ca-2*. Error bars are smaller than symbols, if not visible. **B**, Comparison of germination for wild-type, *pp2ca-1*, *abh1*, and PP2CAox seeds at 1  $\mu\text{M}$  ABA or in the absence of ABA after 5 d. **C**, Germination of seeds from wild-type and 14 individual T1 lines with constitutive 35S::*AtPP2CA* expression at 2.5  $\mu\text{M}$  ABA 5 d after germination.

line, wild-type plants, and the ABA-hypersensitive mutant, *abh1* (Fig. 2A). In the absence of exogenous ABA, *pp2ca-1* mutant seeds germinated as well as wild-type seeds (Fig. 2, A and B). And in the presence of 1  $\mu\text{M}$  ABA, a clear ABA-hypersensitive inhibition of seed germination was observed. Indeed, the ABA hypersensitivity of *pp2ca-1* was almost as strong as that of *abh1* (Fig. 2, A and B). Later, a second T-DNA insertion mutant line WiscDsLox341D03 was released by the Arabidopsis Biological Resource Center (ABRC; stock no. CS851888; Sussman et al., 2000). Plants homozygous for the T-DNA insertion recovered by PCR genotyping were renamed *pp2ca-2* and sequencing of the T-DNA flanking region showed that the insertion lay at the beginning of exon III (position +966; Fig. 3A). As for *pp2ca-1*, RT-PCR failed to amplify a product corresponding to a full-length cDNA in the *pp2ca-2* mutant (Fig. 3B). Seeds from *pp2ca-2* were also tested in germination assays and displayed a strong ABA hypersensitivity very similar to that of *pp2ca-1* (Supplemental Fig. 1A). Together, our data obtained from PP2CAox, as well as the *pp2ca-1* and *pp2ca-2* T-DNA disruption mutants, demonstrate an important role of *AtPP2CA* as a negative regulator of ABA signaling during seed germination.

#### *AtPP2CA* Affects ABA-Promoted Inhibition of Root Growth

To test whether *AtPP2CA* gene disruption or constitutive expression of *AtPP2CA* in plants could affect other ABA responses, and because *AtPP2CA* was shown to be expressed in the stele of the Arabidopsis root system (Cherel et al., 2002), we investigated ABA inhibition of root growth by transferring 6-d-old seedlings on 0.25  $\times$  Murashige and Skoog plates with 0, 2.5, 5, 10, 25, and 50  $\mu\text{M}$  ABA. Elongation of the primary root was measured 6 d after the transfer in three independent experiments (Fig. 3C). Disruption of *AtPP2CA* in *pp2ca-1* and *pp2ca-2* plants exhibits a moderate, but significant, increase in ABA sensitivity compared to wild type during root growth on 0.25  $\times$  Murashige and Skoog media supplemented with 2.5, 5, 10, 25, and 50  $\mu\text{M}$  ABA (Fig. 3C). Interestingly, ABA inhibition of root elongation in the originally isolated PP2CAox plants was the same as in wild type at all ABA concentrations measured ( $P > 0.04$  to 0.36 in all conditions tested; Fig. 3C), suggesting that elevated *AtPP2CA* transcript levels have no dramatic effect on ABA regulation of root elongation. A possible explanation could be that during the ABA inhibition of root elongation, the expression of PP2CA interacting partners is the rate-limiting step rather than PP2CA levels. Moreover, due to the moderate ABA hypersensitivity of *pp2ca-1* and *pp2ca-2* in root assays compared to the strong seed germination phenotype, it is conceivable that partial redundancy with other PP2Cs is more pronounced in roots. When plants were exposed to 50  $\mu\text{M}$  ABA for extended time periods (18 d), *pp2ca-1* plants show



**Figure 3.** Disruption of *AtPP2CA* causes reduced root elongation in response to ABA. A, Schematic representation of the genomic organi-

ABA-hypersensitive inhibition of root elongation as well as retarded growth and chlorosis in aerial parts of plants (Fig. 3D), further exhibiting an enhanced sensitivity to ABA.

#### The *AtPP2CA* Transcript Level Is Up-Regulated by Both ABA and Drought Treatments

*AtPP2CA* is expressed ubiquitously in plant organs with the highest transcript levels in leaves and its expression is up-regulated by stresses, including ABA, cold, drought, and salt treatment (Tahtiharju and Palva, 2001; Cherel et al., 2002). However, the time courses of *AtPP2CA* gene induction in ABA or drought experiments have not yet been reported, to our knowledge. Thus, to assess the stress-related regulation of this gene, we studied the accumulation of *AtPP2CA* transcripts from wild-type plant leaves in response to ABA and drought treatments. The results clearly show that the *AtPP2CA* mRNA is highly and rapidly up-regulated by both treatments (Fig. 4). Significant transcript increases were detected within 30 min of exposure to ABA or drought (Fig. 4). However, whereas the peak of ABA induction occurred at 1 h, induction by drought increased rather progressively before reaching a peak at approximately 6 h.

#### *AtPP2CA* Disruption Causes ABA Hypersensitivity in Stomatal Guard Cells

The control of water loss by ABA is a crucial survival mechanism for plants during drought periods. To investigate the role of *AtPP2CA* in regulating water homeostasis, we measured the loss of fresh weight of detached rosette leaves (Fig. 5A). Overexpression of the *AtPP2CA* cDNA in planta led to an approximately 1.5-fold increase in the water-loss rate (Fig. 5A). In contrast, the gene disruption line *pp2ca-1* did not exhibit significant differences to transpiration


zation of the *AtPP2CA* gene with four exons (black boxes). Positions of the *pp2ca-1* and *pp2ca-2* T-DNA insertions are indicated and orientation of the left-border sequence of the respective T-DNAs is represented by broken arrows. B, RT-PCR analysis shows no full-length transcript in *AtPP2CA* T-DNA disruption lines *pp2ca-1* and *pp2ca-2*. PCR reactions were performed with oligonucleotides PP2CAEx1-F and PP2CAEx4-R (Table I) and samples were withdrawn from the reaction after 28, 32, and 36 cycles. Amplification of *EF1α* cDNA with primers EF1α-F and EF1α-R (Table I) was used for controls. C, Comparison of root elongation of wild-type (white bars), *pp2ca-1* (dark gray bars), *pp2ca-2* (light gray bars), and PP2CAox (black bars) seedlings; 6-d-old seedlings were transferred to plates supplemented with 0, 2.5, 5, 10, 25, and 50 μM ABA, and root elongation was monitored after 6 d. Each data point represents the mean of three independent experiments with eight seedlings each. Asterisks (\*) indicate a significant change between wild-type and *pp2ca-1* or *pp2ca-2* plants ( $P < 0.001$ ). D, Growth of wild type, *pp2ca-1*, and PP2CAox on 0.25 × Murashige and Skoog medium supplemented with 50 μM ABA. The photographs show plants on plates with 0 (top) and 50 μM ABA (bottom) 18 d after transfer of 6-d-old seedlings from 0.25 × Murashige and Skoog medium.

rates of wild-type leaves (Fig. 5A). Similarly, some stomatal ABA response mutants show no detached-leaf water-loss phenotype, including earlier findings on mutant alleles of the *PP2Cs* *ABI1*, *ABI2*, and *AtPP2C-HAB1* (Gosti et al., 1999; Merlot et al., 2001; Saez et al., 2004). This lack of a phenotype in detached-leaf wilting assays may be attributable to the limited resolution of this method.


Therefore, we more directly analyzed stomatal movement responses to ABA in loss- and gain-of-function *AtPP2CA* plants (Fig. 5, B and C). Compared to wild type, guard cells from PP2CAox plants exhibit a clear insensitivity in ABA-induced stomatal closure analyses. Guard cells in these plants show a significantly reduced response to 10  $\mu\text{M}$  ABA, which clearly results in stomatal closure in wild-type plants (Fig. 5B). In contrast, *AtPP2CA* gene disruption results in an ABA-hypersensitive stomatal closure response at 1  $\mu\text{M}$  ABA (Fig. 5B for *pp2ca-1*, and 5C for *pp2ca-2*). These data show that *AtPP2CA* plays an important role in ABA signal transduction events and the regulation of stomatal aperture. Analyses of stomatal aperture responses (Fig. 5, B and C) and ratios of stomatal apertures to stomatal heights illustrate the same findings in *AtPP2CA* disruption and gain-of-function lines (Supplemental Fig. 1, B and C).

#### Analysis of *AtPP2CA* mRNA Splicing in the *abh1* Mutant

Interestingly, the *AtPP2CA* mRNA was previously shown to exhibit a reduced mRNA level in the ABA-hypersensitive *abh1* mutant (Hugouvieux et al., 2001).



**Figure 4.** *AtPP2CA* transcripts are rapidly and highly up-regulated by both ABA and drought treatments. Northern-blot analyses of *AtPP2CA* in wild-type leaves either treated with ABA (50  $\mu\text{M}$ ; left) or excised and subjected to desiccation (right). Total RNA was extracted from leaves at times specified by the number above each lane. Hybridization signals with *ACTIN7* cDNA (*ACT7*) were used for standardization of equal amounts of RNA. Values obtained prior to the indicated treatments were set to 1.



**Figure 5.** *AtPP2CA* modulates the stomatal response to ABA. A, Ectopic expression of *AtPP2CA* in plants causes enhanced leaf evaporation rate compared to wild type. Loss of fresh weight of detached rosette leaves at the same developmental stages was measured for wild-type (black circles), *pp2ca-1* (white triangles), and PP2CAox plants (white circles) at the indicated time points. Data represent the mean of three independent experiments  $\pm$  SEM. B, Stomatal closing is ABA hypersensitive in *pp2ca-1* and ABA insensitive in PP2CAox plants. Stomatal aperture measurements of wild type (white bars), *pp2ca-1* (shaded bars), and PP2CAox (black bars) in response to 0, 1, and 10  $\mu\text{M}$  ABA. Data represent the mean of  $n = 4$  independent experiments  $\pm$  SEM with  $4 \times 50$  stomata per data point. Asterisks (\*) indicate significant changes between the indicated genotype and wild type ( $P < 0.001$ ). C, Stomatal closing is ABA hypersensitive in *pp2ca-2*. Stomatal aperture measurements of wild type (white bars) and *pp2ca-2* (shaded bars) in response to 0, 1, and 10  $\mu\text{M}$  ABA. Data represent the mean of two independent experiments  $\pm$  SEM with  $2 \times 50$  stomata per data point. Asterisks (\*) indicate significant changes between the indicated genotype and wild type ( $P < 0.001$ ). See Supplemental Figure 1, B and C, for stomatal aperture ratios from experiments in Figure 5, B and C.



Transcript levels were analyzed in northern-blot analyses in *abh1* and compared to those in wild-type controls. *AtPP2CA* transcripts normalized to *Actin7* mRNA were 3.3-fold lower in *abh1* compared to wild-type controls (Fig. 1B).


ABH1 is the Arabidopsis homolog of an 80-kD subunit of the dimeric mRNA cap-binding complex, which additionally consists of a 20-kD subunit, AtCBP20 (Hugouvieux et al., 2001; Kmiecik et al., 2002; Papp et al., 2004). In yeast (*Saccharomyces cerevisiae*) and human HeLa cells, the cap-binding complex was shown to participate in pre-mRNA splicing (Izaurralde et al., 1994; Lewis and Izaurralde, 1997; Fortes et al., 1999). We investigated whether the *AtPP2CA* transcript undergoes differential splicing in *abh1* compared to wild type. Differential splicing could contribute to the down-regulation of the *AtPP2CA* mRNA in *abh1* due to a reduced turnover of *AtPP2CA* transcript maturation

(Clark et al., 2002). We designed an RT-PCR approach to analyze qualitative and quantitative differences in *AtPP2CA* transcript maturation (Fig. 6A). Forward primers were selected for amplification of the full reading frame or for amplification of intron sequences with the corresponding reverse primer being located at the 3' part of the most downstream exon IV (Fig. 6A).


RT-PCR on DNase-treated total RNA from leaves yielded similar amplification product qualities and quantities for all four reactions (Fig. 6B). PCR product 1 resulted in a single band corresponding to the full-length reading frame. PCR product 2, with the forward primer location in the first intron, showed faint bands with sizes corresponding to a fully unspliced pre-mRNA (1,028 nucleotides). PCR product 2 also resulted in splice intermediates emerging from intron I independent of removal of intron II or intron III (936 nucleotides) and even removal of both introns II and III (844 nucleotides) from the pre-mRNA without remarkable differences between wild type and *abh1* (Fig. 6B). This holds equally true for PCR products 3 and 4 (Fig. 6B). Identical results were obtained for an analysis on RNA isolated from independently grown plants (data not shown). We conclude from these results that splicing of the *AtPP2CA* pre-mRNA is not affected in the *abh1* background and down-regulation of *AtPP2CA* transcripts is more likely caused by other mechanisms.

To elucidate whether elevated transcript levels of *AtPP2CA* can restore normal ABA sensitivity or even cause ABA insensitivity in *abh1*, we transformed wild-type (Col-0) and *abh1* plants with a *AtPP2CA* cDNA under the control of the cauliflower mosaic virus 35S promoter in a binary vector different from 35SpBARN (see "Materials and Methods"). Forty independent T1 plants were isolated each for wild type and for *abh1* and tested for ABA responses. Individual lines with single-insertion segregation patterns and the strongest ABA insensitivity in seed germination were selected to obtain homozygous lines. ABA germination assays were performed in triplicate and confirmed earlier findings (Fig. 2C) that introduction of a 35S::*AtPP2CA* fusion in the wild type always confers a strong ABA insensitivity in seed germination independent of the binary vector used (data not shown).

In the *abh1* background, ectopic expression of the *AtPP2CA* cDNA had a much weaker effect than in the wild-type background, with only two homozygous single-insertion lines being able to restore wild-type-like ABA responses during germination (Fig. 7A, white diamonds). Most of the 40 *abh1* mutant lines expressing the 35S::*AtPP2CA* construct exhibited a range of ABA sensitivities between *abh1* and wild type in T2 generation germination experiments (data not shown). Based on the hypothesis that *abh1* might affect *AtPP2CA* transcripts in these constitutively cDNA-expressing lines, we tested *AtPP2CA* transcript integrity in RT-PCR experiments (Fig. 7B). Amplification of the full-length reading frame resulted in a single band in all



**Figure 6.** Analysis of *AtPP2CA* intron splicing in wild type and *abh1*. A, Schematic representation of *AtPP2CA* genomic organization with exons (black boxes) and introns (lines between exons). Positions of primers (arrows) used for the RT-PCR analysis are indicated and resulting amplification products (dotted lines) are shown for each PCR reaction performed, numbered from 1 to 4. Product sizes for full-length amplification, corresponding to completely unspliced transcripts, or for smaller fragments, corresponding to fully or partially processed mRNAs, are given for each PCR reaction. B, DNA fragments of fully and partially spliced *AtPP2CA* mRNA amplification products (36 cycles) generated in PCR reactions 1 to 4 (Fig. 6A) for wild-type controls (C) and *abh1* (a) are size fractionated on a 1.8% agarose gel. Amplification (28 cycles) of the *EF1 $\alpha$*  was used as internal control. Sizes of coelectrophoresed DNA standard fragments are given in kilobase pairs.



**Figure 7.** Ectopic expression of *AtPP2CA* partially suppresses the ABA hypersensitivity of *abh1*. **A**, Comparison of germination rates of wild type (black circles), PP2CAox in wild type (white circles), *abh1* (black squares), and seeds from three independent *abh1* 35S::*AtPP2CA* (white diamonds) lines germinated on Murashige and Skoog plates containing 0, 0.3, 0.5, 1, and 2.5  $\mu\text{M}$  ABA after 5 d. Data represent the mean  $\pm$  SEM of three independent experiments. Error bars are smaller than symbols, if not visible. Data for wild type, *abh1*, and PP2CAox in A are the same as in Figure 2A for reference purposes. **B**, RT-PCR analysis of *AtPP2CA* transcript levels in wild type, *abh1*, and the three independent *abh1* 35S::*AtPP2CA* lines displayed in A from the more sensitive to the more ABA resistant. DNA fragments from RT-PCR analyses of *EF1 $\alpha$*  were used for standardization of equal amplification rates and the value obtained for *AtPP2CA* quantity in *abh1* was set to 1. RT-PCR reactions were performed for 20, 24, 28, and 32 cycles to quantify relative mRNA levels and representative images after 28 cycles are shown.

lines investigated, indicating no detectable qualitative impact of *abh1* on the full-length transcripts (Fig. 7B). Ratiometric analysis of the *AtPP2CA* transcript levels, with *elongation factor 1 $\alpha$*  (*EF1 $\alpha$* ) as a reference, indicated that *AtPP2CA* mRNA abundance in 35S::*AtPP2CA*/*abh1* lines was indeed increased up to 1.5-fold compared to *abh1* (Fig. 7B). However, the transcript abundance was persistently lower than in wild-type plants (Fig. 7B). Northern analyses of *AtPP2CA* transcript levels confirmed these results (data not shown). Thus, elevated *AtPP2CA* transcript levels can partially suppress the ABA hypersensitivity of *abh1* and the overall lower levels of *AtPP2CA* transcripts in *abh1* compared to wild type indicate a continued down-regulation of *AtPP2CA* in *abh1* (Fig. 7).

## DISCUSSION

Here we report isolation of a strong dominant ABA response mutant overexpressing the *AtPP2CA* cDNA

during an ABA-insensitive screen of a library of 33,000 35S::cDNA-expressing Arabidopsis lines. We characterize *AtPP2CA* gene disruption and overexpression phenotypes in Arabidopsis. We show that T-DNA insertions in the *AtPP2CA* gene result in a strongly increased sensitivity to the phytohormone ABA during seed germination (Fig. 2, A and B; Supplemental Fig. 1A) and also render guard cells more sensitive to ABA during stomatal closure at 1  $\mu\text{M}$  ABA (Fig. 5, B and C) and affect root elongation in the presence of exogenous ABA (Fig. 3, C and D). On the other hand, overexpression of *AtPP2CA* impairs stomatal closure in response to 10  $\mu\text{M}$  ABA (Fig. 5B). Seed germination of PP2CAox lines displays a greatly decreased sensitivity to ABA (Fig. 2, A and C). Moreover, constitutive expression of *AtPP2CA* in the ABA-hypersensitive mutant *abh1* is shown to partially restore ABA sensitivity in *abh1* (Fig. 7A). Together, these results point to an important function of *AtPP2CA* as a negative regulator of ABA signal transduction events. The identification of a negative regulator in ABA signaling based on a cDNA overexpression screen shows that this approach can be used to isolate mutants in genes that modulate complex signaling networks in plants (Schroeder et al., 2001; Fedoroff, 2002; Finkelstein et al., 2002; Hetherington and Woodward, 2003; Himmelbach et al., 2003; Fan et al., 2004; Cutler and McCourt, 2005).

In an earlier study, *AtPP2CA* was shown to be linked to cold acclimation in Arabidopsis (Tahtiharju and Palva, 2001). It was shown that *AtPP2CA* is highly induced during cold acclimation. Plants with suppression of *AtPP2CA* transcripts by an antisense approach were shown to exhibit an ABA-dependent accelerated development of freezing tolerance (Tahtiharju and Palva, 2001) and an increased ABA sensitivity in seed germination was mentioned. As the antisense construct used shows nucleotide homology to five of the nine group A PP2Cs, and to avoid the possible effects of cosuppression of these related group A PP2C genes, we pursued analyses of T-DNA disruption lines of the *AtPP2CA* gene (Fig. 3A). Accordingly, northern-blot and RT-PCR analyses of the *pp2ca-1* and the *pp2ca-2* lines show absence of mature *AtPP2CA* transcripts (Figs. 1B and 3B).

### *AtPP2CA*, a Negative Regulator of Physiological ABA Responses

We report that *AtPP2CA* gene disruption lines show a strongly increased sensitivity to ABA during seed germination, which appears to be more pronounced than in a *AtP2C-HAB1* disruption line (Leonhardt et al., 2004; Saez et al., 2004). Thus, we investigated the effect of *AtPP2CA* gene disruption and ectopic expression at the whole-plant level.

Application of exogenous ABA is well established to affect root growth as an antagonist of auxin, impairing cell elongation and causing an arrest in mitotic cell cycle activity (Himmelbach et al., 1998). The ABA-insensitive dominant *abi1-1* mutant and sustained ex-



pression of *AtP2C-HAB1* in 35S::*AtP2C-HAB1* plants have been shown to exhibit less sensitivity to ABA inhibition of root growth (Beaudoin et al., 2000; Ghassemian et al., 2000; Saez et al., 2004). However, previous studies have not analyzed *PP2C* gene disruption lines for altered ABA-dependent root elongation. In our study, we show that *AtPP2CA* gene disruption results in a moderate hypersensitive response to ABA in root elongation compared to wild type under the conditions tested (Fig. 3, C and D).

Furthermore, we show that ectopic expression of *AtPP2CA* results in increased transpiration rates of detached rosette leaves (Fig. 5A). It has been shown previously that the dominant ABA-insensitive mutants *abi1-1* and *abi2-1* are sensitive to water stress conditions and impair early ABA signal transduction (Koornneef et al., 1984; Finkelstein, 1994; Leung et al., 1994, 1997; Meyer et al., 1994; Pei et al., 1997; Allen et al., 1999). More recently, plants overexpressing the *AtP2C-HAB1* protein phosphatase were also shown to exhibit increased transpiration rates (Saez et al., 2004). However, we could not observe a difference in the transpiration rates of detached leaves from the *pp2ca-1* plants (Fig. 5A), resembling findings on the *abi1-1R1* to *abi1-1R7* intragenic revertant lines and the T-DNA insertion line of *AtP2C-HAB1* (Saez et al., 2004), even though these mutants show ABA-hypersensitive responses in stomatal movements (Gosti et al., 1999; Leonhardt et al., 2004). Previous leaf water loss analyses from detached leaves have shown that these assays can show phenotypic differences in mutations in which steady-state stomatal apertures already differ from wild-type controls prior to excising leaves (Kwak et al., 2001; Cominelli et al., 2005; Liang et al., 2005). More direct (double-blinded) analyses of ABA-induced stomatal closing show that *AtPP2CA* disruption causes ABA-hypersensitive stomatal closing (Fig. 5, B and C), which also appears to be more pronounced than in the *AtP2C-HAB1* gene disruption line (Leonhardt et al., 2004). Together, these results emphasize the importance of *AtPP2CA* as a strong negative regulator of ABA signal transduction during seed germination and stomatal closure.

#### *AtPP2CA* Shows a Similar Expression Profile to Other Group A *PP2C* Genes

Sixty-nine *PP2C* genes are encoded in the Arabidopsis genome compared to 15 *PP2C* genes in humans (Kerk et al., 2002; Schweighofer et al., 2004). This implies genetic redundancy and a more specific role of plant *PP2Cs* during developmental stages and responses to environmental changes. With the exception of *AtP2C-HAB1*, no gene disruption phenotype for other plant *PP2Cs* has been reported (Leonhardt et al., 2004; Saez et al., 2004). *AtP2C-HAB1* is highly expressed in guard cells in response to ABA and gene disruption causes ABA-hypersensitive stomatal closure (Leonhardt et al., 2004) and increased sensitivity

during seed germination (Leonhardt et al., 2004; Saez et al., 2004).

*AtPP2CA* transcript levels are the second highest expressed of all group A *PP2Cs* in dry seeds of Arabidopsis and are significantly down-regulated upon imbibition (Supplemental Fig. 2; Zimmermann et al., 2004; Nakabayashi et al., 2005). A similar, but overall lower, expression pattern can be seen for *AtP2C-HAB1* (Supplemental Fig. 2). Since other not-yet-characterized *PP2Cs* of group A also show high transcript levels in dry seeds and down-regulation during imbibition, an ABA-related function in seed germination can be anticipated for these genes. Moreover, *AtPP2CA* was found to be among the mRNAs with the highest levels of ABA-mediated induction in guard cells, showing an average 7-fold ABA induction (Leonhardt et al., 2004), again similar to the transcript abundance of *AtP2C-HAB1* (Supplemental Fig. 3). Guard cell expression of *AtPP2CA* and *AtP2C-HAB1* was also shown in independent studies, which investigated their promoter activities in planta with reporter gene fusions (Cherel et al., 2002; Saez et al., 2004).

Despite the large gene family of *PP2Cs* and similar expression patterns of *AtP2C-HAB1* and *AtPP2CA*, the

**Table 1.** Oligonucleotide primers used in this study

Primer Name	Sequence (5'-3')
PP2CAEx1-F	CAAATGGCTGGGATTTGTTGC
PP2CAInt1-F	CTCATGTACGTTAATTCCTCTGTTAG
PP2CAInt2-F	CGAATTGGTTAGTATGAATCAAGATGGC
PP2CAInt3-F	CTGATGATTATTGTTTTGTTTGATAGGT
PP2CAEx4-R	AAGACGACGCTTGATTATTCCTC
SALK-LBa1	TGGTTCACGTAGTGGGCCATCG
p745	AACGTCGCCAATGTGTTATTAAGTTGTC
PP2CAEx1Sa1	TACCGTCGACAAATGGCTGGGATTTGTTGC
PP2CAEx4Spe1	TCCACTAGTTAAGACGACGCTTGATTATCC
EF1a-F	GGCCACGTCGATTCTGGAAA
EF1a-R	GGCTTGGTTGGAGTCATCTT
35S-F	CGCACAATCCCACTATCCTTCGCAAG
NOS-R	GATAATCATCGCAAGACCGGCAACAGG
1'BAR-F	GCGCAAGACGTGACGTAAGTATCCG
1'BAR-R	CCCTCTAGAGTCGACCTGCAGGCATCG
BARN-LB1	GGGCCAGGCGGTGAAGGGCAATC
ACTIN7-F	GGCCGATGGTGAAGATATTCAGCCACTTG
ACTIN7-R	TCGATGGACCTGACTCATCGTACTCACTC
ABI3F1	CGGTTTTAGATTACTTATTAGC
ABI3R1	CCACCCGCTAGTCTTCTTGCC
ABI3F2	GGCGGTGGTAAAGAACGATGAAGC
ABI3R2	CCGAGGTTACCCACGTCGC
ABI3F3	GGGTAACCTCGGAAGGATCG
ABI3R3	CCCATGCATGCACGAGAAG
ABI4F1	GGGATGCTCATCGTATATAATATG
ABI4R1	GGACCCCTTAGCTTCCCAAC
ABI5F1	GGGAACACTAGTAAAGCAG
ABI5R1	CCGCTTTGTAGGAAGACTGTTG
ABI5F2	GCGGAGCTGGAAGTGTCAAAG
ABI5R2	CCGCCTCACCATTATC
ABI5F3	GCGCCATGGACGACTCTACTTCTCGC
ABI5R3	GGTAACGAAAACCTTTATTGG

limited functional redundancy in single gene disruption lines during the process of ABA signal transduction in seeds and guard cells may imply a high degree of specificity toward downstream targets of these two PP2Cs. In plants, our knowledge about PP2C targets is still limited and no target has been identified for AtPP2C-HAB1. However, the inward-rectifying potassium channel AKT2 was shown to interact with AtPP2CA in yeast and AKT2 channel activity is negatively modulated by AtPP2CA in heterologous expression systems (Cherel et al., 2002). Also, the ABI2 protein phosphatase has been found to interact in yeast with the protein kinase PKS3 (Guo et al., 2002). In addition, ABI1 can interact with the ABA-inducible homeodomain transcription factor AtHB6 (Himmelbach et al., 2002). Moreover, phospholipase D $\alpha$ 1 (PLD $\alpha$ 1)-derived phosphatidic acid has been shown to bind and regulate ABI1 (Zhang et al., 2004).

#### Modulation of AtPP2CA in *abh1*

The *abh1* mutation causes ABA hypersensitivity in seed germination and stomatal movements and modulates ion channel activities in guard cells (Hugouvieux et al., 2001, 2002). Down-regulation of the AtPP2CA transcript level was previously reported in the *abh1* mutant (Hugouvieux et al., 2001). Because a protein homologous to ABH1, the 80-kD subunit of the dimeric nuclear cap-binding protein CBP80, has been shown to affect splicing during pre-mRNA maturation in yeast and mammalian cells (Izaurrealde et al., 1994; Lewis and Izaurrealde, 1997; Fortes et al., 1999; Clark et al., 2002), we investigated the hypothesis that the AtPP2CA transcript undergoes differential splicing in *abh1* compared to wild type, therefore causing down-regulation of the transcript. An RT-PCR analysis specifically designed to amplify splice intermediates based on differential intron splicing efficiencies in *abh1* and wild type did not reveal any differences in pre-mRNA splicing of AtPP2CA (Fig. 6). Therefore, pre-mRNA splicing seems unlikely to cause down-regulation of the AtPP2CA transcript, an effect that was observed for some transcripts in the yeast *gcr3* mutant that encodes the CBP80 protein (Clark et al., 2002).

In this study, we investigated the hypothesis that AtPP2CA down-regulation in *abh1* contributes to the ABA hypersensitivity in *abh1*. Constitutive expression of AtPP2CA in the *abh1* mutant suppressed the ABA hypersensitivity of *abh1* plants. Interestingly, however, overexpression did not render *abh1* plants as ABA-insensitive as 35S::AtPP2CA wild-type plants (Fig. 7A). Out of 40 *abh1* plants harboring the 35S::AtPP2CA construct, only two homozygous single-insertion lines resulted in ABA sensitivity similar to wild-type plants in seed germination experiments. The comparison of AtPP2CA transcript levels by RT-PCR in these gain-of-function lines revealed that the AtPP2CA transcript levels were up to 1.5-fold higher than the AtPP2CA transcript level in *abh1*, but still significantly lower than in wild-type plants (Fig. 7B). Because the identi-

fication of strong AtPP2CA gain-of-function phenotypes in the *abh1* background proved substantially more difficult than in wild type, a negative feedback mechanism may limit AtPP2CA expression in *abh1*. With AtPP2CA gene disruption lines being less ABA hypersensitive in seed germination than *abh1* (Fig. 2, A and B) and because AtPP2CA overexpression only partially restores wild-type-like ABA sensitivity in *abh1*, we conclude that additional mechanisms contribute to ABA hypersensitivity in *abh1*.

In conclusion, we demonstrate that the protein phosphatase AtPP2CA acts as a strong negative regulator of ABA signal transduction during seed germination (Fig. 2A) and the regulation of stomatal closure (Fig. 5B). Yoshida et al. (2006) have conducted an independent screen for ABA signaling components in Arabidopsis. They have characterized the same protein phosphatase AtPP2CA also showing ABA hypersensitivity in loss-of-function mutants and insensitivity in AtPP2CA overexpressors. Despite the large number of PP2C genes in the Arabidopsis genome, this study demonstrates that loss- and gain-of-function of AtPP2CA causes strong modulation of ABA responses. With the negative regulatory role of PP2CA in ABA signal transduction, a challenging question for future research will be to uncover the interacting proteins of PP2CA.

## MATERIALS AND METHODS

### Plant Material, Growth Conditions, and Transformation

Arabidopsis (*Arabidopsis thaliana*) plants (Col-0) were grown in a Conviron growth chamber (Controlled Environments Limited) in plastic pots filled with ready-to-use soil (Professional Blend). After sowing, pots were kept at 4°C for 4 to 7 d. Growing conditions were 22°C, 75% humidity, with a 16-h-light/8-h-dark photoperiod regime at approximately 75  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . Seeds used for comparative studies were from plants grown and harvested in parallel.

Seeds of the activation-tagged lines for identification of ABA-insensitive mutants were kindly provided by D. Weigel (Max-Planck-Institute for Developmental Biology, Tuebingen, Germany; Weigel et al., 2000) and by W. Scheible (Carnegie Institution, Stanford, CA; Sedbrook et al., 2004). Seeds from the library of 35S::cDNA Arabidopsis-expressing lines (LeClere and Bartel, 2001) were obtained from ABRC and correspond to the CS84450 stock number. The binary vectors constructed and described below were introduced into *Agrobacterium tumefaciens* strain GV3101 by electroporation, which was then used to transform wild-type (Col-0) or *abh1* plants by floral dipping (Clough and Bent, 1998).

### Molecular Characterization of a New *abi5* Allele and *pp2ca-1* and *pp2ca-2* Insertional Mutants

Given the recessive nature of the strong ABA insensitivity in the homozygous 54.7 mutant, we hypothesized that its phenotype could be mediated by disruption of positive regulators such as *ABI3*, *ABI4*, and *ABI5*. To test this hypothesis, PCR on genomic DNA from 54.7 and wild-type plants with a set of specific primers for *ABI3*, *ABI4*, and *ABI5* (Table I) was carried out. Only one PCR reaction with ABI5-F3/ABI5-R2 primers (Table I) did not lead to any products from 54.7 genomic DNA (data not shown). PCR genotyping reactions from 54.7 genomic DNA were then carried out with ABI5-F3/ABI5-R2 and with BARN-LB1 (Table I; Fig. 1A), a specific primer of the 35SpBARN T-DNA left border. This led to amplification of two PCR products whose sequencing showed that a tandem T-DNA insertion occurred in the first intron of *ABI5* approximately 500 bp before the start codon.

*pp2ca-1* and *pp2ca-2* mutants of the *AtPP2CA* gene (At3g11410) were obtained from ABRC and correspond to the SALK\_028132 and WiscDsLox341D03 lines, respectively. Genotyping PCR reactions for *pp2ca-1* and *pp2ca-2* were performed with PP2CAEx1-F/PP2CAEx4-R primers and with PP2CAEx1-F/SALK-LBa1 primers (SALK\_028132) or with PP2CAEx1-F/p745 primers (WiscDsLox341D03) and amplified products were sequenced (Table I).

### Molecular Characterization of the PP2CAox Mutant and Generation of the Reconstructed 35S::AtPP2CA Lines

The cDNA within the T-DNA of 393.1 plants was PCR amplified from genomic DNA with 35S-F/NOS-R primers (Table I; Fig. 1A) and purified with the QIAEX II kit (Qiagen). The amplified DNA was then cloned into pGEM-T Easy vector (Promega) and sequenced. Because the PCR fragment matched the full-length cDNA of *AtPP2CA* perfectly, it was then excised from pGEM-T Easy with *Sma*I and *Not*I restriction enzymes, subcloned into *Sma*I/*Not*I linearized 35SpBARN vector (Fig. 1A; LeClere and Bartel, 2001), and then used to transform *Arabidopsis* wild-type plants.

Vector PS173 (kindly provided by Professor Jeff F. Harper, University of Nevada, Reno) was used to constitutively express *AtPP2CA* cDNA in wild-type and *abh1* plants. After amplification from total wild-type (Col-0) cDNA (first-strand cDNA synthesis kit; Amersham Biosciences) and cloning the *AtPP2CA* cDNA into the pGEM-T Easy vector (Promega), the sequenced cDNA was excised with *Sa*II and *Spe*I restriction enzymes and subcloned into the PS173 vector previously digested with *Xho*I and *Xba*I restriction enzymes.

### Northern-Blot and RT-PCR Analyses

Total RNA was extracted from leaves using TRIzol reagent (Life Technologies/Gibco-BRL) and quantified by absorption and migration of an aliquot on agarose gel. For ABA and drought treatments, rosette leaves of 3- to 4-week-old wild-type plants were either sprayed with 50  $\mu$ M ABA or excised and subjected to desiccation for 0.5, 1, 2, 3, 6, or 12 h before extraction. Fifteen micrograms of total RNA were separated in a denaturing formaldehyde-agarose gel and blotted to a Hybond-N membrane (Amersham-Pharmacia). Blots were hybridized with random-priming <sup>32</sup>P-labeled probes (Megaprime DNA labeling system; Amersham-Pharmacia). *AtPP2CA* and *ACTIN7* probes were amplified by PCR from cDNA using PP2CAEx1-F/PP2CAEx4-R and ACTIN7-F/ACTIN7-R primers, respectively (Table I). PCR fragments were purified using the QIAEX II kit (Qiagen).

RT-PCR experiments were performed on total RNA isolated as described above after DNase I treatment (DNA-free; Ambion). Reverse transcription (first-strand cDNA synthesis kit, Amersham Biosciences) was performed on 2.5  $\mu$ g of RNA and 2  $\mu$ L were used for PCR reactions (*Ex Taq* DNA polymerase; TaKaRa Mirus Bio). Samples were withdrawn after 20, 24, 28, and 32 cycles (splicing) or 28, 32, and 36 cycles (T-DNA disruption lines) and products were analyzed by agarose gel electrophoresis. Hybridization/PCR signals were quantified using Adobe Photoshop 5.5 software (Adobe Systems) after subtraction of background levels. Expression levels for northern-blot and RT-PCR analyses were normalized against the corresponding *ACTIN7* and *EF1 $\alpha$*  RNA levels, respectively.

### Root Growth and Germination Assays

For ABA germination assays, sterilized seeds were plated on minimal medium (0.25 $\times$  Murashige and Skoog medium, no Suc) supplemented with increasing ABA concentrations. After stratification of 4 d at 4°C, plates were transferred to a Conviron growth chamber (Controlled Environments Limited). To score seed germination, the percentage of seeds that had germinated and developed fully green expanded cotyledons was determined in three independent experiments (36 seeds per genotype and experiment).

Root growth assays to assess ABA sensitivity were carried out by transferring 6-d-old seedlings onto minimal medium (0.25 $\times$  Murashige and Skoog medium, no Suc) supplemented with the indicated ABA concentrations on 0.8% agar (Phytigel; Sigma) plates. Root growth was measured 6 d after the transfer in three independent experiments with eight individuals per genotype and experiment. *t*-Test (one-tailed, homoscedastic) *P*-values are as follows: *pp2ca-1*, *P* = 3.6E-07 (2.5  $\mu$ M ABA); *P* = 3.3E-06 (5  $\mu$ M ABA); *P* = 4.6E-04 (10  $\mu$ M ABA); *P* = 1.6 E-07 (25  $\mu$ M ABA); and *P* = 6.8E-06 (50  $\mu$ M ABA). *pp2ca-2*, *P* = 9.7E-06 (2.5  $\mu$ M ABA); *P* = 6.5E-06 (5  $\mu$ M ABA); *P* = 7.2E-05 (10  $\mu$ M ABA); *P* = 8.9E-13 (25  $\mu$ M ABA); and *P* = 1.2E-07 (50  $\mu$ M ABA).

### Leaf Water Loss and Stomatal Closure Measurements

Time-dependent analyses of loss of fresh weight were performed with detached rosette leaves at the same developmental stage and size from single 3-week-old plants. Three leaves per genotype were excised, kept in the Conviron growth chamber (Controlled Environments Limited), and fresh weight was measured at the indicated periods of time in three independent experiments.

Double-blind stomatal movement assays were performed such that the genotype and applied ABA concentrations were unknown. Stomatal responses were analyzed in 3- to 4-week-old plants grown in a Conviron growth chamber. Leaves were floated for 2.5 h in stomatal opening solution (Pei et al., 1997) containing 50 mM KCl, 50  $\mu$ M CaCl<sub>2</sub>, and 10 mM MES (pH 6.15). After incubation in ABA for 2.5 h, leaves were blended and the stomatal aperture was measured. Control experiments were performed in parallel with no ABA added. *t*-Test (one-tailed, homoscedastic) *P*-values were calculated: *P* = 2.3E-08 for *pp2ca-1* and *P* = 5.7E-13 for *pp2ca-2* at 1  $\mu$ M ABA, and *P* = 2.0E-09 for PP2CAox at 10  $\mu$ M ABA.

Sequence data from this article can be found in the GenBank/EMBL data libraries under accession number NM\_111974.

### ACKNOWLEDGMENTS

We thank David Waner and Majse Nafisi for screening the activation-tagging populations, Bonnie Bartel for providing the 35SpBARN vector as well as helpful discussions, Nadia Robert, Jean Colcombet, and Dongyul Sung for discussions, and Taehoun Kim and Jared Young for comments on the manuscript.

Received August 24, 2005; revised October 13, 2005; accepted October 23, 2005; published December 16, 2005.

### LITERATURE CITED

- Abe H, Urao T, Ito T, Seki M, Shinozaki K, Yamaguchi-Shinozaki K (2003) *Arabidopsis* AtMYC2 (bHLH) and AtMYB2 (MYB) function as transcriptional activators in abscisic acid signaling. *Plant Cell* **15**: 63–78
- Allen GJ, Kuchitsu K, Chu SP, Murata Y, Schroeder JI (1999) *Arabidopsis* *abi1-1* and *abi2-1* phosphatase mutations reduce abscisic acid induced cytoplasmic calcium rises in guard cells. *Plant Cell* **11**: 1785–1798
- Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Cheuk R, et al (2003) Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* **301**: 653–657
- Arabidopsis Genome Initiative (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* **408**: 796–815
- Beaudoin N, Serizet C, Gosti F, Giraudat J (2000) Interactions between abscisic acid and ethylene signaling cascades. *Plant Cell* **12**: 1103–1116
- Brocard-Gifford I, Lynch TJ, Garcia ME, Malhotra B, Finkelstein RR (2004) The *Arabidopsis thaliana* ABSCISIC ACID-INSENSITIVE8 encodes a novel protein mediating abscisic acid and sugar responses essential for growth. *Plant Cell* **16**: 406–421
- Cherel I, Michard E, Platet N, Moulins K, Alcon C, Sentenac H, Thibaud JB (2002) Physical and functional interaction of the *Arabidopsis* K(+) channel AKT2 and phosphatase AtPP2CA. *Plant Cell* **14**: 1133–1146
- Clark TA, Sugnet CW, Ares M Jr (2002) Genomewide analysis of mRNA processing in yeast using splicing-specific microarrays. *Science* **296**: 907–910
- Clough SJ, Bent AF (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* **16**: 735–743
- Cominelli E, Galbiati M, Vavasseur A, Conti L, Sala T, Vuylsteke M, Leonhardt N, Dellaporta SL, Tonelli C (2005) A guard-cell-specific MYB transcription factor regulates stomatal movements and plant drought tolerance. *Curr Biol* **15**: 1196–1200
- Cutler S, Ghassseman M, Bonetta D, Cooney S, McCourt P (1996) A protein farnesyl transferase involved in abscisic acid signal transduction in *Arabidopsis*. *Science* **273**: 1239–1241
- Cutler S, McCourt P (2005) Dude, where's my phenotype? Dealing with redundancy in signaling networks. *Plant Physiol* **138**: 558–559

- Fan LM, Zhao Z, Assmann SM (2004) Guard cells: a dynamic signaling model. *Curr Opin Plant Biol* 7: 537–546
- Fedoroff NV (2002) Cross-talk in abscisic acid signaling. *Sci STKE* 2002: RE10
- Finkelstein RR (1994) Maternal effects govern variable dominance of two abscisic acid response mutations in *Arabidopsis thaliana*. *Plant Physiol* 105: 1203–1208
- Finkelstein RR, Gampala SS, Rock CD (2002) Abscisic acid signaling in seeds and seedlings. *Plant Cell (Suppl)* 14: S15–S45
- Finkelstein RR, Lynch TJ (2000) The *Arabidopsis* abscisic acid response gene *ABI5* encodes a basic leucine zipper transcription factor. *Plant Cell* 12: 599–610
- Finkelstein RR, Wang ML, Lynch TJ, Rao S, Goodman HM (1998) The *Arabidopsis* abscisic acid response locus *ABI4* encodes an APETALA 2 domain protein. *Plant Cell* 10: 1043–1054
- Fortes P, Kufel J, Fornerod M, Polycarpou-Schwarz M, Lafontaine D, Tollervey D, Mattaj IW (1999) Genetic and physical interactions involving the yeast nuclear cap-binding complex. *Mol Cell Biol* 19: 6543–6553
- Ghasseman M, Nambara E, Cutler S, Kawaide H, Kamiya Y, McCourt P (2000) Regulation of abscisic acid signaling by the ethylene response pathway in *Arabidopsis*. *Plant Cell* 12: 1117–1126
- Giraudat J, Hauge BM, Valon C, Smalle J, Parcy F, Goodman HM (1992) Isolation of the *Arabidopsis* *ABI3* gene by positional cloning. *Plant Cell* 4: 1251–1261
- Gosti F, Beaudoin N, Serizet C, Webb AAR, Vartanian N, Giraudat J (1999) ABI1 protein phosphatase 2C is a negative regulator of abscisic acid signaling. *Plant Cell* 11: 1897–1910
- Guo Y, Xiong L, Song CP, Gong D, Halfter U, Zhu JK (2002) A calcium sensor and its interacting protein kinase are global regulators of abscisic acid signaling in *Arabidopsis*. *Dev Cell* 3: 233–244
- Hetherington AM, Woodward FI (2003) The role of stomata in sensing and driving environmental change. *Nature* 424: 901–908
- Himmelbach A, Hoffmann T, Leube M, Hohener B, Grill E (2002) Homeodomain protein ATHB6 is a target of the protein phosphatase ABI1 and regulates hormone responses in *Arabidopsis*. *EMBO J* 21: 3029–3038
- Himmelbach A, Iten M, Grill E (1998) Signaling of abscisic acid to regulate plant growth. *Philos Trans R Soc Lond B Biol Sci* 353: 1439–1444
- Himmelbach A, Yang Y, Grill E (2003) Relay and control of abscisic acid signaling. *Curr Opin Plant Biol* 6: 470–479
- Hoth S, Morgante M, Sanchez JP, Hanafey MK, Tingey SV, Chua NH (2002) Genome-wide gene expression profiling in *Arabidopsis thaliana* reveals new targets of abscisic acid and largely impaired gene regulation in the *abi1-1* mutant. *J Cell Sci* 115: 4891–4900
- Hugouvieux V, Kwak JM, Schroeder JI (2001) An mRNA cap binding protein, ABH1, modulates early abscisic acid signal transduction in *Arabidopsis*. *Cell* 106: 477–487
- Hugouvieux V, Murata Y, Young JJ, Kwak JM, Mackesy DZ, Schroeder JI (2002) Localization, ion channel regulation, and genetic interactions during abscisic acid signaling of the nuclear mRNA cap-binding protein, ABH1. *Plant Physiol* 130: 1276–1287
- Izaurrealde E, Lewis J, McGuigan C, Jankowska M, Darzynkiewicz E, Mattaj IW (1994) A nuclear cap binding protein complex involved in pre-mRNA splicing. *Cell* 78: 657–668
- Kerk D, Bulgrien J, Smith DW, Barsam B, Veretnik S, Gribskov M (2002) The complement of protein phosphatase catalytic subunits encoded in the genome of *Arabidopsis*. *Plant Physiol* 129: 908–925
- Kmieciak M, Simpson CG, Lewandowska D, Brown JWS, Jarmolowski A (2002) Cloning and characterization of two subunits of *Arabidopsis thaliana* nuclear cap-binding complex. *Gene* 283: 171–183
- Koornneef M, Reuling G, Karssen CM (1984) The isolation and characterization of abscisic acid insensitive mutants of *Arabidopsis thaliana*. *Physiol Plant* 61: 377–383
- Kwak JM, Moon JH, Murata Y, Kuchitsu K, Leonhardt N, DeLong A, Schroeder JI (2002) Disruption of a guard cell-expressed protein phosphatase 2A regulatory subunit, RCN1, confers abscisic acid insensitivity in *Arabidopsis*. *Plant Cell* 14: 2849–2861
- Kwak JM, Mori IC, Pei ZM, Leonhardt N, Torres MA, Dangl JL, Bloom RE, Bodde S, Jones JDC, Schroeder JI (2003) NADPH oxidase *AtrbohD* and *AtrbohF* genes function in ROS-dependent ABA signaling in *Arabidopsis*. *EMBO J* 22: 2623–2633
- Kwak JM, Murata Y, Baizabal-Aguirre VM, Merrill J, Wang M, Kemper A, Hawke SD, Tallman G, Schroeder JI (2001) Dominant negative guard cell K<sup>+</sup> channel mutants reduce inward-rectifying K<sup>+</sup> currents and light-induced stomatal opening in *Arabidopsis*. *Plant Physiol* 127: 473–485
- LeClere S, Bartel B (2001) A library of *Arabidopsis* 35S-cDNA lines for identifying novel mutants. *Plant Mol Biol* 46: 695–703
- Leonhardt N, Kwak JM, Robert N, Waner D, Leonhardt G, Schroeder JI (2004) Microarray expression analyses of *Arabidopsis* guard cells and isolation of a recessive abscisic acid hypersensitive protein phosphatase 2C mutant. *Plant Cell* 16: 596–615
- Leung J, Bouvier-Durand M, Morris PC, Guerrier D, Chedford F, Giraudat J (1994) *Arabidopsis* ABA response gene ABI1: features of a calcium-modulated protein phosphatase. *Science* 264: 1448–1452
- Leung J, Merlot S, Giraudat J (1997) The *Arabidopsis* ABSCISIC ACID-INSENSITIVE2 (*ABI2*) and *ABI1* genes encode homologous protein phosphatases 2C involved in abscisic acid signal transduction. *Plant Cell* 9: 759–771
- Lewis JD, Izaurrealde E (1997) The role of the cap structure in RNA processing and nuclear export. *Eur J Biochem* 247: 461–469
- Liang YK, Dubos C, Dodd IC, Holroyd GH, Hetherington AM, Campbell MM (2005) *AtMYB61*, an R2R3-MYB transcription factor controlling stomatal aperture in *Arabidopsis thaliana*. *Curr Biol* 15: 1201–1206
- Lu C, Fedoroff N (2000) A mutation in the *Arabidopsis* *HYL1* gene encoding a dsRNA binding protein affects responses to abscisic acid, auxin, and cytokinin. *Plant Cell* 12: 2351–2366
- Merlot S, Gosti F, Guerrier D, Vavasseur A, Giraudat J (2001) The ABI1 and ABI2 protein phosphatases 2C act in a negative feedback regulatory loop of the abscisic acid signalling pathway. *Plant J* 25: 295–303
- Meyer K, Leube MP, Grill E (1994) A protein phosphatase 2C involved in ABA signal transduction in *Arabidopsis thaliana*. *Science* 264: 1452–1455
- Mustilli AC, Merlot S, Vavasseur A, Fenzi F, Giraudat J (2002) *Arabidopsis* OST1 protein kinase mediates the regulation of stomatal aperture by abscisic acid and acts upstream of reactive oxygen species production. *Plant Cell* 14: 3089–3099
- Nakabayashi K, Okamoto M, Koshiba T, Kamiya Y, Nambara E (2005) Genome-wide profiling of stored mRNA in *Arabidopsis thaliana* seed germination: epigenetic and genetic regulation of transcription in seed. *Plant J* 41: 697–709
- Osakabe Y, Maruyama K, Seki M, Satou M, Shinozaki K, Yamaguchi-Shinozaki K (2005) Leucine-rich repeat receptor-like kinase1 is a key membrane-bound regulator of abscisic acid early signaling in *Arabidopsis*. *Plant Cell* 17: 1105–1119
- Papp I, Mur LA, Dalmadi A, Dulai S, Koncz C (2004) A mutation in the *Cap Binding Protein 20* gene confers drought tolerance to *Arabidopsis*. *Plant Mol Biol* 55: 679–686
- Pei ZM, Ghasseman M, Kwak CM, McCourt P, Schroeder JI (1998) Role of farnesyltransferase in ABA regulation of guard cell anion channels and plant water loss. *Science* 282: 287–290
- Pei ZM, Kuchitsu K, Ward JM, Schwarz M, Schroeder JI (1997) Differential abscisic acid regulation of guard cell slow anion channels in *Arabidopsis* wild-type and *abi1* and *abi2* mutants. *Plant Cell* 9: 409–423
- Rodriguez PL, Benning G, Grill E (1998) ABI2, a second protein phosphatase 2C involved in abscisic acid signal transduction in *Arabidopsis*. *FEBS Lett* 421: 185–190
- Saez A, Apostolova N, Gonzalez-Guzman M, Gonzalez-Garcia MP, Nicolas C, Lorenzo O, Rodriguez PL (2004) Gain-of-function and loss-of-function phenotypes of the protein phosphatase 2C *HAB1* reveal its role as a negative regulator of abscisic acid signalling. *Plant J* 37: 354–369
- Schroeder JI, Allen GJ, Hugouvieux V, Kwak JM, Waner D (2001) GUARD CELL SIGNAL TRANSDUCTION. *Annu Rev Plant Physiol Plant Mol Biol* 52: 627–658
- Schweighofer A, Hirt H, Meskiene I (2004) Plant PP2C phosphatases: emerging functions in stress signaling. *Trends Plant Sci* 9: 236–243
- Sedbrook JC, Ehrhardt DW, Fisher SE, Scheible WR, Somerville CR (2004) The *Arabidopsis* *SKU6/SPIRAL1* gene encodes a plus end-localized microtubule-interacting protein involved in directional cell expansion. *Plant Cell* 16: 1506–1520
- Seki M, Ishida J, Narusaka M, Fujita M, Nanjo T, Umezawa T, Kamiya A, Nakajima M, Enju A, Sakurai T, et al (2002) Monitoring the expression pattern of around 7,000 *Arabidopsis* genes under ABA treatments using a full-length cDNA microarray. *Funct Integr Genomics* 2: 282–291

- Seo M, Koshiba T** (2002) Complex regulation of ABA biosynthesis in plants. *Trends Plant Sci* 7: 41–48
- Sheen J** (1998) Mutational analysis of protein phosphatase 2C involved in abscisic acid signal transduction in higher plants. *Proc Natl Acad Sci USA* 95: 975–980
- Sussman MR, Amasino RM, Young JC, Krysan PJ, Austin-Phillips S** (2000) The *Arabidopsis* knockout facility at the University of Wisconsin-Madison. *Plant Physiol* 124: 1465–1467
- Tahtiharju S, Palva T** (2001) Antisense inhibition of protein phosphatase 2C accelerates cold acclimation in *Arabidopsis thaliana*. *Plant J* 26: 461–470
- Takahashi S, Seki M, Ishida J, Satou M, Sakurai T, Narusaka M, Kamiya A, Nakajima M, Enju A, Akiyama K, et al** (2004) Monitoring the expression profiles of genes induced by hyperosmotic, high salinity, and oxidative stress and abscisic acid treatment in *Arabidopsis* cell culture using a full-length cDNA microarray. *Plant Mol Biol* 56: 29–55
- Uno Y, Furihata T, Abe H, Yoshida R, Shinozaki K, Yamaguchi-Shinozaki K** (2000) *Arabidopsis* basic leucine zipper transcription factors involved in an abscisic acid-dependent signal transduction pathway under drought and high-salinity conditions. *Proc Natl Acad Sci USA* 97: 11632–11637
- Wang XQ, Ullah H, Jones AM, Assmann SM** (2001) G protein regulation of ion channels and abscisic acid signaling in *Arabidopsis* guard cells. *Science* 292: 2070–2072
- Weigel D, Ahn JH, Blazquez MA, Borevitz JO, Christensen SK, Fankhauser C, Ferrandiz C, Kardailsky I, Malancharuvil EJ, Neff MM, et al** (2000) Activation tagging in *Arabidopsis*. *Plant Physiol* 122: 1003–1014
- Wu Y, Sanchez JP, Lopez-Molina L, Himmelbach A, Grill E, Chua NH** (2003) The *abi1-1* mutation blocks ABA signaling downstream of cADPR action. *Plant J* 34: 307–315
- Xiong L, Gong Z, Rock CD, Subramanian S, Guo Y, Xu W, Galbraith D, Zhu JK** (2001a) Modulation of abscisic acid signal transduction and biosynthesis by an Sm-like protein in *Arabidopsis*. *Dev Cell* 1: 771–781
- Xiong L, Lee B, Ishitani M, Lee H, Zhang C, Zhu JK** (2001b) FIERY1 encoding an inositol polyphosphate 1-phosphatase is a negative regulator of abscisic acid and stress signaling in *Arabidopsis*. *Genes Dev* 15: 1971–1984
- Yamaguchi-Shinozaki K, Shinozaki K** (2005) Organization of cis-acting regulatory elements in osmotic- and cold-stress-responsive promoters. *Trends Plant Sci* 10: 88–94
- Yoshida R, Hobo T, Ichimura K, Mizoguchi T, Takahashi F, Aronso J, Ecker JR, Shinozaki K** (2002) ABA-activated SnRK2 protein kinase is required for dehydration stress signaling in *Arabidopsis*. *Plant Cell Physiol* 43: 1473–1483
- Yoshida T, Nishimura N, Kitahata N, Kuromori T, Ito T, Asami T, Shinozaki K, Hirayama T** (2006) ABA-Hypersensitive Germination3 encodes a protein phosphatase 2C (AtPP2CA) that strongly regulates abscisic acid signaling during germination among *Arabidopsis* PP2Cs. *Plant Physiol* 140: 115–126
- Zhang W, Qin C, Zhao J, Wang X** (2004) Phospholipase *Da1*-derived phosphatidic acid interacts with ABI1 phosphatase 2C and regulates abscisic acid signaling. *Proc Natl Acad Sci USA* 101: 9508–9513
- Zhu JK** (2002) Salt and drought stress signal transduction in plants. *Annu Rev Plant Biol* 53: 247–273
- Zimmermann P, Hirsch-Hoffmann M, Hennig L, Gruissem W** (2004) GENEVESTIGATOR. *Arabidopsis* microarray database and analysis toolbox. *Plant Physiol* 136: 2621–2632