

# Nonsteroidal anti-inflammatory drugs repress $\beta$ -secretase gene promoter activity by the activation of PPAR $\gamma$

Magdalena Sastre<sup>\*†</sup>, Ilse Dewachter<sup>‡</sup>, Steffen Rossner<sup>§</sup>, Nenad Bogdanovic<sup>¶</sup>, Evan Rosen<sup>||</sup>, Peter Borghgraef<sup>‡</sup>, Bernd O. Evert<sup>\*</sup>, Lucia Dumitrescu-Ozimek<sup>\*</sup>, Dietmar R. Thal<sup>\*\*</sup>, Gary Landreth<sup>††</sup>, Jochen Walter<sup>\*</sup>, Thomas Klockgether<sup>\*</sup>, Fred van Leuven<sup>‡</sup>, and Michael T. Heneka<sup>\*†††</sup>

Departments of <sup>\*</sup>Neurology and <sup>\*\*</sup>Neuropathology, University of Bonn, Sigmund Freud Strasse 25, 53127 Bonn, Germany; <sup>‡</sup>Experimental Genetics Group, Department of Human Genetics, Katholieke Universiteit Leuven, Campus Gasthuisberg O&N 06, B-3000 Leuven, Belgium; <sup>§</sup>Paul Flechsig Institute for Brain Research, Department of Neurochemistry, University of Leipzig, 04109 Leipzig, Germany; <sup>¶</sup>Karolinska Institute, Neurotec, Geriatric Department and Huddinge Brain Bank KFC, Novum, Plan4, 14186 Stockholm, Sweden; <sup>||</sup>Division of Endocrinology, Beth Israel Deaconess Medical Center, Boston, MA 02215; and <sup>††</sup>Alzheimer Laboratory, Case Western Reserve University, 10900 Euclid Avenue, Cleveland, OH 44106

Edited by Vernon Martin Ingram, Massachusetts Institute of Technology, Cambridge, MA, and approved November 9, 2005 (received for review May 9, 2005)

**Epidemiological evidence suggests that nonsteroidal anti-inflammatory drugs (NSAIDs) decrease the risk for Alzheimer's disease (AD). Certain NSAIDs can activate the peroxisome proliferator-activated receptor- $\gamma$  (PPAR $\gamma$ ), which is a nuclear transcriptional regulator. Here we show that PPAR $\gamma$  depletion potentiates  $\beta$ -secretase [ $\beta$ -site amyloid precursor protein cleaving enzyme (BACE1)] mRNA levels by increasing BACE1 gene promoter activity. Conversely, overexpression of PPAR $\gamma$ , as well as NSAIDs and PPAR $\gamma$  activators, reduced BACE1 gene promoter activity. These results suggested that PPAR $\gamma$  could be a repressor of BACE1. We then identified a PPAR $\gamma$  responsive element (PPRE) in the BACE1 gene promoter. Mutagenesis of the PPRE abolished the binding of PPAR $\gamma$  to the PPRE and increased BACE1 gene promoter activity. Furthermore, proinflammatory cytokines decreased PPAR $\gamma$  gene transcription, and this effect was suppressed by NSAIDs. We also demonstrate that *in vivo* treatment with PPAR $\gamma$  agonists increased PPAR $\gamma$  and reduced BACE1 mRNA and intracellular  $\beta$ -amyloid levels. Interestingly, brain extracts from AD patients showed decreased PPAR $\gamma$  expression and binding to PPRE in the BACE1 gene promoter. Our data strongly support a major role of PPAR $\gamma$  in the modulation of amyloid- $\beta$  generation by inflammation and suggest that the protective mechanism of NSAIDs in AD involves activation of PPAR $\gamma$  and decreased BACE1 gene transcription.**

amyloid | inflammation | Alzheimer's disease | transgenic mice

**F**ormation of amyloid peptides [amyloid- $\beta$  (A $\beta$ )] by neurons is generally thought to be a prime trigger of the pathogenesis of Alzheimer's disease (AD). The generation of A $\beta$  is initiated by a protease that cleaves a larger precursor protein, the  $\beta$ -amyloid precursor protein ( $\beta$ APP), at the N terminus side of the A $\beta$  peptide.  $\beta$ -secretase or  $\beta$ -site APP-cleaving enzyme (BACE1) was cloned and identified as a transmembrane aspartyl protease (1). BACE1 deficiency precludes A $\beta$  formation in transgenic mice (2) and does not cause or promote any neurological or phenotypic abnormalities. Moreover, BACE1 inactivation rescues memory deficits in transgenic mice (3), strongly supporting the importance of BACE1 as a therapeutic target in AD.

Several studies have demonstrated that BACE1 expression can be modulated by various factors. For instance, oxidative stress increases the expression and activity of BACE1 in NT2 neurons (4), whereas it is up-regulated in chronic models of gliosis (5) and experimental traumatic brain injury (6). In addition, BACE1 protein levels and the  $\beta$ -secretase product ( $\beta$ -C-terminal fragment) are increased in brain of sporadic AD patients (7). Although BACE1 expression increases with age in mice (8), there is no consensus regarding BACE1 mRNA levels in AD patients (7, 9). Evidence has been presented for regulation of BACE1 expression at the tran-

scriptional as well as the translational levels. The untranslated 5'-BACE1 transcript leader contains upstream ORFs that can reduce the translation of the main ORF (10). The BACE1 gene promoter region was recently cloned, and several putative transcription factor-binding sites were identified (11–13).

We recently demonstrated that BACE1 mRNA and protein levels are increased by proinflammatory mediators and down-regulated by nonsteroidal anti-inflammatory drugs (NSAIDs) (14). Because certain NSAIDs are agonists for peroxisome proliferator-activated receptor- $\gamma$  (PPAR $\gamma$ ) (15), we hypothesized that the protective mechanisms by which NSAIDs mediate reduction of A $\beta$  via BACE1 could involve activation of PPAR $\gamma$  (14). PPAR $\gamma$  represent ligand-activated transcription factors that belong to a nuclear receptor superfamily, and two isoforms, i.e., PPAR $\gamma$ 1 (16) and PPAR $\gamma$ 2 (17), are formed from the same gene by alternative mRNA splicing. PPAR $\gamma$  forms heterodimers with retinoid X receptors (RXR) and upon ligand activation, the PPAR/RXR heterodimer recruits coactivators and binds to sequence-specific PPRE present in the promoter region of its target genes (18). Alternatively, PPAR $\gamma$  can inhibit specific gene expression without direct binding to the gene promoter, because transrepression of several genes, i.e., iNOS and COX-2, is achieved in part by antagonizing the activities of transcription factors STAT1, NF- $\kappa$ B, and AP-1 (19).

In the present work, we analyzed the promoter region of the BACE1 gene and identified a functional PPRE. Further, we examined whether PPAR $\gamma$  is altered in AD by determination of PPAR $\gamma$  levels in brain of AD patients and AD transgenic mice. We demonstrate that NSAIDs modulate BACE1 transcription by repressing its promoter activity specifically through PPAR $\gamma$  activation.

## Materials and Methods

**Materials and Antibodies.** Immunostimulants and inhibitors tested were IFN- $\gamma$  (Sigma), TNF- $\alpha$  (Roche, Mannheim, Germany), ibuprofen (Sigma), pioglitazone (Takeda, Osaka), indomethacin,

Conflict of interest statement: No conflicts declared.

This paper was submitted directly (Track II) to the PNAS office.

Abbreviations: A $\beta$ , amyloid- $\beta$ ; AD, Alzheimer's disease;  $\beta$ APP,  $\beta$ -amyloid precursor protein; APPsw, APP695 containing the Swedish mutation; BACE,  $\beta$ -site APP cleaving enzyme; MEF, mouse embryonic fibroblasts; NSAIDs, nonsteroidal antiinflammatory drugs; PPAR $\gamma$ , peroxisome proliferator-activated receptor- $\gamma$ ; PPRE, PPAR responsive element; siRNA, small interfering RNA; HEK, human embryonic kidney.

<sup>†</sup>To whom correspondence may be addressed. E-mail: magdalena.sastre@ukb.uni-bonn.de or heneka@uni-muenster.de.

<sup>††</sup>Present address: Department of Neurology, University of Muenster, Mendelstrasse 7, 48149 Muenster, Germany.

© 2006 by The National Academy of Sciences of the USA

naproxen, sulindac sulfide, BAY11-7082 (Alexis Biochemicals, Grünberg, Germany), and the PPAR $\gamma$  antagonist GW0072 (provided by Timothy Willson, Glaxo). The monoclonal antibody 6E10 was obtained from Signet Laboratories (Dedham, MA), polyclonal pan-A $\beta$  antibody was obtained from Calbiochem, polyclonal antibody for PPAR $\gamma$  from Alexis Biochemicals, as well as polyclonal H-100 from Santa Cruz Biotechnology, which is highly specific for PPAR $\gamma$  and does not crossreact with PPAR $\alpha$  or PPAR $\beta$ . The polyclonal antibody 369 for APP was a generous gift from Sam Gandy (Thomas Jefferson University, Philadelphia) and polyclonal antibody against CT-BACE1 7520 from C. Haass (Ludwig-Maximilians-Universität, Munich). Tissue culture reagents were obtained from Invitrogen. All other chemicals were purchased from Sigma.

**Human Postmortem Brain Samples.** Human brains were obtained from routine autopsies at the Huddinge Brain Bank in accordance with the laws and the permission of the ethical committee. The control group included brains from subjects who died either of nonneurological diseases or traffic accidents and had no history of long-term illness or dementia (seven male, three female; mean age was 77  $\pm$  4 years, range 62–91 years). The AD group included the brains from patients with clinically and pathologically confirmed AD (four male, six female, mean age 83  $\pm$  2 years, range 74–93 years).

**Animals.** The transgenic mice used in this study were of the FVB/N genetic background and expressed APPV717I under the control of the mouse thy1 gene promoter. Mice 10 months of age were used, because APPV717I mice begin to deposit amyloid peptides at this age. Mice were fed Purina 5002 rodent chow *ad libitum* supplemented with either ibuprofen or pioglitazone for 7 days. There were six animals per group. The final dosage of drug was computed to be 62.5 mg/kg per day of ibuprofen and 40 mg/kg per day of pioglitazone.

**Cell Lines, Cell Culture, and Transient Transfection.** Mouse embryonic fibroblast (MEF) knockout from PPAR $\gamma$  were obtained as described (20). Mouse neuroblastoma N2a cells stably transfected with APP695 containing the Swedish mutation (K595N/M596L), so-called APP<sup>sw</sup>, were obtained from G. Thinakaran (University of Chicago, Chicago). Human embryonic kidney (HEK)293 cells stably transfected with APP<sup>sw</sup> were kindly supplied by C. Haass (University of Munich, Munich). Transfections were carried out by using Lipofectamine 2000 (Invitrogen) or FuGene (Roche). PPAR $\gamma$ 1 reporter plasmid pGL3- $\gamma$ P3000 was obtained from Johan Auwerx, Strasbourg, France). Cells were incubated for 20 h with OPTIMEM alone or with cytokines IFN- $\gamma$  (1 ng/ml) + TNF- $\alpha$  (30 ng/ml).

**PPAR $\gamma$  Small Interfering (si)RNA.** siRNA duplex to down-regulate PPAR $\gamma$  expression was directed to the target sequence 5'-AAGACCACTCGCATTCTTTG-3' of the mouse PPAR $\gamma$  cDNA and to the sequence 5'-GACCACTCCCACTCCTTTG-3' of the human PPAR $\gamma$  cDNA. As a control, the following target sequence was used: 5'-AAGAGGTGGC-CATCCGAATTT-3', which did not modify PPAR $\gamma$  expression.

**Site-Directed Mutagenesis.** Site-directed mutagenesis in the 1.5-kb BACE promoter fragment cloned in pGL3 vector was carried out by using the QuickChange Site-Directed Mutagenesis Kit (Stratagene).

**Western Blotting and Determination of Secreted A $\beta$  and CTF- $\beta$ .** Determination of A $\beta$  and CTFs and Western blotting were done as described (14).

**Luciferase Assay.** The luciferase reporter assay was performed according to the instructions of the manufacturer (Promega).

**EMSA.** Nuclear proteins (10  $\mu$ g) were incubated for 20 min on ice with double-stranded <sup>32</sup>P-labeled oligonucleotides in a final volume of 15  $\mu$ l. Oligonucleotides with the consensus PPRE site present in BACE1 promoter were: 5'-ATCAGGTGGGTCATGAGG-3' 3'-TAGTCCACCCAGTACTCC-5' and the mutant 5'-ATCAGCAGGCACATGAGG-3' 3'-TAGTCGTCCGTG-TACTCC-5'.

**RNA Preparation and RT-PCR.** Total RNA was extracted by using TRIzol reagent according to the manufacturer's instructions (Sigma). RT-PCR was performed as described (21). The primers used for mouse BACE were: 5'-CCGGCGGGAGTGGTATTATGAAGT-3' and 5'-GATGGTGATGCGGAAGGACTGATT-3'; for murine GAPDH: 5'-ACGACAGTCCATGCCATCAC-3' and 5'-TCCACCACCTGTTGCTGTA-3'; and for murine PPAR $\gamma$ : 5'-ATGCTGGCCTCCCTGATGAATAAA-3' and 3'-ACAA-GCGGTTCCACGAGGTC-5'.

**Immunohistochemistry.** Mouse brain sections were stained with antibody pan-A $\beta$ . For quantitative image analysis of subiculum immunostaining, serial sagittal sections were examined. All images were acquired by using a standard light and immunofluorescence microscope (Nikon, Eclipse E-800) connected to a digital camera (DXC-9100P, Sony, Cologne, Germany) and to a personal computer system with LUCIA imaging software (LUCIA 32G, Version 4.11; Laboratory Imaging, Düsseldorf, Germany).

**Statistical Evaluation.** Data were statistically analyzed by SYSTAT (Systat, Evanston, IL) by using ANOVA followed by the Tukey post hoc test.

## Results


**Effect of PPAR $\gamma$  Suppression on A $\beta$  Secretion and BACE1 Expression.** We sought to determine the specificity of the effect of inflammatory cytokines and ibuprofen (14, 22) in MEFs either wild type or deficient in PPAR $\gamma$  (PPAR $\gamma$ -/- MEF). The latter cells lack PPAR $\gamma$  protein and mRNA as assessed by Western blotting and semiquantitative RT-PCR, respectively (Fig. 1A). Western blotting analysis with two different antibodies showed specificity for PPAR $\gamma$ , because there was no signal detected in PPAR $\gamma$  null cells.

MEF were transiently transfected with APP<sup>sw</sup> (APP695 containing the Swedish mutation) and incubated with a combination of cytokines (TNF- $\alpha$  + IFN- $\gamma$ ). PPAR $\gamma$ -/- MEF did not show any modulation of A $\beta$  secretion and BACE1 protein expression by immunostimulation and/or incubation with ibuprofen (10  $\mu$ M) (Fig. 1B and C), and this effect was reversed by PPAR $\gamma$  transfection. Interestingly, the levels of BACE1 protein in PPAR $\gamma$  knockout cells were increased compared with wild-type cells and cells retransfected with PPAR $\gamma$  cDNA, indicating that PPAR $\gamma$  affected BACE1 cytokine-induced expression (Fig. 1C).

Additionally, we determined whether the suppression of PPAR $\gamma$  by siRNA affected the levels of BACE1 protein and the secretion of A $\beta$ . In N2a cells with stable expression of APP<sup>sw</sup>, transfection of siRNA against mouse PPAR $\gamma$  effectively decreased PPAR $\gamma$  expression (Fig. 1A). Unlike control cells, siRNA-treated cells showed no modulation of A $\beta$  secretion or of BACE1 protein levels by incubation with proinflammatory cytokines or ibuprofen (Fig. 1D and E). We compared our results using another sequence of siRNA, which did not decrease PPAR $\gamma$  levels, and observed no change in BACE1 expression (data not shown). Specific ELISA for A $\beta$ <sub>1-40</sub> and A $\beta$ <sub>1-42</sub> in media from HEK293 cells stably transfected with APP<sup>sw</sup> and transiently transfected with siRNA against human PPAR $\gamma$  showed the same effects (Fig. 1F).

Because PPAR $\gamma$  transcriptional activation depends on ligand binding, we determined whether transfection of N2a-APP<sup>sw</sup> cells





**Fig. 3.** PPAR $\gamma$  modulates BACE1 promoter activity. (A) Luciferase activities of a 1.5-kb BACE1 promoter/luciferase reporter construct transfected in PPAR $\gamma$  wild-type, knockout, and knockout transfected with PPAR $\gamma$  cDNA MEF. Cells were treated with IFN- $\gamma$  (1 ng/ml) + TNF- $\alpha$  (30 ng/ml) with or without ibuprofen (IBU) (10  $\mu$ M),  $n = 5$ . (B) NSAIDs and PPAR $\gamma$  agonists inhibit transcription of BACE1 promoter. N2a-sw cells transfected with BACE1 promoter and incubated with IBU, Pio, Indo, Napro, GW0072X (1  $\mu$ M), BAY11-7082, and sulindac sulfide  $n = 4$ , all at 10  $\mu$ M concentration. (C) PPAR $\gamma$  transfection inhibits BACE1 promoter activity. N2a-sw cells were transfected with BACE1 promoter construct and PPAR $\gamma$ 1, PPAR $\gamma$ 2, and PPAR $\gamma$ 2 E499Q cDNA and incubated with or without IBU (10  $\mu$ M);  $n = 4$ . (D) MEF transfected with BACE1 promoter control and mutated at the PPRE site were incubated with IBU (10  $\mu$ M). (E) N2a cells transfected with BACE1 promoter control and mutated at the PPRE site were incubated with IBU (10  $\mu$ M). Columns represent mean  $\pm$  SEM,  $n = 4$ . Asterisks, significant differences between wild-type cells and treated cells. #, differences between transfected cells treated or untreated. \*,  $P \leq 0.05$ ; \*\*,  $P \leq 0.01$ ; \*\*\*,  $P \leq 0.001$ ; #,  $P \leq 0.05$ ; ##,  $P \leq 0.01$ , ANOVA followed by a Tukey post hoc test. (F) Gel-shift analysis with the BACE1-PPRE probe using nuclear extract from MEF cells. The major PPAR $\gamma$ -containing complex is indicated by the arrow. Lane 1, MEF wild-type cells; lane 2, MEF PPAR $\gamma$  knockout cells; lane 3, MEF PPAR $\gamma$  knockout cells transfected with PPAR $\gamma$ 1 cDNA; lane 4, MEF wild-type cells incubated with an excess of unlabeled BACE1-PPRE probe. (G) Gel shift using HEK293 cells transfected with human PPAR $\gamma$ 2 cDNA. Lane 1, control; lane 2, supershift analysis; lane 3, molar excess of unlabeled BACE1-PPRE probe; lane 4, labeled mutant BACE1-PPRE probe (BACE1-PPREM).

critical for the PPAR $\gamma$  repression, we analyzed truncated constructs by removing sequence segments that appeared potentially important for the regulation of the BACE1 gene (see Fig. 6, which is published as supporting information on the PNAS web site). The promoter activity of BPR.Del (containing residues -1 to -753)


was significantly higher than that of the original 1.5-kb promoter construct, suggesting suppressor elements to be located between -754 and -1,541 bp (11). The activity of this construct was high when tested in cells with or without PPAR $\gamma$ , indicating that the deleted region could contain consensus-binding sites for PPAR $\gamma$  or other transcription factors modulated by PPAR $\gamma$ .

We then tested a number of NSAIDs and PPAR $\gamma$  agonists on BACE1 promoter activity in N2a neuroblastoma cells. Ibuprofen, indomethacin, naproxen, or the PPAR $\gamma$  agonist pioglitazone all repressed BACE1 promoter activity, whereas coincubation with the PPAR $\gamma$  antagonist GW0072 reversed the suppressive effect of ibuprofen (Fig. 3B). We also measured the effect of the NF- $\kappa$ B inhibitor BAY11-7082 and sulindac sulfide, which is an NSAID that does not activate PPAR $\gamma$ , and observed no change in BACE1 promoter activity (Fig. 3B). On the other hand, overexpression of PPAR $\gamma$ 1 and of PPAR $\gamma$ 2 in N2a cells robustly repressed BACE1 gene promoter activity in a ligand-dependent manner (Fig. 3C). Ibuprofen had no effect on BACE1 gene promoter activity in cells transfected with the mutant PPAR $\gamma$ 2-E499Q.

**The Rat BACE1 Gene Promoter Contains a PPAR $\gamma$  Responsive Element PPRE.** The increase in promoter activity of the BPR-Del(-1 to -753) mutant suggested the presence of suppressors of BACE1 transcription between -754 and -1,541 bp. Inspection of the sequence of this region revealed a possible consensus-binding site for PPAR $\gamma$  between -1356 and -1338, 5'-GGTGGGTCATGAG-GTTCA-3', which was experimentally analyzed by introducing point mutations in the BACE1 gene promoter reporter. Mutation of four nucleotides of the putative PPAR $\gamma$  site resulted in up-regulation of BACE1 gene promoter activity in wild-type MEF cells as opposed to unchanged activity in PPAR $\gamma$ -/- MEF (Fig. 3D). Furthermore, ibuprofen failed to decrease the promoter activity of the putative PPRE mutant BACE1 gene promoter construct (Fig. 3D). The same results were observed in N2a cells transfected with wild-type BACE1 gene promoter and mutated in the PPRE (Fig. 3E).

To define further the specificity of the putative PPRE, we performed gel-shift analysis with a double-stranded oligonucleotide probe corresponding to the BACE1 promoter region containing the PPRE. The radiolabeled double-stranded probe (PPREwt) was incubated with nuclear extracts from MEF and HEK293 cells transfected with human PPAR $\gamma$ 2, resulting in the formation of several high-molecular complexes (Fig. 3F and G). The formation of one of the complexes was efficiently reduced by an excess of unlabeled probe, indicating the specificity of the PPAR $\gamma$  bound to this consensus motif (Fig. 3F, lane 4, and G, lane 3). Incubation with an antiserum to PPAR $\gamma$  resulted in the disappearance of the same complex (Fig. 3G, lane 2). The substitution of four bases in the synthetic oligonucleotide (PPREM) abolished the binding (Fig. 3G, lane 4). Nuclear extracts from PPAR $\gamma$ -/- MEF cells and PPAR $\gamma$ -/- MEF transfected with PPAR $\gamma$ 1 cDNA were also used as controls (Fig. 3F, lanes 2 and 3).

**Treatment with PPAR $\gamma$  Agonists Affects the Levels of PPAR $\gamma$ , BACE1, and A $\beta$  Generation in APPV717I Mice.** To prove the effects of PPAR $\gamma$  activation *in vivo*, we performed experiments using the APPV717I mouse model of AD. In wild-type animals, the levels of PPAR $\gamma$  mRNA and protein decreased with age and were higher compared with APPV717I transgenic mice, in agreement with the existence of an inflammatory component in AD mice (Fig. 4A). We have recently observed that ibuprofen and pioglitazone treatment in APP transgenic mice results in a reduction of glial inflammation and A $\beta$  levels in plaques as measured by ELISA (24). Therefore, we sought the underlying mechanism behind these effects and detected increased PPAR $\gamma$  transcript and reduced BACE1 mRNA levels (Fig. 4B) as well as reduced CTF- $\beta$  levels (Fig. 4E). To validate that PPAR $\gamma$  activation decreases total A $\beta$  generation *in vivo*, we determined the intracellular A $\beta$  levels by staining brain sections with



**Fig. 4.** PPAR $\gamma$  in the brain of APPV7171 transgenic mice and in AD patients. (A) Quantification of PPAR $\gamma$  expression in cortex from mouse brain control and APPV7171 of 3 and 16 months of age. (B) Quantification of the mRNA levels of PPAR $\gamma$  and BACE1 APPV7171 mice treated with ibuprofen (IBU) and pioglitazone (PIO). (C) Immunostaining of intracellular A $\beta$  in subiculum from APPV7171 mice treated with IBU and PIO. (D) Quantification of percentage of intracellular A $\beta$  in subiculum from APPV7171 mice treated with IBU and PIO. Bar, 100  $\mu$ M. Columns represent mean  $\pm$  SEM,  $n = 4$ . (E) Representative Western blot for CTF- $\beta$  fragments in cortex from APPV7171 mice treated with IBU and PIO. (F) Representative Western blot for PPAR $\gamma$  expression in brain lysates from two control and two AD patients. As a negative control,  $\beta$ -actin expression was analyzed. (G) Quantification of the levels of  $\beta$ -actin, PPAR $\gamma$ , and BACE1 in the frontal cortex of 10 controls and 10 AD patients. Asterisks, significant differences between control and AD. \*,  $P \leq 0.05$ ; \*\*,  $P \leq 0.01$ . ANOVA followed by a Tukey post hoc test or Student's  $t$  test for human brains.

antibody pan-A $\beta$ . Treatment with ibuprofen and pioglitazone dramatically reduced the levels of intracellular A $\beta$  (Fig. 4 C and D), without affecting the APP levels (data not shown), suggesting that PPAR $\gamma$  activation could regulate A $\beta$  generation.

**PPAR $\gamma$  and BACE1 Expression in AD Brain.** To determine whether PPAR $\gamma$  is important for BACE1 regulation in humans, we examined the levels of expression of PPAR $\gamma$  in postmortem brain sections from AD patients (see Fig. 7, which is published as supporting information on the PNAS web site). We quantified the protein levels of PPAR $\gamma$  and BACE1 in the frontal cortex of 10 AD patients compared with 10 controls. In AD brain, PPAR $\gamma$  protein levels were reduced by 40% compared with controls (Fig. 4 F and G), whereas BACE1 expression was increased (Fig. 4 F). To investigate whether the binding of PPAR $\gamma$  to the BACE1 gene promoter was affected, we performed gel-shift analysis with the PPREwt

probe after incubation with nuclear extracts from brain of five AD patients and five controls. The retarded band corresponding to the PPRE complex was reduced by 50% (see Fig. 7), indicating that PPAR $\gamma$  protein levels and its binding are decreased in AD brain. Combined with all of the other data, these findings strongly point to a direct role of PPAR $\gamma$  in the increased BACE1 transcription and activity in AD, facilitating the generation of A $\beta$ .

## Discussion

Epidemiological and experimental evidence suggested a significant inflammatory component in AD and documented the beneficial effect of NSAIDs in this pathology (25, 26), stating an anti-inflammatory effect. The underlying molecular mechanism was related to decreased secretion of A $\beta$  peptides in cultured cells and in amyloid mouse models (14, 22, 24, 27–29). Some NSAIDs, but not all, have been shown to be able to activate PPAR $\gamma$  (15). Ibuprofen, indomethacin, and naproxen are among the five most-prescribed NSAIDs, which potentially decrease the risk for AD (30). In this paper, we suggest that the NSAIDs that are PPAR $\gamma$  activators reduce A $\beta$  levels by repressing BACE1 promoter activity.

Because it is widely recognized that a local inflammatory response within the brain contributes to the pathophysiology of AD, we performed our assays under inflammatory stimuli. For this reason, our data related to NSAIDs on APP processing are not comparable to those previously reported by using non-inflammatory conditions (28, 29, 31, 32). We recently observed that proinflammatory cytokines activate BACE1 expression, transcription, and activity in cultured neuroblastoma cells (14). The increase was reversed after treatment with NSAIDs or PPAR $\gamma$  agonists. Therefore, we sought to define whether this effect was specific for PPAR $\gamma$  and to determine the molecular mechanisms by which PPAR $\gamma$  decreases BACE1 transcription and thereby A $\beta$  generation. Here we show that the effect of ibuprofen and other NSAIDs on BACE1 expression and A $\beta$  secretion is specifically mediated by PPAR $\gamma$ . Importantly, BACE1 transcription was found to be up-regulated in MEF PPAR $\gamma$  knockout cells compared with wild-type cells. These results were confirmed by reporter gene assays, which demonstrated that lack of endogenous PPAR $\gamma$  facilitates BACE1 promoter activity, suggesting that PPAR $\gamma$  could be a suppressor for BACE1 promoter. Moreover, incubation of N2a cells with NSAIDs that are PPAR $\gamma$  activators showed a decrease in BACE1 gene promoter activity, and this effect was not reproduced by NSAIDs that are not PPAR $\gamma$  agonists, such as sulindac sulfide and aspirin. Equally robust effects were detected by overexpression of PPAR $\gamma$ 1 and PPAR $\gamma$ 2, which significantly reduced BACE1 promoter activity.

The upstream sequences of the BACE1 gene promoter are highly conserved among rat, mouse, and human, indicating that this region contains regulatory elements that modulate BACE1 gene transcription (11–13). We identified a putative PPAR $\gamma$  consensus sequence in the region located between nucleotides –1356 and –1338. The activity of the BACE1 gene promoter mutated in the putative consensus sequence for PPAR $\gamma$  was increased in MEF cells relative to the wild-type promoter and was not affected by ibuprofen, strengthening the hypothesis of the negative modulation of BACE1 transcription by PPAR $\gamma$ . That PPAR $\gamma$  binds to the putative PPRE in the BACE1 gene promoter was demonstrated directly by mobility-shift assays. The BACE1 gene promoter contains other putative transcription factor-binding sites, i.e., for NF- $\kappa$ B, SP-1, AP1, and AP2. Because PPAR $\gamma$  can modulate the activity of different transcription factors, we cannot exclude completely that PPAR $\gamma$  agonists also affect BACE1 gene promoter activity by antagonizing the activity of other transcriptional regulators.

It has been suggested that inflammatory cytokines and oxidative stress decrease PPAR $\gamma$  mRNA in adipocytes (23), and thiazolidinediones reverse this effect (33). Similarly, we demonstrate here that certain combinations of inflammatory cytokines are able to

decrease PPAR $\gamma$  mRNA levels and PPRE activity in neuronal cells, and this effect is suppressed by incubation with NSAIDs. In addition, we show that PPAR $\gamma$  gene transcription is strongly reduced by inflammatory cytokines, and this can be prevented by incubation with ibuprofen or indomethacin. In this line, it has been recently shown in adipocytes that TNF- $\alpha$  suppresses PPAR $\gamma$ 2 transcription by inhibiting the binding of C/EBP $\delta$  to the PPAR $\gamma$ 2 promoter (34). Interestingly, we observed strongly decreased PPAR $\gamma$  protein levels in the frontal cortex of AD brain and in transgenic mice. An increase of the release of inflammatory cytokines by glial cells could be the determinant for the decrease in PPAR $\gamma$  levels in the brain and consequently could alter BACE1 transcription. This would support the hypothesis of the existence of a vicious cycle that accelerates the progression of AD (Fig. 8, which is published as supporting information on the PNAS web site). We believe that PPAR $\gamma$  is herein a major factor, because in the absence of PPAR $\gamma$ , there is no modulation of A $\beta$  secretion by inflammation. The decrease of PPAR $\gamma$  levels detected in the brain of AD patients is not likely to be solely due to neuronal loss or secondary effects, i.e., loss of synapses. We have observed a similar decrease of PPAR $\gamma$  levels in lymphocytes from multiple sclerosis patients compared with healthy controls (35). It is therefore tempting to speculate that patients, even with minor signs of cerebral inflammation, could be predisposed to develop AD earlier due to reduced PPAR $\gamma$  levels that would consequently increase BACE1 expression and A $\beta$  generation.

Our *in vivo* results support the *in vitro* observations, indicating that the PPAR $\gamma$  mRNA levels are up-regulated in transgenic mice after treatment with PPAR $\gamma$  agonists and as a consequence, decreased BACE1 expression leads to reduced total intracellular A $\beta$ . This indicates that PPAR $\gamma$  activation is indeed involved in the regulation of A $\beta$  generation, in contrast with another hypothesis that suggests PPAR $\gamma$  activation could cause an increase of extracellular A $\beta$  degradation (31). Our data are in agreement with previous publications describing that treatment with certain NSAIDs and PPAR $\gamma$  agonists in AD mouse models decreased A $\beta$ <sub>1-40</sub> and A $\beta$ <sub>1-42</sub> levels (24, 27, 29, 36, 37). We do not exclude that some NSAIDs could show a selective

reduction in A $\beta$ <sub>1-42</sub>, but that effect could be independent of PPAR $\gamma$  activation. In addition, the concentrations of NSAIDs required to detect an effect in A $\beta$ <sub>1-40</sub>/A $\beta$ <sub>1-42</sub> ratio are considerably higher (28), whereas the concentrations of NSAIDs to start the activation of PPAR $\gamma$  and decrease BACE1 transcription are in the same range as active concentrations of NSAIDs found in human plasma and cerebrospinal fluid (38–40).

## Conclusion

We have shown that PPAR $\gamma$  represses BACE1 gene promoter activity in response to ligand binding to a PPRE located in the BACE1 gene promoter. This could be a potential mechanism by which NSAIDs have a protective effect against the development of AD. Moreover, we demonstrate that inflammatory cytokines modulate PPAR $\gamma$  transcription and PPRE activity, suppressing the negative regulation of PPAR $\gamma$  on BACE1 gene expression *in vivo* and *in vitro*. These effects may explain the overexpression of BACE1 in the brain under inflammatory conditions and emphasize the hypothesis that neuroinflammatory mechanisms significantly contribute to the pathogenesis of AD.

We thank Dr. Sam Gandy (Thomas Jefferson University, Philadelphia), Dr. Rotraut Mossner (University of Göttingen, Göttingen, Germany), and Dr. Christian Haass (University of Munich, Munich) for generous gifts of antibodies and cells; Dr. Gopal Thinakaran (University of Chicago, Chicago) for providing cells; Dr. Timothy M. Willson (Glaxo Wellcome) for PPAR $\gamma$  antagonist GW0072; Dr. Efrat Levy (Nathan Kline Institute for Psychiatric Research, Orangeburg, NY) for APPsw cDNA; Dr. Ron Evans (The Salk Institute for Biological Studies, San Diego) for PPAR- $\gamma$ 1 cDNA; Dr. Bruce Spiegelman (Harvard Medical School, Boston) for mouse PPAR $\gamma$ 2-E499Q cDNA; Dr. Walter Wahli (University of Lausanne, Lausanne, Switzerland) for human PPAR $\gamma$ 2 cDNA; and Dr. Andreas von Knethen (University of Kaiserslautern, Kaiserslautern, Germany) for the PPAR $\gamma$  responsive element (PPRE) reporter plasmid. This investigation was supported by the Sonderforschungsbereich 400 [SFB 400, Teilprojekt A8 (to M.T.H. and T.K.) and DFG WA1477/3 (to J.W.)], by Fonds voor Wetenschappelijk Onderzoek Vlaanderen, by the Katholieke Universiteit Leuven Research Fund, and by Katholieke Universiteit Leuven Research and Development. I.D. is a postdoctoral fellow at Fonds voor Wetenschappelijk Onderzoek Vlaanderen.

- Vassar, R., Bennett, B. D., Babu-Khan, S., Kahn, S., Mendiaz, E. A., Denis, P., Teplow, D. B., Ross, S., Amarante, P., Loeloff, R., et al. (1999) *Science* **286**, 735–741.
- Luo, Y., Bolon, B., Damore, M. A., Fitzpatrick, D., Liu, H., Zhang, J., Yan, Q., Vassar, R., & Citron, M. (2003) *Neurobiol. Dis.* **14**, 81–88.
- Ohno, M., Sametsky, E. A., Younkin, L. H., Oakley, H., Younkin, S. G., Citron, M., Vassar, R., & Disterhoft, J. F. (2004) *Neuron* **41**, 27–33.
- Tamagno, E., Bardini, P., Obbili, A., Vitali, A., Borghi, R., Zaccheo, D., Pronzato, M. A., Danni, O., Smith, M. A., Perry, G., & Tabato, M. (2002) *Neurobiol. Dis.* **10**, 279–288.
- Hartlage-Rubsamen, M., Zeitschel, U., Apelt, J., Gartner, U., Franke, H., Stahl, T., Gunther, A., Schliebs, R., Penkowa, M., Bigl, V., & Rossner, S. (2003) *Glia* **4**, 169–179.
- Blasko, I., Beer, R., Bigl, M., Apelt, J., Franz, G., Rudzki, D., Ransmayr, G., Kampfl, A., & Schliebs, R. (2004) *J. Neural Transm.* **111**, 523–536.
- Holsinger, R. M., McLean, C. A., Beyreuther, K., Masters, C. L., & Evin, G. (2002) *Ann. Neurol.* **51**, 783–786.
- Fukumoto, H., Rosene, D. L., Moss, M. B., Raju, S., Hyman, B. T., & Irizarry, M. C. (2004) *Am. J. Pathol.* **164**, 719–725.
- Li, R., Lindholm, K., Yang, L. B., Yue, X., Citron, M., Yan, R., Beach, T., Sue, L., Sabbagh, M., Ca, H., et al. (2004) *Proc. Natl. Acad. Sci. USA* **101**, 3632–3637.
- Rogers, G. W., Jr., Edelman, G. M., & Mauro, V. P. (2003) *Proc. Natl. Acad. Sci. USA* **101**, 2794–2799.
- Lange-Dohna, C., Zeitschel, U., Gaunitz, F., Perez-Polo, J. R., Bigl, V., & Rossner, S. (2003) *J. Neurosci. Res.* **73**, 73–80.
- Christensen, M. A., Zhou, W., Qing, H., Lehman, A., Philipsen, S., & Song, W. (2004) *Mol. Cell. Biol.* **24**, 865–874.
- Sambamurti, K., Kinsey, R., Maloney, B., Ge, Y. W., & Lahiri, D. K. (2004) *FASEB J.* **18**, 1034–1036.
- Sastre, M., Dewachter, I., Landreth, G. E., Willson, T. M., Klockgether, T., van Leuven, F., & Heneka, M. T. (2003) *J. Neurosci.* **23**, 9796–9804.
- Lehmann, J. M., Lenhard, J. M., Oliver, B. B., Ringold, G. M., & Kliewer, S. A. (1997) *J. Biol. Chem.* **272**, 3406–3410.
- Kliewer, S. A., Forman, B. M., Blumberg, B., Ong, E. S., Borgmeyer, U., Mangelsdorf, D. J., Umeson, K., & Evans, R. M. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 7355–7359.
- Tontonoz, P., Hu, E., & Spiegelman, B. M. (1994) *Cell* **79**, 1147–1156.
- Tugwood, J. D., Issemann, I., Anderson, R. G., Bundell, K. R., McPheat, W. L., & Green, S. (1992) *EMBO J.* **11**, 433–439.
- Li, A. C., Brown, K. K., Silvestre, M. J., Willson, T. M., Palinski, W., & Glass, C. K. (2000) *J. Clin. Invest.* **106**, 523–531.
- Rosen, E. D., Hsu, C. H., Wang, X., Sakai, S., Freeman, M. W., Gonzalez, F. J., & Spiegelman, B. M. (2002) *Genes Dev.* **16**, 22–26.
- Heneka, M. T., Klockgether, T., & Feinstein, D. L. (2000) *J. Neurosci.* **20**, 6862–6867.
- Blasko, I., Apochal, A., Boeck, G., Hartmann, T., Grubeck-Loebenstien, B., & Ransmayr, G. (2001) *Neurobiol. Dis.* **8**, 1094–1101.
- Xing, H., Northrop, J. P., Grove, J. R., Kilpatrick, K. E., Su, J.-L., & Ringold, G. M. (1997) *Endocrinology* **138**, 2776–2783.
- Heneka, M. T., Sastre, M., Dumitrescu-Ozimek, L., Hanke, A., Dewachter, I., Kuiperi, C., O'Banion, K., Klockgether, T., van Leuven, F., & Landreth, G. E. (2005) *Brain* **128**, 1442–1453.
- Akiyama, H., Barger, S., Barnum, S., Bradt, B., Bauer, J., Cole, G. M., Cooper, N. R., Eikelenboom, P., Emmerling, M., Fiebich, B. L., et al. (2000) *Neurobiol. Aging* **21**, 383–421.
- Szekely, C. A., Thorne, J. E., Zandi, P. P., Ek, M., Messias, E., Breitner, J. C., & Goodman, S. N. (2004) *Neuroepidemiology* **23**, 159–169.
- Lim, G. P., Yang, F., Chu, T., Chen, P., Beech, W., Teter, B., Tran, T., Ubeda, O., Ashe, K. H., Frautschy, S. A., & Cole, G. M. (2000) *J. Neurosci.* **20**, 5709–5714.
- Weggen, S., Eriksen, J. L., Das, P., Sagi, S. A., Wang, R., Pietrzik, C. U., Findlay, K. A., Smith, T. E., Murphy, M. P., Bulter, T., et al. (2001) *Nature* **414**, 212–216.
- Eriksen, J. L., Sagi, S. A., Smith, T. E., Weggen, S., Das, P., McLendon, D. C., Ozols, V. V., Jessing, K. W., Zavitz, K. H., Koo, E. H., et al. (2003) *J. Clin. Invest.* **112**, 440–449.
- In't Veld, B. A., Ruitenbergh, A., Hofman, A., Launer, L. J., van Duijn, C. M., Stijnen, T., Breteler, M. M. B., & Stricker, B. H. C. (2001) *N. Engl. J. Med.* **345**, 1515–1521.
- Samachro, I. E., Serneels, L., Spittaels, K., Merchiers, P., Dominguez, D., & De Strooper, B. (2004) *J. Neurosci.* **24**, 10908–10917.
- Lleo, A., Berezovska, O., Herl, L., Raju, S., Deng, A., Bacskai, B. J., Frosh, M. P., Irizarry, M., & Hyman, B. T. (2004) *Nat. Med.* **10**, 1065–1066.
- Gimble, J. M., Robinson, C. E., Wu, X., Kelly, K. A., Rodriguez, B. R., Kliewer, S. A., Lehmann, J. M., & Morris, D. C. (1996) *Mol. Pharmacol.* **50**, 1087–1094.
- Kudo, M., Sugawara, A., Uruno, A., Takeuchi, K., & Ito, S. (2004) *Endocrinology* **145**, 4948–4956.
- Klotz, L., Schmidt, M., Sastre, M., Giese, T., Knolle, P., Klockgether, T., & Heneka, M. T. (2005) *J. Immunol.* **175**, 4948–4955.
- Yan, Q., Zhang, J., Liu, H., Babu-Khan, S., Vassar, R., Biere, A. L., Citron, M., & Landreth, G. (2003) *J. Neurosci.* **23**, 7504–7509.
- Sung, S., Yang, H., Uryu, K., Lee, E. B., Zhao, L., Shineman, D., Trojanowski, J. Q., Lee, V. M., & Pratico, D. (2004) *Am. J. Pathol.* **165**, 2197–2206.
- Jaradat, M. S., Wongsud, B., Phornchirasilp, S., Rangwala, S. M., Shams, G., Sutton, M., Romstedt, K. J., Noonan, D. J., & Feller, D. R. (2001) *Biochem. Pharmacol.* **62**, 1587–1595.
- Bannwarth, B., Netter, P., Lapique, F., Pere, P., Thomas, P., & Gaucher, A. (1990) *Eur. J. Clin. Pharmacol.* **38**, 343–346.
- Bannwarth, B., Lapique, F., Pehourcq, F., Gillet, P., Schaeverbeke, T., Laborde, C., Dehais, J., Gaucher, A., & Netter, P. (1995) *Br. J. Clin. Pharmacol.* **40**, 266–269.