

Stable isotopes in elephant hair document migration patterns and diet changes

Thure E. Cerling^{*†‡}, George Wittemyer[§], Henrik B. Rasmussen[¶], Fritz Vollrath[¶], Claire E. Cerling[†], Todd J. Robinson[†], and Iain Douglas-Hamilton^{||}

^{*}Department of Geology and Geophysics, 135 South 1460 East, University of Utah, Salt Lake City, UT 84112; [†]Department of Biology, 257 South 1400 East, University of Utah, Salt Lake City, UT 84112; [§]University of California, Berkeley, CA 94720; [¶]Department of Zoology, Oxford University, Oxford OX1 3PS, United Kingdom; and ^{||}Save The Elephants Foundation, Nairobi, Kenya

Contributed by Thure E. Cerling, November 4, 2005

We use chronologies of stable isotopes measured from elephant (*Loxodonta africana*) hair to determine migration patterns and seasonal diet changes in elephants in and near Samburu National Reserve in northern Kenya. Stable carbon isotopes record diet changes, principally enabling differentiation between browse and tropical grasses, which use the C₃ and C₄ photosynthetic pathways, respectively; stable nitrogen isotopes record regional patterns related to aridity, offering insight into localized ranging behavior. Isotopically identified range shifts were corroborated by global positioning system radio tracking data of the studied individuals. Comparison of the stable isotope record in the hair of one migrant individual with that of a resident population shows important differences in feeding and ranging behavior over time. Our analysis indicates that differences are the result of excursions into mesic environments coupled with intermittent crop raiding by the migrant individual. Variation in diet, quantified by using stable isotopes, can offer insight into diet-related wildlife behavior.

13-carbon | 15-nitrogen | chronology | human–elephant conflict

The stable isotope ratios of ¹³C/¹²C in hair records the diet of mammals (1–4). It is particularly useful in distinguishing diets of C₃ browse versus C₄ grass in tropical regions (5–7) because of the large difference in ¹³C/¹²C ratios between plants using the C₃ and C₄ photosynthetic pathways, respectively. In tropical regions, the C₃ pathway is used primarily by trees and shrubs, whereas plants using the C₄ pathway are principally grasses (8, 9).

Hair is a particularly useful indicator of diet change (3, 4) because the isotope turnover of mammal tissues is high enough to resolve short-term diet changes. Recent advances in methodology, progressed through the study of large mammals with controlled diet changes (10, 11), allows detailed reconstruction of the diet history of individual large mammals in wild populations (12, 13).

In this study, we determine the growth rates and stable ¹³C/¹²C and ¹⁵N/¹⁴N ratios in elephant hair collected between 2001 and 2004. We focus on the behavior of a resident population of Samburu National Reserve, Northern Kenya, for the time period of 2000 to 2002. We compare stable isotope results of this resident population with a migrant elephant (B1013) that visited Samburu Reserve up to several times each year. Differences in isotope ratios between the resident individuals and the migrant indicate different behaviors, including rapid migration across long distances by the migrant individual and differences in the fraction C₄ biomass in the diet. The latter may be related to seasonal crop raiding, which can be quantified by using stable isotope ratios.

Materials and Methods

Global Positioning System (GPS) radio collars were fitted to elephants in Samburu National Reserve, Northern Kenya, between 2001 and 2004 (14). Collars were programmed to record positions at hourly intervals, offering detailed records of move-

ment. Tail hairs from 35 elephants were collected over this period during immobilization operations while the collars were being fitted and for some, later, when collar batteries were being changed or when the collars were being removed. Of these 35 elephants, 7 have a secure isotopic dietary chronology over the period from late 2000 and extending through February 2002; GPS spatial information is available for all resident individuals from July 2001 to July 2002 and from February 2002 to July 2002 for the migrant B1013. We focus on comparing the behavior of the migrant B1013 to that of the resident Samburu group during this time interval.

The elephants tracked in this study used both the semiarid region in and around Samburu National Reserve and the mesic Imenti Forest on the flanks of Mt. Kenya. These two distinct ecotones are <60 km apart and are located between 37° and 38° east, just north of the equator. The elevation of the semiarid Samburu region is ≈900 m above sea level and dominated by acacia–comiphora savanna and scrub bushland. Rainfall averages ≈350 mm per year in this lowland region and occurs during biannual rainy seasons, which generally take place in April and November. The elevation of the Imenti Forest is ≈2,000 m above sea level and is dominated by evergreen and broad leaf deciduous tree species. Rainfall in this region also occurs biannually and averages ≈900 mm per year.

In this study, we used 10-day composite normalized differential vegetation index (NDVI) data, available through Satellite Probatoire d’Observation de la Terre (SPOT), to determine changes in season across the study area. NDVI is a remote sensing index value calculated as the ratio between red and near infrared reflection that is highly correlated with green (photosynthetically active) biomass (15, 16). Remotely sensed data provide a direct measure of photosynthetic activity over large spatial regions, offering advantages over the classically used point-sampled rainfall data in areas, like the study region, where weather stations are sparse. Isotope profiles for each elephant were compared with longitudinal 10-day NDVI records to determine the impact of seasonality on diet.

¹³C/¹²C and ¹⁵N/¹⁴N ratios of elephant hair and plant material were measured on an isotope ratio mass spectrometer after combustion in a flow-through modified Carlo–Erba system. Values are reported using the conventional permil (‰) notation, where

$$\delta^{13}\text{C} = \left(\frac{(^{13}\text{C}/^{12}\text{C})_{\text{sample}}}{(^{13}\text{C}/^{12}\text{C})_{\text{standard}}} - 1 \right) \cdot 1,000,$$

and an analogous terminology describes ¹⁵N/¹⁴N ratios. Standards are Vienna Pee Dee Belemnite (VPDB) and air for ^δ¹³C and ^δ¹⁵N, respectively. Uncertainties for average ^δ¹⁵N values for

Conflict of interest statement: No conflicts declared.

Abbreviations: GPS, Global Positioning System; NDVI, normalized differential vegetation index.

[†]To whom correspondence should be addressed. E-mail: tcerling@earth.utah.edu.

© 2006 by The National Academy of Sciences of the USA




Fig. 2. Histogram of hours spent by B1013 outside Imenti Forest. In total, 2,179 h of GPS positions were recorded when B1013 was in or around the Imenti Forest. During this period, 236 h of GPS positions (10.8%) were recorded outside the designated forest boundaries in areas dense with subsistence farming. Being essentially on the equator, the sun rises and sets at 6:30 every day with little variation. All but 2 of the 236 h spent outside the Imenti Forest were during night hours during this interval.

Three of these periods were associated with wet seasons and periods of elevated remotely sensed NDVI values, representing increased vegetative productivity. In general, African elephants have a diet dominated by browse (6) and consume grass only as a windfall when it is in new growth. These periods of grass consumption occurred when B1013 was observed in Samburu National Reserve and when NDVI values exceeded 0.3. This pattern of elevated C_4 consumption during wet seasons was comparable with observed patterns in resident Samburu elephants (Fig. 1C); during these intervals, B1013 had similar C_4 grass consumption as the other six elephants. The single exception is the highest C_4 consumption peak, which occurred during the dry season between June 18 and August 16, 2001. By comparison, the other six elephants had a low fraction of C_4 grass consumption during this dry season. During this interval, while B1013 was in the Imenti Forest, average NDVI was low, yet C_4 grasses made up 36% of the diet integrated over this interval, with peak values approaching 50% C_4 biomass. NDVI in the Imenti region was below the yearly average from June 11 to November 21, 2001, corresponding with the dry season period when vegetative productivity is decreased.

GPS tracking data from the subsequent period spent in Imenti reveals that, while in residence of the Imenti Forest, B1013 was outside the forest reserve boundaries only during nighttime hours (Fig. 2). Most elephant crop raiding occurs at night, apparently to avoid human interaction (17, 18). Subsistence farming occurs adjacent to the Imenti Forest (19), and this region, around the time of the data collection, was one of the major crop raiding conflict zones in Kenya. Thus, it appears that this isotope signal is a quantifiable record of crop raiding (Fig. 3). While in the Imenti Forest, C_3 vegetation was the principal food available, and therefore




Fig. 3. Detail of diet for B1013 from June 1 to October 1, 2001 using ≈ 1 -mm hair segments. The diet estimate of C_4 fraction uses C_3 and C_4 endmember mixing lines for both mesic (upper boundary) and xeric environments (lower boundary). These have average $\delta^{13}C$ values of -28% and -12% for mesic environments and -26% and -13% for xeric environments, respectively. Average interval of time is 1.8 days (maximum, 2.4 days; minimum, 0.9 days).

the C_4 component of diet was most likely obtained from adjacent crops, evidently at night. During the dry season between June and August of 2001, our study indicates that B1013 was supplementing his C_3 forest diet with nocturnal crop raiding of C_4 maize. Fig. 3 shows a detailed stable isotope profile along with the estimated fractionation C_4 intake during this unusual C_4 diet period during the long dry season of 2001.

Quantification of longitudinal diet records from hair in wild animals can offer important information to conservationists and wildlife managers. This article provides an example of one of the many possible uses that stable isotopes can provide in resolving the elephant–human conflict issue. Despite the recording of conflict incidences, it is often difficult to understand elephant decisions made at an individual level and to quantify the dietary importance of crop raiding by elephants. Combining studies of elephant movement patterns and overall diet with incidences of raiding injects additional scientific data into the highly topical discussion of human–elephant conflict alleviation.

We thank J. R. Ehleringer and the Stable Isotope Ratio Facility for Environmental Research (SIRFER) at the University of Utah, the government of Kenya for permission to work in Kenya, the Kenya Wildlife Service for facilitation, Save The Elephants for technical and logistical field support, and the Packard Foundation.

- DeNiro, M. J. & Epstein, S. (1978) *Geochim. Cosmochim. Acta* **42**, 495–506.
- DeNiro, M. J. & Epstein, S. (1981) *Geochim. Cosmochim. Acta* **45**, 341–351.
- White, C. D. (1993) *J. Archaeol. Sci.* **20**, 657–666.
- Macko, S. A., Engel, M. H., Lubec, V. G., O'Connell, T. C. & Hedges, R. E. M. (1999) *Philos. Trans. R. Soc. London B* **354**, 65–76.
- Lee-Thorp, J. & van der Merwe, N. J. (1987) *S. Afr. J. Sci.* **83**, 712–715.
- van der Merwe, N. J., Lee-Thorp, J. A., Thackeray, J. F., Hall-Martin, A., Kruger, F. J., Coetzee, H., Bell, R. H. V. & Lindeque, M. (1990) *Nature* **346**, 744–746.
- Cerling, T. E., Harris, J. M. & Passey, B. H. (2003) *J. Mammal.* **84**, 456–471.
- Bender, M. M. (1971) *Phytochemistry* **10**, 1239–1245.
- Sage, R. F., Li, M. & Monson, R. K. (1999) in *C4 Plant Biology*, eds Sage, R. F. & Monson, R. K. (Academic, San Diego), pp. 551–584.
- Ayliffe, L. K., Cerling, T. E., Robinson, T., West, A. G., Sponheimer, M., Passey, B. H., Hammer, J., Roeder, B., Dearing, M. D. & Ehleringer, J. R. (2004) *Oecologia* **139**, 11–22.
- West, A. G., Ayliffe, L. K., Cerling, T. E., Robinson, T. F., Karren, B., Dearing, M. D. & Ehleringer, J. R. (2004) *Funct. Ecol.* **18**, 616–624.
- Cerling, T. E., Passey, B. H., Ayliffe, L. K., Cook, C. S., Ehleringer, J. R., Harris, J. M., Dhidha, M. B. & Kasiki, S. M. (2004) *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **206**, 367–376.
- Cerling, T. E. & Viehl, K. (2004) *Afr. J. Ecol.* **42**, 88–92.
- Douglas-Hamilton, I., Krink, T. & Vollrath, F. (2005) *Naturwissenschaften* **92**, 163–168.
- Goward, S. N. & Prince, S. D. (1995) *J. Biogeogr.* **22**, 549–564.
- Diallo, O., Diouf, A., Hanan, N. P., Niaye, A. & Prevost, Y. (1991) *Int. J. Remote Sensing* **12**, 1259–1279.
- Thouless, C. R. (1994) *Oryx* **28**, 119–127.
- Hoare, R. E. (1995) *Pachyderm* **19**, 54–63.
- Gathaara, G. N. (1999) *Aerial Survey of the Destruction of Mt. Kenya, Imenti and Ngare Ndare Forest Reserves* (Kenya Wildlife Service, Nairobi, Kenya) p. 33.