

# A novel multifunctional factor involved in *trans*-splicing of chloroplast introns in *Chlamydomonas*

Livia Merendino, Karl Perron, Michèle Rahire, Isabelle Howald,  
Jean-David Rochaix and Michel Goldschmidt-Clermont\*

Department of Molecular Biology and Department of Plant Biology, University of Geneva 30, quai E.  
Ansermet, CH-1211 Geneva 4, Switzerland

Received October 31, 2005; Revised and Accepted December 16, 2005

## ABSTRACT

In the chloroplast of *Chlamydomonas reinhardtii*, *psaA* mRNA is spliced *in trans* from three separate precursors which assemble to form two group II introns. A fourth transcript, *tscA*, completes the structure of the first intron. Of the fourteen nucleus-encoded factors involved in *psaA* splicing, only two are required for splicing of both introns. We cloned and characterized the first of these more general factors, Raa1. Consistently with its role in *psaA* splicing, Raa1 is imported in the chloroplast where it is found in a membrane fraction and is part of a large ribonucleoprotein complex. One mutant, *raa1-L137H*, is defective for splicing of both introns, but another allelic mutant, *raa1-314B*, still expresses the 3' part of the *Raa1* gene and is deficient only in splicing of intron 2. This observation and a deletion analysis indicate the presence of two domains in Raa1. The C-terminal domain is necessary and sufficient for processing of *tscA* RNA and splicing of the first intron, while the central domain is essential for splicing of the second intron. The combination of these two functional domains in Raa1 suggests that this new factor may coordinate *trans*-splicing of the two introns to improve the efficiency of *psaA* maturation.

## INTRODUCTION

Group II introns are typically characterized by small elements of conserved primary sequence and a common secondary

structure involving a central core and six radiating domains with helical stems. They also share many conserved tertiary structure interactions among the domains, as well as between the domains and the exons [reviewed in (1,2)]. Group II introns are prevalent in the organelle genomes of chloroplasts and mitochondria, and are also found in bacteria and archaea [reviewed in (3)]. These introns are thought to be evolutionarily related to the introns in nuclear pre-mRNAs, where catalysis is mediated by the ribonucleoprotein complexes of the spliceosome, and proceeds by a similar mechanism involving two *trans*-esterification reactions and a lariat intermediate (4).

In the green alga *Chlamydomonas reinhardtii*, the *psaA* gene is composed of three exons which are at distant positions in the chloroplast genome and are transcribed separately (5,6). The precursor transcripts are spliced *in trans* to generate the mature *psaA* mRNA (Figure 1D). Flanking the exons, the sequences which constitute the split introns have characteristic features of group II introns. The second intron is assembled from the precursors of exon 2 and of exon 3. However the first intron is actually composed of three transcripts: the precursors of exon 1 and of exon 2, and a small non-coding RNA, transcribed from a separate locus, *tscA* (7). This intron in three pieces can be seen as an intermediate between the usual introns of group II, containing their own catalytic sequences, and the introns of nuclear pre-mRNAs, where part of the structure and catalytic activity have been proposed to reside in the *trans*-acting RNAs of the snRNPs.


While some group II introns are well-studied ribozymes that can self-splice *in vitro*, others have been refractory to self-splicing in the test tube. Both types require accessory protein factors for efficient splicing *in vivo*. Because the catalytic activity is carried by the RNA, these factors may function as RNA chaperones, may assist proper folding of the intron, or may stabilize it in the active conformation (8).

\*To whom correspondence should be addressed. Tel: +41 22 379 6188; Fax: +41 22 379 6868; Email: michel.goldschmidt-clermont@molbio.unige.ch  
Present addresses:

Livia Merendino, Laboratoire Plastiques et Différenciation Cellulaire, UMR 5575, Université Joseph Fourier, BP 53, 38041 Grenoble Cedex 9, France  
Karl Perron, Department of Genetics and Microbiology, CMU, 9 avenue de Champel, CH-1211 Geneva, Switzerland

© The Author 2006. Published by Oxford University Press. All rights reserved.

The online version of this article has been published under an open access model. Users are entitled to use, reproduce, disseminate, or display the open access version of this article for non-commercial purposes provided that: the original authorship is properly and fully attributed; the Journal and Oxford University Press are attributed as the original place of publication with the correct citation details given; if an article is subsequently reproduced or disseminated not in its entirety but only in part or as a derivative work this must be clearly indicated. For commercial re-use, please contact journals.permissions@oxfordjournals.org



**Figure 1.** Characterization of *raa1-314B* and *raa1-L137H*. (A) Immunoblot analysis of total cell extracts from the wild-type (WT), the mutants *raa1-314B* and *raa1-L137H*, and the mutants rescued with the genomic DNA (cosmid #2), or the midigene construct. Top, PsaA; bottom, small subunit of Rubisco. (B) Immunoblot analysis with anti-Raa1 antibodies (affinity purified) of total cell extracts from the WT and the mutants *raa1-L137H* and *raa1-314B*. The arrow labels a non-specific band. (C) RNA blot analysis with probes (indicated on the left) specific for *psaA* exons 1, 2, 3, the 5' part of intron 2 (5'i2), *tscA* RNA or *psbA* transcript as a control. Symbols correspond to the transcripts on the splicing scheme (panel D), except the white square which labels a non-specific hybridization signal. (D) *Trans*-splicing of *psaA* mRNA. The precursors of exon1 (black square, 0.4 kb), exon 2 (black circle, 3.6 kb) and exon 3 (open circle, 2.4 kb) are spliced *in trans* to produce intermediates (exon1-exon2, asterisk, 2.6 kb or exon2-exon3, diamond, 3.8 kb) and mature *psaA* mRNA (black triangle, 2.8 kb). Splicing of the first split intron requires *tscA* RNA (black star, 0.4 kb) which is processed from longer precursors (open stars) containing *chlN*.

In *Chlamydomonas*, a set of at least fourteen nuclear loci are essential for *trans*-splicing the two split group II introns of *psaA* (9). Three of these factors have been characterized more in detail, Raa3, Raa2 and Rat1. Raa3 is necessary for *trans*-splicing of the first intron [class C, (10)]. This novel protein is part of a large complex which also contains the precursor of exon 1 and the *tscA* RNA. Raa2 is essential for *trans*-splicing of the second intron [class A, (11)]. Although Raa2 is related to pseudouridine synthases, this enzymatic activity is not required for its function *in trans*-splicing. Rat1 is required for processing of *tscA* RNA from a larger precursor and for splicing of the first intron (12). Rat1 shows sequence similarity to the NAD<sup>+</sup> binding domain of poly(ADP-ribose) polymerase, but this domain can be mutated without affecting the function of Rat1 in splicing. These factors are highly specific since they are each required for splicing of only one of the introns. However, two nuclear loci encode factors that are required for splicing of both introns, and may thus have a more general role in splicing of group II introns [class B, (9)]. Here we report the characterization of the first of these factors, Raa1, which is required for *trans*-splicing of the two introns of *psaA*. We map two distinct domains of Raa1, one involved in processing of *tscA* and splicing of intron 1, the other in splicing of intron 2. Our results show that Raa1 is a multifunctional protein required, as a component of a large membrane-bound and RNA-containing complex, for the splicing of these group II introns.

**MATERIALS AND METHODS**

**Strains and media**

Procedures for growing *C.reinhardtii* and media were described (13) (TAP: Tris Acetate Phosphate; HSM: High Salt Minimal). The strain *arg2 cwd* also has a plastome mutation conferring spectinomycin resistance and was obtained from Dr R. Loppes (University of Liège, Belgium). Mutant *L137H* was obtained by mutagenesis with 5-fluoro-deoxyuridine and UV, as described previously (6,9).

**Oligonucleotides**

See Supplementary Table I.

**Generation of the 314B mutant strain**

*arg2 cwd* cells were transformed with plasmid pARG7.8 (14) and selected on TAP medium without arginine in the dark. Screening of ca. 4 × 10<sup>3</sup> colonies yielded six strains with the fluorescence induction kinetics of mutants deficient in photosystem I or the b6f complex and a *psaA* RNA profile characteristic of *trans*-splicing mutants (9). Mutant *314B* was chosen for further study because analysis of fourteen progenies from a back-cross to the wild-type showed co-segregation of the pARG7.8 insertion with the *trans*-splicing mutant phenotype, suggesting that the mutation was tagged.

**Transformation of *Chlamydomonas***

For transformation of *raa1-314B*, 3 × 10<sup>7</sup> cells were treated with 1 µg of the appropriate plasmid DNA by the glass bead method (15) and spread on a single agar plate (HSM or TAP plus 10 µg/ml paromomycin).

The *raal-L137H* strain was transformed by electroporation [modified from (16)]. A total of  $10^8$  cells of the mutant were transformed with 1  $\mu$ g of the appropriate plasmid DNA in presence of 50  $\mu$ g of salmon sperm DNA and spread on a single plate as above. The electric pulse was applied in the absence of a shunt resistor with 700 V and 25  $\mu$ F.

### RNA blot hybridization

RNA blot analysis of *psaA* transcripts and the hybridization probes were described previously (11). The *tscA* probe was a 720 bp PCR fragment (7).

### Cloning of the *Raal1* gene

For cloning of *Raal1*, a genomic library of BamHI-digested DNA fragments from *raal-314B* was constructed in bacteriophage lambda EMBL4 (17). The library was screened by hybridization with radiolabelled pBR329, the plasmid vector of pARG7.8 (14). Restriction mapping of one of the positive phage ( $\lambda$ P23) and DNA blot hybridizations with the pARG7.8 probe, or with total genomic DNA from *C.reinhardtii* as a probe, allowed the identification of a 1.8 kb HindIII fragment that contained neither parts of the pARG7.8 plasmid, nor repetitive sequences. This fragment was used to screen a cosmid library of wild-type *C.reinhardtii* DNA (18). Three positive cosmids were obtained and used for transformation of the *raal-314B* mutant and selection on minimal medium (HSM). All three were capable of rescuing phototrophic growth. Cosmid #2 was used for restriction mapping and subcloning. A minimal region was identified spanning the ClaI to ScaI sites. Probes from this region were used to screen cDNA libraries (see below).

*Raal1* was also independently identified by transformation of *L137H* with an ordered cosmid library (19) and selection on HSM in the light. One cosmid [3(G7)] was found to restore the wild-type phenotype. This cosmid (30 kb insert) was used for restriction mapping and subcloning, and was found to contain the *Raal1* gene.

### Complementation analysis in vegetative diploids

Vegetative diploids were obtained by fusion of either *raal-L137H* or *raa2-A18* with *raal-314B*. To allow selection of diploids, paromomycin-resistant derivatives of *raal-L137H* and *raa2-A18* strains were obtained by transformation with the *aphVIII* gene [pSI103, (20)]. The *raal-314B* strain has a chloroplast mutation conferring resistance to spectinomycin.

Removal of the cell wall was achieved by incubating loopfuls of each strain with autolysin in dim light for 1 h at 25°C. Vegetative fusion was induced by spreading the cells on selective plates in fusion solution [83 mM PEG6000, 20 mM CaCl<sub>2</sub>, 20 mM glycine and NaOH (pH 8)] or a control solution [20 mM CaCl<sub>2</sub>, 20 mM glycine and NaOH (pH 8)]. Diploids were selected in dim light on TAP plates containing paromomycin (10  $\mu$ g/ml) and spectinomycin (120  $\mu$ g/ml).

The genotype of *raal-L137H* / *raal-314B* diploids was verified by PCR amplification of the *aphVIII* gene (characteristic of *raal-L137H*) with oligos *para2-5'* and *para2-3'* and of the pARG 7.8 plasmid (characteristic of *raal-314B*) with oligos pBRBam2 and ArgSau2. The genotype of *raa2-A18* / *raal-314B* diploids was verified by amplification of *aphVIII*

and of *Raa2* [which is present in *raal-314B* but partly deleted in *raa2-A18*, (11)] with oligos A18/rbcS and A18/Nci1.

### Immunoblot analysis

Protein samples were analyzed on 5% polyacrylamide gels reinforced with 0.15% linear poly(acrylamide-co-acrylic acid) (Aldrich). Proteins were transferred to nitrocellulose for 16 h at 25 V. Raal rabbit antibodies were obtained by immunization with a mixture of three histidine-tagged polypeptides corresponding to residues 1009–1349, 1510–1829 and 1830–2103 of Raal1. They were affinity purified using the three antigens immobilized on Ni NTA. The monoclonal HA-11 antibody was obtained from Eurogentec.

### Raal1 genomic and midigene constructs

The subclone g $\Delta$ Hi (116b) was obtained by inserting the 12 kb HindIII fragment from cosmid #2 into Bluescript KS (+) digested with HindIII. Subclone g $\Delta$ Sc was obtained by inserting the 11 kb EcoRI-ScaI fragment from 116b into Bluescript KS (+) digested with EcoRI and SmaI. Subclone g $\Delta$ Sa was derived from 116b by partial digestion with Sall and religation.

The cDNA clone C11 (Supplementary Figure 1) was isolated by screening a library constructed in bacteriophage lambda gt10 (21) with the 0.8 kb Sall-BstXI genomic fragment containing exons 11 and 12. The cDNA clone 405 (Supplementary Figure 1) was found by screening a library constructed in lambda NM1149 (H. Sommer, personal communication) with the 1.5 kb Sall-SphI fragment from C11 (exons 6–14). The cDNA clone 551 (Supplementary Figure 1) was obtained from a dedicated cDNA library. This library was prepared by priming synthesis of the first strand of cDNA with oligos i2 and m3 and cloning the double-stranded cDNA with EcoRI-NotI adaptors (Amersham-Pharmacia) into lambda gt10. This library was screened with a 0.68 kb EcoRI-BglII from 405 (exons 4–6). All cDNA inserts were excised with EcoRI and cloned into Bluescript KS (+). The open reading frame (ORF) that the cDNA clones defined encodes a predicted polypeptide of 2103 amino acid. It is preceded by stop codons in every frame, implying that the entire coding sequence has been obtained.

The HA-tagged midigene, mWT (also called pI2), was constructed as follows. Subclone B04 was derived by cloning the 9 kb BscI (ClaI) fragment from g $\Delta$ Hi into Bluescript KS (+) digested with BscI. A 3.4 kb Sall fragment from cDNA C11 was cloned into Sall-digested B04, to yield plasmid 269 (the Sall site used for this fusion is in exon 6). The C11 cDNA (recloned in pET15b, Novagen) was tagged with a His<sub>6</sub>-Myc-BstEII-Kan<sup>r</sup>-*BstEII* sequence introduced in the SmaI site in exon 6 (5'-ctcatgcgcggtcccaccaccaccaccacctcggcgagca-gaagctgatctccgaggagacctggtaacc-Kan<sup>r</sup>-gtaacctgctc-3', the Kan<sup>r</sup> cassette was derived from pUC4K, Amersham-Pharmacia). This tagged cDNA sequence was then transferred as a Sall-StuI fragment into plasmid 269, to yield plasmid 349. The triple HA epitope fragment (22) was then inserted as a NruI-NaeI fragment into 349 digested with BstEII to replace the Kan<sup>r</sup> cassette and obtain plasmid 490. Finally the 3.5 kb SacI fragment of 490 (one SacI site in the polylinker, the other in exon 3) was replaced with the 7 kb fragment from 116b to provide the HA-tagged midigene pI2.

The *aphVIII* gene, conferring paromomycin resistance (20) was amplified by PCR from plasmid pSL17 (S. Lemaire and J. D. Rochaix, unpublished data). The construct mΔBg was derived from pI2 by digestion with BglII (at the intron 6 and exon 7 junction of *Raal1*) and KpnI (in the polylinker) and ligation to the *aphVIII* gene (amplified with Paro Bgl 5' and Paro 3' and digested with BglII and KpnI). The construct mΔKp was derived from pI2 by digestion with KpnI (in exon 16 of *Raal1* and in the poly linker) and ligation to *aphVIII* (amplified with Paro Kpn 5' and Paro 3' and digested with KpnI). The construct mΔAM was derived from pI2 by digestion with AgeI, blunting with mung bean nuclease, digestion with MscI (in exon 2 of *Raal1*) and religation. In a second step, the *aphVIII* gene was amplified with Paro Mlu 5' and Paro Mlu 3', digested with MluI and inserted in the MluI site, upstream of *Raal1*. The clone pI4 was similarly derived from pI2 by inserting the *aphVIII* gene in the MluI site. The construct mΔNS was derived from pI4 by digestion with NruI and SmaI (in exons 2 and 4 of *Raal1*, respectively) and religation.

### RT-PCR analysis of *Raal1* transcripts

Total RNAs were isolated as described (23) and then treated with RQ1 DNase (Promega). Reverse transcription was performed at 50°C with M-MLV reverse transcriptase (Promega). To analyse three different regions (exons 1 and 2, exons 12 and 13 and exons 17 and 18) of the *Raal1* transcript, the corresponding first strand cDNA was synthesized with oligos AUG 2, middle 4 or oligo dT, respectively. PCR amplification was performed with the following pairs of oligos: AUG5'-1 and AUG3'-1; m5'-3 and m3'-3; STOP5'-6 and STOP3'-6, respectively. The PCR products span two exons so that amplification of the cDNA can be distinguished from amplification of any contaminating genomic DNA which would include the intervening intron. As control, RT-PCR analysis was also performed for *Cbl* transcripts (24), using oligo dT to synthesize the first strand cDNA and oligos UP and DOWN for the PCR. The Arg7-*Raal1* fusion transcript in *raal1-314B* was revealed by synthesizing the first strand cDNA with oligo middle 4 (in exon 14 of *Raal1*) and PCR amplification with oligos Arg7ex11 and m3'-3 (in exon 13 of *Raal1*).

### Sequence analysis of *Raal1*

The sequence of the *Raal1* cDNA is deposited in the EMBL database under accession no. AJ605114. The genomic sequence is now available (scaffold 33) in the draft of the *C.reinhardtii* genome, at the Joint Genome Institute of the US Department of Energy (<http://genome.jgi-psf.org/chlre2/chlre2.home.html>). Comparison of the cDNA and genomic sequences shows that the *Raal1* gene is composed of 18 exons with very different sizes (from 60 to 2473 bp). The polyA tail is preceded 11 bp upstream by the TGTA consensus signal typical of *C.reinhardtii* (25).

Transit peptide sequences were predicted using the algorithms ChloroP (26) (<http://www.cbs.dtu.dk/services/ChloroP/>) and Predotar (27) (version 0.5; <http://www.inra.fr/predotar/>). Protein secondary structure was predicted with GORIV (28) ([http://npsa-pbil.ibcp.fr/cgi-bin/npsa\\_automat.pl?page=npsa\\_gor4.html](http://npsa-pbil.ibcp.fr/cgi-bin/npsa_automat.pl?page=npsa_gor4.html))

### Cell fractionation

Cells were lysed with a nebulizer and chloroplasts were purified on Percoll gradients as described previously (10) in the presence of protease inhibitor mix A (PIM A: 1 mM ε-caproic acid, 0.5 mM benzamidine, 0.6 μg/ml leupeptin, 2 μM E64, 2 μM pepstatin and 0.5 mM AEBSF). Chloroplasts were resuspended in hypotonic buffer [10 mM HEPES (pH 7.8), 5 mM MgCl<sub>2</sub>, 1mM DTT] supplemented with protease inhibitor mix B (PIM B: 1 mM ε-caproic acid, 0.5 mM benzamidine, 0.6 μg/ml leupeptin, 2 mM ortho-phenantroline and 1/200 Protease Inhibitor Cocktail P8849 from Sigma). For further fractionation, the chloroplasts (250 μl, 7 mg protein/ml) were lysed by sonication, the solution was adjusted to 20 mM HEPES (pH 7.8), 50 mM KCl and 10 mM MgCl<sub>2</sub> (HepKM) and subjected to centrifugation at 100 000 g for 30 min at 4°C in a TLA45 rotor (Beckman). The supernatant was recovered for analysis, and the pellet was washed by resuspension in HepKMS (0.2 M sucrose, HepKM, PIM B) and centrifugation at 50 000 g for 15 min at 4°C in a TLA45 rotor. The supernatant was discarded and the pellet was resuspended in HepKM plus PIM B.

For analysis of the protein complexes, cells were collected by centrifugation, resuspended at a density of 2.5 10<sup>8</sup> /ml in HepKM plus PIM A, and lysed by freezing and thawing. The lysate (1 ml aliquot) was subjected to centrifugation at 18 000 g for 20 min at 4°C in a TLA45 rotor (Beckman). The pellet containing the membranes was resuspended in HepKM, 0.2 M (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, PIM B. Where needed, 100 μl of RNaseA were added [10 mg/ml in 100 mM Tris-HCl (pH 7.5) and 10 mM Na acetate) or the same amount of buffer without RNase (mock treatment), followed by incubation at 16°C for 20 min. Membranes and insoluble material were removed by two rounds of centrifugation at 120 000 g for 20 min in a TLA45 rotor, and the supernatant was loaded on sucrose gradients (10 ml; 0.1–1.3 M sucrose, HepKM, PIM A). After centrifugation at 115 000 g for 17 h in the SW40 rotor (Beckman), 10 fractions (1.1 ml) were collected from the bottom by puncturing the tube. Sedimentation was calibrated with the High Molecular Weight Calibration Kit from Amersham-Pharmacia.

## RESULTS

### Characterization of the *raal1-314B* mutant deficient in splicing of *psaA* intron 2 and cloning of the *Raal1* gene

Insertional mutants were generated by transformation of a *Chlamydomonas* arginine auxotroph with a plasmid containing the wild-type *Arg7* gene. Mutant *raal1-314B* was unable to grow phototrophically (data not shown) and had no detectable PsaA protein (Figure 1A, lane 2). RNA blot hybridization (Figure 1C) showed that the pattern of transcripts that accumulate in *raal1-314B* is characteristic of a block in trans-splicing of the second intron of *psaA* [class A, (6)]. Probes specific for each of the three exons hybridize to mature *psaA* mRNA (black triangle, see Figure 1D) in the wild-type (Figure 1C, lane 1) but not in the mutant (lane 2). In the *raal1-314B* mutant, the probe for exon 3 only detects an unspliced precursor of exon 3 (open circle), which is also present in the wild-type. In the mutant, probes for exon 1

or for exon 2 reveal a splicing intermediate (asterisk) with exons 1 and 2 spliced together, which migrates slightly faster than mature *psaA* mRNA. This intermediate, with the 5' part of intron 2 still attached to exon 2 (5'i2 probe, lane 9), is not detected in the wild-type (lane 8). The precursor of exon 1 (black square) over accumulates in *raal-314B*, as was observed previously for all other *psaA* splicing mutants (6). Because of these defects in splicing of *psaA* transcripts, we have called the mutant *raal-314B* (RNA maturation of *psaA*).

Using the Arg7 insertion as a molecular tag, we cloned the *raal* mutant locus. We then obtained wild-type *RAA1* cosmids that could rescue the *raal-314B* mutant by transformation. These results also showed that the *raal-314B* mutation is recessive to the wild-type. Additionally, Southern blot analysis of *raal-314B* DNA indicated that the locus was rearranged (data not shown), confirming that the locus that we cloned was indeed *RAA1*. In the strains transformed with cosmid #2, PsaA protein is recovered (Figure 1A, *314B* genomic, lane 3) and the fully spliced mRNA accumulates (Figure 1C, lane 4, black triangle). The splicing intermediate containing the 5' part of intron 2 is not detected any more (Figure 1C, lane 11, asterisk) and the levels of exon 1 precursor are much lower than in the parental mutant (Figure 1C, lanes 2 and 4, black square).

To map the gene within the 34 kb insert in cosmid #2, we tested subclones for their ability to restore phototrophic growth of *raal-314B* by transformation and identified a minimal region of 7.4 kb. We obtained cDNA clones from this region which encode a predicted polypeptide of 2103 amino acid (See Materials and Methods). To ascertain that the appropriate cDNA had been cloned, a 'midigene' was constructed with the genomic sequence containing the 5' part of the gene (including the promoter) fused to the cDNA from the 3' part (Supplementary Figure 1D). This construct transformed the *raal-314B* mutant to prototrophic growth as efficiently as cosmid #2 (data not shown) and restored normal levels of PsaA protein (Figure 1A, lane 4). The midigene also rescued *trans*-splicing of the *psaA* mRNA (black triangle, Figure 1C, lane 6), even if splicing intermediates are still detected with probes for exon 2 (black circle), exon 3 (diamond), exon 1 (black square) and 5'i2 (black circle, lane 13).

#### ***L137H*, defective in splicing of both *psaA* introns, is an allele of *Raal***

In parallel, we have characterized another mutant strain (*L137H*), that was previously shown to be affected in *trans*-splicing of both the *psaA* introns (class B) and was obtained by UV mutagenesis (6). *L137H* was unable to grow phototrophically (data not shown) and did not accumulate detectable levels of the PsaA protein (Figure 1A, lane 6, the mutant is labelled *raal-L137H* in anticipation of evidence presented below which shows that *L137H* is allelic to *raal-314B*). The pattern of transcripts in this mutant (Figure 1C) is characteristic of a block in *trans*-splicing of both introns (6). Probes specific for each of the three exons fail to detect mature *psaA* mRNA in the mutant (black triangle, lane 3), but revealed unspliced precursors of exon 1 (black square), exon 2 (black circle) and exon 3 (white circle). The exon 2 precursor, with the 5' part of intron 2 still attached to exon 2, was also detected with the 5'i2 probe (black circle, lanes 3 and

10). As noted above for *raal-314B* and other *trans*-splicing mutants, the precursor of exon 1 over accumulates.

To clone the mutant locus, we transformed the *L137H* strain with an indexed genomic library (19) and screened the cosmids for their ability to restore phototrophic growth of the mutant cells. Surprisingly, we identified *Raal* as the nuclear gene that complements the photosynthetic defect of the *L137H* strain, suggesting that *L137H* and *raal-314B* are allelic. These results also imply that the *L137H* mutation is recessive to the wild-type. However, Southern blot analysis did not show any alteration in the *RAA1* locus of the *raal-L137H* mutant (data not shown). In the *L137H* cells transformed with either the cosmid containing the *Raal* locus or the midigene, synthesis of the PsaA protein is recovered (Figure 1A, lanes 7 and 8) and fully spliced mRNA accumulates (black triangle, Figure 1C, lanes 5 and 7).

To confirm that *raal-314B* and *L137H* are indeed allelic, we constructed vegetative diploids to test whether the two mutants could genetically complement (Supplementary Data). The observed lack of complementation, together with the fact that transformation with the *Raal* gene can rescue *psaA* splicing in both mutants, indicate that *raal-314B* and *L137H* are allelic. Furthermore full-length *Raal* protein was not detectable by immunoblotting in the two mutant strains, *raal-314B* and *L137H* (Figure 1B, lanes 2 and 3). We thus refer to the two mutants as *raal-314B* and *raal-L137H*.


#### **The *raal-L137H* strain is also defective in maturation of *tscA* RNA**

*HN31*, another mutant defective in *trans*-splicing of both introns of *psaA* (class B), is allelic to *L137H* (9). Interestingly, *HN31* was previously shown to be defective in the maturation of *tscA* (29), the small non-coding RNA required for splicing of intron 1 (Figure 1D). Mature *tscA* RNA could not be detected in *raal-L137H* (Figure 1C, lane 10, black star). In the mutant, several larger transcripts were more prevalent (white stars). These precursor transcripts also contain sequences of *chlN*, which is downstream of *tscA* (29). However, normal levels of mature *tscA* RNA were recovered in the *raal-L137H* strains transformed with the *Raal* cosmid or the 'midigene' (Figure 1C, lanes 12 and 14). Mature *tscA* RNA was observed in *raal-314B* (lane 9), as expected for a strain competent in splicing of intron 1.

#### **A truncated form of *Raal* is expressed as a fusion to *Arg7* in *raal-314B***

Our data indicate that the mutants are allelic, but their splicing defects are different: in *raal-L137H* there is a block in maturation of the *tscA* RNA and in splicing of both *psaA* introns, while in *raal-314B* only splicing of intron 2 is defective. This could be explained if *Raal* is partly functional in *raal-314B*, and more severely impaired in *raal-L137H*. To further investigate the basis for the different phenotypes, we analysed the expression of *Raal* transcripts by RT-PCR (Figure 2).

We designed primers to amplify a 5' region (exons 1 and 2), an intermediate region (exons 12 and 13) and a 3' region (exons 17 and 18) of *Raal* mRNA (Figure 2A). In the wild-type strain the *Raal* transcript could be detected with the three primer pairs (Figure 2B, lane 2). In *raal-L137H* (lane 3) the three parts of the transcript were also amplified,



**Figure 2.** Expression of *Raa1* in the mutants. (A) Schematic representation of the *Raa1* mRNA with the fragments amplified in the RT-PCR analysis. (B) RT-PCR analysis with RNA from the wild-type (WT), *raa1-314B* and *raa1-L137H*, for three parts of *Raa1* (panel A) and the *Cbl* transcript as a control. The + or – signs indicate whether reverse transcriptase was included (lanes 2–4) or omitted as a control (lane 1). (C) The 3' part of *Raa1* is expressed as a fusion to *Arg7* in *raa1-314B*. The structure of the genomic DNA in *raa1-314B* is shown with the last exons of *Arg7* (open bars), a fragment of the cloning vector pBR329 from pARG7.8 (thick black line), the 3' part of *Raa1* intron 9 (thin black line) and the exons of *Raa1* (shaded bars). The sequence of a short hairpin-loop created at the insertion site by duplication of a 15 bp sequence of *Raa1* is indicated above the scheme. A cryptic donor site in exon 13 of *Arg7* (black arrow) and the acceptor site of exon 10 of *Raa1* (black arrow) are used for splicing across the pBR329 sequence to generate an in-frame fusion transcript as indicated at the bottom.


indicating the presence of the *Raa1* mRNA. However, the full-length *Raa1* protein was not detected by immunoblot analysis (Figure 1B, lane 2).

In *raa1-314B* (Figure 2B, lane 4) the intermediate and 3' regions of the *Raa1* transcript could be amplified. However, no amplification of the 5' region (exons 1 and 2) was detected. This qualitative analysis suggested that the 5' part of the gene was altered but that a truncated form of *Raa1* might be expressed. To investigate this further, we sequenced a genomic fragment from the *raa1-314B* mutant ( $\lambda$ P23, Supplementary Figure 1A) and found that the *Arg7* gene and part of the pBR329 vector were joined to intron 9 of the *Raa1* gene (Figure 2C). Indeed, the presence of an *Arg7*-*Raa1* fusion transcript was revealed by RT-PCR with a 5' primer in exon 11 of *Arg7* and a 3' primer in exon 13 of *Raa1*. Its sequence showed that the fused mRNA is spliced using a cryptic 5' donor site within the last exon of *Arg7* (exon 13) and the 3' acceptor site for exon 10 of *Raa1* (Figure 2C). The exons are fused in-frame, such that a polypeptide is predicted with all but the last 14 residues of the *Arg7* polypeptide

(i.e. arginino-succinate lyase) fused to the 644 C-terminal residues of *Raa1*. *Arg7* is a nucleus-encoded plastid protein (30), so the fusion protein is also predicted to be targeted to the chloroplast. It thus appears that in *raa1-314B* the C-terminal domain of *Raa1* is sufficient for processing of *tscA* RNA and splicing of intron 1, but not for splicing of intron 2. The predicted fusion protein of 116 kDa was not detected by immunoblotting probably because of low expression or high turnover, or because it was masked by non-specific bands.

#### Different domains of *Raa1* are required for its functions in *psaA* trans-splicing

Since the C-terminal domain of *Raa1* expressed in *raa1-314B* is sufficient for processing of *tscA* and for splicing of *psaA* intron 1, we sought to determine which parts of the protein are required for these functions or for splicing of intron 2. We conducted an initial deletion analysis using genomic fragments with 3' truncations of *Raa1* (Figure 3C, construct names beginning with g for genomic). We transformed the



**Figure 3.** Deletion analysis of Raa1. (A) Immunoblot analysis with anti-HA antibodies of total cell extracts from transformants of *raa1-L137H* (lanes 1–10) or *raa1-314B* (lanes 11–16) with the midgene constructs depicted in panel C. (B) RNA blot hybridization of total RNA from the wild-type (lanes 1 and 12), the parental mutants (lanes 2, 3, 13 and 14) and the transformants of *raa1-L137H* (lanes 4–11) or *raa1-314B* (lanes 15–20). Symbols are as in Figure 1. (C) Schematic representation of the Raa1 polypeptide predicted for the wild-type or for the deletion derivatives (open bars, see Supplementary Materials and Methods). Names shown on the left begin with 'g' for genomic constructs, with 'm' for the midgene tagged with the HA epitope (white triangle). The part of Raa1 that is fused to Arg7 in *raa1-314B* is also represented as an open bar. Black lines at the bottom represent the domains involved in splicing of intron 1 or intron 2. In the columns marked 'phototrophy', growth of the two mutants on minimal medium is indicated by '+', lack of growth by '-'. Splicing of intron 1 (i1) or of intron 2 (i2) is indicated by '+', lack of splicing by '-', as deduced from the RNA blot analysis in panel B. (nd: not determined, na: not applicable). The position of the restriction sites used for the deletions are shown at the top (they truncate the predicted polypeptide sequence after the following amino acid residues: AgeI, 85; NruI, 300; MscI, 731; SmaI, 944; Sall, 1136; BglII, 1173; Scal, 1549; HindIII, 1785 and KpnI, 1853). Black triangles represent the 38 amino acid repeats (not drawn to scale).

**A**

MRRHPTCGIATPQAYGVAWPRLPPVAPRPASGQLPPRCRRAVLCAARRPAGGTRAASGAD 60  
 AGGSAGEGEGATAAADSHGTRSRTGLEGRSASGSVAVPAGIGGGAGSDSDGDAESAQRSG 120  
 PRRRGRPPGMRQTMVAVPPTPNAGSAPATALAASSAVQQTVGQAANSPGSGSDGEGSGLA 180  
 TRRRGRPRRSTADASGGGCSIGGSTRAVTSAADYEASAAAA PMPGMPLASGRAE PQQPQ 240  
 QPQSQPQLQQQTEAGGSTATAPGGAGAGSSSGSGSTQRDGRSGAGGGAGGPLSRLLSQRS 300  
 RAMGGVGGAAPNATMPGGAAPPDSDLQFASPMSSFEFAPERSSESDETEGGAVGRSGGAT 360  
 SRRQT P P P P P D P Q P P H Q Q A P P Q Q Q Q P P D A A G A A S A G S M A S T D W M L G S M L E D G E A G S G L A F 420  
 GSPGGASSGSDQDFIITADLDVVGSTAPPLNLDLSDLDWGDDTGAGGGWGWGDASTGGSA 480  
 GSAMGALDGVGVGLGAATSGIIGIWGGDASASA AVAGDASALGQTPGSAAAA L D S W P G V L 540  
 QDDEAP P Q P Q P Q P P Q P Q P Q Q A P Q A P Q A P Q A L Q T P Q A L Q A P Q G R P P V L L K G P A A E S G P S R 600  
 PVPAGDGGGAEATQSGSRMDSHPHPGLDSDFDLVSSYDAEMSY P D A V D C L L P E G V A A A A A 660  
 AAGAGAP G G G G W S E G V L D S M A D A G L P D L S Y S G S G G G S G S G G G G S S E G G K D S A H E A R L P V 720  
 AVAGAGRAPVASSSTPIAAAPAVPATAPAAAVPAPAAAKAAAGA A K A A T G A A K G G A Q A A S 780  
 GSSGSSSSGSS IDGSSRTAAAF A E L L A G G S S A H G A G R A D P L V Q R L A R V L S Y V E S E L L G A P E 840  
 E A A A A A A D A A A E L P G V S L P P L P R P L V P A A A Q P L E L F A E P P A T A R L L A A A E A A E A G G L A A P 900  
 PRQVEAPPSADEWALLGGAAPA S A A S A A S A A S A A S A A G T S R A G P G R Y G G G A G Y E D D T D G G 960  
 S S W E R E R E R E R E G D G E A D F E R R L E A G P A Y V A Y L R A V V L G G L G G D S G G G S G G G G G G G R G G G G P 1020  
 VQAGAVQAALLEQVEAAGDWLQLCWLLEAAAA Q G V A L G P R A A A A A F K Q A A G L V A P Q G A L P 1080  
 RSLAAATTTATTSGLEAAAYRQLCDRLAAGVAALGT SV S A A A S T H T A A R A A G G R A S T S R 1140  
 RGVPPGITAGGR G G G G G G A G S H V S A A S G S D V A Q I S Y G M G A L Q I T C P Q L Y G A V L H V S A S Q L 1200  
 AALTARRRAAASLT A A A A A A L R R S A A A V P T D S A G R G G G G G G D G V R R V S D R E I M M A A A A A S 1260  
 SAVQPNHASELP P Q Q L Q Q Q Q Q Q Q A T S G R G P L S G S A A T A K S A L T A A A G G V W T A A D L A A L A W 1320  
 GVANAYAGSAAVAASAAATADSCATASKAQRQQRQQRQ P P P R P P K H R Q L Q G T Q Q P Q P 1380  
 QPPLPLPVPGGDWLRSLV G A S Y E L M A T E G G G G G G G G V G P R Q L W G L A W S F A R L G Y A P S Q E 1440  
 WMLALLSRAEACLSQFDTEGLCRLWLWALAAMDYVPERLWLRVAVAGQLQARARDFTPDQVV 1500  
 TLHCALARLGYAPRPEVCVALHAAAA RL M P L M T G P Q L A A L A H A A A S F A R W R P G P G F L V A A 1560  
 ASATGGAAAPVPSASSPAAAVQMPAGLLMALATRVLVGMQRATTEAAAVASSPAGGEAGA 1620  
 WA Q P Q P Q P T A L A A A A V D G S D L S M A L W A L A L L R R Q Q Q Q Q Q Q P A A D V L Q L M P E W L A A W W A A 1680  
 AAEPVAATFDATCVS Q S L W A L A E L R E T P G L P H S G A A A A A A A A S G G A A S S T Y A D A A R D A 1740  
 GQAAAQAAGAVAALLAALVPQLGQAATADLSTTIAALADLQYRPSDQWMALFTA E A R R 1800  
 RLGTATATATATGTATTAATATNEDHGLIAYGLAVLGWPLSEAWVQELAAGGYRAMAGAS 1860  
 GEG L A L L L W G L S A R G W S T A S G R F W D T V F R E S G S K W D S C G P R G A V L L Y C A V A D M M P P G Q E P 1920  
 PIPWQRQLVKALRLRVRPRPTALLLPAALRTAAGGCLGPLTRAERQQLWQAGAAAAAAR 1980  
 AAGGGAGGIGLVLGAEAYTAVPGSGLAGAVWWSALPQPHRPPSLPGVKGWAWPQGGPQG 2040  
 SGADEVQDPPEVGVLASRVLLPCELCEPEEHPALAAVAEGRS W W A H D V A A E L A R R W G L V 2100  
 RWS\*

**B**

**Raa1**

1423 LWGLAWSFARLGYAPSQEWMLALLSRAEACLSQFDTEG  
 1461 LCRLWLWALAAMDYVPERLWLRVAVAGQLQARARDFTPDQ  
 1499 VVTLHCALARLGYAPRPEVCVALHAAAA RL M P L M T G P Q  
 1772 LSTTIAALADLQYRPSDQWMALFTA E A R R R L G T A T A T A  
 1826 HGLIAYGLAVLGWPLSEAWVQELAAGGYRAMAGASGEG

**Raa1 consensus**

L--L-wALA-LGY-PS--W--AL-A-A-R-----T---

**Tbc2**

\* \* \* \* \* \* \*

L----R-----RFSP-HL--LL--L--LG-RPPPEW--A

**PPR**

-TYNALINAYAK-G--EEA--LY--M---G--PN-

**TPR**

--AY--LG--Y-----YE-A---Y-..KAL-LNPNN

helix A

helix B



two *raal* mutants and selected for phototrophy on minimal medium, an indication of whether *psaA* splicing is restored. However, in this approach, when phototrophy is not rescued, for instance because only one of the two introns is spliced, transformants are not recovered and the effect of the mutations on the excision of the individual introns cannot be analysed in detail. Therefore, we also constructed deletions of the *Raal* midigene carrying an *aphVIII* cassette, allowing selection on paromomycin (20) and included the HA epitope tag (Figure 3C, construct names beginning with m for midigene). Many of the transformants which were initially positive in anti-HA immunoblots gradually ceased to express *Raal* when they were sub-cultured on plates for a few weeks, most likely because of gene silencing. Therefore we used aliquots of the same culture for immunoblotting (Figure 3A) and for RNA analysis (Figure 3B). Some lines were particularly prone to silencing: only low levels of HA-tagged *Raal* could be detected in the transformant *raal-L137H* mΔNS #4, and none in *raal-L137H* mΔNS #7 (Figure 3A, lanes 8 and 9). Likewise only one line stably expressing *Raal* could be recovered for mΔAM in *raal-L137H* (mΔAM #3, lane 10) or for mΔNS in *raal-314B* (mΔNS #1, lane 14), and none for mΔBg in *raal-314B*.

In *raal-314B*, constructs mΔKp, gΔHi and gΔSc rescued phototrophy (Figure 3C). Indeed, spliced *psaA* mRNA was detected in the transformants mΔKp #4 and #7 (Figure 3B, black triangle, lanes 16 and 17). These results define a domain upstream of *ScaI* which is sufficient for splicing of intron 2. This part of *Raal* and the domain expressed as an *Arg7-Raal* fusion in *raal-314B* are thus capable of intragenic complementation. However in *raal-314B*, mΔBg and gΔSa did not rescue phototrophy (Figure 3C), indicating that a region between the *BglIII* and *ScaI* sites is essential for splicing of intron 2. In the *raal-L137H* mutant, even the shortest 3' deletion, mΔKp, failed to rescue phototrophy (Figure 3C). In these *raal-L137H* mΔKp transformants, no *psaA* mRNA was detected (Figure 3B, exon 3, black triangle, lanes 7 and 8), but the intermediate with exon 2 spliced to exon 3 accumulated (exon 3, diamond). This confirms that mΔKp constructs can still support intron 2 splicing and also implies that the deletions at the C-terminal end of *Raal* affect splicing of intron 1 (Figure 3C). In *raal-L137H*, mΔBg and gΔSa did not rescue phototrophy (Figure 3C), and in mΔBg transformants, the pattern was identical to the *raal-L137H* mutant (Figure 3B, lanes 5 and 6 versus lane 2). Thus the domain required for intron 2 splicing is affected in this deletion, confirming the analysis of the *raal-314B* transformants.

To analyse the N-proximal part of the protein, we constructed internal deletions that leave the putative transit peptide intact. The mΔAM deletion rescued phototrophy in both mutants (Figure 3C), and restored splicing of both introns to produce mature *psaA* mRNA (Figure 3B, exon 3, black triangle, lanes 11, 19 and 20). Thus a large domain in the N-terminal part of *Raal* is not essential for splicing of either

intron. Transformation of *raal-314B* with mΔNS does not rescue phototrophy or splicing of intron 2: the mature *psaA* mRNA is not made (Figure 3B, exon 3, black triangle, lane 18) and the exon1–exon2 intermediate is still accumulated (Figure 3B, 5'i2, asterisk, lane 18). Comparison of the mΔNS and mΔAM *raal-314B* transformants indicate that part of the *MscI* to *SmaI* segment is necessary for intron 2 splicing. In the *raal-L137H* mΔNS transformant #4, the intermediate with exon 2 spliced to exon 1 (Figure 3B, 5'i2, asterisk, lane 9 versus lane 2) indicates that splicing of intron 1 is restored, in spite of the low levels of *Raal* protein expressed in this strain (Figure 3A, lane 8). This confirms that the C-terminal domain of *Raal* is sufficient for splicing of intron 1. Only a very faint band is observed for *raal-L137H* mΔNS #7 (Figure 3B, 5'i2, asterisk, lane 10) which does not detectably express *Raal* (Figure 3A, lane 9).


In this analysis, the domain of *Raal* required for splicing of intron 1 is not distinguishable from the domain required for *tscA* processing: in the mΔKpn and mΔBg transformants of *raal-L137H* where intron 1 is not spliced, mature *tscA* fails to accumulate (Figure 3B, *tscA*, black star, lanes 5–8). Conversely intron 1 is spliced and mature *tscA* RNA is formed in the *raal-314B* mutant, or in the mΔNS and mΔAM transformants of *raal-L137H* (Figure 3B, *tscA*, black star, lanes 3, 9 and 11).

Taken together, these results delineate two functional domains of *Raal* (Figure 3C). A central region involved in splicing of intron 2 extends from *MscI* to somewhere between *BglIII* and *ScaI*, with a possible additional contribution from the N-terminal region upstream of *AgeI*. A C-terminal region involved in splicing of intron 1 and *tscA* maturation starts at the breakpoint of the *Arg7-Raal* fusion in *raal-314B* (Figure 3C, last open bar at the bottom) and ends somewhere between *KpnI* and the C-terminus. The two domains may partly overlap in a short region to the left of the *ScaI* site.

### **Raal1, a novel protein with internal repeats**

The Predotar and ChloroP algorithms predict that *Raal* is imported into the chloroplast with an N-terminal transit peptide of 54 residues (Figure 4). This localization was confirmed by cell fractionation experiments (see below). A striking feature of the predicted *Raal* polypeptide (Figure 4A) is the occurrence of numerous stretches of repeated amino acids, such as (Ala)<sub>9</sub>, (Gly)<sub>9</sub>, (Gln)<sub>8</sub> or (Pro)<sub>4</sub>. *Raal* also contains repetitive sequences like (Pro-Gln)<sub>4</sub> or (Ser-Ala-Ala)<sub>5</sub>. Similar repetitive stretches also occur in other *C.reinhardtii* proteins, notably in another *psaA* splicing factor, *Raa3* (10) and in other proteins involved in chloroplast gene expression (see Discussion). *Raal* harbours five copies of a 38 amino acid repeat (Figure 4B). The first three repeats are in tandem arrangement (residues 1423–1536), the other two are separate, further on the C-terminal side (1772–1809 and 1826–1863). They are related to the PPPEW repeats that are found in *Tbc2*, a

**Figure 4.** The predicted *Raal* polypeptide. (A) The putative transit peptide (54 amino acids) is shown in italics and underlined (thin wavy line). Repeats of four or more identical consecutive amino acid residues are shaded, as well as repeats of alternating residues. The 38 amino acid repeats are underlined (thick black line or double line). (B) Alignment of the 38 amino acid repeats. Residues that are conserved in at least three of the five repeats are shaded. Segments predicted to form  $\alpha$ -helices are underlined (28). The consensus shows residues that are conserved in 3 repeats (uppercase), and 4 or 5 repeats (bold uppercase). The lowercase w represents a position where the aromatic residue W is present twice and Y once. The consensus for the 38 amino acid repeat (PPPEW) of *Tbc2* (31) is shown below, with asterisks marking identical residues. The 34 amino acid TPR repeat with its two helical domains (underlined) and the 35 amino acid PPR repeat (presumptive helical domains underlined) are shown for comparison (51).



**Figure 5.** Raa1 is part of a large RNase sensitive complex. (A) Subcellular localization of Raa1. Fractions were prepared from a transformant of *raa1-314B* expressing the HA-tagged Raa1 protein. The four lanes contain total cell extracts (Tot), the chloroplast fraction (Cp) and a chloroplast lysate fractionated by centrifugation into a soluble fraction (Sup) and a crude membrane fraction (Pellet). Immunoblots with antibodies against the HA epitope (Raa1::HA), PsaA (an integral membrane protein of Photosystem I), the small subunit of Rubisco (a soluble stromal protein) and eiF4A (a cytoplasmic translation factor, the nature of the slightly larger cross-reacting band in the chloroplast fraction is not known). Equal amounts of protein were loaded in each lane (60  $\mu$ g). (B) Raa1 is part of a large RNase-sensitive complex. A soluble extract was prepared by salt treatment of a total membrane fraction, and subjected to sucrose gradient sedimentation. Ten fractions were collected and analyzed by immunoblotting with a monoclonal anti-HA antibody. The extract was either untreated, mock treated for 20 min at 16°C, or treated with RNase before loading on the gradient. The sedimentation of protein markers of known size is indicated (in kDa) above the lanes.

nucleus-encoded protein of *C.reinhardtii* involved in translation of the chloroplast *psbC* mRNA [(31); see Discussion]. Four of the 38 amino acid repeats (repeats 2 to 5) are retained in *raa1-314B*, which can process *tscA* RNA and splice intron 1. The last repeat is disrupted in the  $\Delta$ Kpn deletion which can splice intron 2 but not intron 1, and two of the repeats (2 and 3) are in the region where the two functional domains of Raa1 may overlap upstream of the ScaI site. No other significant similarity was found between the deduced Raa1 polypeptide and sequences deposited in public databases.

### Raa1 is a membrane-associated chloroplast protein

To investigate the subcellular localization of Raa1, we used a strain expressing the HA-tagged wild-type midgene. In this construct, the genomic fragment, which extends 3.8 kb upstream of the ATG and also includes the 5' part of the gene with the first five introns, should preserve normal expression and regulation. As observed for the native protein revealed with the Raa1 antibody (Figure 1B), the HA-tagged protein also migrates more slowly than expected for its calculated molecular weight of 205 kDa (Figure 3A, lanes 1 and 11). This may be due to the unusual sequence features of the protein and its large size.

In order to determine the subcellular localization of the Raa1 protein, chloroplasts were prepared by Percoll gradient centrifugation. Raa1 was found in the chloroplast fraction, which also contains the other chloroplast proteins PsaA and Rubisco, but only trace levels of the cytosolic translation factor eiF4A (Figure 5A, lane 2). When isolated chloroplasts were further fractionated, Raa1 was enriched in the crude membrane pellet along with the integral thylakoid protein PsaA (lane 4), and depleted from the supernatant which contains the soluble protein Rubisco (lane 3). Raa1 thus fractionates as a chloroplast protein, as expected from the genetic data, although its presence in other organelles, such as mitochondria, is not ruled out. The membrane association of Raa1 was confirmed by sucrose gradient equilibrium centrifugation (data not shown).

### Raa1 is part of a large RNase sensitive complex

The Raa1 protein can be released from the membrane fraction by treatment with moderate concentrations of salts such as 0.2 M ammonium sulphate (data not shown). An extract was prepared by washing membranes with salt and was fractionated by sedimentation on sucrose gradients (Figure 5B). Part of Raa1 was found in the lower part of the gradient (fractions 2–4), as a large complex which sediments just ahead of the 670 kDa molecular mass marker. Part of Raa1 was also found in a complex that sediments more slowly (fractions 6–8), where it is associated with Raa2 (32).

When the ammonium sulphate extract was incubated with RNase before sucrose gradient sedimentation, the position of the large complex shifted towards the top of the gradient. This shift was not observed when the extract was mock-incubated in the absence of RNase. The sensitivity to RNase shows the presence of RNA in the large Raa1 complex.

## DISCUSSION

### Raa1 is a multifunctional splicing factor

We have characterized a new mutant, *raa1-314B*, defective in *trans*-splicing of only the second intron of *psaA* (class A), and a mutant that fails to splice both introns, *raa1-L137H* (class B). The two mutants are clearly allelic since they fail to complement in genetic tests and both can be rescued by the wild-type *Raa1* gene. Thus *Raa1* has at least two distinct functions, in splicing of intron 1 and of intron 2. In *raa1-HN31*, which is an allele of *raa1-L137H* (9), there is no detectable mature *tscA* (29), the small RNA required for splicing intron 1 (7). The same defect is also observed in *raa1-L137H*, but in contrast *raa1-314B* splices intron 1 and accumulates mature *tscA* RNA. Thus the function of Raa1 in splicing of intron 1 may be direct (in this case Raa1 would have three distinct roles), or may be indirectly mediated by *tscA*. Two other factors which are required for intron 1 splicing and for processing of *tscA* were described recently, Rat1 and Rat2 (12). Like for Raa1, it is still unclear whether they are directly or indirectly involved in intron 1 splicing. The pattern of longer *tscA* transcripts that accumulate in *raa1-L137H* and *raa1-HN31* and in the *rat1* and *rat2* mutants are similar (12,29). Thus a complex processing pathway requiring at least three factors (Raa1, Rat1 and Rat2)

is involved in the maturation of *tscA* RNA from longer precursors.

The partial defect in *raal-314B* can be explained by our observation that in *raal-314B* the 3' part of *Raal1* is expressed as a fusion to *Arg7*: this domain is thus sufficient to promote *tscA* maturation and splicing of intron 1. Our deletion analysis of *Raal1* confirms that this 3'-terminal domain is required for *tscA* maturation and splicing of intron 1, while a more central domain is required for splicing of intron 2. However we did not reveal separate domains required for intron 1 splicing and for *tscA* RNA maturation. This is the first time that in *Chlamydomonas* a factor involved in splicing of the two introns of *psaA* (class B) is cloned and characterized. Interestingly, of the fourteen nuclear loci that are required for maturation of *psaA* transcripts, only two belong to this class. The other factors described to date, *Raa2* (11), *Raa3* (10), *Rat1* and *Rat2* (12) were shown to be required for either one of the two intervening sequences (classes A and C). The presence of the two functional domains in the same factor suggests that *Raal1* may coordinate splicing of the two *psaA* introns. Mutants defective in splicing of either intron 1 or 2 show that the intervening sequences can be excised independently from each other and in either order (6). However, a mechanism that allows coordinate splicing of the two introns may improve the efficiency of maturation of *psaA* transcripts. Chloroplast factors with dual targets were found in *Arabidopsis* where *PGR3* is required for expression of the *petL* operon and of an *ndh* gene (33) and in maize, where *CRP1* functions in processing of *petD* RNA, and in translation of *petA* (34).

### **Raal1 is part of a large RNA-protein complex associated with membranes**

We have cloned the *Raal1* gene, which encodes a large predicted polypeptide of ~206 kDa. We have shown that *Raal1* is found in the chloroplast fraction, as expected from its predicted transit peptide and from its role in plastid gene expression, although its presence in other organelles cannot be ruled out. Surprisingly it is associated with membranes, and shares this property with *Raa2* (formerly *Maa2*), a protein that is required for *trans*-splicing of the second intron of *psaA* (11), and *Rat1*, a factor required for processing of *tscA* and splicing of the first intron (12). This membrane association may be relevant because *PsaA* is a core subunit of photosystem I embedded in the thylakoid membrane. However *Raa3*, a factor required for splicing of the first intron of *psaA*, is mostly a stromal protein (10), like proteins involved in splicing of maize chloroplast group II introns, *CRS1* and *CRS2* (35,36). Chloroplast polyribosomes and some RNA-binding proteins are associated with membranes, and this association may facilitate the assembly of the photosystems and other membrane complexes [reviewed in (37)]. Whether the membrane association of *Raal1*, *Raa2* and *Rat1* reflects a coupling between RNA splicing, translation and protein assembly into the membrane complexes remains a matter of speculation.

When *Raal1* is released from membranes with moderate concentrations of salt, it sediments in large complexes which can be resolved into two peaks in sucrose gradients. *Raal1* is in part associated with *Raa2*, with which it can be co-immunoprecipitated (32). The complex containing *Raal1* and *Raa2* corresponds to the first peak. The faster

sedimenting component of the second peak (~700 kDa) is sensitive to treatment with RNase, implying that this *Raal1* complex contains RNA. This finding fits with the role of *Raal1* in splicing. However the precursor transcripts are broadly distributed in the gradients (data not shown), and we have not been able to conclusively determine whether they are part of the complex. The RNase-resistant fraction still sediments rapidly, indicating that *Raal1* is in a multimeric complex, perhaps in association with other proteins. *Raa3* is also part of a large ribonucleoprotein complex that contains exon 1 and the *tscA* RNA (10). It will be interesting to investigate interactions of *Raal1* with *Rat1* in splicing of intron 1. Although *Raa3* is mainly in the soluble fraction, and *Raal1* and *Rat1* in the membrane fraction, it is possible that they could also interact transiently.

### **Raal1 is a novel splicing factor with repeats**

Because of the limited sequence homology with other known proteins, it is difficult to predict the mode of action of *Raal1*. As already suggested for other proteins found to be essential for group II intron splicing, we expect that *Raal1* may function as an RNA chaperone or a stabilizer of active RNA conformations (8). In maize, the *CRS1* protein is specifically required for splicing of a group II intron in *atpF*, it binds the intron and promotes its folding (38). *CRS2*, together with *CAF1* or *CAF2*, is required for splicing nine other group II introns (35,39). *CRS1*, *CAF1* and *CAF2* contain conserved CRM domains which may be RNA-binding modules (40). In plants, genes for proteins similar to intron-encoded maturases are found in the chloroplast and in the nucleus (41,42). Other proteins which associate in the chloroplast with group II introns have been described, but their role in splicing has not yet been established (43,44).

One of the striking features of the predicted *Raal1* protein is the presence of numerous stretches of repeated amino acids. Such repetitive stretches occur occasionally in other proteins, but in *Raal1* they are particularly long and numerous, and involve diverse residues (Ala, Gln, Ser, Pro and Arg). Several other proteins that share this feature with *Raal1* are also involved in chloroplast gene expression, such as *Raa3* which is also a *psaA* *trans*-splicing factor (10), *Nac2*, *Mbb1* and *Mca1* which are involved in RNA processing and stability (45–47), or *Tbc2* which is implicated in translation (31). The functional significance of these repetitive stretches is unknown. They may play a structural role and may occur in parts of the protein where the precise sequence is not important so that it has drifted during evolution. There are five copies of a 38 amino acid repeat in *Raal1*, related to the 38 amino acid PPPEW repeats of *Tbc2* (Figure 4). Four of the repeats are retained in the *Arg7*–*Raal1* fusion of *raal-314B* which is sufficient for *tscA* processing and intron 1 splicing, and the last repeat is interrupted in the *mΔKpn* deletion which disrupts these activities (Figure 3C). Two of the repeats are in the region of possible overlap with the domain required for intron 2 splicing, just upstream of the *ScaI* site.

We speculate that although they have a different sequence, the 38 amino acid repeats in *Chlamydomonas* might be a counterpart of the repeats in the PPR and TPR proteins, many of which are also implicated in RNA maturation. The PPR proteins form a large family in plants where a majority are

predicted to be located in the organelles and many are known to interact with RNA [reviewed in (48)]. The TPR-like repeats are found in Mbb1 and Nac2 of *C.reinhardtii* (45,46) and in HCF107 of *Arabidopsis* (49). Each TPR repeat unit folds back to form two anti-parallel  $\alpha$ -helices which are stacked with the helices of the adjacent repeats to generate a superhelical structure (50). The related PPR repeats could form a similar structure (51), and one side of the superhelix could provide a surface for protein–RNA interactions, analogous to the protein–RNA interactions between the Puf repeats in Pumilio and their RNA target (52). Thus the repeat-containing chloroplast factors may have been recruited during evolution, by adaptation of a modular protein fold, to recognize new partners or substrates. It will be of interest to determine whether the repeats in Raa1 actually bind RNA, and specifically the *psaA* precursor transcripts or the *tscA* RNA.

In conclusion, in this report we describe a novel factor, Raa1, which is required for *trans*-splicing of both *psaA* introns in *Chlamydomonas*. Distinct domains in Raa1 are responsible for at least two functions, processing of *tscA* (the small RNA which is required for splicing of *psaA* intron 1) and splicing of *psaA* intron 2. Raa1 may thus play a role in coordinating the excision of the two *psaA* introns. Our analysis of Raa1 provides a basis to further investigate how different events of chloroplast RNA maturation are connected.

## SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

## ACKNOWLEDGEMENTS

The authors wish to thank Nicolas Roggli for preparing the figures, the members of the Rochaix laboratory for helpful discussions, Hans Sommer for providing a cDNA library, Cris Kuhlemeier for a gift of eIF4A antibody and Frédy Barneche, Stéphane Bellafiore, Paula Duek, Angela Falciatore and Frédéric Iseni for helpful comments on the manuscript. L.M. was supported by a long-term HFSP fellowship. This work was supported by a grant of the Swiss National Fund for Scientific Research (3100-0667763.02). Funding to pay the Open Access publication charges for this article was provided by the Swiss National Fund for Scientific Research.

*Conflict of interest statement.* None declared.

## REFERENCES

1. Qin,P.Z. and Pyle,A.M. (1998) The architectural organization and mechanistic function of group II intron structural elements. *Curr. Opin. Struct. Biol.*, **8**, 301–308.
2. Bonen,L. and Vogel,J. (2001) The ins and outs of group II introns. *Trends Genet.*, **17**, 322–331.
3. Toro,N. (2003) Bacteria and Archaea Group II introns: additional mobile genetic elements in the environment. *Environ. Microbiol.*, **5**, 143–151.
4. Villa,T., Pleiss,J.A. and Guthrie,C. (2002) Spliceosomal snRNAs: Mg(2+)-dependent chemistry at the catalytic core? *Cell*, **109**, 149–152.
5. Kück,U., Choquet,Y., Schneider,M., Dron,M. and Bennoun,P. (1987) Structural and transcription analysis of the two homologous genes for the P700 chlorophyll a-apoprotein in *Chlamydomonas reinhardtii*: evidence for *in vivo trans*-splicing. *EMBO J.*, **6**, 2185–2195.
6. Choquet,Y., Goldschmidt-Clermont,M., Girard-Bascou,J., Kuck,U., Bannoun,P. and Rochaix,J.D. (1988) Mutant phenotypes support a *trans*-splicing mechanism for the expression of the tripartite *psaA* gene in the *C.reinhardtii* chloroplast. *Cell*, **52**, 903–913.
7. Goldschmidt-Clermont,M., Choquet,Y., Girard-Bascou,J., Michel,F., Schirmer-Rahire,M. and Rochaix,J.D. (1991) A small chloroplast RNA may be required for *trans*-splicing in *Chlamydomonas reinhardtii*. *Cell*, **65**, 135–143.
8. Schroeder,R., Grossberger,R., Pichler,A. and Waldsich,C. (2002) RNA folding *in vivo*. *Curr. Opin. Struct. Biol.*, **12**, 296–300.
9. Goldschmidt-Clermont,M., Girard-Bascou,J., Choquet,Y. and Rochaix,J.D. (1990) *Trans*-splicing mutants of *Chlamydomonas reinhardtii*. *Mol. Gen. Genet.*, **223**, 417–425.
10. Rivier,C., Goldschmidt-Clermont,M. and Rochaix,J.D. (2001) Identification of an RNA–protein complex involved in chloroplast group II intron *trans*-splicing in *Chlamydomonas reinhardtii*. *EMBO J.*, **20**, 1765–1773.
11. Perron,K., Goldschmidt-Clermont,M. and Rochaix,J.D. (1999) A factor related to pseudouridine synthases is required for chloroplast group II intron *trans*-splicing in *Chlamydomonas reinhardtii*. *EMBO J.*, **18**, 6481–6490.
12. Balczun,C., Bunse,A., Hahn,D., Bannoun,P., Nickelsen,J. and Kuck,U. (2005) Two adjacent nuclear genes are required for functional complementation of a chloroplast *trans*-splicing mutant from *Chlamydomonas reinhardtii*. *Plant J.*, **43**, 636–648.
13. Harris,E.H. (1989) *The Chlamydomonas sourcebook. A comprehensive guide to biology and laboratory use.* Academic Press, Inc., San Diego, CA.
14. Debuchy,R., Purton,S. and Rochaix,J.D. (1989) The argininosuccinate lyase gene of *Chlamydomonas reinhardtii*: an important tool for nuclear transformation and for correlating the genetic and molecular maps of the ARG7 locus. *EMBO J.*, **8**, 2803–2809.
15. Kindle,K.L. (1990) High-frequency nuclear transformation of *Chlamydomonas reinhardtii*. *Proc. Natl Acad. Sci. USA*, **87**, 1228–1232.
16. Shimogawara,K., Fujiwara,S., Grossman,A. and Usuda,H. (1998) High-efficiency transformation of *Chlamydomonas reinhardtii* by electroporation. *Genetics*, **148**, 1821–1828.
17. Sambrook,J., Fritsch,E.F. and Maniatis,T. (1989) *Molecular Cloning. A laboratory manual.* Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
18. Purton,S. and Rochaix,J.D. (1994) Complementation of a *Chlamydomonas reinhardtii* mutant using a genomic cosmid library. *Plant Mol. Biol.*, **24**, 533–537.
19. Zhang,H., Herman,P.L. and Weeks,D.P. (1994) Gene isolation through genomic complementation using an indexed library of *Chlamydomonas reinhardtii* DNA. *Plant Mol. Biol.*, **24**, 663–672.
20. Sizova,I., Fuhrmann,M. and Hegemann,P. (2001) A *Streptomyces rimosus* aphVIII gene coding for a new type phosphotransferase provides stable antibiotic resistance to *Chlamydomonas reinhardtii*. *Gene*, **277**, 221–229.
21. Goldschmidt-Clermont,M. and Rahire,M. (1986) Sequence, evolution and differential expression of the two genes encoding variant small subunits of ribulose biphosphate carboxylase/oxygenase in *Chlamydomonas reinhardtii*. *J. Mol. Biol.*, **191**, 421–432.
22. Silflow,C.D., LaVoie,M., Tam,L.W., Tousey,S., Sanders,M., Wu,W., Borodovsky,M. and Lefebvre,P.A. (2001) The Vfl1 Protein in *Chlamydomonas* localizes in a rotationally asymmetric pattern at the distal ends of the basal bodies. *J. Cell. Biol.*, **153**, 63–74.
23. Merendino,L., Falciatore,A. and Rochaix,J.D. (2003) Expression and RNA-binding properties of the chloroplast ribosomal protein S1 from *Chlamydomonas reinhardtii*. *Plant Mol. Biol.*, **53**, 371–382.
24. Schloss,J.A. (1990) A *Chlamydomonas* gene encodes a G protein beta subunit-like polypeptide. *Mol. Gen. Genet.*, **221**, 443–452.
25. Silflow,C. (1998) Organization of the nuclear genome. In Rochaix,J.D., Goldschmidt-Clermont,M. and Merchant,S (1998) *The Molecular Biology of Chloroplasts and Mitochondria in Chlamydomonas*. Kluwer Academic Publishers, Dordrecht, The Netherlands, Vol. 7, pp. 25–40.
26. Emanuelsson,O., Nielsen,H. and von Heijne,G. (1999) ChloroP, a neural network-based method for predicting chloroplast transit peptides and their cleavage sites. *Protein Sci.*, **8**, 978–984.
27. Small,I., Peeters,N., Legeai,F. and Lurin,C. (2004) Predotar: a tool for rapidly screening proteomes for N-terminal targeting sequences. *Proteomics*, **4**, 1581–1590.
28. Garnier,J., Gibrat,J.F. and Robson,B. (1996) GOR method for predicting protein secondary structure from amino acid sequence. *Methods Enzymol.*, **266**, 540–553.

29. Hahn,D., Nickelsen,J., Hackert,A. and Kuck,U. (1998) A single nuclear locus is involved in both chloroplast RNA trans-splicing and 3' end processing. *Plant J.*, **15**, 575–581.
30. Baginsky,S., Siddique,A. and Gruijsem,W. (2004) Proteome analysis of tobacco bright yellow-2 (BY-2) cell culture plastids as a model for undifferentiated heterotrophic plastids. *J. Proteome Res.*, **3**, 1128–1137.
31. Auchincloss,A.H., Zerges,W., Perron,K., Girard-Bascou,J. and Rochaix,J.D. (2002) Characterization of Tbc2, a nucleus-encoded factor specifically required for translation of the chloroplast psbC mRNA in *Chlamydomonas reinhardtii*. *J. Cell. Biol.*, **157**, 953–962.
32. Perron,K., Goldschmidt-Clermont,M. and Rochaix,J.D. (2004) A multiprotein complex involved in chloroplast group II intron splicing. *RNA*, **10**, 704–710.
33. Yamazaki,H., Tasaka,M. and Shikanai,T. (2004) PPR motifs of the nucleus-encoded factor, PGR3, function in the selective and distinct steps of chloroplast gene expression in *Arabidopsis*. *Plant J.*, **38**, 152–163.
34. Fisk,D.G., Walker,M.B. and Barkan,A. (1999) Molecular cloning of the maize gene crp1 reveals similarity between regulators of mitochondrial and chloroplast gene expression. *EMBO J.*, **18**, 2621–2630.
35. Till,B., Schmitz-Linneweber,C., Williams-Carrier,R. and Barkan,A. (2001) CRS1 is a novel group II intron splicing factor that was derived from a domain of ancient origin. *RNA*, **7**, 1227–1238.
36. Jenkins,B.D. and Barkan,A. (2001) Recruitment of a peptidyl-tRNA hydrolase as a facilitator of group II intron splicing in chloroplasts. *EMBO J.*, **20**, 872–879.
37. Zerges,W. (2000) Translation in chloroplasts. *Biochimie*, **82**, 583–601.
38. Ostersetzer,O., Cooke,A.M., Watkins,K.P. and Barkan,A. (2005) CRS1: a chloroplast group II intron splicing factor, promotes intron folding through specific interactions with two intron domains. *Plant Cell*, **17**, 241–255.
39. Ostheimer,G.J., Williams-Carrier,R., Belcher,S., Osborne,E., Gierke,J. and Barkan,A. (2003) Group II intron splicing factors derived by diversification of an ancient RNA-binding domain. *EMBO J.*, **22**, 3919–3929.
40. Ostheimer,G.J., Hadjivasiliou,H., Kloer,D.P., Barkan,A. and Matthews,B.W. (2005) Structural analysis of the group II intron splicing factor CRS2 yields insights into its protein and RNA interaction surfaces. *J. Mol. Biol.*, **345**, 51–68.
41. Mohr,G. and Lambowitz,A.M. (2003) Putative proteins related to group II intron reverse transcriptase/maturases are encoded by nuclear genes in higher plants. *Nucleic Acids Res.*, **31**, 647–652.
42. Liere,K. and Link,G. (1995) RNA-binding activity of the matK protein encoded by the chloroplast trnK intron from mustard (*Sinapis alba* L.). *Nucleic Acids Res.*, **23**, 917–921.
43. Nakamura,T., Ohta,M., Sugiura,M. and Sugita,M. (1999) Chloroplast ribonucleoproteins are associated with both mRNAs and intron-containing precursor tRNAs. *FEBS Lett.*, **460**, 437–441.
44. Bunse,A.A., Nickelsen,J. and Kuck,U. (2001) Intron-specific RNA-binding proteins in the chloroplast of the green alga *Chlamydomonas reinhardtii*. *Biochim. Biophys. Acta.*, **1519**, 46–54.
45. Boudreau,E., Nickelsen,J., Lemaire,S.D., Ossenbuhl,F. and Rochaix,J.D. (2000) The Nac2 gene of *Chlamydomonas* encodes a chloroplast TPR-like protein involved in psbD mRNA stability. *EMBO J.*, **19**, 3366–3376.
46. Vaistij,F.E., Boudreau,E., Lemaire,S.D., Goldschmidt-Clermont,M. and Rochaix,J.D. (2000) Characterization of Mbb1, a nucleus-encoded tetratricopeptide-like repeat protein required for expression of the chloroplast psbB/psbT/psbH gene cluster in *Chlamydomonas reinhardtii*. *Proc. Natl Acad. Sci. USA*, **97**, 14813–14818.
47. Lown,F.J., Watson,A.T. and Purton,S. (2001) *Chlamydomonas* nuclear mutants that fail to assemble respiratory or photosynthetic electron transfer complexes. *Biochem. Soc. Trans.*, **29**, 452–455.
48. Lurin,C., Andres,C., Aubourg,S., Bellaoui,M., Bitton,F., Bruyere,C., Caboche,M., Debast,C., Gualberto,J., Hoffmann,B. et al. (2004) Genome-wide analysis of *Arabidopsis* pentatricopeptide repeat proteins reveals their essential role in organelle biogenesis. *Plant Cell*, **16**, 2089–2103.
49. Felder,S., Meierhoff,K., Sane,A.P., Meurer,J., Driemel,C., Plucken,H., Klaff,P., Stein,B., Bechtold,N. and Westhoff,P. (2001) The nucleus-encoded HCF107 gene of *Arabidopsis* provides a link between intergenic RNA processing and the accumulation of translation-competent psbH transcripts in chloroplasts. *Plant Cell*, **13**, 2127–2141.
50. Das,A.K., Cohen,P.W. and Barford,D. (1998) The structure of the tetratricopeptide repeats of protein phosphatase 5: implications for TPR-mediated protein-protein interactions. *EMBO J.*, **17**, 1192–1199.
51. Small,I.D. and Peeters,N. (2000) The PPR motif: a TPR-related motif prevalent in plant organellar proteins. *Trends Biochem. Sci.*, **25**, 46–47.
52. Wang,X., McLachlan,J., Zamore,P.D. and Hall,T.M. (2002) Modular recognition of RNA by a human pumilio-homology domain. *Cell*, **110**, 501–512.