

## Molecular Characterization of a Serotype O121:H19 Clone, a Distinct Shiga Toxin-Producing Clone of Pathogenic *Escherichia coli*

Cheryl L. Tarr,<sup>1</sup> Teresa M. Large,<sup>1</sup> Chris L. Moeller,<sup>1</sup> David W. Lacher,<sup>1</sup> Phillip I. Tarr,<sup>2</sup>  
David W. Acheson,<sup>3</sup> and Thomas S. Whittam<sup>1\*</sup>

*Microbial Evolution Laboratory, National Food Safety and Toxicology Center, Michigan State University, East Lansing, Michigan 48824<sup>1</sup>; Children's Hospital and Regional Medical Center, University of Washington School of Medicine, Seattle, Washington 98105<sup>2</sup>; and University of Maryland School of Medicine, Baltimore, Maryland 21201<sup>3</sup>*

Received 14 June 2002/Returned for modification 7 August 2002/Accepted 5 September 2002

**Most illnesses caused by Shiga toxin-producing *Escherichia coli* (STEC) have been attributed to *E. coli* serotype O157:H7, but non-O157 STEC infections are now increasingly recognized as public health problems worldwide. The O121:H19 serotype is being isolated more frequently from clinical specimens and has been implicated in one waterborne outbreak. We used multilocus virulence gene profiling, a PCR-based assay, to characterize the virulence gene content of 24 isolates of serotype O121:H19 and nonmotile variants. We also performed multilocus enzyme electrophoresis and multilocus sequencing to establish the clonal relatedness of O121 isolates and to elucidate the relationship of O121 to common STEC clones. The 24 isolates were found to represent a single bacterial clone, as there was no allelic variation across 18 enzyme loci among the isolates. The complete nucleotide sequence of the intimin gene differed by four substitutions from that of the epsilon (Int-ε) allele of O103:H2 strain PMK5. The typical O121 virulence gene profile was similar to the profiles of enterohemorrhagic *E. coli* (EHEC) clones of *E. coli*: it included a Shiga toxin 2 gene (*stx*<sub>2</sub>), two genes on the EHEC plasmid (*tox*B and *ehx*A), and the gene encoding intimin (*eae*). Despite the similarities, putative virulence genes distributed on O islands—large chromosomal DNA segments present in the O157:H7 genome—were useful for discriminating among STEC serotypes and the O121:H19 clone had a composite profile that was distinct from the profiles of the other major EHEC clones of pathogenic *E. coli*. On the basis of sequencing analysis with 13 housekeeping genes, the O121:H19 clone did not fall into any of the four classical EHEC and enteropathogenic *E. coli* groups but instead was closely related to two *eae*-negative STEC strains.**

Shiga toxin-producing *Escherichia coli* (STEC) bacteria are infectious pathogens that have been implicated in outbreaks and sporadic cases of food- and waterborne diarrheal diseases in humans. The clinical symptoms caused by STEC infection include nonbloody diarrhea, hemorrhagic colitis (HC), and the hemolytic-uremic syndrome (HUS) (25, 33). The first recorded STEC outbreak (in 1982) was attributed to *E. coli* O157:H7 (28), a serotype that still accounts for the greatest proportion of STEC disease in North America, Europe, and Japan. However, more than 200 serotypes have been isolated from humans and non-O157 STEC strains may account for 20 to 70 percent of STEC infections overall in different countries (42). Pathogenic non-O157 STEC bacteria are unlike those of the O157:H7 serotype in that they usually ferment sorbitol, so they are more difficult to differentiate from commensal *E. coli*; thus, the incidence of disease caused by non-O157 STEC strains is more difficult to determine (35).

The defining characteristic of STEC is the production of Shiga toxins (Stx), a family of bacteriophage-encoded cytotoxins with diverse effects on eukaryotic cells (18, 20). There are two main antigenic types of Stx, Stx1 and Stx2, whose genes are highly divergent at the nucleotide level (40), and STEC strains may produce either or both toxin variants. Members of a subset of STEC strains called enterohemorrhagic *E. coli* (EHEC)

effect a particular pathology called attaching and effacing (A/E) lesions, which are mediated in part by the adhesin intimin (16). Intimin is encoded by the *eae* gene on a pathogenicity island (the locus of enterocyte effacement [LEE]) whose presence is widespread among enteric bacterial pathogens (14). Other virulence elements are found in STEC strains, but their role in pathogenesis is unclear. For example, an enterohemolysin (*ehx*A) resides on a large plasmid (pO157) found in *E. coli* O157:H7 strains, variants of which are distributed among STEC strains of a variety of serotypes (3). The pO157 plasmid also encodes other factors that may contribute to virulence, including a type II secretion system (29) and accessory adhesins (24, 34, 36). Other virulence gene candidates were uncovered when the genomes of two O157:H7 strains (7, 26) were compared to the genome for the nonpathogenic *E. coli* strain K-12. The comparison revealed large chromosomal segments called O islands that were not present in the K-12 genome (26), and many of these islands contain DNA homologous to virulence factors from other bacterial pathogens. For example, O island 48 encodes an operon for urease activity, a mechanism for acid resistance in *Yersinia* and *Helicobacter*. The extent of occurrence of various O islands and their associated genes in non-O157 STEC strains is presently unknown, and their role in virulence remains to be defined.

Detecting in STEC strains the presence of the accessory virulence genes described above could provide information that is clinically relevant and of epidemiological value (23). For the present investigation, we used multilocus virulence gene profiling (MVGP)—a PCR assay of 15 potential virulence lo-

\* Corresponding author. Mailing address: Microbial Evolution Laboratory, 165 Food Safety and Toxicology Building, Michigan State University, East Lansing, MI 48824. Phone: (517) 432-3588. Fax: (517) 432-2310. E-mail: whittam@msu.edu/whittam@msu.edu.

TABLE 1. Primers and amplicon sizes for loci included in the PCR assay

Locus	Primer designation	Nucleotide sequence of primer	Amplicon size(s) (bp)
<i>bfpA</i>	bfpA-F11 bfpA-R1	5'-GTCTGCGTCTGATTCCAATA-3' 5'-TCAGCAGGAGTAATAGC-3'	408, 414
<i>toxB</i>	toxB.911F toxB.1468R	5'-ATACCTACCTGCTCTGGATTGA-3' 5'-TCTTACCTGATCTGATGCAGC-3'	602
<i>stx<sub>1</sub></i>	1A.251F 1A.832R	5'-GGGATAGATCCAGAGGAAGG-3' 5'-CCGGACACATAGAAGGAACTC-3'	624
<i>stx<sub>2</sub></i>	2A.506F 2A.848R	5'-CTGGCGTAAATGGAGTTCAG-3' 5'-CCTGTCGCCAGTTATCTGAC-3'	383
<i>eae</i>	eae-F1 eae-R3	5'-ACTCCGATTCTCTGGTGAC-3' 5'-TCTTGTGCGCTTTGGCTT-3'	389
<i>terC</i>	terC.106F terC.275R	5'-TATGCACCGTGATGACAAGC-3' 5'-GGCGAACAGGAGAAGATTG-3'	210
<i>ureA</i>	ureA.109F ureA.213R	5'-TAACTATCCCGAATCCGTGG-3' 5'-GGGATCATTCTGGTATGCCT-3'	146
<i>iha</i>	iha.138F iha.282R	5'-ACGCAGCCGCCAGTGT-3' 5'-CCATCAATCAGTATCAGCGTGTA-3'	186
<i>invG</i>	invG.481F invG.572R	5'-TGACCTGGTCTGTTAATGCTG-3' 5'-CGCCACGTAATCATAAAGTCC-3'	133
<i>spaP</i>	spaP.143F spaP.275R	5'-GGACTTCAGCAAGTGCCATC-3' 5'-ACCACTCATGCCTGTCTCAA-3'	173
Z4326	ShET.1137F ShET.1221R	5'-TCATGCAGACGCAATAAAGG-3' 5'-GATACGCAAGAAGTAATCCTGG-3'	125
<i>fyuA</i>	fyuA.924F	5'-GCAGCAGCAGCATTATTCG-3'	495
<i>mdh</i>	mdh.269F mdh.530R	5'-GGTATGGATCGTTCGACCT-3' 5'-GGCAGAATGGTAACACCAGAGT-3'	304

ci—to characterize the diversity and gene content of *E. coli* O121:H19, an STEC serotype that has been implicated in a waterborne outbreak of HUS (13) and is one of the more common non-O157 serotypes found in sporadic cases of STEC infection (39). We also employed multilocus enzyme electrophoresis (MLEE) to establish the clonal relationships among the O121 isolates. Little is known about the genetic relatedness of *E. coli* O121:H19 to other serotypes that cause HC and HUS. To address this issue, we used multilocus sequencing of housekeeping genes to determine the evolutionary relationships of the O121:H19 clone to other pathogenic clones of *E. coli*.

#### MATERIALS AND METHODS

**Bacterial isolates.** A total of 27 of the strains of serogroup O121 that we analyzed were originally isolated from sporadic cases of diarrheal disease. Seventeen strains originated in Montana and were isolated between 1998 and 2000 (11). These included 13 strains of serotype O121:H19 (TW06388, TW08023, TW08032, TW08043, TW08055, TW08062, TW08063, TW08067, TW08068, TW08082, TW08091, TW08096, and TW08105) and 4 for which the flagellar antigen was not determined (TW08036, TW08039, TW08040, and TW08042). An additional five nonmotile strains were included: PT91-4 (TW03508; Washington, 1989), DA-1 (TW07927; District of Columbia, 1998), DA-5 (TW07931; Massachusetts, 1998), DA-37 (TW07972; Ohio, 1998), and DA-69 (TW08004; Ohio, 1998). One O121:H19 isolate, 3056-85, was obtained from the Centers for Disease Control and Prevention (Oklahoma, 1985). For comparisons in the MLEE

analysis, we included two strains of serotype O121:H7 that were isolated from cows (90-0364 and 403-3; Canada, 1992) and ECOR 21, a nonmotile strain isolated from a steer (19). The phylogenetic analysis included 20 pathogenic strains previously characterized for multiple gene sequences (27) and the O103:H2 strain TW08101 (Montana, 2000). Additional strains included for comparisons in the MVGP analysis were as follows: O26:H11 strains TW08100 and TW08104 (Montana, 2000); O103:H2 strains TW07877 (Germany, 1998), TW07882 (Washington, 1999), and PMK5 (21); O145:NM strain TW08087 (Montana, 2000); O111:H8 strains 3214-99 and 3215-99 (Texas, 1999); O157:H7 strains Sakai (7) and EDL-933 (26); and the strain K-12 (2), which was used as a negative control. More information about the strains included in this study can be obtained from the STEC Center website (<http://www.shigatox.net>).

**Enzyme electrophoresis.** Protein extracts were obtained by lysing bacterial cells and were subjected to horizontal starch gel electrophoresis and staining, following an established methodology (31). A total of 18 enzymes were analyzed for mobility variants: PGI, IDH, ACO, G3P, PE2, AK, MDH, PGD, M1P, GOT, ADH, MPI, G6P, CAK, NSP, TDH, SKD, and GLU. A multilocus genotype was determined for each strain, and distinct genotypes were classified as electrophoretic types (ETs) as previously described (41).

**PCR template preparation.** Luria-Bertani agar plates were inoculated from freezer stocks and incubated overnight at 37°C. Genomic DNA was extracted from single colonies with a Puregene DNA isolation kit (Gentra Systems, Minneapolis, Minn.). Templates for MVGP were produced by suspending single colonies in 200 ml of 1× TE buffer (10 mM Tris, 1 mM EDTA [pH 7.0]). The suspension was heated to 95°C for 10 min and centrifuged for 2 min. For each strain, three isolated colonies were processed and analyzed by MVGP.

**PCR assays for virulence factors and O island genes.** The strains were analyzed by PCR with primers specific for a total of 15 genes that potentially contribute to virulence (Table 1). Nine genes have been found on O islands in the

O157:H7 genome; these include the well-characterized virulence factors *stx*<sub>1</sub> (O island 93), *stx*<sub>2</sub> (O island 45), and intimin (O island 148). Three genes occur on O island 48, including a gene involved in tellurite resistance (*terC*), a locus of the urease gene complex (*ureA*), and an adherence-conferring gene (*iha*) encoding a protein homologous to iron-regulated IrgA of *Vibrio cholerae* (34). Two genes occur on O island 115 and have homology to elements of the Inv-Spa system of *Salmonella* (*spaP* and *invG*), and a homologue of the *Shigella* enterotoxin ShET2 (Z4326) on O island 122 was also included. Lastly, we screened for two genes (*fyuA* and *irp2*) on the high-pathogenicity island (HPI) of *Yersinia*, the presence of which has been reported for EHEC O26:H11 strains (10). We also screened for four plasmid-borne genes, including *ehxA*, the bundlin gene (*bfpA*) of enteropathogenic *E. coli* (EPEC), and two recently described adhesins, *toxB* (36, 38) and *saa* (23). PCR amplification of a conserved housekeeping gene (*mdh*) was used as a positive locus control. Previously published primer sequences were used for *ehxA* (8), *saa* (27), and the two HPI genes (34), except the forward primer for *fyuA*, which was redesigned (Table 1).

Synthesized phosphoramidite dye-linked primers (Research Genetics, Huntsville, Ala.) were used at a final concentration of 0.5 μM in a 20-μl reaction mixture with 1 U of AmpliTaq Gold (Applied Biosystems, La Jolla, Calif.) and final concentrations of deoxynucleoside triphosphates of 0.2 mM each and MgCl<sub>2</sub> of 2.5 mM. The loci were simultaneously amplified in separate reactions with the following thermal cycle, which was preceded by a 10-min soak at 94°C and run for 35 cycles: 92°C for 30 s, 55°C for 30 s, and 72°C for 45 s. The cycle was followed by a final extension step of 72°C for 7 min. Each run included two negative controls (including DNA from *E. coli* strain K-12 and no template) and a positive control that was a mixture of strains. Reaction aliquots were pooled for each strain, and the mix was desalted using Multiscreen-HV plates (Millipore, Bedford, Mass.) and a Sephadex G-50 column (Amersham Pharmacia Biotech, Piscataway, N.J.). The amplicon mix was dried under vacuum and suspended in formamide with 0.5 μl of DNA size standard 600 (Beckman Coulter, Fullerton, Calif.). The dye-labeled fragments were electrophoresed for 1 h at 6.0 kV and detected on a Beckman CEQ8000 capillary sequencer. Fragment sizes were calculated automatically, and the presence or absence of fragments was scored for each chromatogram with Fragment Analysis software, version 5.0 (Beckman Coulter).

**Intimin allele typing.** We determined the nucleotide sequence for the entire *eae* gene in O121:H19 strain TW08023 by following previously published protocols (32). The program ClustalX (37) was used to align the O121 *eae* sequence to 20 major *eae* sequences retrieved from GenBank. The relationship of the O121 sequence to those of other intimin alleles was determined by constructing phylogenetic trees with the program MEGA2 (12) by using the neighbor-joining algorithm. Because the mosaic segments of intimin genes can have different evolutionary histories, *eae* sequences were partitioned into a conserved central domain (CCD) (residue 187 through 517) and the extracellular (EC) region (amino acid positions 559 to 951) for phylogenetic analysis (15, 32). Two *eae* sequencing primers (1460F [5'-GGT CTG GAT CGT ATC GTC TG-3'] and 2009R [5'-ATC AGA ACC ATT TGC CTT CG-3']) were used to amplify a 590-bp fragment, and the sequence was determined in one direction with primer 1460F.

**Evolutionary relationships of the O121 clone.** We sequenced ~500 bp of each of 13 housekeeping genes (*arcA*, *aroE*, *aspC*, *clpX*, *cyaA*, *dnaG*, *fadD*, *grpE*, *icdA*, *lysP*, *mdh*, *mtlD*, and *rpoS*) for two O121 isolates (TW08023 and TW07927) and an O103:H2 strain (TW08101) and compared the multilocus sequences to those of 20 pathogenic *E. coli* strains (27). The total length sequenced for each of the gene fragments ranged from 288 bp (*aroE*) to 561 bp (*clpX*), with a total of 6,228 bp sequenced across the thirteen loci. Primer sequences and details of the methodology will be published elsewhere. The program ClustalX was used to align the sequences. The 13 gene sequences were concatenated for each strain, and phylogenetic trees were constructed with the program MEGA2 (12). The trees were based on synonymous distances calculated by using the modified Nei-Gojobori method with the Jukes-Cantor correction (17). Bootstrap confidence values for each node of the tree were calculated over 1,000 replicate trees.

## RESULTS

**MLEE.** A total of three ETs were observed among the 27 strains of the O121 serogroup. All ETs were equally distant from one another, each differing from the other by 8 out of 18 alleles. ECOR 21 was distinct from all other isolates and was designated ET 1. The two O121:H7 strains were identical to each other and were designated ET 2. The remaining O121

strains, including all that express the H19 flagellar antigen, comprised ET 3. We detected no genetic variation within ET 3 at the level of enzyme mobility, which indicates a very close genetic relationship among all O121:H19 strains and related nonmotile variants.

**MVGP.** Strains that produce only Stx2 are associated with serious complications, such as HUS, more often than strains that produce either Stx1 or both toxins (22). Of the 24 isolates that we analyzed, all but one (TW03508) tested positive for *stx* and the 23 positive strains all carried the *stx*<sub>2</sub> gene. One strain (TW08032) also carried *stx*<sub>1</sub>.

Proteins encoded by the large EHEC plasmid are thought to contribute to the virulence of STEC strains. We screened for two genes within the EHEC plasmid (*ehxA* and *toxB*). The two genes always occurred together, and both genes were present in 22 of the ET 3 O121 isolates. Two ET 3 isolates (TW08032 and TW08042) were negative for both genes, suggesting that they lack the pO157 plasmid. None of the O121 strains was positive for the bundlin gene (*bfpA*), the presence of which is a characteristic of the EPEC pathotype, and all O121 strains also lacked *saa*, a gene contained within a large plasmid and usually found in *eae*-negative STEC strains.

The presence of the LEE island is associated with the ability to cause A/E lesions and is one of the major virulence determinants by which some diarrheagenic *E. coli* strains cause disease in humans (16). We screened for the LEE island with primers placed in the CCD of the *eae* gene and detected a copy of *eae* in each of the 24 isolates. The results of amplification and sequencing of three additional genes (*sepZ*, *tir*, and *sepL*) in one isolate (TW08023) suggested that the O121:H19 clone harbors a complete copy of the LEE pathogenicity island (data not shown).

All ET 3 O121 strains were positive for *terC* and *ureA*, two genes found on O island 48 of the *E. coli* O157:H7 genome (Table 2). This observation suggests that the O121:H19 clone has an O island 48 homologue, although the island differs from that of *E. coli* O157:H7 because it lacks the *iha* gene. All ET 3 strains were also positive for Z4326, the gene encoding the ShET2 homologue of O island 122 in *E. coli* O157:H7. Neither the *invG* or *spaP* gene of O island 115 nor either of the two genes (*fyuA* and *irp2*) of the HPI (Table 2) was detected in the O121:H19 clone.

**Evolutionary relationship of the O121:H19 clone to other EHEC strains.** We sequenced segments of 13 housekeeping genes of two ET 3 O121 strains (TW08023 and TW07927) and an STEC O103:H2 strain (TW08101), because O121 and O103 have been reported to have similar intimin sequences (21). The multilocus sequences were aligned and concatenated with homologous sequences from 20 pathogenic strains representing the four groups of major EPEC and EHEC clones (27). There were a total of 281 polymorphic sites among the 23 strains, and the sequences of the two O121 isolates were identical. In the bootstrap analysis of the combined sequences (Fig. 1), the O121:H19 clone did not cluster significantly with either EHEC 1 group (O157:H7) or EHEC 2 group (O26 and O111). The consensus phylogeny indicates that the O121:H19 clonal frame is not significantly allied with STEC O103:H2, clusters at a distant node with the EPEC 2 and EHEC 2 groups, and is most closely related (82% support) to two *eae*-negative STEC strains (serotypes O113:H21 and XO3:NM).

TABLE 2. Molecular virulence gene profiles of *E. coli* isolates

Serotype (reference isolate)	No. of isolates	Locus														
		<i>bfpA</i>	<i>ehxA</i>	<i>toxB</i>	<i>saa</i>	<i>stx<sub>1</sub></i>	<i>stx<sub>2</sub></i>	<i>eae</i>	<i>terC</i>	<i>ureA</i>	<i>iha</i>	<i>invG</i>	<i>spaP</i>	<i>Z4326</i>	<i>irp2</i>	<i>fyuA</i>
O121:H19 (TW08023)	13	-	+	+	-	-	+	+	+	+	-	-	-	+	-	-
O121:H19 (TW08032)	1	-	-	-	-	+	+	+	+	-	-	-	-	+	-	-
O121:NM (DA-1)	5	-	+	+	-	-	+	+	+	-	-	-	-	+	-	-
O121:NM (PT91-4)	1	-	+	+	-	-	+	+	+	-	-	-	-	+	-	-
O121:ND (TW08039)	3	-	+	+	-	-	+	+	+	-	-	-	-	+	-	-
O121:ND (TW08042)	1	-	-	-	-	-	+	+	+	-	-	-	-	+	-	-
O113:H21 (CL-3)	1	-	+	-	+	-	+	-	-	+	-	+	-	-	-	-
XO3:NM (90-1787)	1	-	+	-	+	-	+	-	-	-	+	-	+	-	-	-
O103:H2 (TW08101)	2	-	+	+	-	+	-	+	-	-	-	-	+	+	-	-
O103:H2 (PMK5)	2	-	+	-	-	+	-	+	-	-	-	-	+	+	-	-
O26:H11 (TW08100)	1	-	+	+	-	+	-	+	+	+	-	+	+	+	+	+
O111:H8 (3215-99)	2	-	+	-	-	+	+	+	+	+	-	+	+	-	-	-
O145:NM (TW08087)	1	-	+	+	-	+	-	+	+	+	+	-	+	+	-	-
K-12	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
O127:H6 (E2348/69)	1	+	-	-	-	-	-	+	-	-	-	-	-	+	-	-
O157:H7 (EDL-933)	3	-	+	+	-	+	+	+	+	+	+	+	+	+	-	-

**Intimin allele of the O121:H19 clone.** The sequence for the *eae* gene of O121:H19 strain TW08023 was 2,847 nucleotides in length, which is indicative of a protein of 948 amino acids. A gene phylogeny based on sequences of the CCD indicates that the intimin allele of the O121:H19 clone is virtually the same as the epsilon allele (Int-ε) of *E. coli* O103:H2 (Fig. 2). There are only four nucleotide differences between the two sequences (Fig. 3): two nonsynonymous substitutions (H209R in the CCD and N587D in domain 0 [D0]) and two synonymous substitutions. To confirm this finding, we used primers flanking the replacement location in D0 and found that all 24 isolates of the O121:H19 clone have the Int-ε variant allele. Interestingly, Int-ε is most similar to Int-β (found in EHEC 2 O26:H11 strains) in the transmembrane region, where the divergence at the amino acid level is less than 1%, but is more than 30% divergent in the EC domains (Fig. 3). Overall, the level of amino acid divergence between Int-ε and other intimin alleles ranges from 34% to 40% in the EC domains (Fig. 3).

## DISCUSSION

A key observation from this study is that *E. coli* O121:H19 represents an independent clone that has acquired particular virulence factors in parallel with the EHEC lineages. The O121:H19 clone has in common with bacteria of the EHEC 1 group (O157:H7 and relatives) and EHEC 2 group (O26, O111, and relatives) at least three virulence characteristics: a Shiga toxin gene, an intimin allele, and genes found on the EHEC plasmid (pO157). However, the phylogenetic tree constructed from the housekeeping gene sequences places the O121:H19 clonal frame on a separate branch distinct from the EHEC 1 and EHEC 2 lineages. Similarly, the EHEC 1 and EHEC 2 clones are not closely related to each other but instead have evolved in parallel through a repeated series of gene acquisitions (27). The results presented here support the hypothesis that *E. coli* O121:H19 represents a third EHEC clone that has independently accumulated the virulence genes that contribute to the EHEC pathovar.

The comparison of *E. coli* strain O121:H19 to strains of the other major EHEC serotypes showed that the fragment profiles of the major EHEC serotypes differed in composition

when all of the genes were examined. There were few genes that could serve as diagnostic markers for a single clone or serotype (Table 2). The exception was the presence of the genes on the HPI (*irp2* and *fyuA*), which differentiates bacteria of the EHEC O26:H11 clone from those of the other serotypes included here. The HPI is prevalent in *E. coli* (30) but thus far has not been found in EHEC serotypes other than O26:H11 strains (10). Bacterial clones associated with other serotypes, including O121:H19, were differentiated on the basis of the composite profile rather than that of a single diagnostic gene. It appears that the genomic island markers are more stable than plasmid or phage markers; the latter tended to differ within a serotype and occasionally among the single-colony picks for each strain. The loss of plasmids and phages during




FIG. 1. Gene phylogeny inferred from 13 housekeeping gene sequences. Trees were constructed from synonymous distances by using the neighbor-joining algorithm. The number at each branching point indicates the percentage of trees that support a particular node out of 1,000 bootstrapped replicates.






FIG. 2. Phylogeny of intimin alleles inferred from the CCD. Trees were constructed from synonymous distances by using the neighbor-joining algorithm. The scale at the lower left shows the horizontal distance that corresponds to two synonymous substitutions per 100 synonymous sites.

subculture is documented (9, 16) and could explain in part the variation in the presence of EHEC plasmid and Stx phage genes within a serotype. The stability of the O island genes makes them useful for differentiating STEC serotypes by PCR.

Intimin and other genes on the LEE pathogenicity island have been shown to be crucial for the full virulence of enteropathogenic *E. coli* (5), as they confer upon bacterial strains the ability to cause A/E lesions in humans. The intimin gene of the O121:H19 clone is nearly identical in sequence to Int-ε of *E. coli* O103:H2, a strain of the EPEC 2 clonal group (members of which typically carry the Int-β allele) (Fig. 1). Int-ε is highly similar to Int-β in the 5' half of the gene but is highly divergent in the region encoding the EC domains (Fig. 2); it is possible that segments of Int-β and Int-ε are ancestral and derive from a similar LEE backbone. To address this hypothesis, we sequenced *sepL* (a gene downstream of *eae* on the LEE island) from O121:H19 strain TW08023 and from O103:H2 strain TW08101. The *sepL* sequences are identical and form a tight cluster (average divergence, 0.4%) with the *sepL* sequences from strains that carry Int-β (data not shown). These results suggest that the O121:H19 and O103:H2 clones carry a LEE that is closely related by recent descent to the

LEE of EPEC2 strains but one lineage of the LEE has undergone recombination with a divergent sequence that converted the 3' half of the gene.

The O121:H19 clone comprises a single ET, as no polymorphic loci were detected by MLEE. Additional ETs might be present within the clone but were not detected within our limited sample. Even so, *E. coli* O121:H19 appears to be more homogeneous than other serotypes. In the study by Whittam et al. (41), six serotypes had sample sizes comparable to that of our study (excluding nonmotile strains); among these serotypes, two to five ETs were detected in samples of 10 to 31 isolates and 5 to 70% of the loci were polymorphic. Also, the average genetic diversity (*H*)—a measure that is not correlated with sample size—was lowest in O121:H19 (where it was zero). The *H* averaged across 11 serotypes was 0.107, with a range of 0.005 to 0.183. As a comparison, O157:H7 has an *H* of 0.008 and 95% of the isolates belong to the same ET. The low level of diversity in O121:H19 indicates a relatively recent emergence of this clone from a single ancestral cell.

Recognition of the impact of non-O157 STEC strains on human health is increasing worldwide, but unlike O157:H7, these strains are difficult to differentiate from commensal *E. coli*. Thus, the prevalence of non-O157 STEC is presently underestimated (8, 35). In 1998, the World Health Organization cited O26, O111, O103, and O145 as the most important non-O157 STEC pathogens in humans. Some of the serotypes appear to have emerged more recently than O157:H7. For example, prior to 1995, O103 was not detected by routine surveillance in a lab in Germany and accounted for less than 2% of the STEC infections in Italy (4). However, in 1996, the frequency of detection of O103 had increased to 18.2% in a German laboratory and accounted for 11% of the HUS cases in Italy in 1996 (4). Similarly, the frequency of infection by O26 STEC increased dramatically in 1996 in Germany and Italy (4).

The O121:H19 clone also may be on a trajectory of emergence and spread. In 1996, Beutin and colleagues reported that among 89 non-O157 STEC strains implicated in human disease in Germany, 40% of the strains were of common STEC serotypes (O26, O103, O111, and O145) but none of the remaining isolates belonged to the O121 serogroup (1). Fey et al. (6) examined 335 specimens from patients in Nebraska in 1998 and isolated four non-O157 serotypes, but there were no cases of disease in which an STEC O121 strain was implicated (6). More recently, however, cases of HC and HUS have been attributed to *E. coli* O121:H19. In 1999, bacteria of this serotype were implicated as the cause of a waterborne outbreak of HC in Connecticut (13). Also, a characterization of 67 EHEC strains from Swedish pa-




FIG. 3. Plot of the variable nucleotide sites in intimin alleles. Each vertical line represents a site that differs between the O121:H19 epsilon sequence and the allele labeled in the figure. The domain structure of intimin is shown at the bottom of the figure, with the CCD shaded in gray.

tients collected between 1997 and 1999 (39) found that about one-third of the strains were non-O157 and that of these, 17% were O121:H19. In a surveillance study of STEC strains in diarrheal disease, members of the O121:H19 clone group were the third most common STEC serotype (11), accounting for ~25% of the cases of diarrheal disease caused by non-O157 STEC infections (P. I. Tarr, unpublished data). The molecular data and the recent increase in the incidence of O121:H19 infections suggest that this clone is an emerging pathogen, and surveillance methods should be able to detect this as well as other non-O157 STEC serotypes.

#### ACKNOWLEDGMENTS

This research was supported by the National Institutes of Health (grant AI-47499) (P.I.T.) and the Enteric Pathogen Research Unit (grant N01-AI-65299) at the University of Maryland Medical School.

J. Paton kindly provided primer sequences for *saa* prior to publication.

#### REFERENCES

- Beutin, L., D. Geier, S. Zimmermann, S. Aleksic, H. A. Gillespie, and T. S. Whittam. 1997. Epidemiological relatedness and clonal types of natural populations of *Escherichia coli* strains producing Shiga toxins in separate populations of cattle and sheep. *Appl. Environ. Microbiol.* **63**:2175–2180.
- Blattner, F. R., G. Plunkett III, C. A. Bloch, N. T. Perna, V. Burland, M. Riley, J. Collado-Vides, J. D. Glassner, C. K. Rode, G. F. Mayhew, J. Gregor, N. W. Davis, H. A. Kirkpatrick, M. A. Goeden, D. J. Rose, B. Mau, and Y. Shao. 1997. The complete genome sequence of *Escherichia coli* K-12. *Science* **277**:1453–1462.
- Brunder, W., H. Schmidt, M. Frosch, and H. Karch. 1999. The large plasmids of Shiga-toxin-producing *Escherichia coli* (STEC) are highly variable genetic elements. *Microbiology* **145**:1005–1014.
- Caprioli, A., A. E. Tozzi, G. Rizzoni, and H. Karch. 1997. Non-O157 Shiga toxin-producing *Escherichia coli* infections in Europe. *Emerg. Infect. Dis.* **3**:578–579.
- Donnenberg, M. S., and J. B. Kaper. 1992. Enteropathogenic *Escherichia coli*. *Infect. Immun.* **60**:3953–3961.
- Fey, P. D., R. S. Wickert, M. E. Rupp, T. J. Safraneck, and S. H. Hinrichs. 2000. Prevalence of non-O157:H7 Shiga toxin-producing *Escherichia coli* in diarrheal stool samples from Nebraska. *Emerg. Infect. Dis.* **6**:530–533.
- Hayashi, T., K. Makino, M. Ohnishi, K. Kurokawa, K. Ishii, K. Yokoyama, C. G. Han, E. Ohtsubo, K. Nakayama, T. Murata, M. Tanaka, T. Tobe, T. Iida, H. Takami, T. Honda, C. Sasakawa, N. Ogasawara, T. Yasunaga, S. Kuhara, T. Shiba, M. Hattori, and H. Shinagawa. 2001. Complete genome sequence of enterohemorrhagic *Escherichia coli* O157:H7 and genomic comparison with a laboratory strain K-12. *DNA Res.* **8**:11–22.
- Jaeger, J. L., and D. W. Acheson. 2000. Shiga toxin-producing *Escherichia coli*. *Curr. Infect. Dis. Rep.* **2**:61–67.
- Karch, H., T. Meyer, H. Rüssmann, and J. Heesemann. 1992. Frequent loss of Shiga-like toxin genes in clinical isolates of *Escherichia coli* upon subcultivation. *Infect. Immun.* **60**:3464–3467.
- Karch, H., S. Schubert, D. Zhang, W. Zhang, H. Schmidt, T. Olschlager, and J. Hacker. 1999. A genomic island, termed high-pathogenicity island, is present in certain non-O157 Shiga toxin-producing *Escherichia coli* clonal lineages. *Infect. Immun.* **67**:5994–6001.
- Klein, E. J., J. R. Stapp, C. R. Clausen, D. R. Boster, J. G. Wells, X. Qin, D. L. Swerdlow, and P. I. Tarr. 2002. Shiga toxin-producing *Escherichia coli* in children with diarrhea: a prospective point-of-care study. *J. Pediatr.* **141**:172–177.
- Kumar, S., K. Tamura, I. Jakobsen, and M. Nei. 2000. MEGA 2: molecular evolutionary genetics analysis program. Version 2.0. Pennsylvania State University, University Park, Pa.
- McCarthy, T. A., N. L. Barrett, J. L. Hadler, B. Salisbury, R. T. Howard, D. W. Dingman, C. D. Brinkman, W. F. Bibb, and M. L. Cartter. 2001. Hemolytic-uremic syndrome and *Escherichia coli* O121 at a lake in Connecticut, 1999. *Pediatrics* **108**:E59. [Online.] <http://www.ncbi.nlm.nih.gov>.
- McDaniel, T. K., K. G. Jarvis, M. S. Donnenberg, and J. B. Kaper. 1995. A genetic locus of enterocyte effacement conserved among diverse enterobacterial pathogens. *Proc. Natl. Acad. Sci. USA* **92**:1664–1668.
- McGraw, E. A., J. Li, R. K. Selander, and T. S. Whittam. 1999. Molecular evolution and mosaic structure of  $\alpha$ ,  $\beta$ , and  $\gamma$  intimins of pathogenic *Escherichia coli*. *Mol. Biol. Evol.* **16**:12–22.
- Nataro, J. P., and J. B. Kaper. 1998. Diarrheagenic *Escherichia coli*. *Clin. Microbiol. Rev.* **11**:142–201.
- Nei, M., and S. Kumar. 2000. Molecular evolution and phylogenetics. Oxford University Press, New York, N.Y.
- O'Brien, A. D., V. L. Tesh, A. Donohue-Rolfe, M. P. Jackson, S. Olsnes, K. Sandvig, A. A. Lindberg, and G. T. Keusch. 1992. Shiga toxin: biochemistry, genetics, mode of action, and role in pathogenesis. *Curr. Top. Microbiol. Immunol.* **180**:65–94.
- Ochman, H., and R. K. Selander. 1984. Standard reference strains of *Escherichia coli* from natural populations. *J. Bacteriol.* **157**:690–693.
- O'Loughlin, E. V., and R. M. Robins-Browne. 2001. Effect of Shiga toxin and Shiga-like toxins on eukaryotic cells. *Microbes Infect.* **3**:493–507.
- Oswald, E., H. Schmidt, S. Morabito, H. Karch, O. Marches, and A. Caprioli. 2000. Typing of intimin genes in human and animal enterohemorrhagic and enteropathogenic *Escherichia coli*: characterization of a new intimin variant. *Infect. Immun.* **68**:64–71.
- Paton, A. W., and J. C. Paton. 1998. Detection and characterization of Shiga toxinogenic *Escherichia coli* by using multiplex PCR assays for *stx*<sub>1</sub>, *stx*<sub>2</sub>, *eaeA*, enterohemorrhagic *E. coli* *hlyA*, *rfb*<sub>O111</sub>, and *rfb*<sub>O157</sub>. *J. Clin. Microbiol.* **36**:598–602.
- Paton, A. W., and J. C. Paton. 2002. Direct detection and characterization of Shiga toxinogenic *Escherichia coli* by multiplex PCR for *stx*<sub>1</sub>, *stx*<sub>2</sub>, *eae*, *ehxA*, and *saa*. *J. Clin. Microbiol.* **40**:271–274.
- Paton, A. W., E. Voss, P. A. Manning, and J. C. Paton. 1998. Antibodies to lipopolysaccharide block adherence of Shiga toxin-producing *Escherichia coli* to human intestinal epithelial (Henle 407) cells. *Microb. Pathog.* **24**:57–63.
- Paton, J. C., and A. W. Paton. 1998. Pathogenesis and diagnosis of Shiga toxin-producing *Escherichia coli* infections. *Clin. Microbiol. Rev.* **11**:450–479.
- Perna, N. T., G. Plunkett, V. Burland, B. Mau, J. D. Glasner, D. J. Rose, G. F. Mayhew, P. S. Evans, J. Gregor, H. A. Kirkpatrick, G. Posfai, J. Hackett, S. Klink, A. Boutin, Y. Shao, L. Miller, E. J. Grobeck, N. W. Davis, A. Lim, E. T. Dimalanta, K. D. Potamousis, J. Apodaca, T. S. Anantharaman, J. Lin, G. Yen, D. C. Schwartz, R. A. Welch, and F. R. Blattner. 2001. Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature* **409**:529–533.
- Reid, S. D., C. J. Herbelin, A. C. Bumbaugh, R. K. Selander, and T. S. Whittam. 2000. Parallel evolution of virulence in pathogenic *Escherichia coli*. *Nature* **406**:64–67.
- Riley, L. W., R. S. Remis, S. D. Helgerson, H. B. McGee, J. G. Wells, B. R. Davis, R. J. Hebert, E. S. Olcott, L. M. Johnson, N. T. Hargrett, P. A. Blake, and M. L. Cohen. 1983. Hemorrhagic colitis associated with a rare *Escherichia coli* serotype. *N. Engl. J. Med.* **308**:681–685.
- Schmidt, H., B. Henkel, and H. Karch. 1997. A gene cluster closely related to type II secretion pathway operons of gram-negative bacteria is located on the large plasmid of enterohemorrhagic *Escherichia coli* O157 strains. *FEMS Microbiol. Lett.* **148**:265–272.
- Schubert, S., A. Rakin, H. Karch, E. Carniel, and J. Heesemann. 1998. Prevalence of the “high-pathogenicity island” of *Yersinia* species among *Escherichia coli* strains that are pathogenic to humans. *Infect. Immun.* **66**:480–485.
- Selander, R. K., D. A. Caugant, H. Ochman, J. M. Musser, M. H. Gilmour, and T. S. Whittam. 1986. Methods of multilocus enzyme electrophoresis for bacterial population genetics and systematics. *Appl. Environ. Microbiol.* **51**:873–884.
- Tarr, C. L., and T. S. Whittam. 2002. Molecular evolution of the intimin gene in O111 clones of pathogenic *Escherichia coli*. *J. Bacteriol.* **184**:479–487.
- Tarr, P. I. 1995. *Escherichia coli* O157:H7: clinical, diagnostic, and epidemiological aspects of human infection. *Clin. Infect. Dis.* **20**:1–10.
- Tarr, P. I., S. S. Bilge, J. C. Vary, Jr., S. Jelacic, R. L. Habeeb, T. R. Ward, M. R. Baylor, and T. E. Besser. 2000. Iha: a novel *Escherichia coli* O157:H7 adherence-conferring molecule encoded on a recently acquired chromosomal island of conserved structure. *Infect. Immun.* **68**:1400–1407.
- Tarr, P. I., and M. A. Neill. 1996. Perspective: the problem of non-O157:H7 Shiga toxin (Verocytotoxin)-producing *Escherichia coli*. *J. Infect. Dis.* **174**:1136–1139.
- Tatsuno, I., M. Horie, H. Abe, T. Miki, K. Makino, H. Shinagawa, H. Taguchi, S. Kamiya, T. Hayashi, and C. Sasakawa. 2001. *toxB* gene on pO157 of enterohemorrhagic *Escherichia coli* O157:H7 is required for full epithelial cell adherence phenotype. *Infect. Immun.* **69**:6660–6669.
- Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
- Tobe, T., T. Hayashi, C. G. Han, G. K. Schoolnik, E. Ohtsubo, and C. Sasakawa. 1999. Complete DNA sequence and structural analysis of the enteropathogenic *Escherichia coli* adherence factor plasmid. *Infect. Immun.* **67**:5455–5462.
- Welinder-Olsson, C., M. Badenfors, T. Cheasty, E. Kjellin, and B. Kaijser. 2002. Genetic profiling of enterohemorrhagic *Escherichia coli* strains in re-

- lation to clonality and clinical signs of infection. *J. Clin. Microbiol.* **40**:959–964.
40. **Whittam, T. S.** 1998. Evolution of *Escherichia coli* O157:H7 and other Shiga toxin-producing *E. coli* strains, p. 195–209. *In* J. B. Kaper and A. D. O'Brien (ed.), *Escherichia coli* O157:H7 and other Shiga toxin-producing *E. coli* strains. American Society for Microbiology, Washington, D.C.
41. **Whittam, T. S., M. L. Wolfe, I. K. Wachsmuth, F. Ørskov, I. Ørskov, and R. A. Wilson.** 1993. Clonal relationships among *Escherichia coli* strains that cause hemorrhagic colitis and infantile diarrhea. *Infect. Immun.* **61**:1619–1629.
42. **World Health Organization.** 1999. Zoonotic non-O157 Shiga toxin-producing *Escherichia coli* (STEC). Report of a WHO Scientific Working Group meeting, 23–26 June 1998, Berlin, Germany. [Online.] <http://www.who.int/emc-documents/zoonoses/whocsgraph988c.html>.

---

*Editor:* J. T. Barbieri