

mechanisms remains to be fully elucidated, but its widespread distribution, numerous effects, and subtle interrelations with other physiological control systems, particularly in controlling oxygen balance, suggest that it may be a most important homeostatic metabolite.

ANDREW H WATT
Clinical research officer

PHILIP A ROUTLEDGE
Senior lecturer

Department of Pharmacology and Therapeutics,
University of Wales College of Medicine,
Cardiff CF4 4XN

- 1 Anonymous. 3rd International Symposium on adenosine. *Pflügers Arch* 1986;407(suppl 1):1-57S.
- 2 Newby AC. Adenosine and the concept of "retaliatory metabolites." *Trends in Biochemical Sciences* 1984;9:42-4.
- 3 Schrader J, Haddy FJ, Gerlach E. Release of adenosine, inosine and hypoxanthine from the isolated guinea pig heart during hypoxia, flow-autoregulation and reactive hyperemia. *Pflügers Arch* 1977;369:1-6.
- 4 Achterberg PW, Stroewe RJ, DeJong JW. Myocardial adenosine cycling rates during normoxia and under conditions of stimulated purine release. *Biochem J* 1986;235:13-7.
- 5 Berne RM. The role of adenosine in the regulation of coronary blood flow. *Circ Res* 1980;47:807-13.
- 6 Berne RM. Criteria for involvement of adenosine in the regulation of blood flow. In: Paton DM, ed. *Methods in adenosine research*. New York: Plenum Press, 1986:331-6.
- 7 Watkinson WP, Foley DH, Rubio R, Berne RM. Myocardial adenosine formation with increased cardiac performance in the dog. *Am J Physiol* 1979;236:H13-21.
- 8 Katori M, Berne RM. Release of adenosine from anoxic hearts. *Circ Res* 1966;19:410-25.
- 9 Olsson RA. Changes in content of purine nucleoside in canine myocardium during coronary occlusion. *Circ Res* 1970;26:301-6.
- 10 Manfredi JP, Sparks HV, Jr. Adenosine's role in coronary vasodilation induced by atrial pacing and norepinephrine. *Am J Physiol* 1982;243:H536-H545.
- 11 Hanley FL, Grattan MT, Stevens MB, Hoffman JI. Role of adenosine in coronary autoregulation. *Am J Physiol* 1986;250:H558-66.
- 12 Merrill GF, Downey HF, Jones CE. Adenosine deaminase attenuates canine coronary vasodilatation during systemic hypoxia. *Am J Physiol* 1986;250:H579-83.
- 13 Watt AH, Penny WJ, Singh H. The effect of adenosine on coronary flow and left ventricular function in man. *Pflügers Arch* 1986;407(suppl 1):54S.
- 14 Sollevi A, Torssell L, Öwall A, Edlund A. Levels and effects of adenosine in humans. *Pflügers Arch* 1986;407(suppl 1):25S.
- 15 Szentmiklosi AJ, Nemeth M, Szegi J, Papp JGy, Szekeres L. Effect of adenosine on sinoatrial and ventricular automaticity of the guinea pig. *Arch Pharmacol* 1980;311:147-9.
- 16 DiMarco JP, Sellers TD, Berne RM, West GA, Belardinelli L. Adenosine: electrophysiologic effects and therapeutic use for terminating paroxysmal supraventricular tachycardia. *Circulation* 1983;68:1254-63.
- 17 Belardinelli L, Mattos EC, Berne RM. Evidence for adenosine mediation of atrioventricular block in the ischemic canine myocardium. *J Clin Invest* 1981;68:195-205.
- 18 Adgey AAJ, Allen JD, Geddes JS, et al. Acute phase of myocardial infarction. *Lancet* 1971;ii:501-4.
- 19 DiMarco JP, Sellers TD, Lerman BB, Greenberg ML, Berne RM, Belardinelli L. Diagnostic and therapeutic use of adenosine in patients with supraventricular tachyarrhythmias. *J Am Coll Cardiol* 1985;6:417-25.
- 20 Watt AH, Bernard MS, Webster J, Passani SL, Stephens MR, Routledge PA. Intravenous adenosine in the treatment of supraventricular tachycardia: a dose-ranging study and interaction with dipyridamole. *Br J Clin Pharmacol* 1986;21:227-30.
- 21 Klabunde RE. Dipyridamole inhibition of adenosine metabolism in human blood. *Eur J Pharmacol* 1983;93:21-6.
- 22 Sollevi A, Lagerkranser M, Irestedt L, Gordon E, Lindquist C. Controlled hypotension with adenosine in cerebral aneurysm surgery. *Anesthesiology* 1984;61:400-5.
- 23 Fukunaga AF, Flacke WE, Bloor BC. Hypotensive effects of adenosine and adenosine triphosphate compared with sodium nitroprusside. *Anesth Analg* 1982;61:273-8.
- 24 Sollevi A, Lagerkranser M, Andreen M, Irestedt L. Relationship between arterial and venous adenosine levels and vasodilatation during ATP- and adenosine-infusion in dogs. *Acta Physiol Scand* 1984;120:171-6.
- 25 Spielman WS. Antagonistic effect of theophylline on the adenosine-induced decrease in renin release. *Am J Physiol* 1984;247:F246-51.
- 26 Winn HR, Morii S. The role of adenosine in autoregulation of cerebral blood flow. *Ann Biomed Eng* 1985;13:321-8.
- 27 Dobson JG Jr, Rubio R, Berne RM. Role of adenine nucleotides, adenosine and inorganic phosphate in the regulation of skeletal muscle blood flow. *Circ Res* 1971;29:375-84.
- 28 Arend LJ, Haramati A, Thompson CI, Spielman WS. Adenosine-induced decrease in renin release: dissociation from hemodynamic effects. *Am J Physiol* 1984;247:F447-52.
- 29 Spielman WS, Thompson CI. A proposed role for adenosine in the regulation of renal hemodynamics and renin release. *Am J Physiol* 1982;242:F423-35.
- 30 Burnstock G. Purinergic transmission. In: Iversen LL, ed. *Handbook of psychopharmacology*. Vol 5. New York: Plenum Press, 1975:131-94.
- 31 Stone TW. Physiological roles for adenosine and adenosine 5'-triphosphate in the nervous system. *Neuroscience* 1981;6:523-55.
- 32 Daly JW, Bruns RF, Snyder SH. Adenosine receptors in the central nervous system: relationship to the central actions of methylxanthines. *Life Sci* 1981;28:2083-97.
- 33 Dragunow M. Adenosine: the brain's natural anticonvulsant? *Trends in Pharmacological Sciences* 1986;7:128-30.
- 34 Phillis JW. Interactions of the anticonvulsants diphenylhydantoin and carbamazepine with adenosine on cerebral cortical neurones. *Epilepsia* 1984;25:765-72.
- 35 Born GVR, Honour AJ, Mitchell JRA. Inhibition by adenosine and by 2-chloroadenosine of the formation and embolization of platelet thrombi. *Nature* 1964;202:761-5.
- 36 Sollevi A, Torssell L, Fredholm BB, Settergren G, Blombäck M. Adenosine spares platelets during cardiopulmonary bypass in man without causing systemic vasodilatation. *Scand J Thorac Cardiovasc Surg* 1985;19:155-9.
- 37 Pritchard JB, Chavez-Peon F, Berlin RD. Purines: supply by liver to tissues. *Am J Physiol* 1970;219:1263-8.
- 38 Lerner MH, Lowy BA. The formation of adenosine in rabbit liver and its possible role as a direct precursor of erythrocyte adenine nucleotides. *J Biol Chem* 1974;249:959-66.
- 39 Watt AH, Routledge PA. Adenosine stimulates respiration in man. *Br J Clin Pharmacol* 1985;20:503-6.

- 40 Watt AH, Reid PG, Stephens MR, Routledge PA. Adenosine-induced respiratory stimulation depends on site of infusion. Evidence for an action on the carotid body? *Br J Clin Pharmacol* 1986;22:238-39P.
- 41 Reid PG, Watt AH, Routledge PA, Smith AP. Adenosine not inosine stimulates respiration in man. *Pflügers Arch* 1986;407(suppl 1):57S.
- 42 Biaggioni I, Olafsson B, Robertson R, Hollister AS, Robertson D. Characteristics of adenosine-induced stimulation of respiration in man. *Pflügers Arch* 1986;407(suppl 1):55S.
- 43 Buss DC, Routledge PA, Watt AH. Respiratory effects of intravenous adenosine in anaesthetised rabbits before and after carotid nerve section. *Br J Pharmacol* 1986;88:413P.
- 44 Monteiro EC, Ribeiro JA. Adenosine modulation of respiration mediated by carotid body chemoreceptors in the rat. In: Ribeiro JA, Pallot DJ, eds. *Chemoreceptors in respiratory control*. London: Croom Helm (in press).
- 45 Cushley MJ, Tattersfield AE, Holgate ST. Inhaled adenosine and guanosine on airway resistance in normal and asthmatic subjects. *Br J Clin Pharmacol* 1983;15:161-5.
- 46 Cushley MJ, Tattersfield AE, Holgate ST. Adenosine-induced bronchoconstriction in asthma. Antagonism by inhaled theophylline. *Am Rev Respir Dis* 1984;129:380-4.

Primary pulmonary hypertension

Pulmonary hypertension is usually secondary to cardiac or pulmonary disease, and only rarely is it "primary" or "unexplained." At sea level for a cardiac output of 5-6 litres systolic pulmonary arterial pressure is about 20 mm Hg and diastolic about 12 mm Hg. In pulmonary hypertension the mean arterial pressure is above 19-20 mm Hg. Although rare,¹ primary pulmonary hypertension is important because it affects young people and has a 10 year survival of only 25%.²

The diagnosis is often delayed³ as the signs of raised pulmonary vascular resistance without a cardiac cause, respiratory insufficiency, or collagen vascular disease may be missed. The diagnosis must be considered in breathless patients, particularly when associated with syncope and angina.^{2,4} Right heart failure develops late,⁴ but a loud pulmonary second heart sound is often heard.^{2,4} Signs of raised pulmonary vascular resistance on chest radiograph and electrocardiogram are found in 95% of cases.² Spirometric values are normal, but gas transfer for carbon monoxide may be low.⁵ Doppler M mode echocardiography shows characteristic signs and allows an assessment of pulmonary artery pressure and exclusion of a left to right shunt.^{6,8} Ventilation and perfusion lung scintigraphy can exclude proximal vessel thromboemboli.⁹

Right heart catheterisation confirms the diagnosis and provides prognostic information: the lower the cardiac output the worse the prognosis.^{2,4} The most discriminating test is the mixed venous oxygen saturation: when it is less than 63% survival at five years is 17%, whereas when it is greater than 63% survival is as high as 55%.² "Spontaneous" recovery may occur in those with high cardiac output.^{4,10}

Surgical endarterectomy can improve the haemodynamic measurements and symptoms in patients with longstanding proximal pulmonary artery obstruction from thromboembolism.^{11,12} Diagnosis is with ventilation and perfusion lung scintigraphy and preoperative assessment requires pulmonary angiography.⁹

In patients without proximal vessel obstruction the haemodynamic disturbance is a result of a "restriction" of the peripheral pulmonary vasculature, in contrast to the "vasoconstriction" of hypoxia.¹³ Idiopathic pulmonary hypertension is associated with a generalised loss in the precapillary non-muscular vessels,^{13,14} whereas in patients with primary pulmonary hypertension and peripheral thrombosis muscular and non-muscular arteries are obstructed in a patchy distribution.¹³ Adaptive dilatational lesions

and smooth muscle hypertrophy with intimal fibrosis are seen in vessels upstream from these restrictions.¹⁵⁻¹⁷

Treatment is aimed at reducing the consequences of the low cardiac output. By preventing secondary thromboembolism anticoagulants improve survival in both idiopathic and peripheral thrombotic forms.² By raising cardiac output, and therefore oxygen tissue delivery, vasodilators relieve the symptoms.¹⁸ A trial of vasodilators during catheterisation identifies responders.¹⁹ A titrated dose of intravenous prostacyclin is the most effective acute vasodilator²⁰ and because of its short half life²¹ avoids the dangers of a sustained fall in systemic arterial pressure.²² Successful vasodilatation is shown by a greater than 30% rise in cardiac output²³ without an increase in pulmonary artery pressure and a less than 20% fall in systemic arterial pressure.

Patients with clear ventilation and perfusion scans and a mixed venous oxygen saturation greater than 63% should be given anticoagulants and an oral vasodilator (hydralazine,²⁴ captopril,²⁵ or diltiazem²⁶) if they have responded to prostacyclin at catheterisation. If the mixed venous oxygen saturation is less than 63% and prostacyclin was successful at catheterisation a long term infusion of prostacyclin, though expensive, can improve the quality of life.^{18, 27} As prostacyclin inhibits platelet aggregation²⁸ anticoagulants are theoretically not needed. The optimal long term dose of vasodilators can be calculated from monitoring systemic arterial pressure and by measuring maximum rate of oxygen consumption during exercise testing.²⁹ When the mixed venous oxygen saturation is less than 63% and there is no response to vasodilators at catheterisation heart-lung transplantation may be the treatment.³⁰ The 5-10% of patients with primary pulmonary hypertension and veno-occlusive disease can probably be managed according to the same system.

More effective treatments will come only when we learn more of the homeostatic mechanisms governing the interface between the blood and endothelium. Acute experimental injury of pulmonary endothelium may lead to obliteration of the precapillary arteries.^{31, 32} Clinical conditions associated with thrombosis—for example, with lupus anticoagulant—may cause primary pulmonary hypertension.³³ This protein (an IgM or IgG) may act by interfering with prostacyclin formation.^{34, 35} Other inherited thrombotic disorders—for example, deficiencies of antithrombin III³⁶ or heparin cofactor II³⁷—may account for some familial cases of primary pulmonary hypertension³⁸ and for some people developing the disease after taking aminorex fumarate³⁹ or fenfluramine.⁴⁰

TIM HIGENBOTTAM

Consultant Physician and Director,
Regional Pulmonary Function Laboratory,
Papworth Hospital,
Cambridge CB3 8RE

- 11 Moser KM, Spragg RG, Utley J, Daily PO. Chronic thrombotic obstruction of major pulmonary arteries. Results of thromboendarterectomy in 15 patients. *Ann Intern Med* 1983;99:299-305.
- 12 Daily PO, Johnsson GG, Simmons CJ, Moser KM. Surgical management of chronic pulmonary embolism: surgical treatment and late results. *J Thorac Cardiovasc Surg* 1980;79:523-31.
- 13 Reid LM. Structure and function in pulmonary hypertension: new perceptions. *Chest* 1986;89:279-88.
- 14 Anderson EG, Simon G, Reid L. Primary and thrombo-embolic pulmonary hypertension: a quantitative pathological study. *J Pathol* 1973;110:273-93.
- 15 Wagenvoort CA, Wagenvoort N. Primary pulmonary hypertension. A pathological study of the lung vessels in 156 clinically diagnosed cases. *Circulation* 1970;42:1163-84.
- 16 Heath D, Whitaker W, Brown JW. Idiopathic pulmonary hypertension. *Br Heart J* 1957;19:83-92.
- 17 Bjornsson J, Edwards WD. Primary pulmonary hypertension: a histopathological study of 80 cases. *Mayo Clin Proc* 1985;60:16-25.
- 18 Higenbottam TW, Wheelton D, Wells F, Wallwork J. Long-term treatment of primary pulmonary hypertension with continuous intravenous epoprostenol (prostacyclin). *Lancet* 1984;ii:1046-7.
- 19 Packer M, Greenberg M, Massie B, Dash H. Deleterious effects of hydralazine in patients with pulmonary hypertension. *N Engl J Med* 1982;306:1326-31.
- 20 Simonneau G, Herve P, Petitpretz P, et al. Detection of a reversible component in primary pulmonary hypertension. *Am Rev Respir Dis* 1986;133:223A.
- 21 Rubin LJ, Graves BM, Reaves JT, Frosolano M, Handel F, Cato AE. Prostacyclin induced acute pulmonary vasodilation in pulmonary hypertension. *Circulation* 1982;66:334-8.
- 22 Wade EG, Rowlands DJ. Diazoxide in treatment of primary pulmonary hypertension. *Br Heart J* 1979;42:362-3.
- 23 Rich S, D'Alonzo DR, Dantzker PS, Levy L. Magnitude and implication of spontaneous haemodynamic variability in primary pulmonary hypertension. *Am J Cardiol* 1985;55:159-63.
- 24 Rubin LJ, Peter RH. Oral hydralazine therapy for primary pulmonary hypertension. *N Engl J Med* 1980;302:69-73.
- 25 Ikram H, Maslowski AH, Nicholls MG, Espiner EA, Hull FTL. Haemodynamic and hormonal effects of captopril in primary pulmonary hypertension. *Br Heart J* 1982;48:541-5.
- 26 Crevey BJ, Dantzker DR, Bower JS, Popat KD, Walker SD. Haemodynamic and gas exchange effects of intravenous diltiazem in patients with pulmonary hypertension. *Am J Cardiol* 1982;49:578-83.
- 27 Jones DK, Higenbottam TW, Wallwork J. Practical experience of long-term prostacyclin with primary pulmonary hypertension. *Am Rev Respir Dis* 1985;31:85.
- 28 Moncada S, Gryglewski R, Bunting S, Vane JR. An enzyme isolated from arteries transforms prostaglandin endoperoxides to an unstable substance that inhibits platelet aggregation. *Nature* 1976;263:663-5.
- 29 D'Alonzo GE, Gianotti L, Dantzker DR. Non-invasive assessment of haemodynamic improvement during chronic vasodilator therapy in obliterative pulmonary hypertension. *Am Rev Respir Dis* 1986;133:380-4.
- 30 Burke CM, Baldwin JC, Morris AJ, et al. Twenty-eight cases of human heart-lung transplantation. *Lancet* 1986;ii:518-9.
- 31 Meyrick B, Gamble W, Reid L. Development of crotalaria pulmonary hypertension: haemodynamic and structural study. *Am J Physiol* 1980;230:692-702.
- 32 Coffesky JT, Jones RC, Reid LM, Evans JN. Mechanical properties and structures of isolated pulmonary arteries following prolonged hyperoxia. *Am Rev Respir Dis* 1985;135:405-12.
- 33 Asherson RA, Mockworth-Young CG, Boey ML, et al. Pulmonary hypertension in systemic lupus erythematosus. *Br Med J* 1983;287:1024-5.
- 34 Carreras LO, Defreyne G, Machin S. Arterial thrombosis, intra-uterine death, and lupus anticoagulant: detection of an immunoglobulin interfering with prostacyclin formation. *Lancet* 1981;ii:244-6.
- 35 Carreras LO, Veringhen JG. "Lupus" anticoagulant and thrombosis: possible role of inhibition of prostacyclin formation. *Thromb Haemost* 1982;48:38-40.
- 36 Eigeberg O. Inherited antithrombin deficiency causing thrombophilia. *Thrombosis et Diathesis Haemorrhagica* 1965;13:516-30.
- 37 Tran TH, Marbet GA, Duckert F. Association of hereditary heparin Co-factor II deficiency with thrombosis. *Lancet* 1983;ii:413-6.
- 38 Loyd JE, Primm RK, Newman JH. Familial primary pulmonary hypertension: clinical patterns. *Am Rev Respir Dis* 1984;129:194-7.
- 39 Gurthier HP, Gertsch M, Salzmann C, Shepper M, Stucki P, Wyss F. Haufen sich die primär vasculären Formen des chronischen cor pulmonale? *Schweiz Med Wochenschr* 1986;98:1579-89.
- 40 Douglas JG, Munro JF, Kitchin AH, Muir AL, Proudfoot AT. Pulmonary hypertension and fenfluramine. *Br Med J* 1981;283:881-3.

Long live health promotion

Health promotion is an intensely political issue. Twenty years ago people naively imagined that a whiff of advertising and a few leaflets would be enough to solve the smoking problem. Now most people—although the minister currently responsible for prevention would seem to be an exception—recognise that exhorting people to eat better, pull themselves together, "get on their bikes," or "just say no" is not enough. Information is certainly needed (and it must be carefully composed), but in addition those who want to improve the health of the people will have to do battle with the immensely powerful organisations that have a vested interest in, for example, keeping people smoking, drinking large quantities of alcohol, and eating unhealthy food. As Peter Taylor clearly described in *Smoke Ring: the Politics of Tobacco*¹ and as the television programme *Yes, Prime Minister*² has so humorously but so chillingly illustrated, those campaigners are also likely

- 1 Loogen F, Both A. Primare pulmonare hypertone. *Z Kardiol* 1976;65:1-14.
- 2 Fuster V, Steele PM, Edwards WD, Gersh BJ, McGoon MD, Frye RL. Primary pulmonary hypertension: a natural history and importance of thrombosis. *Circulation* 1984;70:580-7.
- 3 Voelkel N, Reeves JT. Primary pulmonary hypertension. In: Moser KM, ed. *Pulmonary vascular disease*. New York: Marcel Dekker, 1979:573-628.
- 4 Rozkovec A, Montanes P, Oakley CM. Factors that influence the outcome of primary pulmonary hypertension. *Br Heart J* 1986;55:449-58.
- 5 Alpert JS, Braunwald E. In: Braunwald E, ed. *Heart disease*. Philadelphia: Saunders, 1980: vol 2, p 1638.
- 6 Goodman DJ, Harrison DC, Papp RL. Echocardiographic features of primary pulmonary hypertension. *Am J Cardiol* 1974;33:438-43.
- 7 Hatle L, Angelsen BAJ, Transdal A. Non-invasive estimation of pulmonary artery systolic pressure with doppler ultrasound. *Br Heart J* 1981;45:157-65.
- 8 Kitabatake A, Inoue M, Asao MI, et al. Non-invasive evaluation of the ratio of pulmonary to systemic flow in ASD by Duplex Doppler echocardiography. *Circulation* 1984;69:73-9.
- 9 Moser KM, Fedullo PF, Shure D, Elliot G. In what patients with apparent primary pulmonary hypertension is pulmonary angiography indicated? *Am Rev Respir Dis* 1986;133:1221A.
- 10 Bourdillon PDV, Oakley CM. Regression of primary pulmonary hypertension. *Br Heart J* 1976;38:264-70.