

Research Article

JNK1-dependent antimitotic activity of thiazolidin compounds in human non-small-cell lung and colon cancer cells

F. Teraishi, S. Wu, J. Sasaki, L. Zhang, J. J. Davis, W. Guo, F. Dong and B. Fang*

Department of Thoracic and Cardiovascular Surgery, Unit 445, University of Texas M. D. Anderson Cancer Center, 1515 Holcombe Blvd., Houston, Texas 77030 (USA), Fax: +1713 794 4901, e-mail: bfang@mdanderson.org

Received 9 August 2005; received after revision 23 August 2005; accepted 24 August 2005
Online First 22 September 2005

Abstract. We recently identified two thiazolidin compounds, 5-[(4-methylphenyl)methylene]-2-(phenylamino)-4(5H)-thiazolone (MMPT) and 5-(2,4-dihydroxybenzylidene)-2-(phenylimino)-1,3-thiazolidin (DBPT), that inhibit the growth of human non-small-cell lung and colon cancer cells independent of P-glycoprotein and p53 status. Here we further investigated the mechanism by which these thiazolidin compounds mediate their anticancer effects. Treatment of cancer cells with MMPT and DBPT led to a time-dependent accumulation of cells arrested in the G2/M phase with modulation of the expression of proteins such as cyclin

B1, cdc25C, and phosphorylated histone H3. Moreover, treatment with MMPT and DBPT increased M-phase arrest with abnormal spindle formation. DBPT-mediated G2/M phase arrest and phosphorylation of cdc25C and histone H3 were abrogated when JNK activation was blocked either with SP600125, a specific JNK inhibitor, or a dominant-negative JNK1 gene. Moreover, DBPT-mediated microtubule disruption was also blocked by SP600125 treatment. Our results demonstrate that thiazolidin compounds can effectively induce G2/M arrest in cancer cells and that this G2/M arrest requires JNK activation.

Key words. Cancer therapy; mitotic arrest; microtubule disruption; non-small-cell lung cancer; cell cycle.

Microtubules have crucial roles in intracellular transport, morphogenesis, and cell division. Various agents targeting tubulin, the major protein component of microtubules, have been reported to cause cell cycle arrest at mitosis and induce apoptosis in cancer cells [1, 2]. These microtubule-interacting drugs can be classified into two major groups: the first, which includes drugs such as vinca alkaloids, colchicine, and nocodazole, interferes with mitosis and changes the balance of the tubulin in microtubules to a depolymerized state; the second group, which includes agents such as taxanes, changes the balance of tubulin to a polymerized state. Vinca alkaloids such as vinorelbine and vincristine have been widely used as cytotoxic agents

for the treatment of hematopoietic neoplasia and some solid tumors [3, 4]. The taxanes, including paclitaxel and docetaxel, are newer antitumor agents that are used clinically to treat several solid tumors of the head and neck, breast, lung, ovary, and bladder [5–7]. Despite their opposing effects on tubulin polymerization, all of these agents cause cell cycle arrest at the metaphase-to-anaphase transition [8], and all induce apoptosis.

Although microtubule-interacting drugs have been used clinically to inhibit the progression of some cancers, multiple cycles of treatment usually result in acquired resistance that abolishes drug efficacy [9]. One of the mechanisms by which neoplastic cells acquire resistance is the overexpression of efflux pumps such as P-glycoprotein, a transmembrane pump that binds to many anti-tumor drugs and increases their efflux out of cells. Many

* Corresponding author.

microtubule-interacting drugs, such as the taxanes and vinblastine, are well-known substrates of P-glycoprotein [10–13]. The development of new compounds that are effective against drug-resistant cells while sparing normal cells is therefore important for cancer therapy.

To this end, we recently identified two thiazolidin compounds, 5-[(4-methylphenyl)methylene]-2-(phenylamino-4(SH)-thiazolone) (MMPT) and 5-(2,4-dihydroxybenzylidene)-2-(phenylimino)-1,3-thiazolidin (DBPT), as antineoplastic agents [14, 15]. These compounds have selective cytotoxic effects against human lung and colon cancer cells *in vitro* and *in vivo*, and these effects are independent of P-glycoprotein expression. In addition, the compounds have no obvious toxic effects on normal human fibroblasts and mesenchymal stem cells at the 50% inhibitory concentration for cancer cell lines. We also found that MMPT and DBPT induce apoptosis through caspase activation that is dependent on JNK activation. However, JNK activation alone does not induce apoptosis, suggesting that other molecular events are involved in the antitumor activity of these compounds.

The goal of the present study was to investigate the mechanism of the antitumor action of MMPT and DBPT on non-small-cell lung and colon cancer cells. We found that MMPT and DBPT induced G2/M phase arrest evidenced by up-regulation of G2/M checkpoint proteins such as cyclin B1, cdc25C, Bcl-2, and histone H3. Treatment with MMPT and DBPT also caused cell cycle arrest at the mitotic phase, with abnormal spindle formation in cancer cells. Moreover, DBPT-mediated G2/M phase arrest and microtubule disruption were blocked by inhibition of JNK activation, suggesting that JNK activation also plays a critical role in G2/M arrest by thiazolidin compounds.

Materials and methods

Cells and culture conditions. The human non-small-cell lung cancer cell lines H1299 and H460 and the human colon cancer cell lines DLD-1 and HCT116 (p53 wild type and p53^{-/-}; generously provided by Dr. B. Vogelstein, The Johns Hopkins University, Baltimore, MD.) were routinely propagated in monolayer culture in RPMI 1640 medium supplemented with 10% heat-inactivated fetal calf serum, 25 mM HEPES, 100 units/ml penicillin, and 100 mg/ml streptomycin. Paclitaxel-resistant H460/TaxR cells and vinorelbine-resistant H460/VinR cells were generated from parental H460 cells by initially adding 5 nM paclitaxel or vinorelbine to the culture medium and thereafter escalating the dose of paclitaxel or vinorelbine to 100 nM. The resistance indexes (i.e., IC₅₀ in resistant cells/IC₅₀ in parental cells) were 80.0 in H460/TaxR cells and 213.9 in H460/VinR cells. Both H460/TaxR and H460/VinR expressed high levels of P-glycoprotein [15]. All cell lines were maintained in the presence of 5% CO₂ at 37°C.

Chemicals. MMPT and DBPT were obtained from ChemBridge Corporation (San Diego, Calif.), dissolved in dimethylsulfoxide (DMSO) to a concentration of 10 mM, and stored at 4°C as described previously [14, 15]. Nocodazole and vinorelbine were purchased from Sigma-Aldrich (St. Louis, MO.) and dissolved in DMSO to a concentration of 10 mM. Paclitaxel was purchased from Bristol-Myers Squibb (Princeton, N. J.). The JNK-specific inhibitor SP600125 was purchased from Calbiochem (La Jolla, Calif.), dissolved in DMSO, stored at -20°C, and protected from light. An equal volume of DMSO (<0.2%) had no obvious effects on cell viability and was used as a control.

Flow cytometry analysis. For analysis of intracellular DNA content, lung cancer and colon cancer cells were treated with various concentrations (0.3–20 μM) of MMPT or DBPT, collected in 0.125% trypsin, washed twice in PBS, and fixed in 70% methanol overnight at -20°C. The cells were then resuspended in PBS containing 10 μg/ml propidium iodide (Roche Diagnostics, Indianapolis, Ind.) and 10 μg/ml RNase A (Sigma-Aldrich) at 37°C for 30 min. Cell cycle analysis was performed using an EPICS Profile II flow cytometer (Coulter, Hialeah, Fla.) with MultiCycle Phoenix Flow Systems software (Phoenix Flow Systems, San Diego, Calif.). All experiments were repeated three times.

For detection of cells in mitosis, fixed cells were resuspended in PBS containing 5 μg/ml mouse monoclonal antibody against MPM-2 (Upstate Biotechnology, Lake Placid, N. Y.). The cells were then incubated with fluorescein isothiocyanate-conjugated goat anti-mouse secondary antibody (BD Biosciences PharMingen, San Diego, Calif.), 10 μg/ml propidium iodide, and 10 μg/ml RNase A at 37°C for 1 h in the dark. Stained cells were analyzed with the EPICS Profile II flow cytometer using MultiCycle Phoenix Flow Systems software. All experiments were repeated at least twice.

Western blot analysis. Western blot analysis was performed with primary antibodies against cyclin A, cdc2p34, cdc25C, and Bcl-2 (Santa Cruz Biotechnology, Santa Cruz, Calif.); cyclin B1 (NeoMarkers, Fremont, Calif.); phosphorylated histone H3 (Upstate Biotechnology), and β-actin (Sigma). Cells were washed twice in cold PBS, collected, and lysed in lysis buffer [62.5 mM Tris (pH 6.8), 2% sodium dodecyl sulfate, and 10% glycerol] containing 1x proteinase inhibitor cocktail (Roche Diagnostics). The lysates were spun at 14,000 x g in a microcentrifuge at 4°C for 10 min, and the supernatants were used as whole-cell extracts. Protein concentrations were determined using a BCA protein assay kit (Pierce, Rockford, Ill.). Equal amounts (30–50 μg) of proteins were subjected to electrophoresis under reducing conditions on 10–12.5% (w/v) polyacrylamide gels and then

electrophoretically transferred to nitrocellulose transfer membranes (Amersham Biosciences, Piscataway, N. J.). The membranes were incubated with primary antibody followed by peroxidase-linked secondary antibody. An electrochemiluminescence Western blotting system (Amersham) was used to detect secondary probes.

Immunofluorescence microscopy. Cells were grown on a glass chamber slide (Becton Dickinson, Franklin Lakes, N. J.) and treated with the indicated chemicals for 14 h. The cells were then washed with PBS and permeabilized with microtubule-stabilizing buffer [80 mM PIPES-KOH (pH 6.8), 5 mM egtazic acid, 1 mM MgCl₂, and 0.5% Triton X-100] for 5 min at room temperature. The permeabilized cells were fixed with chilled absolute methanol for 10 min at -20°C as described previously [16]. After being washed, cells were incubated with a mouse anti- α -tubulin monoclonal antibody (Sigma) for 1 h at ambient temperature and then incubated with a fluorescein isothiocyanate-conjugated anti-mouse secondary antibody (BD Biosciences PharMingen). After extensive washing, cells were rinsed once in water, mounted with Vectashield (Vector Laboratories, Burlingame, Calif.), and examined under a Nikon BX61 microscope (Melville, N. Y.).

Statistical analysis. Differences among the treatment groups were assessed by analysis of variance using StatSoft statistical software (Tulsa, Okla.). $p < 0.05$ was considered significant.

Results

Treatment with thiazolidin compounds causes induction of G2/M phase arrest in cancer cells. We recently found that MMPT and DBPT effectively induce JNK-dependent apoptosis in non-small-cell lung and colon cancer cells through caspase activation [14, 15]. To further elucidate the anticancer mechanisms of thiazolidin compounds, we examined the effects of MMPT and DBPT on cell cycle progression. H1299 and H460 cells were treated with 0.3–10 μ M MMPT and analyzed for cell cycle distribution by flow cytometry. Unexpectedly, G2/M phase arrest was elicited by 3 μ M MMPT after 12 h of treatment (fig. 1A). The accumulation of cells in the G2/M phase was accompanied by synchronous decreases in the numbers of cells in the G1 and S phases. Treatment with 10 μ M MMPT also resulted in accumulation of cells in the G2/M phase, beginning as early as 6 h and peaking around 80% after 12 h of treatment (fig. 1B). Because p53 is the wild type in H460 cells but homologously deleted in H1299 cells, this result suggested that the action of MMPT on G2/M phase arrest is p53 independent. Treatment of DLD-1, HCT116 wild type, and HCT116




Figure 1. Effect of thiazolidin compounds on cell cycle progression in cancer cells. (A) Flow cytometry analysis of propidium iodide-stained H1299 and H460 cells treated with various concentrations of MMPT for 12 h. (B) Percentages of H1299 and H460 cells in G1, S, and G2/M phases after treatment with 10 μ M MMPT for the indicated times. (C) Flow cytometry analysis of propidium iodide-stained DLD-1 cells, wild-type HCT116 cells (HCT116^{+/+}), and p53-deficient HCT116 cells (HCT116^{-/-}) were treated with 3 or 5 μ M DBPT, respectively, for the indicated time periods. (D) Percentages of DLD-1, HCT116^{+/+}, and HCT116^{-/-} cells in G1, S, and G2/M phases after treatment with 3 or 5 μ M DBPT for the indicated times.

p53-deficient colon cancer cells with 3 or 5 μ M DBPT, respectively, also resulted in a time-dependent increase

in cells arrested in G2/M phase (fig. 1C). Accumulation of cells in G2/M phase started as early as 6 h and peaked around 80% at 12 h after treatment in all three cell lines (fig. 1D), indicating that DBPT-mediated G2/M phase arrest is also p53 independent. In contrast, G2/M phase arrest was not observed in normal human fibroblasts treated with MMPT or DBPT at the 50% inhibitory concentration for cancer cells (data not shown).

Thiazolidin compounds affect the regulation of G2/M checkpoint proteins. We next used Western blotting to examine the effects of MMPT and DBPT on the expression of G2/M checkpoint proteins in cancer cells. Cyclin A expression was noticeably reduced after 24 h of treatment with MMPT and DBPT, and cyclin B1 expression was transiently increased after 6–12 h of treatment (fig. 2A, B). No dramatic change was observed for the level of cdc2p34. We also detected a slower-migrating form of the cdc25C phosphatase and a phosphorylated form of histone H3 after 6–24 h of MMPT treatment. Similar changes in the migration of cdc25C and phosphorylation of histone H3 were observed in DLD-1 cells 6–24 h after treatment with 3 μ M DBPT (fig. 2B).

Bcl-2 phosphorylation is a common event during M phase [17, 18]. To determine whether thiazolidin compounds use this mechanism to induce G2/M phase arrest, we examined time-dependent changes in Bcl-2 phosphorylation in H460, H460/TaxR, and H460/VinR cells treated with 10 μ M MMPT. Western blotting revealed an obvious increase in Bcl-2 phosphorylation after 12–24 h of MMPT treatment (fig. 2C). This result indicated that the effect of MMPT on Bcl-2 phosphorylation is independent of P-glycoprotein expression.

Induction of M phase arrest by thiazolidin compounds in cancer cells. To further pinpoint whether cell cycle arrest by thiazolidin compounds occurred at the G2 or M phase, we conducted a flow cytometry analysis of MMPT-treated H1299 and H460 cells stained with propidium iodide and an antibody against MPM-2, a marker for the onset of mitosis [19]. Anti-mouse secondary antibody was used as a control, and H460 cells treated with 50 nM paclitaxel were used as a positive control for MPM-2 (data not shown). We observed a marked accumulation of MPM-2-positive cells 12 h after treatment with 10 μ M MMPT (fig. 3A, B). A similar accumulation of MPM-2-positive cells was observed in DLD-1 cells treated with 3 μ M DBPT (fig. 3C). These results indicate that treatment of cancer cells with thiazolidin compounds results in an accumulation of cells arrested in M phase.

Effect of thiazolidin compounds on mitotic spindles in cancer cells. In addition to inducing M phase arrest and Bcl-2 phosphorylation, most microtubule-interacting agents inhibit mitosis by disrupting the organization




Figure 2. Thiazolidin compounds regulate G2/M checkpoint proteins. H1299 and H460 cells (A) and DLD-1 cells (B) were treated with 10 μ M MMPT or 3 μ M DBPT, respectively, for the indicated time periods. Expression and phosphorylation status of G2/M checkpoint proteins in whole-cell lysates were analyzed by immunoblotting with the indicated antibodies. Arrowheads indicate the phosphorylated form of cdc25C. β -Actin was used as the loading control. p-, phosphorylated. (C) H460, H460/TaxR, and H460/VinR cells were treated with 10 μ M MMPT for the indicated time periods. The Phosphorylation status of Bcl-2 in whole-cell lysates was analyzed by immunoblotting with anti-Bcl-2 antibody. Arrowhead indicates phosphorylated form of Bcl-2. β -actin was used as the loading control.

of mitotic spindles. To examine whether thiazolidin compounds share this ability, we evaluated the status of the mitotic spindles in H460, H460/TaxR, and H460/VinR cells after 14 h of treatment with DMSO, vinorelbine, MMPT, or paclitaxel (fig. 4A). Normal spindle formation was confirmed in dividing cells that had been treated with DMSO. Vinorelbine treatment of parental H460 cells resulted in the accumulation of many mitotic cells with depolymerized mitotic spindles. A similar accumulation was not observed in vinorelbine-treated H460/TaxR or H460/VinR cells. Monoaster spindles




Figure 3. Induction of M phase arrest by thiazolidin compounds. H1299 (A) and H460 (B) cells were treated with 10 μ M MMPT, and DLD-1 cells (C) were treated with 3 μ M DBPT for the indicated time periods. Cells were then stained with anti-MPM-2 antibody and propidium iodide (PI) and analyzed by flow cytometry. Histograms show the DNA content of cells; dot plots show proportions of mitotic and 4N cells after MMPT or DBPT treatment. Boxes indicate cells staining positively for MPM-2. Percentages of MPM-2-positive cells are shown.

were observed in all three cell lines after treatment with MMPT, and some of the H460/TaxR and H460/VinR cells had a two- or three-spindle polar organization. This result suggested that the effect of MMPT on microtubule disruption is independent of P-glycoprotein expression.

Similar changes were observed in DLD-1 cells after 14 h of treatment with 3 μ M DBPT (fig. 4B). In contrast, the increased microtubule polymerization caused by paclitaxel treatment resulted in destruction of the mitotic spindles and aggregated spindle formation in H460 and DLD-1 cells. This phenomenon was, however, not observed in paclitaxel-treated H460/TaxR and H460/VinR cells (fig. 4A, B).

JNK activation is required for DBPT-induced G2/M phase arrest and microtubule disruption. Our previ-




Figure 4. Effect of thiazolidin compounds on cellular microtubule organization. H460, H460/TaxR, and H460/VinR cells (A) and DLD-1 cells (B) were treated with DMSO (control), 100 nM vinorelbine, 15 μ M MMPT/3 μ M DBPT, or 100 nM paclitaxel. After 14 h, cellular microtubules were visualized by indirect immunofluorescence with anti- α -tubulin antibody. Normal spindle formation was observed in control (inside left upper box). Abnormal monoaster spindles were observed in cells treated with MMPT or DBPT but not in control cells.

ous study showed that activation of JNK but not ERK or p38 is required for MMPT- and DBPT-induced apoptosis [14, 15]. To further investigate the possible role of JNK activation on induction of G2/M phase arrest, we treated DLD-1 cells with DBPT in the presence or absence of the JNK inhibitor SP600125 for 12 h and then evaluated the cell cycle progression by flow cytometry. Treatment with 20 μ M SP600125 alone, which was enough to block DBPT-mediated JNK activation [14], resulted in a slight increase of cells at G2/M phase. Interestingly, SP600125 dramatically blocked DBPT-induced G2/M phase arrest (fig. 5A). Whereas the percentage of G2/M phase cells was 40.8% with 20 μ M SP600125 alone and 86.0% with 3 μ M DBPT alone, only 44.5% of cells were in the G2/M phase with the combination of SP600125 and DBPT. This result indicates that SP600125 inhibits DBPT-induced G2/M phase arrest ($p < 0.001$). Our previous data showed that JNK1 activation is crucial for DBPT-




Figure 5. The effects of JNK inhibition on DBPT-mediated G2/M phase arrest and microtubule disruption. (A) Inhibition of JNK activation blocks DBPT-mediated G2/M phase arrest. DLD-1 cells were treated with 3 μM DBPT in the presence or absence of 5 or 20 μM SP600125 for 12 h. The cell cycle progression of stained cells with PI was determined by flow cytometry. Data represent the mean ± SD from three independent experiments performed in triplicate. *p < 0.001, as compared with DBPT treatment alone. (B) Effect of JNK1 activation on DBPT-mediated G2/M phase arrest. DLD-1 cells stably transfected with dnJNK1, dnJNK2, or an empty vector (control) were treated with 3 μM DBPT for 6 h. The cell cycle progression of stained cells with PI was determined by flow cytometry. Data represent the mean ± SD of two independent experiments. *p < 0.001 and **p < 0.01, as compared with DBPT-treated parental DLD-1 cells. (C) SP600125 blocks DBPT-mediated histone H3 phosphorylation. DLD-1 cells were treated with 3 μM DBPT in the presence or absence of 20 μM SP600125 for 6 h. Whole-cell lysates were analyzed by immunoblotting with cdc25C and phosphorylated histone H3 antibodies. β-actin was used as the loading control. (D) Effect of JNK inhibition on DBPT-mediated microtubule disruption. DLD-1 cells were treated with DMSO (control), 3 μM DBPT, or 100 nM paclitaxel in the presence or absence of 20 μM SP600125 for 14 h. Cellular microtubules were visualized by indirect immunofluorescence with anti-α-tubulin antibody. The data represent one of two experiments with similar results.

mediated apoptosis and that different JNK isoforms may have different roles in DBPT-induced apoptosis [14]. To further test the role of JNK1 and JNK2 activation in DBPT-mediated G2/M phase arrest, we evaluated the effect of DBPT in DLD-1 cells stably transfected with plasmid expressing the dominant-negative mutants, dnJNK1 or dnJNK2 [14]. Cells were treated with 3 μM DBPT for 6 h, and cell cycle progression was analyzed by flow cytometry. The expression of dnJNK1 in DLD-1 cells significantly blocked DBPT-mediated G2/M phase arrest after 6 h (p < 0.001). In contrast, the expression of dnJNK2 enhanced DBPT-mediated G2/M phase arrest when compared with DBPT-treated parental DLD-1 cells (p < 0.01) (fig. 5B). These results indicate that JNK1 and JNK2 may also have different roles in DBPT-mediated

G2/M-phase arrest. We also evaluated the expression of cdc25C and phosphorylated histone H3 in DLD-1 cells treated with DBPT in the presence of SP600125 for 6 h. Treatment with 20 μM SP600125 alone had no effect on the phosphorylated form of histone H3 and cdc25C expression when compared with the control. Interestingly, SP600125 dramatically blocked DBPT-mediated changes in the migration of cdc25C and phosphorylation of histone H3 (fig. 5C). To further examine whether inhibition of JNK affects DBPT-caused microtubule disruption, DLD-1 cells were treated with DBPT or paclitaxel in the presence or absence of 20 μM SP600125 for 14 h and the mitotic spindles were visualized by immunofluorescence microscopy (fig. 5D). Central spindle formation was detected in DLD-1 cells treated with DMSO and

SP600125. Interestingly, although abnormal monoaster spindle formations were observed in DLD-1 cells treated with DBPT alone, they were not seen in DLD-1 cells treated with DBPT in the presence of SP600125. In contrast, SP600125 had little impact on microtubule depolymerization of DLD-1 cells treated with paclitaxel. These results suggest that JNK1 activation is required for DBPT-mediated microtubule disruption.

Discussion

In the present study, we found that thiazolidin compounds can induce G2/M arrest and disruption of mitotic spindles in non-small-cell lung and colon cancer cells. Our flow cytometry and immunoblotting analyses showed that thiazolidin compounds elicit changes in the expression of G2/M checkpoint proteins and increase the phosphorylation of MPM-2, cdc25C, and histone H3 in non-small-cell lung and colon cancer cells. Consistently, DBPT also induced G2/M-phase arrest with increased phosphorylation of MPM-2, cdc25C, and histone H3 in prostate cancer cell lines [our unpublished data]. These changes in expression and phosphorylation are consistent with mitotic arrest [19–21]. In addition, the result of immunofluorescence microscopy showed that thiazolidin compounds interfere with normal spindle formation, as do nocodazole and vinorelbine. Together, these data indicate that thiazolidin compounds induce mitotic arrest with microtubule disruption. A recent report showed that p53 activation plays a key role in the induction of G2/M phase arrest by several chemotherapeutic agents [22, 23]. However, microtubule-interacting agents such as paclitaxel, nocodazole, and vinorelbine cause G2/M phase arrest independent of p53 status, despite increasing levels of transcriptionally active p53 [24, 25]. Similarly, we found that although MMPT and DBPT up-regulate p53 expression [our unpublished data], they are still able to cause G2/M phase arrest in p53-deficient H1299 and HCT116^{-/-} cells.

Most microtubule-interacting agents induce apoptosis in cancer cells, however, the mechanisms by which they do so remain uncharacterized. Nevertheless, a previous report showed that endonucleolytic cleavage of DNA, a characteristic of apoptosis, is observed only in cells arrested in G2/M phase after treatment with epothilone and paclitaxel [26]. It has also been reported that paclitaxel-induced cell death in cancer cells might result from apoptosis directly after mitotic arrest [2, 27, 28]. Our data showed that both MMPT- and DBPT-mediated G2/M phase arrest occurred as early as 6 h and peaked around 80% after 12 h of treatment, whereas MMPT- and DBPT-induced apoptosis mainly started after 12 h of treatment [14, 15]. Thus, G2/M phase arrest by thiazolidin compounds occurred earlier than apoptosis. Moreover, DBPT-mediated G2/M phase arrest and phosphorylation

of cdc25C and histone H3 were both blocked by inhibition of JNK activation with either a specific JNK inhibitor or a dominant-negative JNK1 gene. Because DBPT-induced G2/M-phase arrest precedes apoptosis and because our previous study showed that DBPT-induced apoptosis was also blocked by inhibition of JNK activation [14], these two events (G2/M arrest and apoptosis) are very likely coupled and persistent G2/M arrest probably leads to apoptosis. Indeed, several reports have shown that Bcl-2 phosphorylation may be involved in one of the pathways of antimitotic drug-induced apoptosis [29–31]. Our study showed that Bcl-2 phosphorylation increased in H460, H460/TaxR, and H460/VinR cells after MMPT treatment. Phosphorylated Bcl-2 appeared to accumulate at the same time as the phosphorylated forms of MPM-2, cdc25C, and histone H3. However, G2/M phase arrest and apoptosis induction by thiazolidin compounds may occur independently, although both events are blocked by inhibition of JNK activation. Therefore, molecular mechanisms of apoptotic induction and G2/M arrest caused by thiazolidin compounds remain to be further delineated. Although this study has revealed the effects of thiazolidin compounds in cell cycle arrest and microtubule disruption, the direct targets of thiazolidin compounds remain to be identified. Whereas inhibition of JNK activation blocked DBPT-caused G2/M phase arrest and microtubule disruption, the mechanisms by which the JNK signaling pathway triggers G2/M arrest and microtubule disruption are not yet clear. Further characterization of the biological function of thiazolidin compounds may shed light on the roles of JNK activation in G2/M arrest and apoptosis induction.

Acknowledgements. We thank P. Lo for editorial review and K. M. Ramirez for technical assistance with flow cytometry analysis. This study was supported by National Cancer Institute grants RO1 CA 092487-01A1 (B. Fang), RO1 CA 098582-01A1 (B. Fang), Core Grant CA 16672, and a Lung SPORE (CA 70907) development award.

- 1 Correia J. J. and Lobert S. (2001) Physicochemical aspects of tubulin-interacting antimitotic drugs. *Curr. Pharm. Design* **7**: 1213–1228
- 2 Jordan M. A., Wendell K., Gardiner S., Derry W. B., Copp H. and Wilson L. (1996) Mitotic block induced in HeLa cells by low concentrations of paclitaxel (Taxol) results in abnormal mitotic exit and apoptotic cell death. *Cancer Res.* **56**: 816–825
- 3 Masters G. A., Hoffman P. C., Drinkard L. C., Samuels B. L., Golomb H. M. and Vokes E. E. (1998) A review of ifosfamide and vinorelbine in advanced non-small cell carcinoma of the lung. *Semin. Oncol.* **25**: 8–14
- 4 Ngan V. K., Bellman K., Panda D., Hill B. T., Jordan M. A. and Wilson L. (2000) Novel actions of the antitumor drugs vinflunine and vinorelbine on microtubules. *Cancer Res.* **60**: 5045–5051
- 5 Fossella F. V., Lee J. S., Murphy W. K., Lippman S. M., Calayag M., Pang A. et al. (1994) Phase II study of docetaxel for recur-

- rent or metastatic non-small-cell lung cancer. *J. Clin. Oncol.* **12**: 1238–1244
- 6 Rowinsky E. K. (1997) The development and clinical utility of the taxane class of antimicrotubule chemotherapy agents. *Annu. Rev. Med.* **48**: 353–374
- 7 Sandercock J., Parmar M. K. and Torri V. (1998) First-line chemotherapy for advanced ovarian cancer: paclitaxel, cisplatin and the evidence. *Br. J. Cancer* **78**: 1471–1478
- 8 Sorger P. K., Dobles M., Tournebise R. and Hyman A. A. (1997) Coupling cell division and cell death to microtubule dynamics. *Curr. Opin. Cell Biol.* **9**: 807–814
- 9 Drukman S. and Kavallaris M. (2002) Microtubule alterations and resistance to tubulin-binding agents. *Int. J. Oncol.* **21**: 621–628
- 10 Bosch I. and Croop J. (1996) P-glycoprotein multidrug resistance and cancer. *Biochim. Biophys. Acta* **1288**: F37–F54
- 11 Goldman B. (2003) Multidrug resistance: can new drugs help chemotherapy score against cancer? *J. Natl. Cancer Inst.* **95**: 255–257
- 12 Krishna R. and Mayer L. D. (2000) Multidrug resistance (MDR) in cancer: mechanisms, reversal using modulators of MDR and the role of MDR modulators in influencing the pharmacokinetics of anticancer drugs. *Eur. J. Pharm. Sci.* **11**: 265–283
- 13 Smyth M. J., Krasovskis E., Sutton V. R. and Johnstone R. W. (1998) The drug efflux protein, P-glycoprotein, additionally protects drug-resistant tumor cells from multiple forms of caspase-dependent apoptosis. *Proc. Natl. Acad. Sci. USA* **95**: 7024–7029
- 14 Teraishi F., Wu S., Zhang L., Guo W., Davis J. J., Dong F. et al. (2005) Identification of a novel synthetic thiazolidin compound capable of inducing c-Jun NH2-terminal kinase-dependent apoptosis in human colon cancer cells. *Cancer Res.* **65**: 6380–6387
- 15 Teraishi F., Wu S., Sasaki J., Zhang L., Zhu H. B., Davis J. J. et al. (2005) P-glycoprotein-independent apoptosis induction by a novel synthetic compound, MMPT [5-[(4-Methylphenyl)methylene]-2-(phenylamino)-4(5H)-thiazolone]. *J. Pharmacol. Exp. Ther.* **314**: 355–362
- 16 Sasaki J., Ramesh R., Chada S., Gomyo Y., Roth J. A. and Mukhopadhyay T. (2002) The anthelmintic drug mebendazole induces mitotic arrest and apoptosis by depolymerizing tubulin in non-small cell lung cancer cells. *Mol. Cancer Ther.* **1**: 1201–1209
- 17 Ling Y. H., Tornos C. and Perez-Soler R. (1998) Phosphorylation of Bcl-2 is a marker of M phase events and not a determinant of apoptosis. *J. Biol. Chem.* **273**: 18984–18991
- 18 Scatena C. D., Stewart Z. A., Mays D., Tang L. J., Keefer C. J., Leach S. D. et al. (1998) Mitotic phosphorylation of Bcl-2 during normal cell cycle progression and Taxol-induced growth arrest. *J. Biol. Chem.* **273**: 30777–30784
- 19 Vandre D. D. and Borisy G. G. (1989) Anaphase onset and dephosphorylation of mitotic phosphoproteins occur concomitantly. *J. Cell Sci.* **94**: 245–258
- 20 Giet R. and Glover D. M. (2001) *Drosophila* aurora B kinase is required for histone H3 phosphorylation and condensin recruitment during chromosome condensation and to organize the central spindle during cytokinesis. *J. Cell Biol.* **152**: 669–682
- 21 Morris M. C., Heitz A., Mery J., Heitz F. and Divita G. (2000) An essential phosphorylation-site domain of human cdc25C interacts with both 14-3-3 and cyclins. *J. Biol. Chem.* **275**: 28849–28857
- 22 Bunz F., Dutriaux A., Lengauer C., Waldman T., Zhou S., Brown J. P. et al. (1998) Requirement for p53 and p21 to sustain G2 arrest after DNA damage. *Science* **282**: 1497–1501
- 23 Taylor W. R. and Stark G. R. (2001) Regulation of the G2/M transition by p53. *Oncogene* **20**: 1803–1815
- 24 Stewart Z. A., Tang L. J. and Pietenpol J. A. (2001) Increased p53 phosphorylation after microtubule disruption is mediated in a microtubule inhibitor- and cell-specific manner. *Oncogene* **20**: 113–124
- 25 Tishler R. B., Lamppu D. M., Park S. and Price B. D. (1995) Microtubule-active drugs taxol, vinblastine, and nocodazole increase the levels of transcriptionally active p53. *Cancer Res.* **55**: 6021–6025
- 26 Wang T. H., Wang H. S. and Soong Y. K. (2000) Paclitaxel-induced cell death: where the cell cycle and apoptosis come together. *Cancer* **88**: 2619–2628
- 27 Basu A. and Haldar S. (1998) Microtubule-damaging drugs triggered bcl2 phosphorylation – requirement of phosphorylation on both serine-70 and serine-87 residues of bcl2 protein. *Int. J. Oncol.* **13**: 659–664
- 28 Woods C. M., Zhu J., McQueney P. A., Bollag D. and Lazarides E. (1995) Taxol-induced mitotic block triggers rapid onset of a p53-independent apoptotic pathway. *Mol. Med.* **1**: 506–526
- 29 Blagosklonny M. V., Giannakakou P., el Deiry W. S., Kingston D. G., Higgs P. I., Neckers L. et al. (1997) Raf-1/bcl-2 phosphorylation: a step from microtubule damage to cell death. *Cancer Res.* **57**: 130–135
- 30 Srivastava R. K., Mi Q. S., Hardwick J. M. and Longo D. L. (1999) Deletion of the loop region of Bcl-2 completely blocks paclitaxel-induced apoptosis. *Proc. Natl. Acad. Sci. USA* **96**: 3775–3780
- 31 Wang S., Wang Z., Boise L., Dent P. and Grant S. (1999) Loss of the bcl-2 phosphorylation loop domain increases resistance of human leukemia cells (U937) to paclitaxel-mediated mitochondrial dysfunction and apoptosis. *Biochem. Biophys. Res. Commun.* **259**: 67–72



To access this journal online:
<http://www.birkhauser.ch>
