

EARLY FLOWERING3 Encodes a Novel Protein That Regulates Circadian Clock Function and Flowering in Arabidopsis

Karen A. Hicks,¹ Tina M. Albertson,² and D. Ry Wagner^{3,4}

Institute of Molecular Biology, University of Oregon, Eugene, Oregon 97403

Higher plants use photoperiodic cues to regulate many aspects of development, including the transition from vegetative to floral development. The *EARLY FLOWERING3 (ELF3)* gene is required for photoperiodic flowering and normal circadian regulation in Arabidopsis. We have cloned *ELF3* by positional methods and found that it encodes a novel 695–amino acid protein that may function as a transcriptional regulator. *ELF3* transcript level is regulated in a circadian manner, as is expected of a *zeitnehmer* input pathway component. Overexpression of the *LATE ELONGATED HYPOCOTYL* gene, which has been proposed to function as a clock component, did not abolish circadian regulation of *ELF3* transcription, providing further evidence that *ELF3* is a circadian clock input pathway component.

INTRODUCTION

Both plants and animals use seasonal cues to synchronize their reproductive development with the external environment. The primary seasonal cue used by organisms is day-length, or photoperiod. Higher plants use photoperiodic cues to regulate many aspects of development, including the transition from vegetative to floral development. Arabidopsis is a facultative long-day (LD) plant, flowering earlier in LD conditions than in short-day (SD) conditions.

Mutational analysis of Arabidopsis has led to the discovery of a number of genes that are required for the photoperiodic regulation of flowering. Mutations in the *CONSTANS (CO)* and *GIGANTEA (GI)* genes result in late flowering in LD conditions but have little or no effect in SD conditions, suggesting that the wild-type genes are required for the promotion of flowering in inductive (LD) conditions. *CO* encodes a zinc finger protein that likely functions as a transcription factor, and *GI* encodes a novel putative membrane protein (Putterill et al., 1995; Fowler et al., 1999). In contrast to *co* and *gi* mutants, *flowering locus t (ft)* mutants flower late in both LD and SD conditions (Ruiz-García et al., 1997). *FT* belongs to a family of putative membrane-associated proteins that can bind hydrophobic ligands (Kardailsky et al., 1999; Kobayashi et al., 1999). One proposed molecular function of *FT* is the generation of peptide molecules that could act as transmissible signals (Kobayashi et al., 1999). *FT* appears to

function partially downstream of *CO*. Although the identification of genes that control flowering has begun to reveal a great deal about the regulation of cell fate during plant development, these genes have not revealed how circadian clock function influences the vegetative-to-floral transition.

Circadian clocks appear to be involved in photoperiodic responses in cyanobacteria, fungi, plants, and animals. Although a molecular feedback loop required for circadian regulation in *Neurospora* and animals has been described in detail, the molecular mechanisms of plant circadian clocks remain largely unknown. Genetic approaches have resulted in the isolation of a number of intriguing recessive mutations that alter the free-running period of the Arabidopsis circadian clock, including *TIMING OF CAB (TOC) 1*, *CIRCADIAN CLOCK ASSOCIATED (CCA) 1*, and *ZEITLUPE (ZTL)* (Somers et al., 1998, 2000; Green and Tobin, 1999). *ZTL* encodes a novel F-box-containing protein that may be involved in light-dependent proteolysis. *TOC1* encodes a histidine kinase similar to those of bacterial two-component signaling systems (Strayer et al., 2000). *CCA1* encodes a single MYB domain-containing transcription factor whose expression is regulated in a circadian manner. The overexpression of *CCA1*, or of the closely related gene *LATE ELONGATED HYPOCOTYL (LHY)*, results in photoperiod-insensitive early flowering and arrhythmicity of circadian clock-associated gene expression (Schaffer et al., 1998; Wang and Tobin, 1998). In addition, both *CO* and *GI* are regulated in a circadian manner.

Mutations in the *EARLY FLOWERING3 (ELF3)* locus also result in the loss of both photoperiod sensitivity and circadian regulation, making *ELF3* a candidate for linking circadian clock function with the photoperiodic induction of flowering. *elf3* mutant plants flower early and at the same developmental time in both LD and SD light conditions (Zagotta et al., 1996). The long hypocotyl phenotype of *elf3*

¹Current address: Department of Biology, Kenyon College, Gambier, OH 43022.

²Current address: Department of Pediatrics, University of Colorado Health Sciences Center, Denver, CO 80262.

³Current address: Exelixis Plant Sciences, 16160 S.W. Upper Boones Ferry Rd., Portland, OR 97224.

⁴To whom correspondence should be addressed. E-mail rwagner@exelixis.com; fax 503-670-7703.

mutant plants suggests a defect in light reception or the transduction of light signals (Zagotta et al., 1996). In addition, leaf movements and circadian clock-regulated gene expression are arrhythmic in *elf3* mutants in constant light conditions but not in constant dark conditions, suggesting that a circadian clock remains functional in the absence of wild-type *ELF3* function (Hicks et al., 1996). On the basis of these results, the *ELF3* gene product was proposed to function in a light input pathway to the circadian oscillator. The absence of *ELF3* was hypothesized to alter the coordination

of light and circadian regulatory pathways, resulting in the altered flowering time and photoperiodic insensitivity observed in *elf3* mutants. This model is supported by recent results showing that *ELF3* is required to gate light input to the circadian oscillator, altering the sensitivity of the central oscillator to light at a particular point in the circadian cycle (McWatters et al., 2000).

To elucidate the molecular mechanism of *ELF3* function, we have isolated the *ELF3* gene by positional cloning methods. *ELF3* encodes a novel protein of 695 amino acids that




Figure 1. Map-Based Cloning of *ELF3*.

(A) Physical mapping of *ELF3*. Molecular markers are shown above the line with relevant recombination distances, and the YAC contig of ~500 kb of Arabidopsis genomic DNA is shown below the line. YAC 6D6 is chimeric, with the unlinked region represented by a dashed line. cM, centimorgan.

(B) YAC end clone 10A10L was used as a starting point for identifying λ and cosmid clones. RFLP analysis localized *ELF3* to the 60-kb minimal recombinant interval shown. The closed boxes indicate the RFLPs. BAC clones spanning this region were identified, and cosmid subclones of BAC 4D15 were used for transformation rescue. Cosmid B8 complemented *elf3-1* and *elf3-3* mutant phenotypes (see Figure 2).

(C) The positions and structures of two transcription units within cosmid B8 are shown. ORF, open reading frame.




Figure 2. Complementation of *elf3* Mutants.

(A) Ten-day-old seedlings grown in SD conditions. Left to right: wild-type Columbia, *elf3-1*, transgenic *elf3-1* containing cosmid B8, and transgenic *elf3-1* containing cosmid E11.
 (B) Hypocotyl lengths of 10-day-old seedlings grown in SD conditions. Col, wild-type Columbia; Ws, Wassilewskija. Error bars indicate \pm SD ($n = 20$).
 (C) Twenty-five-day-old plants grown in LD conditions. Left to right: wild-type Columbia, *elf3-1*, transgenic *elf3-1* containing cosmid B8, and transgenic *elf3-1* containing cosmid E11.
 (D) Flowering time expressed as vegetative leaves produced before flowering. Plants were grown in either SD or LD conditions. Error bars indicate \pm SD ($n \geq 16$).

may function as a transcriptional regulator. *ELF3* mRNA level is regulated in a cyclic manner, peaking at \sim 14 to 16 hr after sunrise regardless of daylength. Continuation of cyclic expression in constant conditions shows that *ELF3* is regulated by a circadian clock. Constitutive expression of *LHY* does not abolish the circadian rhythm of *ELF3* gene expression, suggesting that additional genes, and potentially additional feedback loops, are involved in the regulation of *ELF3*.

RESULTS

Positional Cloning of *ELF3*

The *ELF3* gene was isolated on the basis of its chromosomal map position. *ELF3* was mapped initially to the mid-

dle of chromosome 2, between molecular markers g6842 and GPA1 (Zagotta et al., 1996). A yeast artificial chromosome (YAC) contig covering this region (H. Goodman, personal communication) was refined, and several restriction fragment length polymorphisms (RFLPs) in the region were identified using YAC end clones as probes (Figure 1A). Genomic cosmid clones (Figure 1B) were isolated from this region and used to identify RFLPs distinguishable from *ELF3* by recombination. This defined an \sim 60-kb region of chromosome 2 that contained the *ELF3* gene.

Bacterial artificial chromosomes (BACs) that spanned this minimal recombinant interval were identified from the Texas A&M University BAC library (Figure 1B). A cosmid mini-library of BAC 4D15 was constructed, and members of a group of cosmid clones that spanned the region of interest were transformed individually into *elf3* mutant plants. Transformation with an 11-kb cosmid clone, B8, rescued all *elf3* mutant

A

TGAAACTCACTTTGGTTTGGTTGATCCCTTTAGTCTGTTTTTCGGATTTTCCTGATTCGTTTGGTGGTGGATGATCTATCCAGTTGGTCCCTTGGGTTAAGATATTTTCATTTGGTGGT
 TTGTTTTATGAAGCTATTGTTGCGAAAGTTGGAGTCTTTTCAGTCTTTTAGTGTGAATATTAAGGAAAGGGAGACTTTTGGTGGTGAAGTTAGGTATTATTTGGGTTTGGAGAAGTTGCAAGTTGAAAGGTTGTAATTTGAGTGT
 ATGAAGAGGGGAAAGATGAGGGAAGATATGGAACCTATCTTCTCGGCTTCAATGTAATGATCCAGATAAAGAGGGCCTAGAGCTCCTCTAGAAACAAGATGGCTTTATGAGCAGCTTAGTATTCTCTCAGAGGTTTGGT
 MKRGRKDEEKILEPMPFPRRLHVNDADKGGPRAPPRNKMALYEQLSIPSRFGE 50
 Δ T
 GATCATGACAGGATGATTCCTAGTAAACAACAAGCACTTTGGTTTCATCTGGACATCTAGTCCAGCTTTGTTGGTGGAAAGAACTTATCTGTCACATCTTGATTTCTCAGCCCAACCAAGCACTGAGAAGTTGCTCC
 DHGTFMNSRSNNTSTLTVHPGFPSSQPCGVERNLSVQHLDDSSAANQATEKFFVS 100
 CAATGCTCTCATGAAAATGTGAGATCTTCGGCACAGCATGATCAGAGAAAATGGTGGAGAGGAGAGATTTTCAGTTCAGTATATTAATTAATCAAGAAGATCTCAGTCTCATGGCAGAACAAGAGTGGTATTGAGAAGAA
 QMSFMEHNVRSSEAQH DQRKMVREEEEDFAVPPVYINSRRSQSHGRTKKSGLIEKE 150
 AAACACACCCAAATGTCACCTAGCTCTCATCTCCATTCGATTTCAAGAAGTGAATCAGACAGCTCAAAAGCAAAAGCATGTTGGCTACTGTTCAAAACCTGAAGTTAGGGATCAGTCAAGGCGAATCCAGGTCAGTGGC
 KH T P M V A P S S H H S I R F Q E V N Q T G S K Q N V C L A T C S K P E V R D Q V K A N A R S G G 200
 TTTGTAATCTCTTAGATGTATCAGTACAGAGGAGATTGATCTCGAAAATCAGCATCAAGTCATGATAGAGTAAATGATTATAATGCTTCCTTGAGACAGAGCTAGAAAATCGGTTATACCGAGTCTGCCAAAACCTGTCGAAG
 F V I S L S L G D V T E E I D L E K S A S S H D R V N D Y N A S L R Q E S R N R L Y R D G G K T R L K 250
 Δ T
 GACACTGATAATGGCTGAATCTCACTTGGCAACGGAAAATCATTACAGAGGGTCAATGGCAGTCTCGAAGCATGTAATGATCTGAATCAGCAAAAGCAGAGCATGGCCCTCTCCAGCAGATAAATGAGGCAAGTGT
 D T D N G A K S H L A T E N H S Q E G H G S E E D I D N D R E Y S K S R A C A S L Q Q I N E K A S D 300
 Δ T
 GACCTTCTGATGATTCGATGGTGATTTCTATATCCAGCATAGATCTCTCCGAGTATGTTGGTGATATATAGGTCAAAAACGTTCTGGAGAGCAAGAAAGCCATTCGCAATCAAGAAGATATTGCTGTTCAACTATTGAG
 D V S D D S M V D S I S S I D V S P D D V V G I L G Q K R F W R A R K A I A N Q Q R V F A V Q L F E 350
 Δ T
 TTGCACAGACTGATTAAGTTCAAAAATTTATTCGTCATCCCGGATCTCTCTGCTGATGAGATCAGTTTCTGGAAAAGTTCTGCTAAAAGCTATCCAGTGAAGAAGCTCCTCCATCAGAAATTCGTGTAAGCCCTCTACCA
 L H R L I K V Q K L I A A S P D L L L D E I S F L G K V S A K S Y P V K K L L P S E F L V K P P L P 400
 CATGTTGCTCAACAAGGGGTGACTCGGAGAAGACTGACCAACATAAATGGAAGTCTCAGCTGAGACAGTGTGGGAGTTGTCAAACTAAGTCAATCAACAATCCAATACATGCTTTTCAACAACCCACCGCTTCA
 H V V V K Q R G D S E K T D Q H K M E S S A E N V V G R L S N Q G H H Q Q S N Y M P P A N N P P A S 450
 CCGCTCAAAATGGATATGCTTTCTCCTCAGCCTCTCTTCCAGAAATCAGCAATGGTGTATGCTCCCTCGAAGGACTGATATACAGCTCACCAGCTATGCAACACAGCGGGCATATGAGGATATAT
 P A P N G Y C F P P Q P P P S G N H Q W L I P V M S P S E G L I Y K P H P G M A H T G H Y G G Y Y 500
 GTCATATATGCTACACCAATGTAAGCCCAATATCACCCGGATGATTCACCACCTGCTGGTAAAGTCTTCCCTCATATGGAATGCCCACCAATGAAGCCATATTTCCAGCCCAACAACAACAACA
 G H Y M P T P M V M P Q Y H P G M G F P P P G N G Y F P P Y G M M P T I M N P Y C S Q Q Q Q Q Q 550
 CCCAATGACAAATGACCAAGTTGGACATCTGGAATCTCAGAACCCCAACAACAAGAGATCTGATAATGAACCTCCACAGCAACAGCAGCCAAAGCTTATCCCGGAGCAAGAGAGCGGACAGGGAGC
 P N E Q M N Q F G H P G N L O N T Q Q Q Q Q R S D N E P A P Q Q Q Q P T K S Y P R A R K S R Q G S 600
 ACAGAGCAGTCCAGTGGCCCAAGGATCTCTGATGACAGTCTTCGGCCATTCGACGCGCTGATGAGGACGCAACATCAACAATGACCTGAGCAACAGATGACACCAACCAACAGCAGCAAGCAACTGTTACTCAG
 T G S S P S P S P Q G I S G S K S F R P F A V D E D S N I N N A P E Q T M T T T T T R T T V T Q 650
 ACAACAGAGATGGGGAGGAGTACGAGGTGATAAGGTGCTACCAACAACCAAGCTCCGAGTGAAGATCTCCAGCAATTTCCAGTCAATCAAGAAGAGCTAAACGCTATGCTCTAGCTTATCTCTATGCT
 T R D G G G V T R V I K V V P H N A K L A S E N A A R I F Q S I Q E R K R Y D S S K P * 695
 GTATTGCTACTGATATGATTTTCAAAAATGAAAATTTGATGATGATATCTCAATACCAATGAAAGCTATTATGGTGAAGCTCATTATAT

B

Arabidopsis	1	MKRGRKDEEKILEPMPFPRRLHVNDADKGGPRAPPRNKMALYEQLSIPSRFGE	49
Tomato		MKRGTGEEIKVMGPMFPRRLHVNDADKGGPRAPPRNKMALYEQLSIPSRFGE	
Rice		GGGKAKKVMGPFPRRLHVNDADKGGPRAPPRNKMALYEQRFVPSRF	
Arabidopsis	300	DDVSDDSMVDSISSIDVSPDDVGIIGKRFWARAIANQQRVFAVQFELHRLIKVQKLIASPDLLLEDISFLGKVSAKS	382
Tomato		DDVSDDSMRVESISGDISPDDVGIIGLRFWKARRAIINQQRVFAVQVFEHRLIKVQKLIASPNSSLEDPAYLKGK-PLKS	
Rice		DDVSDDS-VECTIGWISPDKIVGATGTRFWKARRAIINQQRVFAVQVFEHLIKVQKLIASPE	
Arabidopsis	470	QWLIPVMS-PSEGLV-----YKPHPGM	490
Tomato		QWLIPVMS-PSEGLV-----YKPHSGP	
Soybean		QWIPVMS-SSPSEQSPTSLTWTVSCSTLRGSLKSRKSYSLGTTKPIIGP	
Rice		QWLIPVMS-PLEGLV-----YKPHVSGP	
Arabidopsis	526	YFPPGCMPTIMNPHYCSQQQQQ	549
Soybean		YFPPGCM-PVMNQATSGSAVEQV	
Maize		YFPPFSM-EAV-----SGSAVEQV	
Rice		YFPPFSI-EVNMPTAP-PVVECG	
Arabidopsis	591	PRARKSRQGSIGSSPSG-PQGI	611
Soybean		ALKERGIQGSIRSSPSMAQGI	
Maize		ASRDSEIQGSASSP-----	
Rice		ASRDSEAIQSSASSPFRFCS	
Arabidopsis	658	VTRVIKVVPHNATASENAARIEQSIQERKRYD	691
Soybean		QTQAIKVVPHNRKSAEESAARIVQSIQERKQHD	
Maize		QPRVIKVVPHNATASENAARIEQSIQERKQND	
Rice		QTNVIKVVPHNATASENAARIEQSIQERKQND	

Figure 3. DNA Coding and Amino Acid Sequence of *ELF3* and Sequence Comparison with Putative Orthologs.

(A) The DNA coding corresponding to *ELF3* cDNA clone 8.2 is shown, along with the predicted amino acid sequence of the longest open

phenotypes assayed (Figures 1B and 2). Cosmids E1, E11, and E39 did not rescue the *elf3* mutant (Figure 2 and data not shown).

A 20-kb genomic region including cosmid B8 was subcloned and sequenced. Sequence analysis revealed two potential transcription units contained within cosmid B8, neither of which showed significant similarity to genes of known function. The molecular nature of *elf3* mutant alleles was used to determine that the more distal gene contained on cosmid B8 was *ELF3*. The EcoRI fragment from the distal end of B8 identified a 1- to 2-kb deletion in *elf3-2*, which was generated by fast neutron mutagenesis (data not shown). Sequence analysis of eight additional *elf3* alleles identified further mutations within this transcription unit (Figure 3A).

ELF3 Encodes a Novel Protein

The *ELF3* gene has four exons and three introns and is predicted to encode a novel soluble protein of 695 amino acids that is particularly rich in serine, proline, and glutamine (Figure 3A). Database searches revealed strong sequence similarity between *ELF3* and predicted proteins from tomato, soybean, rice, and maize (Figure 3B) and weaker similarity between *ELF3* and an Arabidopsis protein of unknown function on chromosome 3 (GenBank accession number BAB01726).

ELF3 contains a proline-rich region between amino acids 440 and 540 that consists of ~25% proline residues. This region is shared by both the tomato expressed sequence tag and the Arabidopsis ELF3-like protein from chromosome 3. An acidic region is located from amino acids 206 to 320, and part of this region is shared by ELF3-like proteins from tomato and rice. A threonine-rich stretch of amino acids is located near the C-terminal end of the protein from amino acids 636 to 652, and three short runs of glutamine residues are found between amino acids 544 and 585. The length of the first glutamine repeat is polymorphic between the Columbia and Wassilewskija ecotypes: the Wassilewskija gene contains 16 glutamine codons, whereas the Columbia gene contains seven (data not shown). The proline-rich region, the acidic region, and the threonine/glutamine-rich region could play a role in transcriptional activation. A potential nuclear targeting signal beginning at amino acid

591 was predicted by analysis with PROSORT2 (Figure 3A), which is consistent with a role for ELF3 in transcriptional regulation. Potential nuclear targeting signals also were predicted in the ELF3-like proteins from tomato and rice. Although a nuclear targeting signal was not identified in the partial sequences available from soybean and maize ELF3-like proteins, these were predicted to be nuclear proteins on the basis of their generally basic nature. Finally, *ELF3* contains a large number of potential phosphorylation sites, suggesting that phosphorylation may be involved in ELF3 regulation.

The *elf3-1*, *elf3-3*, *elf3-4*, *elf3-5*, and *elf3-9* alleles all contain single base changes that result in premature stop codons (Figure 3A). *elf3-6* and *elf3-8* are independently derived alleles that contain identical single base changes at the exon 4 splice acceptor site. This mutation leads to the inclusion of 28 amino acids from a different reading frame before a stop codon is encountered. All of these alleles result in a strong mutant phenotype (Hicks et al., 1996; Zagotta et al., 1996) and are likely to be either null alleles or strong reduced-function alleles.

elf3-7 is a weak allele that causes early flowering and long hypocotyl phenotypes. However, *elf3-7* homozygous mutants remain sensitive to photoperiod, flowering earlier in LD than in SD (J. Reed, personal communication; K.A. Hicks, unpublished results). The *elf3-7* allele contains a single base change at the exon 1 splice donor site (Figure 3A). Analysis of *elf3-7* transcripts by reverse transcriptase-polymerase chain reaction (PCR) (data not shown) indicated the use of several cryptic splice sites that resulted in several different transcripts, all but one of which contain premature stop codons. On the basis of similarity to the strong alleles described above, we deduce that transcripts containing a premature stop codon are unlikely to confer the weak phenotype observed in the *elf3-7* allele. However, the use of one in-frame cryptic splice site was observed. The use of this splice site is predicted to result in the loss of eight amino acids, a molecular change that could be consistent with the observed partial function phenotype.

ELF3 Transcript Level Cycles Daily

As shown in Figures 4A and 4B, *ELF3* transcript level varied during a 24-hr period, with a trough in the early day and a peak in the early night. The maximal transcript level was

Figure 3. (continued).

reading frame. Amino acid numbering is shown at right, and intron positions are marked with inverted triangles. Molecular changes in eight *elf3* alleles are shown above the DNA sequence. The closed inverted triangle is a cryptic splice site used in some *elf3-7* transcripts (see text). A potential nuclear targeting signal is shown in boldface. Translation stop codon is indicated by an asterisk.

(B) Black boxes indicate identical residues, and gray boxes indicate conserved residues between ELF3 and ELF3-like predicted proteins in other plant species. Dashes indicate gaps in the sequences. Tomato, soybean, and maize ELF3-like proteins were predicted from partial sequences from expressed sequence tag clones. The rice ELF3-like protein was predicted from genomic sequence.




Figure 4. *ELF3* Transcript Levels in Wild-Type and Mutant Backgrounds.

Ten micrograms of total RNA was hybridized with antisense riboprobes for *ELF3* or *CCR2* and then subjected to RNase digestion, as described

observed ~16 hr after dawn in 12-hr-light/12-hr-dark cycles. In the same RNA samples, expression of the circadian clock-regulated gene *CCR2* was highest in the late day, as has been reported (Kreps and Simon, 1997). Diurnal cycling of the *ELF3* transcript was observed in a variety of adult shoot tissues (Figure 4C). A very low level of *ELF3* expression also was detected in seedling root tissue and followed a similar temporal pattern of expression (data not shown). Diurnal cycling of *ELF3* transcript was observed in *elf3* mutant alleles (Figure 4D and data not shown).

Phenotypic analysis suggested that *ELF3* plays a role in light signal transduction in addition to its role in regulating flowering time. In an effort to determine the potential upstream regulators of *ELF3*, we measured the level of *ELF3* transcript in known mutants that show defects in light signal transduction and/or floral induction. None of the mutations tested had a profound effect on the level or gross temporal pattern of *ELF3* expression. Expression of *ELF3* in the *cop1-6* background was somewhat altered from that of the wild type in SD conditions, with higher expression than in the wild type observed at T4 and lower expression observed at T20; this effect on *ELF3* expression by the *cop1-6* mutation was less pronounced in LD conditions.

***ELF3* Transcript Level Shows a Circadian Rhythm in Constant Conditions**

The observation of a diurnal rhythm of *ELF3* transcript level suggested that an endogenous circadian clock might regulate *ELF3* expression. One hallmark of circadian rhythms is that they continue in constant conditions. Therefore, *ELF3* expression was analyzed in wild-type seedlings entrained to 12-hr-light/12-hr-dark cycles and transferred to constant conditions. Figure 5A shows that *ELF3* transcript level con-

tinued to cycle for at least 4 days after transfer to constant light conditions, indicating that *ELF3* is regulated by a circadian clock. The rhythmic expression of a previously described circadian clock-regulated gene, *CCR2*, also is shown in Figure 5A. Although *ELF3* and *CCR2* expression was not in phase in 12-hr-light/12-hr-dark cycles (Figures 4A and 4B), expression of these genes appeared to be relatively in phase after 48 hr in constant light conditions (Figure 5A). This altered phasing of gene expression was consistently observed. A similar experiment was conducted in which wild-type seedlings were transferred to constant dark after entrainment in light/dark cycles, and *ELF3* transcript level was measured. Cycling of *ELF3* transcript level was observed during the first day of constant dark conditions; however, rhythmic expression of *ELF3* was not maintained at detectable levels (Figure 5C). Damping of circadian oscillations also has been observed for a number of other circadian clock-regulated genes, including *CAB*, *CAT2*, *CCA1*, and *CCR2* (Millar and Kay, 1991; Kreps and Simon, 1997; Zhong et al., 1997; Wang and Tobin, 1998).

ELF3* Transcript Cycling Requires *ELF3

Previously, we showed that *ELF3* is required for the circadian regulation of the *cab2* promoter and of leaf sleep movements in constant light (Hicks et al., 1996). To determine whether *ELF3* function is required for the circadian regulation of its own expression, we analyzed *ELF3* gene expression in *elf3* mutant seedlings that were first entrained to 12-hr-light/12-hr-dark cycles and then transferred to constant light. Figure 5B shows that *ELF3* transcript levels did not cycle in the *elf3-1* mutant after transfer to constant light conditions, indicating that *ELF3* is required for its own circadian regulation. The loss of rhythmicity in *CCR2* expression

Figure 4. (continued).

in Methods. β -ATPase served as an internal control, and tRNA was used as a negative control. Time 0 (T0) indicates dawn. Quantitative representation of expression data normalized to β -ATPase is shown in all panels except (A). Open and closed bars along x-axis represent light and dark photoperiods, respectively.

(A) Wild-type seedlings were grown in 12-hr-light/12-hr-dark conditions for ~1 week, and samples were collected every 4 hr. The specific probe was complementary to *ELF3* at top and to *CCR2* at bottom.

(B) Quantification of *ELF3* and *CCR2* mRNA levels shown in (A) after normalization to β -ATPase mRNA. The experiment was repeated at least three times.

(C) *ELF3* transcript level in mature wild-type plants grown in 9-hr-light/15-hr-dark SD conditions. Tissue samples were collected at T3, T13, and T21.

(D) *ELF3* transcript level in *elf3* mutant seedlings grown in 12-hr-light/12-hr-dark conditions. Samples were collected every 4 hr. The experiment was repeated five times for *elf3-1*.

(E) *ELF3* transcript level in wild-type seedlings grown in either 9-hr-light/15-hr-dark SD conditions (9L:15D) or 18-hr-light/6-hr-dark LD conditions (18L:6D). Samples were collected every 2 hr. The experiment was repeated twice.

(F) and (G) *ELF3* transcript level in wild-type and mutant seedlings grown in 9-hr-light/15-hr-dark SD conditions (F) or 18-hr-light/6-hr-dark LD conditions (G). Samples were collected at T4 and T20. Col, Columbia; Ler, Landsberg; *phy*, phytochrome; Ws, Wassilewskija.




Figure 5. *ELF3* Transcript Level Continues to Cycle in Constant Conditions.

Wild-type (**A**) and (**C**) and *elf3* mutant (**B**) seedlings were entrained under 12-hr-light/12-hr-dark conditions and then shifted to either constant light (**A**) and (**B**) or constant dark (**C**) at time 0, which corresponds to normal dawn. Samples were taken every 4 hr and analyzed by RNase protection. *ELF3* and *CCR2* transcript levels were normalized to β -ATPase. Experiments were repeated at least three times. Col, Columbia.

in the *elf3-1* mutant is shown as well. Similar results were observed for two additional mutant alleles, *elf3-2* and *elf3-3* (data not shown).

Constitutive Expression of *LHY* Does Not Abolish the Circadian Rhythm of *ELF3* Gene Expression

The rhythmic expression of two MYB-related transcription factors, *CCA1* and *LHY*, has been shown to be required for circadian regulation in Arabidopsis. Constitutive overexpression of either of these genes disrupts the circadian regulation of a number of genes and results in arrhythmic leaf sleep movements (Schaffer et al., 1998; Wang and Tobin, 1998). Therefore, we anticipated that overexpression of *LHY* also would lead to a loss of rhythmic *ELF3* gene expression. However, as shown in Figure 6A, the constitutive overexpression of *LHY* did not abolish the circadian rhythm of *ELF3* gene expression. Although the overexpression of *LHY* resulted in a reduction in *ELF3* transcript levels, gene expression was rhythmic for at least 2 days in constant light. In contrast, Figure 6B shows that the overexpression of *LHY* resulted in low levels of *CCR2* transcript that did not show a circadian rhythm, as reported previously (Schaffer et al., 1998).

DISCUSSION

ELF3 Encodes a Novel Protein

The *ELF3* gene is predicted to encode a novel protein. The *ELF3* sequence was previously identified as *pyk20* in an unrelated promoter tagging approach (Puzio et al., 1999). Searches of current databases revealed significant sequence similarity to *ELF3* only in other higher plants. *ELF3* may function as a transcription factor on the basis of the presence of sequence features that are commonly found in transcriptional regulators: a proline-rich region, an acidic region, and a threonine/glutamine-rich region (Figure 3). *ELF3*-like proteins from tomato and rice share these sequence features in part, strengthening the possibility that *ELF3* encodes a novel transcription factor. Although none of these proteins appears to contain a DNA binding domain, it is possible that *ELF3* acts to regulate transcription in concert with other factors. Given the pleiotropic nature of the *elf3* mutant phenotypes, many of which are related directly to defects in light perception, it is likely that *ELF3* functions in photoreceptor-mediated signal transduction pathways. It is now clear that some higher plant photoreceptors, such as the phytochromes, function in the nucleus; thus, the *ELF3* protein may interact with these photoreceptors to regulate gene transcription.

There are a number of examples of altered gene expression in *elf3* mutants that suggest potential downstream tar-




Figure 6. *ELF3* Transcript Level Continues to Cycle in *lhy* Mutant Seedlings.

Wild-type Landsberg (Ler) and *lhy* mutant seedlings were entrained under 12-hr-light/12-hr-dark conditions and then shifted to constant light at time 0, which corresponds to normal dawn. Samples were taken every 4 hr and analyzed by RNase protection. *ELF3* (A) and *CCR2* (B) transcript levels were normalized to β -ATPase. Scales for *ELF3* (A) differ between wild type and *lhy*. The experiment was repeated twice.

gets of transcriptional regulation by ELF3. The rhythmic expression of *Gl*, *LHY*, *CAB*, and *CCR2* is abolished in constant light conditions in *elf3* null mutants (Hicks et al., 1996; Schaffer et al., 1998; Fowler et al., 1999) (Figure 5). In addition, the expression of *Gl* was higher in *elf3* null mutants in LD cycles, particularly during the later half of the cycle, when *ELF3* expression normally peaks (Fowler et al., 1999). The question remains whether ELF3 functions to regulate the transcription of some or all of these genes directly, or in contrast, whether ELF3 affects their expression indirectly by way of the circadian clock.

ELF3 Transcript Is Regulated in a Circadian Manner

On the basis of the early flowering phenotype of *elf3* null alleles, ELF3 likely functions as a repressor of flowering. In wild-type Arabidopsis, flowering is more delayed in SD conditions than in LD conditions, presumably because of the increased level or activity of a floral repressor in SD conditions. However, no difference in *ELF3* transcript level was observed between LD and SD conditions (Figure 4E). Diurnal cycling of *ELF3* transcript occurred in both conditions, with the peak level of transcript occurring ~16 to 18 hr after dawn. Although this corresponds to the middle of the dark period in 9-hr-light/15-hr-dark SD conditions, maximal transcript levels were observed at the light-to-dark transition in 18-hr-light/6-hr-dark LD conditions.

Significantly, *elf3-1* mutants display a semidominant early flowering phenotype in LD conditions but a fully recessive early flowering phenotype in SD conditions (data not shown). This observation, together with the observation of the partial mutant phenotype in the *elf3-7* splice site mutant, suggests that the ELF3 protein itself functions in a quantitative manner. Thus, the relative abundance of functional ELF3 protein in the nucleus may serve as a mechanism to measure the length of day and to transmit this information to the circadian clock, with a high level of active ELF3 protein being found during the long night in SD-grown plants.

We propose that although the level of *ELF3* transcript was similar in LD and SD conditions, the activity of ELF3 protein may be decreased by LD conditions. This decrease could be caused at the molecular level by ELF3 interaction with, or modification by, another protein or ligand that is strictly regulated by light. This interaction or modification would occur only in the presence of light. Thus, maximum ELF3 activity would be observed in SD conditions, when the peak level of *ELF3* transcript overlaps with the dark period. One candidate for directing such interactions with the ELF3 protein is phytochrome B, which was found to interact with ELF3 in yeast two-hybrid experiments and in vitro (Liu et al., 2001). This proposed interaction mechanism could account for the quantitative aspect of photoperiodism in Arabidopsis.

Our results on the nature of *ELF3* transcriptional regulation, namely, that *ELF3* transcripts accumulate in a circadian manner and that this accumulation is dependent on the presence of functional ELF3 protein in constant light conditions (Figure 5), strengthen the conclusion that ELF3 functions within a *zeitnehmer* ("time-taker") feedback loop (McWatters et al., 2000) and suggest that the input pathway is regulated rhythmically by feedback from the central oscillator to *ELF3* transcription.

Constitutive Expression of *LHY* Does Not Abolish the Circadian Rhythm of *ELF3* Gene Expression

The rhythmic expression of central oscillator components is required for circadian function in systems such as Neurospora

and animals, in which the molecular feedback loop has been well characterized (Dunlap, 1999). Therefore, the constitutive expression of an *Arabidopsis* oscillator component is expected to result in a loss of rhythmicity. Two MYB-related transcription factors, *CCA1* and *LHY*, have been proposed as oscillator components on the basis of the disruption of circadian regulation when either factor is expressed constitutively (Schaffer et al., 1998; Wang and Tobin, 1998). However, the continued rhythmic expression of *ELF3* in the presence of constitutive *LHY* expression suggests that *LHY* is not a component of the *ELF3*-regulating central oscillator. It remains possible that multiple clocks function within *Arabidopsis* and that the *ELF3*-related clock is separate from the clock that requires *LHY* function.

ELF3 Gene Function in Flowering Plants

Database searches for gene sequences related to *ELF3* have revealed significant homology only with sequences from other higher plant species. This leaves open the possibility that the *ELF3* protein, and the homologous proteins of other plant species, function in a regulatory hierarchy that has evolved specifically in the plant kingdom. If this is the case, then plants may have developed novel molecular mechanisms for the regulation of input signals to the circadian clock, or there may exist a type of circadian clock that is specific to plants.

One possible explanation for such a novel system in multicellular plants is that higher plants are uniquely constrained within their environment. For example, some species of flowering plants quite accurately measure slight changes in daylength to promote or repress critical developmental events such as the initiation of flowering. The precise seasonal timing of such events is crucial for plant reproduction and population dispersal. It will be of great interest to determine whether the putative *ELF3* homologs identified in other flowering plant species play similar roles in regulating circadian clock function and aspects of plant development such as the initiation of flowering. Although there is clear evidence that *ELF3* functions to repress floral initiation in the LD plant *Arabidopsis*, what function does the tomato homolog of *ELF3* have in day-neutral varieties of that species? To investigate questions such as this, it will be essential to elucidate the molecular function of the *ELF3* protein in *Arabidopsis* and to determine if the homologous proteins in species such as tomato have equivalent functions. In addition, it will be critical to determine whether the components of the circadian clock and the pathways that regulate the initiation of flowering are conserved among these higher plant species. It remains a possibility that not only are circadian components and regulatory mechanisms divergent among bacteria, fungi, plants, and animals but that even within the plant kingdom multiple molecular mechanisms exist that control circadian clock function and, in turn, related developmental processes.

METHODS

Plant Material

elf3-1, *elf3-2*, *elf3-3*, and *elf3-4* have been described (Hicks et al., 1996; Zagotta et al., 1996). *elf3-5*, *elf3-7*, *elf3-8*, and *elf3-9* were isolated in the *Arabidopsis thaliana* Columbia ecotype by using ethyl methanesulfonate mutagenesis. *elf3-7* and *elf3-9* were kindly provided by Jason Reed (University of North Carolina, Chapel Hill), and *elf3-8* was kindly provided by Andrew Millar (University of Warwick, Coventry, UK). *elf3-6* was isolated in the Wassilewskija ecotype from the publicly available Feldmann T-DNA insertion populations (*Arabidopsis* Biological Resource Center, Columbus, OH).

pif was kindly provided by José Martínez-Zapater (Instituto Valenciano de Investigaciones Agrarias, Valencia, Spain). *aux1*, *hy4* (2.23N), *hy5* (Ci88), *co*, *det1-1*, and *cop1-6* were obtained from the *Arabidopsis* Biological Resource Center. *phyA-201* and *phyB-5* were kindly provided by Joanne Chory (Salk Institute for Biological Studies, La Jolla, CA) (Nagatani et al., 1993; Reed et al., 1994). *cry2-1*, *cry2-2*, and *CRY2OX* were kindly provided by Chentao Lin (University of California, Los Angeles) (Guo et al., 1998).

Cloning and Sequencing of *ELF3*

Meiotic recombination events were generated near *ELF3* by screening for recombination events between *elf3-1* and the flanking markers *erecta* and *pif*. Crosses were analyzed between *elf3-1* (Columbia ecotype) and *erecta aux1* (Landsberg ecotype) or *erecta pif1* (Landsberg ecotype). Restriction fragment length polymorphism (RFLP) mapping was performed by DNA gel blot analysis according to standard protocols.

The yeast artificial chromosome (YAC) contig spanning the region between g6842 and *GPA1* was provided by Howard Goodman (Massachusetts General Hospital, Boston). YAC end clones were obtained by plasmid rescue or inverse polymerase chain reaction (PCR). End clones were used to refine the contig by DNA gel blot hybridization and to map RFLPs between the Columbia and Landsberg ecotypes.

Genomic clones were isolated from the Mulligan and Davis λ genomic library, the Olszewski cosmid library (Olszewski et al., 1988), the Texas A&M University (TAMU) bacterial artificial chromosome (BAC) library (Choi et al., 1995), and a cosmid library constructed from TAMU BAC 4D15 by using the pOCA28 vector. Cosmid clones were introduced into *Agrobacterium tumefaciens* strain GV3101 by electroporation, and *elf3-1* and *elf3-3* were transformed by vacuum infiltration (Bechtold et al., 1993; Bent et al., 1994).

DNA fragments were subcloned into the pBluescript SK⁻ vector (Stratagene) for sequence analysis. The *ELF3* gene was amplified by PCR from genomic DNA of mutant alleles and subcloned into pBluescript SK⁻ or pCR2.1 (Invitrogen, Carlsbad, CA) for sequence analysis. Mutations were confirmed by direct sequencing of a pool of at least 10 PCR products. *elf3-7* transcripts were isolated with reverse transcriptase-PCR by using the cDNA cycle kit (Invitrogen), and PCR products were subcloned as described above.

RNA Analysis

Tissue samples were harvested as described above, and total RNA was extracted essentially as described by Nagy et al. (1988). RNA

used for the experiment shown in Figure 6 was kindly provided by Isabelle Carré (University of Warwick). RNase protection assays were performed according to standard protocols. Templates were prepared with PCR amplification from DNA subclones by using one primer complementary to the vector sequence and a second primer complementary to the insert. The *ELF3* primer sequence was 5'-TGG-CACCTAGCTCTCATC-3', the *CCR2* primer sequence was 5'-CGC-TTGTATGCTTCTACTTGG-3', and the β -ATPase primer sequence was 5'-TTCCTTGAGAGCTACGAGATG-3'. The *CCR2* plasmid was provided by J. Kreps (Torrey Mesa Research Institute, La Jolla, CA). β -ATPase was obtained as an expressed sequence tag clone (GenBank accession number N96685) from the Arabidopsis Biological Resource Center. The intensity of protected fragments was measured by PhosphorImager analysis (Molecular Dynamics, Sunnyvale, CA).

ACKNOWLEDGMENTS

We are grateful to I. Carré for providing RNA samples and to R. Formosa and H. Foss for technical assistance. This research was supported by National Science Foundation Grant MCB-9808208 and by U.S.–Israel Binational Agricultural Research and Development Agency Grant US-2964-97 to D.R.W. and by a National Institutes of Health postdoctoral fellowship to K.A.H.

REFERENCES

- Bechtold, N., Ellis, J., and Pelletier, G. (1993). *In planta* Agrobacterium mediated gene transfer by infiltration of adult *Arabidopsis thaliana* plants. C. R. Acad. Sci. Paris Life Sci. **316**, 1194–1199.
- Bent, A.F., Kunkel, B.N., Dahlbeck, D., Brown, K.L., Schmidt, R., Giraudat, J., Leung, J., and Staskawicz, B.J. (1994). *RPS2* of *Arabidopsis thaliana*: A leucine-rich repeat class of plant disease resistance genes. *Science* **265**, 1856–1860.
- Choi, S., Creelman, R.A., Mullet, J.E., and Wing, R.A. (1995). Construction and characterization of a bacterial artificial chromosome library of *Arabidopsis thaliana*. *Weeds World* **2**, 17–20.
- Dunlap, J.C. (1999). Molecular bases for circadian clocks. *Cell* **96**, 271–290.
- Fowler, S., Lee, K., Onouchi, H., Samach, A., Richardson, K., Morris, B., Coupland, G., and Putterill, J. (1999). GIGANTEA: A circadian clock-controlled gene that regulates photoperiodic flowering in *Arabidopsis* and encodes a protein with several possible membrane-spanning domains. *EMBO J.* **18**, 4679–4688.
- Green, R.M., and Tobin, E.M. (1999). Loss of the circadian clock-associated protein 1 in *Arabidopsis* results in altered clock-regulated gene expression. *Proc. Natl. Acad. Sci. USA* **96**, 4176–4179.
- Guo, H., Yang, H., Mockler, T.C., and Lin, C. (1998). Regulation of flowering time by *Arabidopsis* photoreceptors. *Science* **279**, 1360–1363.
- Hicks, K.A., Millar, A.J., Carré, I.A., Somers, D.E., Straume, M., Meeks-Wagner, D.R., and Kay, S.A. (1996). Conditional circadian dysfunction of the *Arabidopsis early-flowering 3* mutant. *Science* **274**, 790–792.
- Kardailsky, I., Shukla, V.K., Ahn, J.H., Dagenais, N., Christensen, S.K., Nguyen, J.T., Chory, J., Harrison, M.J., and Weigel, D. (1999). Activation tagging of the floral inducer *FT*. *Science* **286**, 1962–1965.
- Kobayashi, Y., Kaya, H., Goto, K., and Iwabuchi, M. (1999). A pair of related genes with antagonistic roles in mediating flowering signals. *Science* **286**, 1960–1962.
- Kreps, J.A., and Simon, A.E. (1997). Environmental and genetic effects on circadian clock-regulated gene expression in *Arabidopsis*. *Plant Cell* **9**, 297–304.
- Liu, X.L., Covington, M.F., Fankhauser, C., Chory, J., and Wagner, D.R. (2001). *ELF3* encodes a circadian clock-regulated nuclear protein that functions in an *Arabidopsis PHYB* signal transduction pathway. *Plant Cell* **13**, 1293–1304.
- McWatters, H.G., Bastow, R.M., Hall, A., and Millar, A.J. (2000). The *ELF3 zeitnehmer* regulates light signalling to the circadian clock. *Nature* **408**, 716–720.
- Millar, A.J., and Kay, S.A. (1991). Circadian control of *cab* gene transcription and mRNA accumulation in *Arabidopsis*. *Plant Cell* **3**, 541–550.
- Nagatani, A., Reed, J.W., and Chory, J. (1993). Isolation and initial characterization of *Arabidopsis* mutants that are deficient in phytochrome A. *Plant Physiol.* **102**, 269–277.
- Nagy, F., Kay, S.A., and Chua, N.-H. (1988). Analysis of gene expression in transgenic plants. In *Plant Molecular Biology Manual*, S. Gelvin and R. Schilperoort, eds (Dordrecht, The Netherlands: Kluwer Academic Publishers), pp. 1–29.
- Olszewski, N., Martin, F., and Ausubel, F. (1988). Specialized binary vector for plant transformation: Expression of the *Arabidopsis thaliana AHAS* gene in *Nicotiana tabacum*. *Nucleic Acids Res.* **16**, 10765–10782.
- Putterill, J., Robson, F., Lee, K., Simon, R., and Coupland, G. (1995). The *CONSTANS* gene of *Arabidopsis* promotes flowering and encodes a protein showing similarities to zinc finger transcription factors. *Cell* **80**, 847–857.
- Puzio, P.S., Lausen, J., Almeida-Engler, J., Cai, D., Gheysen, G., and Grundler, F.M.W. (1999). Isolation of a gene from *Arabidopsis thaliana* related to nematode feeding structures. *Gene* **239**, 163–172.
- Reed, J.W., Nagatani, A., Elich, T.D., Fagan, M., and Chory, J. (1994). Phytochrome A and phytochrome B have overlapping but distinct functions in *Arabidopsis* development. *Plant Physiol.* **104**, 1139–1149.
- Ruiz-García, L., Madueno, F., Wilkinson, M., Haughn, G., Salinas, J., and Martínez-Zapater, J.M. (1997). Different roles of flowering-time genes in the activation of floral initiation genes in *Arabidopsis*. *Plant Cell* **9**, 1921–1934.
- Schaffer, R., Ramsay, N., Samach, A., Corden, S., Putterill, J., Carré, I.A., and Coupland, G. (1998). The *late elongated hypocotyl* mutation of *Arabidopsis* disrupts circadian rhythms and the photoperiodic control of flowering. *Cell* **93**, 1219–1229.

Somers, D.E., Webb, A.A.R., Pearson, M., and Kay, S.A. (1998). The short-period mutant, *toc1-1*, alters circadian clock regulation of multiple outputs throughout development in *Arabidopsis thaliana*. *Development* **125**, 485–494.

Somers, D.E., Schultz, T.F., Milnamow, M., and Kay, S.A. (2000). *ZEITLUPE* encodes a novel clock-associated PAS protein from *Arabidopsis*. *Cell* **101**, 319–329.

Strayer, C., Oyama, T., Schultz, T.F., Raman, R., Somers, D.E., Mas, P., Panda, S., Kreps, J.A., and Kay, S.A. (2000). Cloning of the *Arabidopsis* clock gene *TOC1*, an autoregulatory response regulator homolog. *Science* **289**, 768–771.

Wang, Z.-Y., and Tobin, E.M. (1998). Constitutive expression of the *CIRCADIAN CLOCK ASSOCIATED 1 (CCA1)* gene disrupts circadian rhythms and suppresses its own expression. *Cell* **93**, 1207–1217.

Zagotta, M.T., Hicks, K.A., Jacobs, C.I., Young, J.C., Hangarter, R.P., and Meeks-Wagner, D.R. (1996). The *Arabidopsis* *ELF3* gene regulates vegetative photomorphogenesis and the photoperiodic induction of flowering. *Plant J.* **10**, 691–702.

Zhong, H., Resnick, A., Straume, M., and McClung, C.R. (1997). Effects of synergistic signaling by phytochrome A and cryptochrome1 on circadian clock-regulated catalase expression. *Plant Cell* **9**, 947–955.