

A Ca²⁺-Binding Domain in RyR1 that Interacts with the Calmodulin Binding Site and Modulates Channel Activity

Liangwen Xiong, Jia-Zheng Zhang, Rong He, and Susan L. Hamilton

Department of Molecular Physiology and Biophysics, Baylor College of Medicine, Houston, Texas

ABSTRACT A fragment of RyR1 (amino acids 4064–4210) is predicted to fold to at least one lobe of calmodulin and to bind Ca²⁺. This fragment of RyR1 (R4064–4210) was subcloned, expressed, refolded, and purified. Consistent with the predicted folding pattern, R4064–4210 was found to bind two molecules of Ca²⁺ and undergo a structural change upon binding Ca²⁺ that exposes hydrophobic amino acids. R4064–4210 also binds to RyR1, the L-type Ca²⁺ channel (Cav_{1.1}), and several synthetic calmodulin binding peptides. Both R4064–4210 and a peptide representing the calmodulin-binding region of RyR1 (R3614–3643) alter the Ca²⁺ dependence of (³H)ryanodine binding to RyR1, suggesting that they may both be interfering with an intramolecular interaction between amino acids 4064–4210 and amino acids 3614–3643 in the native RyR1 to alter or regulate the response of the channel to changes in Ca²⁺ concentration. The finding that a domain within RyR1 binds Ca²⁺ and interacts with calmodulin-binding motifs may provide insights into the mechanism for calcium- and calmodulin-dependent regulation of this channel and perhaps for its regulation by the L-type Ca²⁺ channel.

INTRODUCTION

Ca²⁺ release channels, also known as ryanodine receptors (RyRs), regulate the release of Ca²⁺ from sarcoplasmic reticulum (SR) stores. The released Ca²⁺ triggers muscle contraction and activates a cascade of Ca²⁺-dependent signal transduction pathways. Ca²⁺ and calmodulin (CaM) are important as *in vivo* modulators of RyRs. RyR1 activity displays a bell-shaped dependence on Ca²⁺ concentration, with enhancement of channel activity in the low μ M range, but inhibition of the channel in the high μ M to mM range (1,2). Several motifs within the primary amino-acid sequence of RyRs have been suggested to resemble Ca²⁺-binding E-F hand motifs in Ca²⁺-binding proteins (3,4). Putative E-F hands have been identified in RyR1 at amino acids 4253–4264, 4407–4416, and 4489–4499 (5), in RyR2 at amino acids 1136–1347 and 2010–2021 (6), and in RyR3 at amino acids 3934–3945 (7). Fragments of RyR1 containing some of these E-F hand motifs have been shown to bind Ca²⁺ (8–10). The low-affinity inhibitory Ca²⁺ binding site was suggested to be within the negatively charged region between residues 1872–1923 (11,12); however, the replacement of this glutamate-rich sequence with the corresponding, less-acidic sequence from RyR2 did not change low-affinity Ca²⁺ inactivation of RyR1 (12). Du and MacLennan suggested that the Ca²⁺ binding sites for both activation and inactivation of RyR1 are located at the C-terminus between residues 3726 and 5037 (13). This region has three potential E-F hands (5) and several other possible Ca²⁺-binding sequences (14–16). Mutation of some amino

acids in this region alters Ca²⁺ binding and Ca²⁺ regulation of RyR1 (17). To determine if any of these putative E-F hands are involved in Ca²⁺ regulation of the channel, Fessenden et al. (18) created mutant RyRs in which the putative E-F hands were scrambled either singly or in combination with other E-F hands. These investigators found that, in intact myotubes, the mutations did not affect functional responses to depolarization, caffeine, or 4-chloro-m-cresol (4-CmC) (18). However, (³H)ryanodine binding and its Ca²⁺ dependence were altered by the E-F-1 mutations and abolished by the E-F-2 mutations, suggesting an involvement of these E-F hands in Ca²⁺ regulation of the channel. Why would this be different in the myotubes? One possibility is that a Ca²⁺-binding protein such as CaM, which can itself both activate and inhibit the channel in a Ca²⁺-dependent manner, can override the need for Ca²⁺ binding directly to the channel in the myotubes. This unanswered question emphasizes the need for further investigation.

CaM can function as either an activator or an inhibitor of RyR1, depending on whether it has Ca²⁺ bound (19). At sub- μ M Ca²⁺ concentrations, CaM is an activator of RyR1, but at higher Ca²⁺ concentrations, it becomes an inhibitor (19–21). Both Ca²⁺-free and Ca²⁺-bound CaMs bind to RyR1 at a site close to amino acids 3614–3643 (22,23). This sequence is likely to represent the binding site for the C-lobe of CaM (24), whereas the N-lobe may bind to an adjacent subunit within the RyR1 tetramer between amino acids 1975 and 1999 (25). We previously have shown that a carboxy-terminal tail fragment of the skeletal L-type channel (Ca_v1.1) α_1 -subunit, which contains both Ca²⁺ and CaM binding sites, binds to RyR1. This interaction is blocked by Ca²⁺CaM (26). We have also shown that a peptide representing the CaM binding site on RyR1 (amino acids 3614–3643) binds to both the Ca_v1.1 and an expressed fragment of the carboxy-terminal tail of its α_1 -subunit (amino acids 1393–1527) (26).

Submitted May 11, 2005, and accepted for publication September 27, 2005.

Address reprint requests to Susan L. Hamilton, Dept. of Molecular Physiology and Biophysics, Baylor College of Medicine, 1 Baylor Plaza, Houston, TX 77030. Tel.: 713-798-3894; Fax: 713-798-5441; E-mail: susanh@bcm.tmc.edu.

© 2006 by the Biophysical Society

0006-3495/06/01/173/10 \$2.00

doi: 10.1529/biophysj.105.066092

The interaction of R3614–3643 with the $\text{Ca}_v1.1$ is also blocked by Ca^{2+} CaM (26). Three sequences on the carboxy-terminal tail of the cardiac L-type Ca^{2+} channel ($\text{Ca}_v1.2$), designated *A* (1609–1628, numbering of the human cardiac channel), *C* (1627–1652), and *IQ* (1665–1685) have been implicated in CaM binding (27–30). Synthetic peptides matching these sequences or their skeletal muscle counterparts do not interact with R3614–3643 (unpublished observation), suggesting that there is another site within $\text{Ca}_v1.1$ that can bind to R3614–3643.

Since the CaM binding domains on $\text{Ca}_v1.1$ and RyR1 do not directly interact, we began to search for other interaction sites. Using 3D-PSSM (<http://www.sbg.bio.ic.ac.uk>), we identified a sequence within RyR1 (amino acids 4064–4210) and one within the carboxyterminal tail of the $\text{Ca}_v1.1$ α_1 -subunit (amino acids 1395–1540) predicted to fold like CaM and bind Ca^{2+} . One possibility is that these CaM-like domains can interact with the CaM binding sites of these proteins and CaM would compete for this interaction. To assess this possibility, we expressed the CaM-like domain of RyR1 and examined its ability to bind Ca^{2+} and to interact with RYR1, the L-type channel, and several CaM-binding peptides. (See Table 1 for a list of all peptides used.)

EXPERIMENTAL PROCEDURES

Materials

The peptides were synthesized in the core facility at Baylor College of Medicine (Houston, TX) under the direction of Dr. Richard Cook. SR membranes were prepared from rabbit hind leg and backstrap skeletal muscle as previously described (31). Protein concentrations were determined by the Lowry method (32) using bovine serum albumin as the standard. (^3H)ryanodine (70–80 Ci/mmol) was purchased from Du Pont New England Nuclear (Boston, MA); unlabeled ryanodine was purchased from Calbiochem (La Jolla, CA).

Creation of the construct for expressing R4064–4210

To subclone the fragment of RyR1 cDNA coding for the region of RyR1 predicted to fold like CaM, we used PCR primers containing *NdeI* and *HindIII* restriction sites, with a stop-codon in the reverse primer. PCR

reactions amplified the desired DNA fragment by selective priming of the full-length rabbit skeletal RyR1 cDNA. The amplified PCR products were agarose gel-purified and then ligated into pCR-Blunt vector (Invitrogen, San Diego, CA). The ligates were transformed into One-Shot TOP-10 Competent Cells (Invitrogen) for amplification. After confirmation by DNA sequencing, the desired DNA fragments were excised using the enzymes whose restriction sites were pre-incorporated into primers. The released DNA fragment was subcloned into pET23a(+) or pET28a(+) vectors (Novagen, Madison, WI) between *NdeI* and *HindIII* sites. The subcloned products were transformed into DH5 α competent cells (Invitrogen) to amplify. DNA sequencing was performed again for verification of the desired DNA sequences in the expression vectors.

Expression, purification, and refolding of R4064–4210

The expression vectors containing the DNA construct for R4064–4210 were transformed into BL21(DE3) competent cells (Novagen). The fragment R4064–4210 was found almost exclusively in inclusion bodies when expressed either with or without a 20-amino acid His tag (MGSSHHHHHSSGLVPRGSH) at the N-terminus of the fragment (in pET28a(+) or pET23a(+) vector). To purify this fragment from inclusion bodies, the cytoplasmic proteins were first extracted using Bacterial Protein Extraction Reagent (B-PER; Pierce Biotechnology, Rockford, IL) and then the bacterial membrane proteins were extracted by B-PER Bacterial Protein Extraction Reagent with lysozyme and nuclease. The remaining insoluble material, which is greatly enriched in R4064–4210, was washed with inclusion body wash buffer (50 mM Tris-HCl pH 7.4, 1% Triton or CHAPS, 5 mM DTT, and 5 mM EDTA). R4064–4210 was extracted with 50 mM Tris-HCl pH 7.4 and 2 M urea. The His-tagged protein was purified with a chelating sepharose column (Amersham Pharmacia Biotech, Piscataway, NJ) with a buffer containing 2 M urea. The chelating sepharose-purified R4064–4210 was dialyzed against 20-mM Tris-HCl pH 7.4 in a 10,000 MWCO cassette (Pierce). The dialysis buffer was changed twice to remove the urea and refold the protein. The urea-extracted untagged protein was refolded in a manner similar to the His-tagged protein and purified by anion exchange (HiTrap Q HP) and phenyl sepharose chromatography (Amersham Pharmacia Biotech).

Circular dichroism spectroscopy

To demonstrate the refolding of R4064–4210, circular dichroism (CD) spectroscopy was used to analyze its secondary structure. Untagged R4064–4210 (0.1 mg/ml) was incubated in 5 mM Tris-HCl pH7.9, 1 mM EGTA, or 1 mM Ca^{2+} for 10 min at room temperature. CD spectra were recorded on an Aviv CD instrument, Model #62A DS (Aviv, Lake Wood, NJ) from 188 to 250 nm with 1 nm/step, using a 2-mm quartz cell. Data were processed by subtracting the CD signal of buffer/additives and by averaging the data obtained from three independent experiments.

Analysis of ^{45}Ca binding to R4064–4210 by equilibrium dialysis

R4064–4210 (15 μM , 100 μl) in 30 mM HEPES pH 7.0, 0.2 mM PMSF, and 5 mM β -mercaptoethanol was injected into one of two chambers of an equilibrium dialysis cell (Bel-Art Products, Pequannock, NJ), separated by a 3500 MWCO membrane. The same amount of the buffer containing different concentrations of $^{45}\text{Ca}^{2+}$ (5–200 μM , 1 $\mu\text{Ci}/\mu\text{mol}$) was injected into the other cell. The equilibrium dialysis cells were incubated at 20°C overnight with rotation, after which 50 μl was removed from each chamber for analysis of $^{45}\text{Ca}^{2+}$ by liquid scintillation counting. $^{45}\text{Ca}^{2+}$ bound to R4064–4210 was calculated from the difference in $^{45}\text{Ca}^{2+}$ radioactivity in the two chambers. The data were fit with a four-parameter Hill equation in SigmaPlot (SPSS, Chicago, IL).

TABLE 1 Peptides used in this study

R4064–4210	MFLKLDK IVGSEAFQDYVTDPRGLISK KDFQKAMDSQKQF TGPEIQFLLSCEADENEMINFEEFAN RFQEPARDIGFNVAV LLTN LSEHVPHDPRLRNFLLELAESI LEYFRPYLGRIEIMGAS RRRIERYF EISETNRAQWEMPQV
R3614–3627	KSKKAVVHKLLSKQ
R3614–3634	KSKKAVVHKLLSKQRRRAVVA
R3614–3643	KSKKAVVHKLLSKQRRRAVVACFRMTPLYN
R3625–3644	SKQRRRAVVACFRMTPLYN
Skeletal IQ	KFYATFLIQEHFRKFMKRQEE
C-peptide	EQANEELRAIIKKIWKRTSMKLLDQV
CaMKII peptide	MHRQEAVDCLKKFNARRKLG

ANSA fluorescence analysis probes conformational changes of R4064–4210 induced by Ca^{2+}

To assess the ability of the RyR1 fragment to undergo Ca^{2+} -induced conformational changes, we used 8-anilino-1-naphthalenesulfonic acid ammonium (ANSA) (Sigma-Aldrich, St. Louis, MO) fluorescence analysis. ANSA (1 μM) was incubated in 30 mM MOPS (pH 7.2) and either 5 mM EGTA (low Ca^{2+}) or 2 mM Ca^{2+} (high Ca^{2+}) for 5 min at room temperature in the presence or absence of untagged R4064–4210 (0.4 μM). Fluorescence emission spectra from 430 to 600 nm were collected with 370-nm excitation using an ISS PC1 Photon Counting Spectrofluorometer (ISS, Champaign, IL), with a 0.5-mm slit for excitation and 2-mm slit for emission, 1 nm/step. Relative fluorescence was corrected for buffer contribution. All experiments were repeated at least three times.

R4064–4210 binding to CaM-binding peptides by ANSA fluorescence analysis

R4064–4210 (0.5 μM) and ANSA (2 μM) were incubated in 30 mM MOPS (pH 7.2) containing 5 mM EGTA, 50 μM Ca^{2+} , 500 μM Ca^{2+} , or 5 mM Ca^{2+} for 5 min at room temperature; 5 μM of different peptides (R3614–3643, C-peptide (27), CaM kinase II Inhibitor (Calbiochem) or the IQ-peptide (30)) were then added. ANSA fluorescence emission data at 470 nm were collected on an SLM 8000C Spectrofluorometer (SLM Instruments, Urbana, IL) with a 370-nm excitation, 8-nm bandpass for excitation, and 16-nm bandpass for emission. Data were processed by subtracting the background of buffer and/or additives where appropriate and the data were fit with a four-parameter Hill equation in SigmaPlot (SPSS).

Identification of the amino acids interacting with R4064–4210

To determine which amino acids in R3614–3643 were involved in R4064–4210 binding, we used overlapping peptide fragments from amino acids 3614–3644 to interact with R4064–4210 in the presence of ANSA. R4064–4210 (0.5 μM) and ANSA (2 μM) were incubated in 30 mM MOPS (pH 7.2) containing 5 mM EGTA (low Ca^{2+}) or 500 μM Ca^{2+} (high Ca^{2+}) for 5 min at room temperature. Peptides (R3614–3627, R3614–3634, R3614–3643, and R3625–3644) were then added and the ANSA fluorescence data were collected on an SLM 8000C Spectrofluorometer (SLM Instruments).

The effect of R4064–4210 and synthetic peptides on (^3H)ryanodine binding

SR membranes (10 μg) were incubated with (^3H)ryanodine (5 nM) for 16 h at room temperature in 100 mM NaCl, 30 mM MOPS (pH 7.2), 100 $\mu\text{g}/\text{ml}$ BSA, and 0.1% CHAPS, containing increasing CaCl_2 with 10 μM different proteins or peptides. Bound radiolabel was separated from free by filtration through Whatman GF/F filters with 5×3 ml washes of ice cold buffer (Whatman, Brentford, Middlesex, UK). Nonspecific binding was determined in the presence of 10- μM unlabeled ryanodine.

Assessing the interaction of R4064–4210 with $\text{Ca}_v1.1$ by pulldown assays

His-tagged R4064–4210 (3 nM) or a synthetic His tag alone were incubated with 300 μl of a Ni charged chelating sepharose bead slurry in binding buffer consisting of 50 mM MOPS (pH 7.4), 20 mM imidazole, 1 mM CaCl_2 (designated high Ca^{2+} buffer) or 2 mM EGTA (designated low Ca^{2+} buffer) for 30 min. The IQ-peptide (3 nM, sequence: KFYATFLIQEHFRKFMKR-QEE) from the C-terminus of the skeletal L-type Ca^{2+} channel ($\text{Ca}_v1.1$, α_1 -

subunit) or digitonin-solubilized T-tubule membranes (24 μg) were added to this bead slurry. After 1-h incubation, the beads were washed twice with 300- μl binding buffer and proteins were eluted with SDS sample buffer for electrophoresis. The pulldown of the IQ-peptide by R4064–4210 was assessed by Coomassie staining of the SDS gel (in duplicate and repeated three times). $\text{Ca}_v1.1$ in the pulldown was determined by Western blotting with an anti- $\text{Ca}_v1.1$ α_{1s} antibody, MA3-920 (Affinity BioReagents, Golden, CO), and Alexa Fluor 680 (Molecular Probes, Eugene, OR) as secondary antibody (in triplicate and repeated twice). An Odyssey Infrared Imaging System (LI-COR, Lincoln, NE) was used to analyze the Western blots. To verify the specific interaction of R4064–4210 with $\text{Ca}_v1.1$, parallel Western blotting experiments were conducted, but with an anti-sodium/potassium ATPase α -1 antibody, MA3-929 (Affinity BioReagents).

RESULTS

R4064–4210 is predicted by 3D-PSSM to fold similar to calmodulin

We have previously shown that a synthetic peptide matching the CaM binding site on RyR1 bound to $\text{Ca}_v1.1$, and, conversely, that a CaM binding peptide from $\text{Ca}_v1.1$ bound to RyR1 (26). Since the CaM binding peptides from these two channels do not themselves interact (unpublished observation), we began to look for sequences within RyR1 that could bind to the carboxy-terminal tail of the $\text{Ca}_v1.1$ α_1 -subunit. A region of $\text{Ca}_v1.1$ (amino acids 1395–1540) that contains the putative E-F hands was previously predicted by 3D-PSSM (<http://www.sbg.bio.ic.ac.uk>) to fold like CaM and was shown to interact with CaM binding peptides from RyR1 and $\text{Ca}_v1.1$ (26). The CaM binding peptide from RyR1 has been shown to bind both Ca^{2+}CaM and apoCaM (22,23). We found a region of RyR1 (amino acids 4064–4210) that was also predicted to fold in a manner similar to CaM (>95% confidence level, 21% sequence identity, Fig. 1 A).

Extraction and refolding of R4064–4210 from inclusion bodies after expression in *Escherichia coli*

A fragment of RyR1 representing this putative CaM-like domain and designated R4064–4210 was expressed in BL21(DE3) cells. The protocol used for the expression generated 30–50 mg of R4064–4210 per liter of culture. The R4064–4210 was found almost exclusively in inclusion bodies but could be solubilized in 2 M urea (*lane 5* of Fig. 1 B). The urea-solubilized R4064–4210 was refolded by dialysis into buffer without urea and was soluble at concentrations as high as 10 mg/ml (*lane 7* of Fig. 1 B).

Assessing the α -helical content of R4064–4210

R4064–4210 appeared to be both homogenous and structured as assessed by its CD signal at both low (*solid line* of Fig. 1 C) and high Ca^{2+} concentrations (*dashed line* of Fig. 1 C). There is a strong positive CD maximum at 192 nm and

molecules of $^{45}\text{Ca}^{2+}$ bound per mol of R4064–4210 (1.9 ± 0.3 pmol $^{45}\text{Ca}^{2+}$ bound per pmol of R4064–4210, with an apparent affinity of $60 \pm 12 \mu\text{M}$, $n = 3$, and a Hill coefficient of 1.6 ± 0.4 ; see Fig. 2 A). To confirm the ability of R4064–4210 to bind Ca^{2+} we also used terbium fluorescence. We found that terbium fluorescence was increased in the presence of R4064–4210 and that the increase is inhibited by Ca^{2+} (data not shown), suggesting that Ca^{2+} competes with terbium for a common binding site on this fragment of RyR1.




FIGURE 2 R4064–4210 binds Ca^{2+} . (A) Analysis of $^{45}\text{Ca}^{2+}$ binding to R4064–4210 by equilibrium dialysis. R4064–4210 ($15 \mu\text{M}$, $100 \mu\text{l}$) in 30 mM HEPES pH 7.0 was injected into one of two chambers of an Equilibrium Dialysis cell, separated by a 3500-MWCO membrane. The same amount of the buffer containing different concentrations of $^{45}\text{Ca}^{2+}$ was injected into the other chamber. After overnight incubation, $50\text{-}\mu\text{l}$ aliquots were removed from each chamber for determination of $^{45}\text{Ca}^{2+}$ by liquid scintillation counting. $^{45}\text{Ca}^{2+}$ bound to R4064–4210 was determined from the difference in $^{45}\text{Ca}^{2+}$ radioactivity in the two chambers. By a four-parameter Hill fitting, R4064–4210 apparent affinity for Ca^{2+} (EC_{50}) = $60 \pm 12 \mu\text{M}$, maximum binding (B_{max}) = 1.9 ± 0.3 pmol $^{45}\text{Ca}^{2+}$ bound per pmol of R4064–4210, and the Hill coefficient = 1.6 ± 0.4 . (B) Ca^{2+} binding to R4064–4210 alters ANSA fluorescence. R4064–4210 ($0.4 \mu\text{M}$) was incubated with ANSA ($1 \mu\text{M}$) in 30 mM MOPS (pH 7.2), with either 5 mM EGTA (low Ca^{2+}) or 2 mM Ca^{2+} (high Ca^{2+}) for 5 min at room temperature. Fluorescence emission spectra from 430 to 600 nm were collected with a 370-nm excitation wavelength. 1, R4064–4210 (low Ca^{2+}); 2, R4064–4210 (high Ca^{2+}); 3, ANSA (low Ca^{2+}); 4, ANSA (high Ca^{2+}); 5, R4064–4210 + ANSA (low Ca^{2+}); and 6, R4064–4210 + ANSA (high Ca^{2+}).

Ca^{2+} -induced changes in R4064–4210 assessed by ANSA fluorescence probe analysis

To determine if R4064–4210 undergoes a structural change upon binding Ca^{2+} , we used ANSA, a probe that increases its fluorescence when bound to the hydrophobic sites on proteins (34). We found that Ca^{2+} increased ANSA fluorescence in the presence of R4064–4210 (Fig. 2 B), suggesting that Ca^{2+} binds to R4064–4210 and induces exposure of hydrophobic regions.

R4064–4210 binds to the CaM binding motif (R3614–3643) of RyR1

R4064–4210 binds directly to a peptide (R3614–3643) representing the CaM binding site on RyR1. We found a very dramatic increase in ANSA fluorescence with increasing concentrations of R3614–3643 (Fig. 3). If the R4064–4210 region of RyR1 is important for Ca^{2+} -dependent regulation of RyR1 activity, the interactions should be in some way regulated by the binding of Ca^{2+} . To assess this, we examined the interaction of R3614–3643 with R4064–4210 at different Ca^{2+} concentrations (Fig. 3). We found that R3614–3643 apparently interacts with R4064–4210 at all of the Ca^{2+} concentrations tested, but that the affinity and the apparent magnitude of the ANSA fluorescence changes were different at different Ca^{2+} concentrations. The apparent affinities of R3614–3643 for R4064–4210 at $<10 \text{ nM}$, $50 \mu\text{M}$ Ca^{2+} , $500 \mu\text{M}$ Ca^{2+} , and 5 mM Ca^{2+} were $803 \pm 47 \text{ nM}$, $230 \pm 7 \text{ nM}$, $268 \pm 8 \text{ nM}$, and $181 \pm 10 \text{ nM}$, respectively. The magnitude of the ANSA fluorescence change was also different at the different Ca^{2+} concentrations, with the smallest change occurring at 5 mM and $<10 \text{ nM}$ Ca^{2+} (concentrations where RyR1 is less active). An interaction of the CaM-like domain (R4064–4210) at both high and low Ca^{2+} concentrations is consistent with R3614–3643 being a binding site for both apoCaM and Ca^{2+} CaM (22,23).

Amino-acid determinants in R3614–3643 required for R4064–4210 binding

To determine which amino acids within the CaM binding site of RyR1 were required for the interaction with R4064–4210, we tested the ability of a series of overlapping peptides from this region for the ability to interact with R4064–4210 using ANSA fluorescence to assess the interactions. We found that a peptide corresponding to amino acids 3614–3634 (*solid circles* in Fig. 4 A) had an apparent affinity ($\text{EC}_{50} = 251 \pm 17 \text{ nM}$) that was similar to R3614–3643 ($268 \pm 8 \text{ nM}$) (*solid squares* in Fig. 4 A) at high Ca^{2+} ; however, the magnitude of the fluorescence change induced by R3614–3634 was much less than that induced by R3614–3643. This finding suggests that determinants between amino acids 3614–3634 were important for affinity, but that determinants between amino acids 3635 and 3643 are needed to induce the conformational




FIGURE 3 R4064–4210 interaction with R3614–3643 at different Ca^{2+} concentrations by ANSA fluorescence probe. $0.5\mu\text{M}$ R4064–4210 and $2\mu\text{M}$ ANSA was incubated in 30 mM MOPS ($\text{pH } 7.2$) containing 5 mM EGTA, \bullet ; $50\mu\text{M}$ Ca^{2+} , \blacksquare ; $500\mu\text{M}$ Ca^{2+} , \blacktriangle ; or 5 mM Ca^{2+} , \blacktriangledown for 5 min at room temperature. Then the incubated materials were titrated with R3614–3643 and the ANSA fluorescence data were collected. Ca^{2+} concentrations affect significantly the interaction of R4064–4210 with R3614–3643 ($\text{EC}_{50} = 803 \pm 47, 230 \pm 7, 268 \pm 8, \text{ and } 181 \pm 10\text{ nM}$; and the Hill coefficient = $1.4 \pm 0.1, 2.8 \pm 0.2, 3.1 \pm 0.2, \text{ and } 1.9 \pm 0.2$ for 5 mM EGTA, $50\mu\text{M}$ Ca^{2+} , $500\mu\text{M}$ Ca^{2+} , and 5 mM Ca^{2+} , respectively).

change at high Ca^{2+} . The R3614–3634 peptide had a higher apparent affinity for R4064–4210 ($320 \pm 21\text{ nM}$, $n = 3$) (solid circles in Fig. 4 B) than R3614–3643 ($803 \pm 47\text{ nM}$, $n = 3$) (solid squares in Fig. 4 B) at low Ca^{2+} , but again the maximal fluorescence change was much less than that produced by R3614–3643. These data suggest that the amino acids between 3635 and 3643 did not contribute greatly to the affinity but were responsible for the greater change in ANSA fluorescence with R4064–4210. A peptide corresponding to amino acids 3625–3644 produced a maximal fluorescence change comparable to R3614–3643 but with a twofold decrease in affinity ($\text{EC}_{50} = 525 \pm 27\text{ nM}$) at high Ca^{2+} (solid diamonds in Fig. 4 A). The affinities of the two peptides were similar at low Ca^{2+} (835 ± 23 vs. $803 \pm 47\text{ nM}$ for 3614–3643) (solid diamonds in Fig. 4 B). A peptide corresponding to amino acids 3614–3627 (solid up-triangles in Fig. 4, A and B) had little effect on the ANSA fluorescence at either high or low Ca^{2+} concentrations.

Effects of R4064–4210 on (^3H)ryanodine binding to RyR1

If the interaction between amino acids 3614–3643 and 4064–4210 occurs within the native RyR1, and if this interaction in some way regulates the functional state of RyR1, both R4064–4210 and R3614–3643 would be predicted to modulate RyR1 activity by interfering with interactions within RyR1. To test this possibility, we examined the effects of R4064–4210 and R3614–3643 on the (^3H)ryanodine binding to RyR1 at different Ca^{2+} concentrations. For comparison, the effects of CaM on (^3H)ryanodine binding are also shown.




FIGURE 4 Mapping the amino acids in R3614–3643 that bind to R4064–4210. R4064–4210 ($0.5\mu\text{M}$) and ANSA ($2\mu\text{M}$) were incubated in 30 mM MOPS ($\text{pH } 7.2$) containing $500\mu\text{M}$ Ca^{2+} (A, high Ca^{2+}) or 5 mM EGTA (B, low Ca^{2+}) for 5 min at room temperature. Peptides R3614–3627, R3614–3634, R3614–3643, and R3625–3644 were added at the indicated concentrations, and the ANSA fluorescence were collected. ANSA fluorescence data in the presence of R4064–4210 per peptide were processed by subtracting the effect of R4064–4210 and peptide on the fluorescence of ANSA. (\bullet , R3614–3634 with $\text{EC}_{50} = 251 \pm 17\text{ nM}$, the Hill coefficient = 1.8 ± 0.2 for high Ca^{2+} and $320 \pm 21\text{ nM}$, 1.7 ± 0.2 for low Ca^{2+} ; \blacksquare , R3614–3643 with $\text{EC}_{50} = 268 \pm 8\text{ nM}$, the Hill coefficient = 3.1 ± 0.2 for high Ca^{2+} and $803 \pm 47\text{ nM}$, 1.4 ± 0.1 for low Ca^{2+} ; \blacklozenge , R3625–3644 with $\text{EC}_{50} = 525 \pm 27\text{ nM}$, the Hill coefficient = 1.5 ± 0.1 for high Ca^{2+} and $835 \pm 23\text{ nM}$, 2.6 ± 0.2 for low Ca^{2+} ; and \blacktriangle , R3614–3627.)

(^3H)Ryanodine binding is frequently used to assess functional effects on RyR1 since ryanodine preferentially binds to the open state of the channel (35). Ca^{2+} titration of (^3H)ryanodine binding to SR membranes in the presence and absence of R4064–4210, R3614–3643, and CaM is shown in Fig. 5. (^3H)Ryanodine binding to SR membranes shows a bell-shaped dependence on Ca^{2+} concentration similar to the Ca^{2+} dependence of channel activity: binding is low at nM Ca^{2+} concentrations, increases in sub- μM Ca^{2+} , and reaches maximum at $\sim 10\text{-}\mu\text{M}$ Ca^{2+} , and then decreases (Fig. 5, curve with solid circles). Our surprising finding were that both R4064–4210 and R3614–3643 (Fig. 5, curves with solid up-triangles and solid down-triangles, respectively) slightly enhanced (^3H)ryanodine binding at $< 10\text{ nM}$ Ca^{2+} , inhibited (^3H)ryanodine at intermediate Ca^{2+} and prevented Ca^{2+} inhibition of (^3H)ryanodine binding at high Ca^{2+} . For comparison, CaM activates the channel at low Ca^{2+} but




FIGURE 5 Peptides alter (^3H)ryanodine binding to SR membranes. SR membranes ($10\ \mu\text{g}$) were incubated with $5\ \text{nM}$ (^3H)ryanodine in $100\ \text{mM}$ NaCl, $30\ \text{mM}$ MOPS (pH 7.2) containing increasing CaCl_2 concentration with or without $10\ \mu\text{M}$ CaM, R4064–4210, or R3614–3643 for 16 h at room temperature. The samples were filtered through Whatman GF/F filters with five washes and the radioactivity determined by liquid scintillation counting. Control, \bullet ; CaM, \blacksquare ; R4064–4210, \blacktriangle ; and R3614–3643, \blacktriangledown .

inhibits the channel at high Ca^{2+} (Fig. 5, curve with solid squares). Since these peptides and fragments would be expected to compete with intramolecular interactions of these same sequences within RyR1, these findings suggest that an interaction between amino acids 3614–3643 and 4064–4210 within RyR1: 1), somewhat inhibits activity in low Ca^{2+} (perhaps stabilizing a resting state of the channel); and 2), enhances channel activity at intermediate Ca^{2+} and inhibits channel activity at high Ca^{2+} . CaM might activate the channel at low Ca^{2+} and inhibit the channel at intermediate Ca^{2+} by competing with these intramolecular interactions.

R4064–4210 binds to CaM binding motifs

Since R4064–4210 is predicted to bind Ca^{2+} and interact with CaM binding motifs within RyR1, we asked whether it could bind to other CaM binding motifs. The CaM binding sites on voltage-dependent calcium channels and RyR1 are complex sites. In both types of protein, more than one motif has been implicated in CaM binding (22,25,27–30). We examined the ability of R4064–4210 to interact with synthetic peptides representing CaM-binding motifs from these channels. Also, to determine if this is a more general protein-protein interaction motif, we assessed the ability of R4064–4210 to bind to the CaM binding peptide from CaM kinase II (aa 281–309, Calbiochem). We used changes in ANSA fluorescence in the presence of R4064–4210 to assess the interactions. We found that ANSA fluorescence in the presence of R4064–4210 increased in the presence of CaM binding peptides, R3614–3643, calmodulin kinase II inhibitor peptide (CaMKII peptide), the $\text{Ca}_v1.2$ C-peptide, and the $\text{Ca}_v1.2$ IQ-peptide (Fig. 6), suggesting that all of these peptides could bind to the CaM-like region of RyR1 and the interactions again occurs at both high and low Ca^{2+}

concentrations. Peptides that did not bind CaM had no effects on the ANSA fluorescence in the presence of R4064–4210 (data not shown). These interactions were confirmed by tryptophan fluorescence changes and pull-down assays with His-tagged R4064–4210 (data not shown).

R4064–4210 binds to the $\text{Ca}_v1.1$ by pull-down analysis

If the R4064–4210 sequence can bind to the CaM binding site on $\text{Ca}_v1.1$, it should be able to pull down native $\text{Ca}_v1.1$. We tested this using His-tagged R4064–4210 and digitonin-solubilized T-tubule membranes containing the $\text{Ca}_v1.1$. Fig. 7 shows that R4064–4210 pulls down the $\text{Ca}_v1.1$ protein. Consistent with our previous findings, the pull-down of $\text{Ca}_v1.1$ by R4064–4210 does not require Ca^{2+} binding to either R4064–4210 or $\text{Ca}_v1.1$ (Fig. 7 A for low Ca^{2+} and Fig. 7 B for high Ca^{2+}). To confirm that the pull-down is specific for $\text{Ca}_v1.1$, pull-down data of Na/K ATPase by R4064–4210 are also shown in Fig. 7. At either low Ca^{2+} (Fig. 7 C) or high Ca^{2+} condition (Fig. 7 D), R4064–4210 does not pull-down Na/K ATPase.

DISCUSSION

RyRs are activated by the binding of Ca^{2+} at μM concentrations, but are inhibited by higher concentrations (1,2). Our data suggest that an expressed fragment containing two putative E-F hands (R4064–4210) and a synthetic peptide (R3614–3643) that represents the CaM binding motif of RyR1 interfere with an interaction that regulates the response of RyR1 to Ca^{2+} . These findings raise questions as to whether Ca^{2+} binding to sites between amino acids 4064 and 4210 is involved in the Ca^{2+} regulation of the channel and, if so, how? One possibility is that Ca^{2+} binding to R4064–4210 establishes which determinants within both the 3614–3643 CaM binding site and amino acids 4064–4210 are actually interacting, and as a result, the functional outcomes of the interaction. We have previously shown that Ca^{2+} binding to CaM causes an N-terminal shift in its interaction within amino acids 3614–3643 and converts CaM from an activator to an inhibitor of the channel (20,21). It is conceivable that a similar shift in the interaction of amino acids 4064–4210 with amino acids 3614–3643 could occur when Ca^{2+} binds to the two sites between amino acids 4064 and 4210. It is, however, also possible that Ca^{2+} binding to other sites within RYR1 alters the aa4064–4210 to aa3614–3643 interaction. The effects of R4064–4210 and R3614–3643 on the Ca^{2+} dependence of (^3H)ryanodine binding supports the possibility that this intramolecular interaction is important for both Ca^{2+} activation and inhibition of the channel. The location of the Ca^{2+} activation and inhibition sites remain to be determined, but could conceivably be the two sites within amino acids 4064–4210. Scrambling the sequence of putative E-F hand 1 and E-F hand 2 within aa4064–4210 does not alter




FIGURE 6 R4064–4210 binds to CaM binding motifs. R4064–4210 (0.5 μM) and ANSA (2 μM) were incubated in 30 mM MOPS (pH 7.2), 5 mM EGTA (low Ca^{2+}), or 500 μM Ca^{2+} (high Ca^{2+}) for 5 min at room temperature. CaM-binding peptides (5 μM , R3614–3643 peptide, A; CaMKII peptide, B; C-peptide, C; and IQ-peptide, D) were added and ANSA fluorescence emission spectra were collected from 430 nm to 560 nm with a 370-nm excitation wavelength. (Curve 1, ANSA + EGTA; curve 2, ANSA + Ca^{2+} ; curve 3, ANSA + peptide + EGTA; curve 4, ANSA + peptide + Ca^{2+} ; curve 5, ANSA + R4064–4210 + EGTA; curve 6, ANSA + R4064–4210 + Ca^{2+} ; curve 7, ANSA + R4064–4210 + peptide + EGTA; and curve 8, ANSA + R4064–4210 + peptide + Ca^{2+} .)

the functional responses to either depolarization or activators (caffeine or 4-CmC) when compared with wtRyR1 expressed in myotubes (19). The question is whether 4-CmC or caffeine can accurately assess alterations in the affinity of Ca^{2+} activation or inhibition sites. Another question is whether the presence of endogenous CaM (which we propose competes for the interaction of aa3614–3643 with aa4064–4210) can obscure apparent changes in Ca^{2+} activation or inhibition of RyR1. Fessenden et al. (36) found that RyR1 expressed in dyspedic 1B5 myotubes was activated by 4-CmC, whereas RyR3 was not. Given that RyR1 and RyR3 are both activated by Ca^{2+} with only an approximately twofold difference in Ca^{2+} affinity (37), but have the same apparent affinity for caffeine, one has to question whether altered responses to caffeine and 4-CmC can be used to indirectly assess RyR1 responses to Ca^{2+} . In addition, caffeine and 4-CmC not only increase Ca^{2+} affinity of RyR1 but also alter the maximal activity. Additional data obtained by Fessenden et al. (18) with SR membranes argues that these two E-F hands play an important role in Ca^{2+} regulation of the channel. Scrambling of the sequence of EF1 altered the Ca^{2+} dependence of (³H)ryanodine binding, with respect to both enhancement and inhibition. Even more surprisingly, scrambling of EF2

abolished high-affinity (³H)ryanodine binding to membranes. Without a direct analysis of the binding of Ca^{2+} to these E-F hand mutants, these studies do not rule out E-F hands 1 and 2 as sites of Ca^{2+} regulation of the channel. The sequence from 4064 to 4210 binds Ca^{2+} and modulates the Ca^{2+} dependence of channel activity and we cannot, therefore eliminate the possibility that the Ca^{2+} sites within this region are regulatory sites.

If the region containing E-F-1 and E-F-2 represents a site at which Ca^{2+} regulates channel activity, the interaction of the CaM binding motif (R3614–3643) with this sequence suggests that CaM alters channel activity by interfering with these interactions in a Ca^{2+} -dependent fashion, activating when it is in its apoCaM form and inhibiting in its Ca^{2+} bound form, both of which bind to R3614–3643. We suggest that CaM activates the channel at low Ca^{2+} and inhibits the channel at intermediate Ca^{2+} by competing with the intramolecular interactions between amino acids 4064–4210 and 3614–3643.

Other laboratories have also suggested that CaM might be altering RyR1 activity by interfering with an intramolecular interaction or by inducing a conformational change in the CaM binding site. Zhu et al. (38) demonstrated that a peptide




FIGURE 7 R4064–4210 binds to the native $\text{Ca}_v1.1$. His-tagged R4064–4210 or the synthetic His-tag alone were incubated with Ni-charged chelating sepharose beads and digitonin solubilized T-tubule membranes in either low, $<10 \text{ nM Ca}^{2+}$ (A), or high, 1 mM Ca^{2+} (B). After centrifugation, the pellets and supernatants were electrophoresced on 10% SDS polyacrylamide gels. After transfer to Immobilon (Millipore, Billerica, MA), the Western blots were developed with an anti- $\alpha 1$ antibody from Affinity BioReagents. For comparison, Western blots were also developed with an anti-Na/K ATPase $\alpha 1$ antibody, with C for low Ca^{2+} and D for high Ca^{2+} . (Lanes 1–3, pull-down with His-tagged R4064–4210; lanes 4–6, pull-down with His tag only; lanes 7–9, supernatant from the pull-down with His-tagged R4064–4210; lanes 10–12, supernatants from the pull-downs with His-tag only; and lane 13, starting T-tubule preparation for the pull-down assay.)

representing the CaM binding site (amino acids 3614–3643) could either activate or inhibit RyR1, depending on the dose and the Ca^{2+} concentration. They proposed that the CaM binding site was an important modulatory site within RyR1. Consistent with this, Rodney et al. (39) showed that the R3614–3643 peptide injected in frog skeletal muscle fibers increased the occurrence of Ca^{2+} sparks in a dose-dependent manner, but not by displacing endogenous CaM. These studies also support the existence of an important intramolecular interaction. Gangopadhyay et al. (40) used an environment-sensitive fluorescent probe, 6-bromoacetyl-2-dimethylaminonaphthalene (badan), to study the interaction between badan-labeled calmodulin (CaM) and the CaM-binding peptide of the ryanodine receptor. They found that the interaction interface and the global conformation of the CaM-CaM binding peptide were altered by the binding of Ca^{2+} to CaM.

Other amino acids are also likely to contribute to the ability of CaM to regulate RyR1 activity. Yamaguchi et al. (41) created chimeras of RyR1 and RyR2 and found that five nonconserved amino acid residues (RyR1 aa3680 and 3682–3685, and RyR2 aa3647 and 3649–3652), by differentially affecting RyR helical probability, played a key role in the ability of CaM to inhibit RyR1.

We have previously proposed that the CaM binding sequence on RyR1 can interact with determinants on the $\text{Ca}_v1.1$ and that the CaM binding sequence on the $\text{Ca}_v1.1$ can bind to determinants on RyR1 (26). We now show that the expressed fragment of RyR1 from amino acids 4064–4210

can bind to the intact $\text{Ca}_v1.1$ and to peptides representing the CaM binding sites of both RyR1 and the $\text{Ca}_v1.1$ (both the C-peptide and the IQ-peptide). We propose that the CaM binding sites on RyR1 and the $\text{Ca}_v1.1$ are more general protein-protein interaction motifs that bind to regions that contain E-F hands. The significance of this for excitation-contraction coupling is that it may provide a unique mechanism for interactions between these two ion channels: the CaM binding site on the $\text{Ca}_v1.1$ binding to the RyR1 CaM-like domain and the RyR1 CaM binding site interacting with a CaM-like domain on the $\text{Ca}_v1.1$. Since R4064–4210 of RyR1 also interacts with the CaM binding site (R3614–3643) on RyR1 to stabilize the closed channel, the interaction with the carboxyterminal tail of the $\text{Ca}_v1.1 \alpha_1$ -subunit might disrupt an inter- or intramolecular interaction, thereby regulating RyR1 activity.

The studies were supported in part by a grant from the Muscular Dystrophy Association, and National Institutes of Health grants No. AR41802 and No. AR44864 (to S.L.H.).

REFERENCES

- Liu, W., D. A. Pasek, and G. Meissner. 1998. Modulation of Ca^{2+} -gated cardiac muscle Ca^{2+} -release channel (ryanodine receptor) by mono- and divalent ions. *Am. J. Physiol.* 274:C120–C128.
- Meissner, G., E. Rios, A. Tripathy, and D. A. Pasek. 1997. Regulation of skeletal muscle Ca^{2+} release channel (ryanodine receptor) by Ca^{2+} and monovalent cations and anions. *J. Biol. Chem.* 272:1628–1638.
- Kretsinger, R. H. 1976. Calcium-binding proteins. *Annu. Rev. Biochem.* 45:239–266.
- Kretsinger, R. H. 1976. Evolution and function of calcium-binding proteins. *Int. Rev. Cytol.* 46:323–393.
- Takekuma, H., S. Nishimura, T. Matsumoto, H. Ishida, K. Kangawa, N. Minamino, H. Matsuo, M. Ueda, M. Hanoaka, T. Hirose, and S. Numa. 1989. Primary structure and expression from complementary DNA of skeletal muscle ryanodine receptor. *Nature.* 339:439–445.
- Nakai, J., T. Imagawa, Y. Hakamat, M. Shigekawa, H. Takeshima, and S. Numa. 1990. Primary structure and functional expression from cDNA of the cardiac ryanodine receptor/calcium release channel. *FEBS Lett.* 271:169–177.
- Hakamata, Y., J. Nakai, H. Takeshima, and K. Imoto. 1992. Primary structure and distribution of a novel ryanodine receptor/calcium release channel from rabbit brain. *FEBS Lett.* 312:229–235.
- Treves, S., P. Chiozzi, and F. Zorzato. 1993. Identification of the domain recognized by anti-(ryanodine receptor) antibodies which affect Ca^{2+} -induced Ca^{2+} release. *Biochem. J.* 291:757–763.
- Chen, S. R., L. Zhang, and D. H. MacLennan. 1992. Characterization of a Ca^{2+} binding and regulatory site in the Ca^{2+} release channel (ryanodine receptor) of rabbit skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.* 267:23318–23326.
- Chen, S. R., L. Zhang, and D. H. MacLennan. 1993. Antibodies as probes for Ca^{2+} activation sites in the Ca^{2+} release channel (ryanodine receptor) of rabbit skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.* 268:13414–13421.
- Zorzato, F., J. Fujii, K. Otsu, M. Phillips, N. M. Green, F. A. Lai, G. Meissner, and D. H. MacLennan. 1990. Molecular cloning of cDNA encoding human and rabbit forms of the Ca^{2+} release channel (ryanodine receptor) of skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.* 265:2244–2256.
- Hayek, S. M., X. Zhu, M. B. Bhat, J. Zhao, H. Takeshima, H. H. Valdivia, and J. Ma. 2000. Characterization of a calcium-regulation

- domain of the skeletal-muscle ryanodine receptor. *Biochem. J.* 351: 57–65.
13. Du, G. G. and D. H. MacLennan. 1999. Ca^{2+} inactivation sites are located in the COOH-terminal quarter of recombinant rabbit skeletal muscle Ca^{2+} release channels (ryanodine receptors). *J. Biol. Chem.* 274:26120–26126.
 14. Chen, S. R. W., and D. H. MacLennan. 1994. Identification of calmodulin, Ca^{2+} , and ruthenium red-binding domains in the Ca^{2+} release channel (ryanodine receptor) of rabbit skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.* 269:22698–22704.
 15. Xiong, H., X. Feng, L. Gao, L. Xu, D. A. Pasek, J. H. Seok, and G. Meissner. 1998. Identification of a two EF-hand Ca^{2+} binding domain in lobster skeletal muscle ryanodine receptor/ Ca^{2+} release channel. *Biochemistry.* 37:4804–4814.
 16. Hamada, T., Y. Sakube, J. Ahn, H. Kim, and H. Kagawa. 2002. Molecular dissection, tissue localization and Ca^{2+} binding of the ryanodine receptor of *Caenorhabditis elegans*. *J. Mol. Biol.* 324:123–135.
 17. Fessenden, J. D., L. Chen, Y. Wang, C. Paolini, C. Franzini-Armstrong, P. D. Allen, and I. N. Pessah. 2001. Ryanodine receptor point mutant E4032A reveals an allosteric interaction with ryanodine. *Proc. Natl. Acad. Sci. USA.* 98:2865–2870.
 18. Fessenden, J. D., W. Feng, I. N. Pessah, and P. D. Allen. 2004. Mutational analysis of putative calcium binding motifs within the skeletal ryanodine receptor isoform, RyR1. *J. Biol. Chem.* 279:53028–53035.
 19. Tripathy, A., L. Xu, G. Mann, and G. Meissner. 1995. Calmodulin activation and inhibition of skeletal muscle Ca^{2+} release channel (ryanodine receptor). *Biophys. J.* 69:106–119.
 20. Rodney, G. G., B. Y. Williams, G. M. Strasburg, K. Beckingham, and S. L. Hamilton. 2000. Regulation of RyR1 activity by Ca^{2+} and calmodulin. *Biochemistry.* 39:7807–7812.
 21. Rodney, G. G., C. P. Moore, B. Y. Williams, J. Z. Zhang, J. Krol, S. E. Pedersen, and S. L. Hamilton. 2001. Calcium binding to calmodulin leads to an N-terminal shift in its binding site on the ryanodine receptor. *J. Biol. Chem.* 276:2069–2074.
 22. Moore, C. P., J.-Z. Zhang, and S. L. Hamilton. 1999. A role for Cysteine 3635 of RyR1 in redox modulation and calmodulin binding. *J. Biol. Chem.* 274:36831–36834.
 23. Rodney, G. G., J. Krol, B. Williams, K. Beckingham, and S. L. Hamilton. 2001. The carboxy-terminal calcium binding sites of calmodulin control calmodulin's switch from an activator to an inhibitor of RyR1. *Biochemistry.* 40:12430–12435.
 24. Xiong, L. W., R. A. Newman, G. G. Rodney, O. Thomas, J. Z. Zhang, A. Persechini, M. A. Shea, and S. L. Hamilton. 2002. Lobe-dependent regulation of ryanodine receptor Type 1 by calmodulin. *J. Biol. Chem.* 277:40862–40870.
 25. Zhang, H., J. Z. Zhang, C. I. Danila, S. L. Hamilton, and A. Noncontiguous. 2003. Inter-subunit binding site for calmodulin on the skeletal muscle Ca^{2+} release channel. *J. Biol. Chem.* 278:8348–8355.
 26. Sencer, S., R. V. Papineni, D. B. Halling, P. Pate, J. Krol, J. Z. Zhang, and S. L. Hamilton. 2001. Coupling of RyR1 and L-type calcium channels via calmodulin binding domains. *J. Biol. Chem.* 276:38237–38241.
 27. Pate, P., J. Mochca-Morales, Y. Wu, J. Z. Zhang, G. G. Rodney, I. I. Serysheva, B. Y. Williams, M. E. Anderson, and S. L. Hamilton. 2000. Determinants for calmodulin binding on voltage-dependent Ca^{2+} channels. *J. Biol. Chem.* 275:39786–39792.
 28. Pitt, G. S., R. D. Zuhlke, A. Hudmon, H. Schulman, H. Reuter, and R. W. Tsien. 2001. Molecular basis of calmodulin tethering and Ca^{2+} -dependent inactivation of L-type Ca^{2+} channels. *J. Biol. Chem.* 276: 30794–30802.
 29. Soldatov, N. M., M. Oz, K. A. O'Brien, D. R. Abernethy, and M. Morad. 1998. Molecular determinants of L-type Ca^{2+} channel inactivation. segment exchange analysis of the carboxyl-terminal cytoplasmic motif encoded by exons 40–42 of the human $\alpha 1\text{C}$ -subunit gene. *J. Biol. Chem.* 273:957–963.
 30. Zühlke, R. D., and H. Reuter. 1998. Ca^{2+} -sensitive inactivation of L-type Ca^{2+} channels depends on multiple cytoplasmic amino acid sequences of the $\alpha 1\text{C}$ -subunit. *Proc. Natl. Acad. Sci. USA.* 95:3287–3294.
 31. Hawkes, M. J., M. Díaz-Muñoz, and S. L. Hamilton. 1989. A procedure for the purification of the ryanodine receptor from skeletal muscle. *Membr. Biochem.* 8:133–145.
 32. Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the folin phenol reagent. *J. Biol. Chem.* 193:265–275.
 33. Venyaminov, S. Y., and J. T. Yang. 1996. Circular Dichroism and the Conformational Analysis of Biomolecules. G. D. Fasman, editor. Plenum Publishing, New York. 69–108.
 34. Muesing, R. A., and T. Nishida. 1971. Disruption of low- and high-density human plasma lipoproteins and phospholipid dispersions by 1-anilinonaphthalene-8-sulfonate. *Biochemistry.* 10:2952–2962.
 35. Chu, A., M. Diaz-Munoz, M. J. Hawkes, K. Brush, and S. L. Hamilton. 1990. Ryanodine as a probe for the functional state of the skeletal muscle sarcoplasmic reticulum Ca^{2+} release channel. *Mol. Pharmacol.* 37:735–741.
 36. Fessenden, J. D., C. F. Perez, S. Goth, I. N. Pessah, and P. D. Allen. 2003. Identification of a key determinant of ryanodine receptor Type 1 required for activation by 4-chloro-m-cresol. *J. Biol. Chem.* 278: 28727–28735.
 37. Murayama, T., and Y. Ogawa. 2004. RyR1 exhibits lower gain of CICR activity than RyR3 in the SR: evidence for selective stabilization of RyR1 channel. *Am. J. Physiol. Cell Physiol.* 287:C36–C45.
 38. Zhu, X., J. Ghanta, J. W. Walker, P. D. Allen, and H. H. Valdivia. 2004. The calmodulin binding region of the skeletal ryanodine receptor acts as a self-modulatory domain. *Cell Calcium.* 35:165–177.
 39. Rodney, G. G., G. M. Wilson, and M. F. Schneider. 2005. A calmodulin binding domain of RyR increases activation of spontaneous Ca^{2+} sparks in frog skeletal muscle. *J. Biol. Chem.* 280:11713–11722.
 40. Gangopadhyay, J. P., Z. Grabarek, and N. Ikemoto. 2004. Fluorescence probe study of Ca^{2+} -dependent interactions of calmodulin with calmodulin-binding peptides of the ryanodine receptor. *Biochem. Biophys. Res. Commun.* 323:760–768.
 41. Yamaguchi, N., L. Xu, K. E. Evans, D. A. Pasek, and G. Meissner. 2004. Different regions in skeletal and cardiac muscle ryanodine receptors are involved in transducing the functional effects of calmodulin. *J. Biol. Chem.* 279:36433–36439.