



Published in final edited form as:

Pediatr Res. 2006 March ; 59(3): 471–477.

Maternal IQ, Child IQ, behavior and achievement in urban 5–7 year olds

Aimin Chen, Donald Schwarz, Jerilynn Radcliffe, and Walter J. Rogan

Epidemiology Branch, National Institute of Environmental Health Sciences, National Institutes of Health, Department of Health and Human Services [A.C., W.J.R.], Research Triangle Park, North Carolina 27709, USA Department of Adolescent Medicine, Children's Hospital of Philadelphia [D.S.], Philadelphia, Pennsylvania 19104, USA Department of Psychology, Children's Hospital of Philadelphia, and University of Pennsylvania School of Medicine [J.R.], Philadelphia, Pennsylvania 19104, USA

Abstract

In one study of children in 27 families with maternal retardation, those children with higher IQ were more likely to have multiple behavior problems than those with lower IQ. If true, this result would affect clinical practice, but it has not been replicated. Since the setting of the initial observation is similar to the setting of childhood lead poisoning, we attempted a replication using data from the Treatment of Lead-exposed Children (TLC) study, in which 780 children aged 12–33 months with blood lead level 20–44 µg/dL were randomized to either succimer treatment or placebo and then followed up to 5 years. Of 656 mothers of TLC children with IQ measured, 113 demonstrated mental retardation (IQ < 70). Whether maternal IQ was less than 70 or over 70, children with IQ over 85 were rated more favorably on cognitive tests and behavioral questionnaires than children with IQ less than 85; these measures included Conners' Parent Rating Scale-Revised at age 5, the Developmental Neuropsychological Assessment at ages 5 and 7, and the Behavioral Assessment System for Children at age 7. Among children of mothers with IQ < 70, those with IQ over 85 did not show more severe clinical behavioral problems, nor were they more likely to show multiple behavior problems. Children with higher IQ have fewer behavior problems, irrespective of the mother's IQ. In the special setting of mothers with IQ < 70, children with higher IQ are not at greater risk of behavior problems.

Keywords

mothers; children; intelligence; mental retardation; behavior

Abbreviations

BASC-PRS: Behavioral Assessment System for Children-Parent Rating Scale; BASC-TRS: Behavioral Assessment System for Children-Teacher Rating Scale; CPRS-R: Conners' Parent Rating Scale-Revised; CPT: Conners' Continuous Performance Test; CVLTC: California Verbal Learning Test for Children; NEPSY: Developmental Neuropsychological Assessment; NESS: Neurological Examination for Subtle Signs; TLC: Treatment of Lead-exposed Children study; WLPB-R: Woodcock Language Proficiency Battery-Revised

Introduction

The relation of maternal IQ with child IQ and child behavior is not well studied in families with maternal mental retardation. In 1997, Feldman and Walton-Allen reported that children of mothers with mental retardation living in poverty had more behavior problems and lower IQ than children from similar families but without maternal mental retardation (1). In addition, they advanced the hypothesis that, among studied children in families with maternal mental retardation, those with higher IQ were more likely to have multiple behavior problems than those with lower IQ. They based this conclusion on their observation that, among the 27 families with maternal mental retardation, none of the 11 children with IQ < 85 scored above the clinical threshold on all 3 sub-scales (conduct, hyperactivity, and emotional) of the Ontario version of the Child Behavior Checklist, whereas 5 of the children with IQ of 85 or greater did. The authors speculated that, since parents with mental retardation may not set reasonable limits for and communicate well with their more intelligent children, these children may take advantage of, and rebel against, less competent parents (1). Early anti-social acts against authorities may also develop (2).

If this hypothesis were true, it would influence clinical practice by focusing attention more on brighter children living in poverty and with mothers with cognitive limitations, since children with higher IQ are not usually thought to be at high risk for behavioral or academic problems (3–5). Because of these clinical implications, we sought to replicate Feldman's finding in another setting in which many relatively disadvantaged families are found, that of urban lead poisoning. We had done a large clinical trial of the treatment of lead poisoning in which we measured maternal IQ, and did multiple measures of child IQ and behavior over 5 years of follow up. We use data from that trial to attempt a direct replication of Feldman's finding, and to explore further the inter-relationships among maternal IQ, child IQ, and child behavior.

Methods

The Treatment of Lead-exposed Children (TLC) study was a multi-center randomized placebo-controlled clinical trial to investigate the effect of treatment with succimer, an oral lead chelator, on IQ, neuropsychological, and behavioral measurements 36 months after the initiation of treatment (6). Altogether 780 children aged 12–33 months with blood lead concentrations of 20–44 µg/dL were randomized to succimer or placebo between August 1994 and January 1997 at four clinical centers: Baltimore MD, Newark NJ, Philadelphia PA, and Cincinnati OH. 396 children were randomly assigned to succimer and 384 to placebo. Depending on the blood lead, children in the succimer group could receive up to 3 26-day courses of treatment; the number of courses given to placebo children was frequency matched to that in the succimer group. The succimer group had a lower blood lead concentration for about 6 months, but did not show any difference in IQ, neuropsychological or behavior tests at 36 months of follow-up (7). An extension of the study to 60 months of follow-up confirmed the main results (8). Since succimer treatment did not affect test scores, we combined across treatment groups for the analysis of maternal IQ, child IQ, behavior, achievement, and neuropsychological function. The study was approved by the institutional review boards at the clinical centers, the Harvard School of Public Health, the CDC, and the National Institute of Environmental Health Sciences. The parents of all the children provided written informed consent at enrollment.

Of 780 children who participated in the trial, 745 had caregiver's IQ measured with the Wechsler Adult Intelligence Scale-Revised (9), among whom were 656 mothers. In this analysis, only data from the 656 children of these mothers were used. The child's full scale IQ was measured with the Wechsler Preschool and Primary Scales of Intelligence-Revised (WPPSI-R) (10) at 36 months of follow-up (about age 5) and with the Wechsler Intelligence Scale for Children-III (WISC-III) (11) (at age 7 years). The neuropsychological and behavioral

tests used at age 5 included the Developmental Neuropsychological Assessment (NEPSY) (12) and Conners' Parent Rating Scale-Revised (CPRS-R) (13), and that used at age 7 were NEPSY (12), Conners' Continuous Performance Test (CPT) (14), California Verbal Learning Test for Children (CVLTC) (15), Woodcock Language Proficiency Battery-Revised (WLPB-R) (16), Neurological Examination for Subtle Signs (NESS) (17), Behavioral Assessment System for Children teacher rating scale (BASC-TRS) and BASC parent rating scale (BASC-PRS) (18). The administration of CPRS-R at age 5 and BASC-PRS at age 7 by parents was completed by trained examiners' reading aloud questions to parents and providing clarification of items as necessary; examiners were blind to maternal IQ levels. Blood lead concentrations were measured by atomic absorption spectrometry based on the methods described by Miller et al (19).

As a first step in using data from the trial to test whether children with higher IQ who had mothers with mental retardation showed more behavior problems, we replicated Feldman's approach, comparing the neuropsychological and behavioral test scores of children with higher (≥ 85) or lower (< 85) IQ scores, given maternal IQ less than 70. We did a similar analysis among children with maternal IQ ≥ 70 . We then compared percentage of children being "at risk" for "clinically significant" problems (CPRS-R indices or BASC problem scores ≥ 60) and percentage of children having "clinically significant" problems (CPRS-R indices or BASC problem scores ≥ 70) across child IQ categories. In addition, we examined the more general finding that children with mentally retarded mothers had lower IQ, poorer school achievement, and more behavior problems. The covariates used for adjustment, determined a priori based on the literature (20–22), were clinic center (Baltimore, Newark, Philadelphia, and Cincinnati), race (black, white and others), gender (male, female), language (English or Spanish), parent's education (under 12 years, 12 years, over 12 years), parents' employment (neither working, either working), single parent (yes, no), concurrent blood lead level (5 or 7 years) and maternal IQ.

Finally, we attempted to assess the relative contributions of maternal IQ, child's IQ, and both on neuropsychological, behavioral, and achievement scores. We made 3 sets of statistical models, all of which included behavior, neuropsychological, or achievement scores as dependent variables. As independent variables, the models utilized either maternal IQ (model 1), child's IQ (model 2), or both (model 3), and all included the covariates: clinical center, race, sex, language, parent's education, parent's employment, single parent, and concurrent blood lead level. SAS 9.0 (SAS Institute Inc. Cary, NC) was used for all statistical analyses.

Results

The mothers and children in these data were predominantly from low socioeconomic status families (Table 1). The 656 children with maternal IQ tested did not differ in baseline characteristics from the other children of the total 780 recruited. Their mean age was about 2 years at baseline, thus 5 years at 36 months of follow-up and 7 years at last follow-up.

The mean (\pm standard deviation, SD) of maternal IQ in the 656 children's mothers was 80 ± 11 ; 113 (17%) had IQs less than 70 (mean \pm SD: 65 ± 4) and 543 (83%) had IQs 70 and over (mean \pm SD: 83 ± 10). At about age 5, 624 children had IQ tested, with mean \pm SD of 81 ± 13 ; at age 7, 549 children had IQ tested, with mean \pm SD of 86 ± 13 . The correlation coefficient of maternal IQ and child IQ was 0.31 at age 5 and 0.41 at age 7; that of child IQ at age 5 and at age 7 was 0.78. The p values of the correlation tests are all < 0.001 .

The test scores of NEPSY and CPRS-R at age 5 stratified by maternal IQ (< 70 or ≥ 70) and child IQ at age 5 (< 85 or ≥ 85) as well as the difference of scores between two child IQ categories are shown in Table 2. Analogously, scores from NEPSY, CPT, CVLTC, WLPB-R, NESS,

BASC-TRS and BASC-PRS at age 7 are shown in Table 3. Not all children completed all tests and so the numbers vary slightly. The tables show the adjusted differences between child IQ categories. The difference between the maternal IQ categories can be obtained by subtracting the comparable columns under each maternal IQ category. If we look at the CPRS-R in the 5 year olds of the lower IQ mothers, the higher IQ children show better mean behavior scores than the lower IQ children. The pattern is the same for the higher IQ mothers, and the size of the differences between the children with different IQ scores is the same, independent of their mother's IQ. At age 7, the pattern is the same, with one exception – “internalizing problems” on the BASC parent scale is at the same level for higher and lower IQ children stratified by maternal IQ category. In these data, then, higher IQ children generally had fewer behavior problems than lower IQ children in both maternal IQ categories.

The percentages of children with behavior test scores over 60 or 70 are shown in Table 4. With either of the clinical cut-off points, children with IQ less than 85 generally had more behavior problems (some comparisons achieve statistical significance) at a clinical level than their higher IQ peers, given equivalent maternal IQ. Among children with maternal IQ <70, the percentage of having both externalizing and internalizing problems, an indicator similar to “multiple behavior problems” in the Feldman paper (1), did not differ statistically between children with IQ < 85 and IQ ≥85; this is also true for children with maternal IQ over 70 (table 4).

BASC scores from both the teacher rating scale and the parent rating scale were checked to see if agreement between teachers and parents was similar for mothers with IQ <70 and mothers with IQ ≥70. The correlation coefficients between teacher and parent reports were 0.26 for adaptive skills, 0.21 for behavioral symptoms index, 0.35 for externalizing problems, and -0.10 for internalizing problems in mothers with IQ < 70, and in mothers with IQ ≥70, these correlation coefficients were 0.33, 0.29, 0.36, and 0.16, respectively. The mean differences between teacher- and parent-rated scores (parent score minus teacher score) ranged from -2.32 to 3.19, all less than 0.3 standard deviation of these scores, with no statistically significant difference between mothers with IQ < 70 and mothers with IQ ≥70. For the percentage of children with BASC scores ≥ 60 or ≥ 70 in the behavioral symptoms index and externalizing problems, the kappa statistics between teacher and parent report were similar for mothers with IQ < 70 and mothers with IQ ≥70 (kappa 0.13 to 0.28). However, for BASC scores ≥60 or ≥70 in internalizing problems and both externalizing and internalizing problems, the kappa statistics between teacher and parent reports were lower (kappa -0.05 to 0.06) in mothers with IQ < 70 than in mothers with IQ ≥70 (kappa 0.10 to 0.14).

The results of general linear models on the association of maternal IQ and child IQ with the other test scores at ages 5 and 7 are shown in Table 5 and Table 6, respectively. For tests at 5 years, maternal IQ was positively and significantly correlated to NEPSY domains but not to CPRS-R indices. The child's own IQ had an effect about twice as large on the NEPSY domains, and children with higher IQ had significantly fewer behavior problems on the CPRS-R. When both maternal and child IQ are considered (model 3), child IQ remained statistically significant, with regression estimates largely similar to those in model 2, but maternal IQ only had significance on NEPSY language and memory scores and then with smaller regression estimates. At age 7, there is a similar picture, with small effects from maternal IQ that are marginally or not statistically significant, larger positive effects from child IQ, and a reduction of the maternal effect with a persistence of the child effect when both are in the model. This finding holds consistently for neuropsychological testing (NEPSY), behavior (the BASCs) and achievement (WLPB-R Broad reading). Additional regression models of maternal IQ and child IQ show the estimate (SE) of 10 point maternal IQ on child IQ was 3.4 (0.5) points at age 5 and 3.9 (0.5) points at age 7 after adjustment for the covariates used in Table 3/4. It is unlikely that maternal IQ accounts for a large proportion of the correlation between child IQ and non-IQ outcomes. Therefore, even considering the association of maternal IQ with child IQ, child

IQ remained a strong factor that was correlated with child non-IQ outcomes at both age 5 and age 7. Higher child IQ was accompanied by better attention and executive function, language proficiency, adaptive skills and by fewer teacher- or parent-reported behavioral problems.

Because these data come from a trial in which half of the children received succimer, an oral chelating drug, for their lead poisoning, we re-analyzed the data by treatment group, and the main results were similar for succimer and for placebo group. We also did an analysis stratified by gender, and the main results did not differ between boys and girls. In no circumstance did the higher IQ children of mothers with lower IQ have significantly more behavior problems than lower IQ children from similar mothers.

Discussion

The link between maternal cognitive ability and child cognition has been found to be both a genetic and an environmental one (23–26). More recently, the link between maternal cognitive ability and early child behavior has been explored as well (1). The few studies that have looked at children of mentally-retarded parents and those of parents of normal intelligence have suggested that parents with mental retardation are more likely to have children with lower cognitive ability, and that those children with higher cognitive ability are more likely to have behavioral problems than children of normal intelligence with parents of normal intelligence (1,2,27).

We found that children with higher IQ were less likely to be “at risk for” or to present with clinically significant behavior problems, irrespective of maternal IQ. This is true at both age 5 and age 7, and in various behavior domains measured by different behavioral tests. We thus failed to replicate Feldman’s finding (1) that higher IQ children of lower IQ mothers have more behavior problems. We also found that mothers with lower IQ are more likely to have children with lower IQ and more behavior problems than mothers of higher intelligence, consistent with previous studies (27,28).

Our study differs from Feldman’s in sample size, which may have influenced outcomes. Feldman studied only 27 children from mothers with IQ<70, but our study had 108 children with such mothers for CPRS-R at age 5, 85 for BASC-TRS and 101 for BASC-PRS at age 7. Small clinical samples such as Feldman’s may not be representative. Feldman’s studied families were referred from community agencies providing advocacy and support services to adults with mental retardation; the mothers were enrolled first. On the other hand, our group of children was selected because of their elevated blood lead concentration. Despite the difference in enrollment strategy, mothers in the Feldman study with mental retardation had mean IQ of 64 (SD 6) while those mothers in this analysis had mean IQ of 65 (SD 4); they were both of lower IQ and not fundamentally different in intelligence level. Nonetheless, it may be that the relationship between maternal IQ, child IQ, and the other psychological tests is different in the lead-exposed children in our study. In the literature, lead exposure was associated with child IQ deficits (29–31) and, though less studied, behavioral problems (20–22). We have multiple measures of their blood lead concentration, and those measures are associated with many of the test scores reported here. We do include the blood lead concentration measured closest to the time of psychological testing, which in these data bear the strongest relationship to test scores, and so the result we see should not be due simply to the effect of lead on both IQ and behavior. Lead exposure is only one of the factors that affect IQ and behavior, and it seems unlikely that lead exposure would bias the IQ and behavior relationship in TLC study because the observed relationship was similar to that in another report in literature (3). It is also plausible that Feldman’s subjects were also lead-exposed. The children in Feldman study were older (10 years old) and were tested by the Ontario version of

the Child behavior Checklist (32), but that seems unlikely to cause different results in the two studies.

Another issue is that the accuracy of parent rating scale for those mothers with $IQ < 70$ may impact results. When we checked the agreement between teacher and parent BASC 14 ratings for scores at age 7, we found mothers with $IQ < 70$ and mothers with $IQ \geq 70$ had similar levels of agreement with teachers for reporting behavior scores except for internalizing problems. The correlation coefficients and kappa statistics between teacher and parent reports in this study are similar to literature reports of parent-teacher behavior ratings of children (33–35). It is unlikely that these mothers (with IQ score of 65) completely lack the ability to report child behavioral problems. The results of teacher rated scores also confirmed the findings from parent rated scores in our study. Nevertheless, our results suggest that clinicians need to be mindful of the discrepancies between teacher and parent report of internalizing problems in the setting of low maternal IQ. In such situations, clinicians need to ask more specific questions of lower IQ mothers to inquire about symptoms of a child's mental health problems, particularly internalizing problems such as anxiety and depression. Clinicians may also need to ask directly children of mothers with lower IQ about mental health symptoms.

We can speculate about why mother's intelligence level was not as influential as child's IQ in shaping child's behavior patterns. It may be true that lower IQ mothers cannot handle mother-child interaction in an ideal way, but the influence from other family members, neighborhood, community and school will gain more strength as the child grows, if the child is sufficiently intelligent to allow this societal interaction to take effect. Similarly, among children of parents suffering from psychiatric problems, higher IQ is associated with better personal and social adaptation (36,37).

The relative contributions of maternal IQ and child IQ to child behavior outcomes merit further consideration. In this analysis, the effect of maternal IQ is always smaller than the effect of child IQ on behavior, and when both variables are included in the model, the effect of maternal IQ is very small and not significant. Maternal IQ per se might have some effect on both child IQ and child behavior. In a small data set with two variables that are highly correlated, the analytical model has very few discordant observations to work with, and thus the individual parameters are unstable when both are present in a model. At the extreme, if there are no higher IQ children of lower IQ mothers, the model cannot estimate independent effects. The data set we use, however, is relatively large, and the correlation between maternal and child IQ relatively modest, and thus the model's partitioning of the effect to the child is plausible. It would still be useful to see if this holds in other data sets. In a sample of 411 13-year-old twins of normal intelligence, Goodman et al (3) found lower child IQ was associated with more behavioral problems, while lower parental IQ (average of maternal and paternal IQ) did not significantly predict behavioral problems. Compared with parental IQ, child IQ may be more powerful in the inter-relationship of parental IQ, child IQ and child behavior. While it may also be that behavioral problems (e.g., hyperactivity) leads to lower IQ (38), the observation that remission of child psychiatric disorder is not associated with a change in measured IQ suggests child IQ is more likely to be a cause rather than consequence of child behavioral problems (3), probably via poorer ability in dealing with distress or lower self-esteem (39, 40).

Overall we conclude that given any maternal IQ, the higher IQ child is always at lower risk for behavior and other problems than the lower IQ child. Among children with similar IQ, maternal IQ does not explain much of the variation in child behavior, neuropsychological function or achievement. These findings support continued efforts to investigate and address factors influencing child cognitive outcomes, such as preventing lead exposure and providing

supports to children with lower cognitive ability in an effort to prevent poor behavioral outcomes.

Acknowledgements

The Treatment of Lead-exposed Children Trial was supported by NIEHS in cooperation with the NIH Office of Minority Health, and by the Centers for Disease Control and Prevention. Succimer and placebo capsules were gifts from McNeil Labs, Fort Washington, PA.

Part of this work was presented at Pediatric Academic Societies' Annual Meeting, May 14–17, 2005, Washington, DC

References

1. Feldman MA, Walton-Allen N. Effects of maternal mental retardation and poverty on intellectual, academic, and behavioral status of school-age children. *Am J Ment Retard* 1997;101:352–364. [PubMed: 9017082]
2. O'Neill AM. Normal and bright children of mentally retarded parents: the Huck Finn syndrome. *Child Psychiatry Hum Dev* 1985;15:255–268. [PubMed: 4042738]
3. Goodman R, Simonoff E, Stevenson J. The impact of child IQ, parent IQ and sibling IQ on child behavioural deviance scores. *J Child Psychol Psychiatry* 1995;36:409–425. [PubMed: 7782405]
4. Andersson HW, Sommerfelt K. The relationship between cognitive abilities and maternal ratings of externalizing behaviors in preschool children. *Scand J Psychol* 2001;42:437–444. [PubMed: 11771813]
5. Anselmi L, Piccinini CA, Barros FC, Lopes RS. Psychosocial determinants of behaviour problems in Brazilian preschool children. *J Child Psychol Psychiatry* 2004;45:779–788. [PubMed: 15056309]
6. TLC. The Treatment of Lead-exposed Children (TLC) trial: design and recruitment for a study of the effect of oral chelation on growth and development in toddlers. *Paediatr Perinat Epidemiol* 1998;12:313–333. [PubMed: 9690266]
7. Rogan WJ, Dietrich KN, Ware JH, Dockery DW, Salganik M, Radcliffe J, Jones RL, Ragan NB, Chisolm JJ Jr, Rhoads GG. The effect of chelation therapy with succimer on neuropsychological development in children exposed to lead. *N Engl J Med* 2001;344:1421–1426. [PubMed: 11346806]
8. Dietrich KN, Ware JH, Salganik M, Radcliffe J, Rogan WJ, Rhoads GG, Fay ME, Davoli CT, Denckla MB, Bornschein RL, Schwarz D, Dockery DW, Aubato S, Jones RL. Effect of chelation therapy on the neuropsychological and behavioral development of lead-exposed children after school entry. *Pediatrics* 2004;114:19–26. [PubMed: 15231903]
9. Silverstein AB. Two- and four-subtest short forms of the WAIS-R: a closer look at validity and reliability. *J Clin Psychol* 1985;41:95–97. [PubMed: 3973047]
10. Wechsler D 1989 *The Wechsler Preschool and Primary Scales of Intelligence-Revised*. The Psychological Corporation, San Antonio, TX.
11. Wechsler D 1991 *Wechsler Intelligence Scale for Children-III*. The Psychological Corporation, San Antonio, TX.
12. Korkman M, Kirk U, Kemp S 1998 *NEPSY: a developmental neuropsychological assessment: manual*. Rev. The Psychological Corporation, San Antonio, TX.
13. Conners CK 1997 *Conners' Rating Scales: technical manual*. Rev. Multi-Health systems, North Tonawanda, NY.
14. Conners CK 1995 *Conners' Continuous Performance Test*. Multi-Health System, North Tonawanda, NY.
15. Delis DC, Kramer JH, Kaplan E, Ober BA 1994 *California Verbal Learning Test-Children's Version*. The Psychological Corporation, San Antonio, TX.
16. Woodcock RW 1991 *Woodcock Language Proficiency Battery-Revised*. Riverside Publishing, Itasca, IL.
17. Denckla MB. Revised Neurological Examination for Subtle Signs (1985). *Psychopharmacol Bull* 1985;21:773–800. [PubMed: 4089106]

18. Reynolds CR, Kamphaus RW 1992 Behavior Assessment System for Children. American Guidance Service, Circle Pines, MN.
19. Miller DT, Paschal DC, Gunter EW, Stroud PE, D'Angelo J. Determination of lead in blood using electrothermal atomisation atomic absorption spectrometry with a L'vov platform and matrix modifier. *Analyst* 1987;112:1701–1704. [PubMed: 3445938]
20. Burns JM, Baghurst PA, Sawyer MG, McMichael AJ, Tong SL. Lifetime low-level exposure to environmental lead and children's emotional and behavioral development at ages 11–13 years. The Port Pirie Cohort Study. *Am J Epidemiol* 1999;149:740–749. [PubMed: 10206624]
21. Wasserman GA, Staghezza-Jaramillo B, Shrout P, Popovac D, Graziano J. The effect of lead exposure on behavior problems in preschool children. *Am J Public Health* 1998;88:481–486. [PubMed: 9518990]
22. Sciarillo WG, Alexander G, Farrell KP. Lead exposure and child behavior. *Am J Public Health* 1992;82:1356–1360. [PubMed: 1415859]
23. Bacharach VR, Baumeister AA. Effects of maternal intelligence, marital status, income, and home environment on cognitive development of low birthweight infants. *J Pediatr Psychol* 1998;19:361–375.
24. Plomin R. Genetics and general cognitive ability. *Nature* 1999;402:C25–29. [PubMed: 10591222]
25. Stoolmiller M. Implications of the restricted range of family environments for estimates of heritability and nonshared environment in behavior-genetic adoption studies. *Psychol Bull* 1999;125:392–409. [PubMed: 10414224]
26. Dickens WT, Flynn JR. Heritability estimates versus large environmental effects: the IQ paradox resolved. *Psychol Rev* 2001;108:346–369. [PubMed: 11381833]
27. Gillberg C, Geijer-Karlsson M. Children born to mentally retarded women: a 1-21 year follow-up study of 41 cases. *Psychol Med* 1983;13:891–894. [PubMed: 6665104]
28. Feldman MA, Case L, Towns F, Betel J. Parent Education Project. I: Development and nurturance of children of mentally retarded parents. *Am J Ment Defic* 1985;90:253–258. [PubMed: 2417482]
29. Schwartz J. Low-level lead exposure and children's IQ: a meta-analysis and search for a threshold. *Environ Res* 1994;65:42–55. [PubMed: 8162884]
30. Pocock SJ, Smith M, Baghurst P. Environmental lead and children's intelligence: a systematic review of the epidemiological evidence. *BMJ* 1994;309:1189–1197. [PubMed: 7987149]
31. Lanphear BP, Hornung R, Khoury J, Yolton K, Baghurst P, Bellinger DC, Canfield RL, Dietrich KN, Bornschein R, Greene T, Rothenberg SJ, Needleman HL, Schnaas L, Wasserman G, Graziano J, Roberts R. Low-level environmental lead exposure and children's intellectual function: an international pooled analysis. *Environ Health Perspect* 2005;113:894–899. [PubMed: 16002379]
32. Offord DR, Boyle MH, Szatmari P, Rae-Grant NI, Links PS, Cadman DT, Byles JA, Crawford JW, Blum HM, Byrne C, et al. Ontario Child Health Study. II. Six-month prevalence of disorder and rates of service utilization. *Arch Gen Psychiatry* 1987;44:832–836. [PubMed: 3498458]
33. Deng S, Liu X, Roosa MW. Agreement between parent and teacher reports on behavioral problems among Chinese children. *J Dev Behav Pediatr* 2004;25:407–414. [PubMed: 15613989]
34. Kumpulainen K, Rasanen E, Henttonen I, Moilanen I, Piha J, Puura K, Tamminen T, Almqvist F. Children's behavioural/emotional problems: a comparison of parents' and teachers' reports for elementary school-aged children. *Eur Child Adolesc Psychiatry* 1999;8(Suppl 4):41–47. [PubMed: 10654132]
35. Gagnon C, Vitaro F, Tremblay RE. Parent-teacher agreement on kindergarteners' behavior problems: a research note. *J Child Psychol Psychiatry* 1992;33:1255–1261. [PubMed: 1400707]
36. Harder DW, Greenwald DF. Parent, family interaction, and child predictors of outcome among sons at psychiatric risk. *J Clin Psychol* 1992;48:151–164. [PubMed: 1573013]
37. Greenwald DF. Child functioning predictors of outcome among boys at risk for psychological disorder. *J Genet Psychol* 1990;151:139–151. [PubMed: 2388050]
38. Schachar R, Rutter M, Smith A. The characteristics of situationally and pervasively hyperactive children: implications for syndrome definition. *J Child Psychol Psychiatry* 1981;22:375–392. [PubMed: 7053050]
39. Broman S, Nichols PL, Shaughnessy P, Kennedy W 1987 Retardation in young children: a developmental study of cognitive deficit. Lawrence Erlbaum Associates, Hillsdale, NJ, pp 24–37.

40. Renick MJ, Harter S. Impact of social comparisons on the developing self-perceptions of learning disabled students. *J Educ Psychol* 1989;81:631–638.

Table 1
Demographic characteristics and blood lead levels of 656 TLC children at baseline and follow-ups

Characteristics	% or mean±SD
Clinic center	
Baltimore	29
Newark	26
Philadelphia	20
Cincinnati	24
African-American	77
Boys	54
English-speaking	95
Parent's education less than 12 years	40
Neither parent working	59
Living with single Parent	74
Succimer (treatment) group	51
Blood lead level (µg/dL)	
Baseline (n=656)	26±5
36 months follow-up (n=628)	12±5
60 months follow-up (n=530 [*])	8±4
Age (years)	
Baseline	2.0±0.5
36 months follow-up blood lead and IQ	5.0±0.5
36 months follow-up CPRS-R	5.1±0.5
36 months follow-up NEPSY	5.2±0.5
60 months follow-up blood lead	7.0±0.2
60 months follow-up IQ, CVLTC and BASC-PRS	7.1±0.2
60 months follow-up BASC-TRS	7.3±0.4
60 months follow-up NEPSY, CPT, WLPB-R and NESS	7.5±0.2

* One child with unexpectedly high blood lead level of 51 µg/dL at 60 months follow-up was excluded from the analysis of 7 year outcomes.

Table 2
Neuropsychological and behavioral test scores at age 5 by maternal IQ and child 5y IQ

Neuropsychological and behavioral test at age 5	Maternal IQ <70						Maternal IQ ≥ 70					
	Child 5y IQ <85			Child 5y IQ ≥ 85			Child 5y IQ <85			Child 5y IQ ≥ 85		
	n	Mean±SD	Difference* β (SE)	n	Mean±SD	Difference* β (SE)	n	Mean±SD	n	Mean±SD	Difference* β (SE)	
NEPSY												
Attention and executive	72	80±19		25	93±17	13.7 (4.5)	245	83±18	222	94±16	9.9 (1.7)	
Language	66	74±10		25	87±16	13.1 (3.1)	236	79±12	217	92±13	11.7 (1.3)	
Sensorimotor	75	80±15		24	93±16	11.7 (3.8)	257	81±15	217	93±14	10.7 (1.4)	
Visuospatial	76	84±12		25	96±10	10.0 (2.9)	260	85±11	225	97±11	10.8 (1.1)	
Memory	77	77±14		24	93±13	13.4 (3.3)	252	84±14	219	97±12	11.0 (1.3)	
CPRS-R												
Oppositional index	83	61±15		25	56±14	-4.8 (3.5)	282	59±14	226	55±13	-4.4 (1.3)	
Hyperactivity index	83	67±14		25	62±13	-4.0 (2.8)	282	65±14	226	61±14	-2.6 (1.2)	
ADHD index	83	62±13		25	56±11	-5.5 (2.7)	282	63±13	226	57±12	-5.8 (1.1)	
Behavioral index	83	63±13		25	58±11	-4.8 (2.6)	282	62±12	226	58±12	-4.3 (1.1)	

* Difference is the difference of child 5y IQ≥85 minus child 5y IQ<85, adjusted for clinic center, race, sex, language, parent's education, parent's employment, single parent, concurrent blood lead level, and maternal IQ

Table 3
Neuropsychological and behavioral test scores at age 7 by maternal IQ and child 7y IQ

Neuropsychological and behavioral test at age 7	Maternal IQ <70						Maternal IQ ≥ 70					
	Child 7y IQ <85			Child 7y IQ ≥85			Child 7y IQ <85			Child 7y IQ ≥85		
	n	Mean±SD	Difference* β (SE)	n	Mean±SD	Difference* β (SE)	n	Mean±SD	Difference* β (SE)	n	Mean±SD	Difference* β (SE)
NEPSY												
Attention/executive functions	63	75±16	17.7 (3.8)	29	94±13	17.7 (3.8)	154	76±13	17.7 (3.8)	257	95±15	17.0 (1.4)
CPT d-Prime	56	59±8	-4.7 (2.4)	28	53±12	-4.7 (2.4)	150	58±10	-4.7 (2.4)	250	54±10	-3.8 (1.1)
CVL/TC												
List A memory	68	38±10	5.1 (2.4)	32	43±13	5.1 (2.4)	170	40±10	5.1 (2.4)	277	47±11	5.9 (1.1)
Learning Slope	68	-0.6±1.1	0.2 (0.3)	32	-0.4±1.5	0.2 (0.3)	170	-0.5±1.1	0.2 (0.3)	277	-0.3±1.1	0.2 (0.1)
WLPB-R												
Broad Reading score	64	83±17	17.3 (4.4)	28	99±18	17.3 (4.4)	157	83±16	17.3 (4.4)	258	103±15	18.1 (1.6)
CPT Hit response time	56	42±15	3.2 (3.8)	28	45±10	3.2 (3.8)	150	41±13	3.2 (3.8)	250	44±12	2.5 (1.4)
NESS												
Sequential movement time	57	1.1±1.2	0.2 (0.3)	28	1.3±1.3	0.2 (0.3)	146	1.4±1.4	0.2 (0.3)	250	0.7±1.1	-0.7 (0.1)
BASC-TRS												
Adaptive skills	59	41±8	7.1 (2.2)	26	48±8	7.1 (2.2)	142	44±8	7.1 (2.2)	221	50±10	5.4 (1.1)
Behavioral Symptoms Index	58	60±12	-7.5 (3.3)	26	51±11	-7.5 (3.3)	146	57±11	-7.5 (3.3)	227	51±11	-4.9 (1.2)
Externalizing problems	59	61±15	-5.6 (4.1)	26	53±14	-5.6 (4.1)	146	57±12	-5.6 (4.1)	228	53±12	-3.2 (1.3)
Internalizing problems	59	54±9	-2.4 (2.7)	26	51±9	-2.4 (2.7)	146	54±10	-2.4 (2.7)	228	51±9	-3.2 (1.1)
School problems	58	65±13	-9.9 (3.6)	26	53±11	-9.9 (3.6)	147	60±12	-9.9 (3.6)	228	52±11	-7.5 (1.2)
BASC-PRS												
Adaptive skills	69	42±11	5.1 (2.9)	32	45±13	5.1 (2.9)	169	44±11	5.1 (2.9)	278	48±10	3.4 (1.1)
Behavioral Symptoms Index	69	58±15	-3.4 (3.6)	32	53±13	-3.4 (3.6)	169	58±16	-3.4 (3.6)	278	54±15	-3.7 (1.6)
Externalizing problems	69	60±17	-3.0 (3.9)	32	55±12	-3.0 (3.9)	169	62±16	-3.0 (3.9)	278	56±14	-5.1 (1.5)
Internalizing problems	69	51±14	1.0 (3.2)	32	51±10	1.0 (3.2)	169	50±13	1.0 (3.2)	278	50±12	0.2 (1.3)

* Difference is the difference of child 7y IQ ≥ 85 minus child 7y IQ <85, adjusted for clinic center, race, sex, language, parent's education, parent's employment, single parent, concurrent blood lead level, and maternal IQ

Table 4
Percentage of behavior test scores ≥ 60 and ≥ 70 by maternal IQ and child IQ at the age of behavior tests

	Maternal IQ <70						Maternal IQ ≥ 70					
	Child IQ <85			Child IQ ≥ 85			Child IQ <85			Child IQ ≥ 85		
	n	% score ≥ 60	% score ≥ 70	n	% score ≥ 60	% score ≥ 70	n	% score ≥ 60	% score ≥ 70	n	% score ≥ 60	% score ≥ 70
CPRS-R												
Oppositional index	83	49	29	25	40	20	282	41	25	226	32	17
Hyperactivity index	83	64	37	25	60	20	282	61**	32*	226	51	26
ADHD index	83	51	33	25	36	12	282	55*	28**	226	34	15
Behavioral index	83	57*	29	25	32	20	282	50	27**	226	40	14
BASC-TRS												
Behavioral Symptoms Index	58	50	26	26	23	8	146	34*	15	227	20	8
Externalizing problems	59	49*	24	26	19	12	146	32	16	228	23	11
Internalizing problems	59	29	7	26	15	8	146	25	7	228	18	4
School problems	58	72**	45**	26	31	4	147	52**	22**	228	22	6
Both externalizing and internalizing problems	59	27	3	26	8	4	145	14	4	227	10	2
BASC-PRS												
Behavioral Symptoms Index	69	39	20	32	25	16	169	41	22*	278	29	13
Externalizing problems	69	42	26	32	31	16	169	49*	30**	278	34	16
Internalizing problems	69	23	6	32	19	3	169	20	5	278	17	7
Both externalizing and internalizing problems	69	18	4	32	16	3	169	17	5	278	12	3

* P<0.05,

** P<0.01 for comparison with children with IQ ≥ 85 under the same maternal IQ category, after adjustment for clinic center, race, sex, language, parent's education, parent's employment, single parent, and concurrent blood lead level.

Table 5
Neuropsychological and behavioral test scores at age 5 by maternal IQ, child IQ or both*

Neuropsychological and behavioral test at age 5	n	Mean±SD	Model 1		Model 2		Model 3	
			10 point Maternal IQ β (SE)	10 point Child 5y IQ β (SE)	10 point Maternal IQ β (SE)	10 point Child 5y IQ β (SE)		
NEPSY								
Attention and executive	565	87±18	2.5 (0.8)	5.7 (0.6)	0.7 (0.7)	5.6 (0.7)		
Language	545	84±14	3.2 (0.6)	6.6 (0.4)	1.2 (0.5)	6.3 (0.5)		
Sensorimotor	574	86±16	1.3 (0.6)	5.9 (0.5)	-0.6 (0.6)	6.0 (0.5)		
Visuospatial	587	90±13	1.5 (0.5)	5.8 (0.4)	-0.5 (0.4)	5.9 (0.4)		
Memory	573	88±15	3.5 (0.6)	6.8 (0.4)	1.3 (0.5)	6.5 (0.5)		
CPRS-R								
Oppositional index	619	58±14	-1.2 (0.6)	-1.4 (0.4)	-0.8 (0.6)	-1.2 (0.4)		
Hyperactivity index	619	64±14	-0.9 (0.5)	-1.4 (0.4)	-0.4 (0.5)	-1.3 (0.4)		
ADHD index	619	60±13	-0.9 (0.5)	-2.6 (0.4)	0.1 (0.5)	-2.6 (0.4)		
Behavioral index	619	61±12	-1.0 (0.5)	-1.8 (0.4)	-0.4 (0.5)	-1.7 (0.4)		

* Model 1, model 2 and model 3 all adjusted for clinic center, race, sex, language, parent's education, parent's employment, single parent, and concurrent blood lead level.

Table 6
Neuropsychological and behavioral test scores at age 7 by maternal IQ, child IQ or both*

	n	Mean±SD	Model 1		Model 2		Model 3	
			10 point Maternal IQ β (SE)	10 point Child 7y IQ β (SE)	10 point Maternal IQ β (SE)	10 point Child 7y IQ β (SE)		
NEPSY								
Attention/executive functions	505	87±17	3.3 (0.7)	8.3 (0.5)	0.1 (0.6)	8.3 (0.5)	8.3 (0.5)	
CPT d-Prime	487	56±10	-0.5 (0.5)	-2.1 (0.4)	0.4 (0.5)	-2.2 (0.4)	-2.2 (0.4)	
CVLTC								
List A memory	551	44±11	1.9 (0.5)	3.9 (0.4)	0.5 (0.5)	3.7 (0.4)	3.7 (0.4)	
Learning Slope	551	-0.4±1.1	0.0 (0.1)	0.1 (0.0)	0.0 (0.1)	0.1 (0.0)	0.1 (0.0)	
WLPB-R								
Broad Reading score	510	94±19	4.7 (0.8)	9.4 (0.5)	1.0 (0.7)	9.1 (0.5)	9.1 (0.5)	
CPT Hit response time	487	43±13	0.3 (0.6)	1.2 (0.5)	-0.2 (0.6)	1.2 (0.5)	1.2 (0.5)	
NESS								
Sequential movement time	483	1±1.3	-0.1 (0.1)	-0.3 (0.0)	0.0 (0.1)	-0.3 (0.1)	-0.3 (0.1)	
BASC-TRS								
Adaptive skills	452	46±10	1.4 (0.5)	2.8 (0.4)	0.5 (0.5)	2.6 (0.4)	2.6 (0.4)	
Behavioral Symptoms Index	461	54±12	-1.4 (0.5)	-2.4 (0.4)	-0.5 (0.6)	-2.3 (0.4)	-2.3 (0.4)	
Externalizing problems	463	55±13	-0.8 (0.6)	-1.3 (0.5)	-0.3 (0.6)	-1.2 (0.5)	-1.2 (0.5)	
Internalizing problems	463	52±10	-0.8 (0.5)	-1.5 (0.4)	-0.2 (0.5)	-1.5 (0.4)	-1.5 (0.4)	
School problems	463	56±13	-2.8 (0.6)	-4.2 (0.4)	-1.2 (0.6)	-3.9 (0.4)	-3.9 (0.4)	
BASC-PRS								
Adaptive skills	553	46±11	1.8 (0.5)	2.4 (0.4)	1.0 (0.5)	2.2 (0.4)	2.2 (0.4)	
Behavioral Symptoms Index	553	56±15	-1.7 (0.7)	-2.3 (0.5)	-0.8 (0.7)	-2.1 (0.6)	-2.1 (0.6)	
Externalizing problems	553	58±15	-1.0 (0.7)	-2.2 (0.5)	-0.2 (0.7)	-2.2 (0.6)	-2.2 (0.6)	
Internalizing problems	553	50±12	-1.2 (0.6)	-0.6 (0.5)	-1.0 (0.6)	-0.3 (0.5)	-0.3 (0.5)	

* Model 1, model 2 and model 3 all adjusted for clinic center, race, sex, language, parent's education, parent's employment, single parent, and concurrent blood lead level.