

Norwalk-like viruses: When the runs can slow you down

Background and epidemiology: In 2002, outbreaks of gastroenteritis caused by Norwalk-like viruses (NLVs) occurred on a cruise ship bound for Alaska,¹ a trans-Canada VIA Rail train² and a Christian camp northwest of Vancouver.³ In the case of the cruise ship, 13% of the estimated 1266 passengers and crew were affected, and the ship had to be taken out of service for aggressive sanitation and disinfection procedures.¹

Norwalk virus was first identified in 1972 after an outbreak of gastrointestinal illness in Norwalk, Ohio. Other viruses of the Caliciviridae family have subsequently been identified as causing similar Norwalk virus illnesses, and the term “Norwalk-like viruses” refers to any members of this family.

NLV infections occur year-round and are the leading cause (in over 65% of cases) of nonbacterial outbreaks of gastroenteritis occurring in Canada and the United States.⁴ NLVs are highly contagious (a person can become infected with fewer than 100 viral particles). Virus transmission is primarily fecal–oral. Affected individuals typically drink contaminated water (including poorly maintained municipal water supplies, wells, recreational lakes, swimming pools and water stored aboard cruise ships) or eat food that has been handled by an infected food service worker or harvested or washed in con-

taminated water (e.g., shellfish, fruit and salad ingredients).

Symptoms of nonbloody diarrhea, nausea, vomiting, abdominal cramps, myalgias, malaise, headache and low-grade fever appear 24–48 hours after infection and last 12–60 hours. During this acute stage and until about 48 hours after the symptoms resolve, the disease is believed to be communicable.⁵ Secondary person-to-person spread is facilitated by enclosed living quarters and reduced levels of personal hygiene and commonly occurs through hand-to-hand contact with fecally contaminated skin, fomites (any contaminated object such as a cutting board, door handle or toys) or spattering and aerosols of infectious material from soiled linens and clothes. Anyone can be infected by NLV, but outbreaks typically occur in institutional settings such as nursing homes, day-care centres and schools, and other group settings such as banquet halls, cruise ships, dormitories and campgrounds.

NLVs are very stable and withstand freezing, heating to 60°C and disinfection with chlorine (at concentrations below 10 ppm).⁶ They are also resistant to acidic conditions, vinegar, alcohol and high sugar concentrations, which enables their survival on fruits and prepared foods.

Clinical management: In isolated cases of acute gastroenteritis other etiologic agents need to be considered, including other viruses (rotaviruses, adenoviruses, caliciviruses and astroviruses), bacteria (*Salmonella*, *Shigella* and *Escherichia coli*) and parasites (*Giardia*). Rotavirus is the most common cause of diarrhea in infants and children less than 5 years old, whereas adenoviruses and astroviruses are common causes of diarrhea in young children; NLVs typically affect older children and adults.⁷ Vomiting is more common than diarrhea in children with NLV infection.⁸ The diagnosis of NLV outbreaks is usually based on a patient’s clinical presentation, and specific diagnostic tests are often not

needed. During suspected outbreaks refrigerated stool or vomit specimens collected during the acute phase of the illness can be examined by means of electron microscopy for NLV particles. Serologic testing can also be useful to detect a specific viral strain. Seroconversion, defined as more than a 4-fold rise in IgG antibody titre during the acute and convalescent phases, is indicative of recent infection.⁸ NLVs are not routinely detected in water, food or environmental specimens, but during an outbreak such specimens should be tested by means of electron microscopy. Nucleic acid hybridization assays and polymerase chain reaction techniques are not yet widely used for outbreak investigation.⁸

No specific antiviral medications are currently indicated, and the emphasis for treatment should be on avoiding dehydration by means of fluid and electrolyte replacement. Oral rehydration is preferred over parenteral except in cases of severe dehydration. Complications of NLV infection, including severe dehydration and electrolyte imbalances, are more common in elderly people, malnourished children and immunocompromised people, but they are rarely fatal.⁸ Long-term sequelae of NLV infection have not been reported. Although acute symptoms typically last 1–3 days, patients may feel debilitated for 2–3 weeks after becoming ill.

Prevention: No vaccine against NLVs is currently available. Outbreaks are preventable and controlled by practising good sanitation and hygiene. Raw fruits and vegetables should always be washed thoroughly with clean water, and shellfish should be thoroughly cooked (a temperature of 85°C–90°C for 90 seconds is required to destroy viruses).⁹

Once an outbreak has occurred, the challenge of implementing control measures must be tackled.¹⁰ Food handlers with symptoms of vomiting or diarrhea should be excluded from work immediately and should not return to work until at least 48 hours after the



Concentrated living quarters of cruise ships can facilitate large-scale outbreaks of gastroenteritis caused by Norwalk-like viruses.

symptoms stop. Epidemic diarrhea occurring in an institutional setting or other group settings is a reportable disease in most jurisdictions, and in the event of an outbreak the local medical officer of health should be notified immediately. Affected individuals should be isolated from group activities until they are symptom free for 48 hours. Infection control precautions, including the use of gloves, masks and gowns, should be used by staff when caring for ill people or cleaning contaminated areas. Aggressive sanitization and disinfection of commonly touched surfaces should be implemented. Soiled clothes and linens should be handled with minimal agitation (to prevent aerosolization of viral particles) and laundered with detergent on a hot water cycle. Immunity to NLV lasts only up to 14 days, which makes people vulnerable to repeat infections throughout their lifetime and creates a challenge for the development of future vaccines.

Allison Chris
Ottawa, Ont.

This article has been peer reviewed.

Allison Chris is a medical student at the University of Ottawa with a special interest in public health.

References

1. *Outbreak of Norwalk virus: Alaska/British Columbia (update)* [Infectious Diseases News Brief]. Ottawa: Population and Public Health Branch, Health Canada; 2002 Aug 9. Available: www.hc-sc.gc.ca/pphb-dgspsp/bid-bmi/dsd-dsm/nb-ab/2002/nb3202_e.html (accessed 2002 Nov 22).
2. *Norwalk-like virus: Canada* [Infectious Diseases News Brief]. Ottawa: Population and Public Health Branch, Health Canada; 2002 Aug 2. Available: www.hc-sc.gc.ca/pphb-dgspsp/bid-bmi/dsd-dsm/nb-ab/2002/nb3102_e.html (accessed 2002 Nov 25).
3. *Virus hits kids' camp*. Guelph (ON): Food Safety Network, University of Guelph; 2002 July 26. Available: 131.104.232.9/fsnet/2002/7-2002/fsnet_july_26.htm#VIRUS%20HITS (accessed 2002 Nov 25).
4. Bidawid S. *Viruses in food contamination* [telelecture]. Ottawa: Food Directorate, Health Canada; June 2002. Available (PDF format): www.alphaweb.org/userfiles/filespublication/file_291611_ALPHA02F.pdf (accessed 2002 Nov 21).
5. Thornhill TS, Kalica AR, Wyatt RG, Kapikian AZ, Chanock RM. Pattern of shedding of the Norwalk particles in stools during experimentally induced gastroenteritis in volunteers as determined by immune electron microscopy. *J Infect Dis* 1975;132:28-34.
6. Keswick BH, Satterwhite TK, Johnson PC, DuPont HL, Secor SL, Bitsura JA, et al. Inactivation of Norwalk virus in drinking water by chlorine. *Appl Environ Microbiol* 1985;50:261-4.
7. *Viral gastroenteritis* [fact sheet]. Respiratory and Enteric Viruses Branch, National Center for Infectious Diseases. Available: www.cdc.gov/ncidod/dvrd/revb/gastro/faq.htm (revised 2001 Aug 20; accessed 2002 Nov 21).
8. Norwalk-like viruses: public health consequences and outbreak management. *MMWR Morb Mortal Wkly Rep* 2001;50(RR-9). Available (PDF format): www.cdc.gov/mmwr/pdf/rr/rr5009.pdf (accessed 2002 Nov 21).
9. Institute of Food Science and Technology: *Food-borne viral infections*. London (UK): The Institute; 2001 Jan 23. Available: www.ifst.org/hotspot12.htm (accessed 2002 Nov 21).
10. *Norwalk-like-virus outbreak interventions*. Vancouver: BC Centre for Disease Control; July 2002. Available (PDF format): www.bccdc.org/download.php?item=779 (accessed 2002 Nov 21).

Public Health: Authors Wanted

We're interested in receiving well-written submissions from public health experts for CMAJ's Public Health page. Submissions should adhere to the format presented above and should provide information applicable to the primary care setting. We encourage potential authors to contact us at pubs@cma.ca to discuss possible topics. — The editors

PHYSICIAN MANAGER INSTITUTE

A five level credit program exclusively for physicians designed to develop superior leadership and management skills

Approved for RCPSC, CFPC, CCHSE credits

cma.ca

FOR INFORMATION:
tel 800 663-7336 or 613 731-8610
x2319 (PMI) or x2261 (In-house PMI)
professional_development@cma.ca

2003 WORKSHOP SCHEDULE

PMI-I / PMI-II

Feb. 23-25 / Feb. 26-28	Victoria, BC
Mar. 23-25 / Mar. 26-28	Toronto, ON
June 1-3 / June 4-6	Ottawa, ON
Sept. 21-23 / Sept. 24-26	Montréal, QC

PMI-III / PMI-IV


Apr. 27-29 / Apr. 30-May 2	Niagara-on-the-Lake, ON
Nov. 2-4 / Nov. 5-7	Vancouver, BC

PMI Refresher

Oct. 24-26	Toronto, ON
------------	-------------

In-house PMI

A practical, cost effective and focused training opportunity held on-site for medical leaders and managers



ASSOCIATION
MÉDICALE
CANADIENNE



CANADIAN
MEDICAL
ASSOCIATION