

Dual role of the exocyst in AMPA receptor targeting and insertion into the postsynaptic membrane

Nashaat Z Gerges¹, Donald S Backos¹,
Chamila N Rupasinghe², Mark R Spaller²
and José A Esteban^{1,*}

¹Department of Pharmacology, University of Michigan Medical School, Ann Arbor, MI USA and ²Department of Chemistry, Wayne State University, Detroit, MI, USA

Intracellular membrane trafficking of glutamate receptors at excitatory synapses is critical for synaptic function. However, little is known about the specialized trafficking events occurring at the postsynaptic membrane. We have found that two components of the exocyst complex, Sec8 and Exo70, separately control synaptic targeting and insertion of AMPA-type glutamate receptors. Sec8 controls the directional movement of receptors towards synapses through PDZ-dependent interactions. In contrast, Exo70 mediates receptor insertion at the postsynaptic membrane, but it does not participate in receptor targeting. Thus, interference with Exo70 function accumulates AMPA receptors inside the spine, forming a complex physically associated, but not yet fused with the postsynaptic membrane. Electron microscopic analysis of these complexes indicates that Exo70 mediates AMPA receptor insertion directly within the postsynaptic density, rather than at extrasynaptic membranes. Therefore, we propose a molecular and anatomical model that dissects AMPA receptor sorting and synaptic delivery within the spine, and uncovers new functions of the exocyst at the postsynaptic membrane.

The EMBO Journal (2006) 25, 1623–1634. doi:10.1038/sj.emboj.7601065; Published online 6 April 2006

Subject Categories: membranes & transport; neuroscience

Keywords: Exo70; hippocampus; membrane trafficking; Sec8; synaptic plasticity

Introduction

The exocyst is a multiprotein complex originally described in yeast to be required for exocytosis (TerBush *et al.*, 1996). It consists of eight different subunits (Sec3, Sec5, Sec6, Sec8, Sec10, Sec15, Exo70 and Exo84), whose mammalian counterparts have already been identified (Ting *et al.*, 1995; Hsu *et al.*, 1996; Kee *et al.*, 1997). The exocyst is localized at the plasma membrane and intracellular compartments (Shin *et al.*, 2000; Yeaman *et al.*, 2001), and it associates with cytoskeletal

elements involved in vesicular trafficking (Hsu *et al.*, 1998; Shin *et al.*, 2000; Vega and Hsu, 2001; Wang *et al.*, 2004). The exocyst is broadly expressed in adult brain and is enriched in synaptic membranes (Hsu *et al.*, 1996; Sans *et al.*, 2003; Vik-Mo *et al.*, 2003). However, its role in neurons has always been associated with membrane trafficking events required for neuronal development, such as neurite outgrowth (Vega and Hsu, 2001), synaptogenesis (Hazuka *et al.*, 1999) and dendritic trafficking of newly synthesized NMDA receptors (NMDARs) (Sans *et al.*, 2003). In contrast, the functional role of the exocyst in established synapses remains unclear.

Synaptic function is critically dependent on the targeting and delivery of a large number of different proteins at the postsynaptic membrane, including neurotransmitter receptors, ion channels, anchoring and cell adhesion molecules, and a variety of signal transduction modulators (Garner *et al.*, 2000; McGee and Brecht, 2003; Yamagata *et al.*, 2003; Kim and Sheng, 2004). AMPA-type glutamate receptors (AMPA receptors) mediate most excitatory synaptic transmission in the brain. Nevertheless, they are highly dynamic components of excitatory synapses. AMPARs can be added to or removed from synapses in an activity-dependent manner, leading to long-term potentiation (LTP) or depression (LTD) of synaptic transmission (for recent reviews, see Sheng and Lee, 2001; Barry and Ziff, 2002; Malinow and Malenka, 2002; Brecht and Nicoll, 2003; Esteban, 2003). In addition to this regulated trafficking, AMPARs continuously cycle in and out of synapses in an activity-independent manner, which requires NSF (Nishimune *et al.*, 1998; Song *et al.*, 1998; Luscher *et al.*, 1999), Hsp90 (Gerges *et al.*, 2004b) and NEEP21 (Steiner *et al.*, 2005) function. Multiple proteins have been described to interact with AMPAR subunits and affect their synaptic localization (for recent reviews, see Scannevin and Huganir, 2000; Henley, 2003; Kim and Sheng, 2004); however, we still know very little about the membrane trafficking mechanism mediating AMPAR targeting and delivery into synapses.

Here we have explored a potential role for the exocyst in the dynamic behavior of AMPARs at the postsynaptic membrane. Using a combination of molecular biology, electrophysiology, fluorescence imaging and electron microscopy, we have found that different subunits of the exocyst mediate AMPAR synaptic trafficking at several distinct levels. Intracellular receptor targeting within the spine is mediated by the Sec8 subunit of the exocyst, possibly through PDZ-dependent interactions with synaptic scaffolding molecules. Independently from vesicle targeting, receptor insertion into the postsynaptic membrane requires the Exo70 subunit. By separately manipulating vesicle targeting and receptor insertion, we now show that AMPAR synaptic delivery occurs from intracellular compartments within the spine directly into the postsynaptic density. Therefore, in addition to characterizing the exocytic machinery that operates at the

*Corresponding author. Department of Pharmacology, University of Michigan Medical School, 1150 W Medical Center Drive, Ann Arbor, MI 48109-0632, USA. Tel.: +1 734 615 2686; Fax: +1 734 763 4450; E-mail: estebanj@umich.edu

Received: 13 December 2005; accepted: 3 March 2006; published online: 6 April 2006

postsynaptic membrane, these results delineate the subcellular pathways by which AMPARs reach their synaptic targets.

Results

Different components of the exocyst play distinct roles in synaptic transmission

The C-terminus of the exocyst subunit Sec8 contains a PDZ motif that interacts with the synaptic scaffolding proteins PSD95 and SAP102 (Riefler *et al*, 2003; Sans *et al*, 2003) and is required for NMDAR trafficking (Sans *et al*, 2003). Therefore, as a first step to evaluate the role of the exocyst in synaptic transmission, we expressed the C-terminus (last 16 amino acids) of Sec8 as a GFP-fusion protein in CA1 hippocampal neurons from organotypic slice cultures using the Sindbis virus system (see Supplementary data). This construct may act as a dominant negative by competing with the endogenous exocyst for its PDZ synaptic partners. The effect of Sec8 C-terminus (Sec8-Ct) on synaptic transmission was evaluated by simultaneous double whole-cell recordings from neurons expressing the recombinant protein and nearby uninfected neurons (Methods). As shown in Figure 1A, Sec8-Ct significantly reduced AMPAR and NMDAR responses in hippocampal CA1 neurons, while GABA_A receptor-mediated transmission remained unaffected. The depression of NMDA responses and lack of effect on GABA_A responses are consistent with previously published observations using another Sec8 dominant negative construct (Sans *et al*, 2003). As control, overexpression of the full-length Sec8 protein (Sec8-wt) fused to GFP did not affect AMPA or NMDA transmission (Figure 1B). Importantly, expression of a truncated form of Sec8-Ct lacking its last four amino acids (Sec8-Ct-Δ4), which are required for its PDZ interactions (Sans *et al*, 2003), did not alter AMPA or NMDA responses either (Figure 1C). These results suggest that the dominant negative effect of Sec8-Ct is indeed due to competition for PDZ-binding domains, which would be required for the synaptic delivery or stability of both AMPARs and NMDARs.

Although both AMPARs (Malinow and Malenka, 2002; Bredt and Nicoll, 2003) and NMDARs (Nong *et al*, 2004; Perez-Otano and Ehlers, 2005; Prybylowski *et al*, 2005) undergo regulated synaptic trafficking, NMDARs are considered to be fairly stable components of the synapse, as compared with AMPARs (Sheng, 2001). Therefore, interference with a potential role of the exocyst specifically in receptor delivery is more likely to affect AMPARs to a larger extent than NMDARs. A truncated form of the exocyst subunit Exo70, lacking its C-terminal half (Exo70-N), has been recently shown to block the exocytic delivery of the glucose transporter Glut4 into the adipocyte cell surface (Inoue *et al*, 2003). Importantly, this truncated Exo70 is still able to interact with endogenous exocyst subunits (see Supplementary Figure 1 and Inoue *et al*, 2003). We therefore evaluated the effect on synaptic transmission of Exo70-N as a GFP fusion protein in CA1 hippocampal neurons. As shown in Figure 1D, AMPAR-mediated responses were significantly reduced in Exo70-N-expressing neurons. Importantly, NMDAR responses were not affected, in contrast to the result obtained with Sec8-Ct. This result is also confirmed by the decrease in the ratio of AMPA to NMDA responses in Exo70-N-expressing neurons (Figure 1D; to note, AMPA/NMDA ratios are intrinsically very variable across slices; therefore, comparisons

between infected and uninfected cells were always drawn from recordings within the same slice, and in most cases, as simultaneous double recordings). As control, overexpression of full-length Exo70 (Exo70-wt) fused to GFP did not affect AMPA or NMDA synaptic responses (Figure 1E). The differing effects of Exo70-N on AMPA and NMDA synaptic transmission suggest that the Exo70 subunit of the exocyst mediates receptor insertion at synapses, which is more likely to affect the continuous synaptic delivery of AMPARs than the stable presence of NMDARs at synapses. (To note, none of these wild-type and dominant negative constructs affected passive membrane properties of the infected cell, such as input resistance and holding current; see Supplementary Figure 2).

Sec8 PDZ-dependent interactions are required for the constitutive synaptic cycling of AMPA receptors

The activity-independent, continuous delivery of AMPARs into synapses was originally inferred from intracellular peptide infusion experiments, in which AMPAR-mediated synaptic transmission rapidly ran down when a specific peptide competing GluR2-NSF interactions was loaded in the recorded cell (Nishimune *et al*, 1998; Luscher *et al*, 1999). We have used the same approach to determine whether the PDZ motif of Sec8 participates in the constitutive cycling of AMPARs. To this end, we recorded AMPAR-mediated synaptic responses from CA1 hippocampal neurons while infusing them with one of three peptides: Sec8 C-t (last eight amino acids of Sec8), NR2B C-t (last eight amino acids of the NMDAR subunit NR2B, which also contains a PDZ ligand motif (Kornau *et al*, 1995; Niethammer *et al*, 1996)) and Sec8-mut C-t (last eight amino acids of Sec8 with the PDZ ligand sequence -TTV mutated to -ATA) (see Supplementary data and Supplementary Figure 3). These experiments were carried out in a blind fashion, that is, the person carrying out the electrophysiological recordings and the analysis did not know which peptide was used in the experiment. As shown in Figure 2, the peptide corresponding to the Sec8 C-terminus produced a fast run-down of synaptic transmission, similar to the one observed for the GluR2-NSF peptide (Nishimune *et al*, 1998; Luscher *et al*, 1999; Gerges *et al*, 2004b). Importantly, the peptide corresponding to the PDZ ligand sequence of NR2B did not affect AMPAR-mediated responses, suggesting that specific PDZ-dependent interactions of Sec8 are required for the continuous synaptic cycling of AMPARs. As control, the mutated Sec8 C-t peptide lacking the PDZ motif did not affect synaptic transmission either. Figure 2B indicates that the differences in run-down of synaptic transmission obtained with the three peptides are not due to systematic changes in access resistance during the recordings.

The exocyst associates with AMPA receptors in hippocampal synaptosomes

Our data shown above suggest that the exocyst may be directly involved in AMPAR synaptic trafficking. Therefore, we tested whether an exocyst/AMPA complex is present in the brain. To this end, we carried out immunoprecipitation experiments from hippocampal synaptosomes after Triton solubilization (see Supplementary data; all immunoprecipitations were replicated in at least three independent experiments). As shown in Figure 3A, immunoprecipitation of the exocyst subunits Sec8 or Exo70 co-precipitated




Figure 1 Effect of different exocyst subunits on synaptic transmission. (A–E) Insets, sample trace of evoked AMPA-receptor-mediated synaptic responses recorded at -60 mV, from uninfected and infected cells. Scale bars, 20 pA and 40 ms. Left: average AMPAR-mediated current amplitude (i.e. the peak of the response recorded at -60 mV) from infected (inf) neurons expressing Sec8-Ct (A), Sec8-wt (B), Sec8-Ct- $\Delta 4$ (C), Exo70-N (D) or Exo70-wt (E), and control neighboring cells not expressing the recombinant protein (uninf). n represents the number of cell pairs. Middle: average NMDAR-mediated current amplitude (recorded at $+40$ mV at a latency when AMPA responses are fully decayed, 60 ms) from uninfected and infected cells (n also represents the number of cell pairs). Right: average GABA_A receptor-mediated current amplitude measured at 0 mV (A) or average AMPA/NMDA ratios for uninfected and infected cells (B–E).

AMPA receptors but not metabotropic (mGluR) receptors. As previously published (Riefler *et al*, 2003; Sans *et al*, 2003), the exocyst also co-precipitated the PDZ protein PSD95 (Figure 3A).

The exocyst complex has been recently shown to associate with NMDARs through its interaction with synaptic anchoring proteins, such as PSD95 and SAP102 (Sans *et al*, 2003). However, AMPA and NMDA receptors do not appear to associate in hippocampal extracts (Wyszynski *et al*, 1999; Sans *et al*, 2000) (see also Figure 3B). Therefore, we inves-

tigated whether the exocyst forms separate complexes with AMPARs and NMDARs at synaptic terminals. To this end, Triton-solubilized synaptosomal extracts from hippocampus were immunoprecipitated with anti-GluR2/3 (AMPA receptors) or anti-NR1 (NMDARs) antibodies (see Supplementary data). As shown in Figure 3B, the exocyst subunits Exo70, Sec8 and Sec6 co-precipitated predominantly with AMPARs, but only to low levels with NMDARs. This result supports the notion that the exocyst forms separate complexes with AMPARs and NMDARs.




Figure 2 Interference of the constitutive synaptic cycling of AMPA receptors with a Sec8 C-t peptide. (A) Time course of AMPAR-mediated responses recorded from CA1 neurons during whole-cell pipette infusion of Sec8 C-t, NR2B C-t or Sec8-mut C-t peptides, as indicated. Responses are normalized to a 2-min baseline from the beginning of the recording. The statistical significance of the run-down of AMPA transmission was calculated from the last 5 min of the recordings. (B) Time course of the pipette access resistance during the recordings shown in (A). There was no systematic change in access resistance with any of the peptides.




Figure 3 Co-immunoprecipitation of exocyst subunits and AMPA receptors from hippocampal synaptosomes. Exocyst subunits Exo70 and Sec8 (A), AMPA receptor subunits GluR2/3 (GR2/3) and NMDA receptor subunit NR1 (B) were immunoprecipitated from Triton-solubilized synaptosomal preparation of hippocampus. IP indicates antibodies used for immunoprecipitation. Non-immune (N.I.) antibodies were used as control. Total solubilized protein (10% of the input) and the immunoprecipitated fractions were analyzed by immunoblotting using the indicated antibodies.

A potential role for Sec8 but not for Exo70 in the dendritic trafficking of AMPA receptors

It has been recently described that the exocyst mediates the export into dendrites of newly synthesized NMDARs in primary neuronal cultures (Sans *et al*, 2003). In order to test whether the exocyst plays a similar role in the trafficking of AMPARs in hippocampal slices, we coexpressed a GFP-tagged




Figure 4 Exo70 is not involved in the dendritic trafficking of newly synthesized AMPA receptors in hippocampal slices. Hippocampal CA1 neurons were transfected with GluR2-GFP alone or cotransfected with GluR2-GFP plus Exo70-N-RFP, Sec8-Ct-RFP or NR2B-Ct-RFP (these constructs are equivalent to their GFP counterparts, only with RFP in the place of GFP; the NR2B construct includes the last 16 amino acids of the NR2B C-terminus). (A) Representative examples of neurons expressing GluR2-GFP alone or coexpressing GluR2-GFP together with Exo70-N-RFP or with Sec8-Ct-RFP, as indicated. Scale bar: 20 μ m. The efficiency of GluR2 transport along dendrites was quantified from the intensity of GFP fluorescence at the main apical dendrite 170–200 μ m away from the cell soma. Fluorescence intensity is normalized to the peak value at the soma after background subtraction (see arrows on left panel and representative scheme in Supplementary Figure 4A). (B) GluR2-GFP presence at distal dendrites (170–200 μ m away from the cell soma) in the presence of Exo70-N, Sec8-Ct or NR2B-Ct, as indicated. Data are presented as mean \pm s.e.m. *n* represents the number of cells.

AMPA subunit (GluR2-GFP), together with Exo70-N, Sec8 C-t or NR2B C-t (as control) tagged with a red fluorescence protein (Exo70-N-RFP, Sec8-Ct-RFP or NR2B-Ct-RFP; see Supplementary data). The efficiency of AMPAR transport along dendrites was quantified from the intensity of GFP fluorescence along the main apical dendrite, after background subtraction (see Supplementary Figure 4A). As shown in Figure 4, the trafficking of GluR2-GFP into distal dendrites was virtually identical with or without coexpressed Exo70-N, indicating that the transport of AMPARs along dendrites does not require a fully functional Exo70 subunit. In contrast, the dendritic transport of GluR2-GFP was significantly impaired when co-expressing Sec8-Ct or NR2B-Ct (Figure 4; see Supplementary Figure 4B for a complete quantification of AMPAR dendritic distribution in the presence of Exo70-N, Sec8-Ct and NR2B-Ct). These results reinforce the interpretation that Exo70 and Sec8 play different roles in AMPAR trafficking: Exo70 probably mediates a local step in AMPAR synaptic delivery, whereas PDZ-dependent interactions of

Sec8 or other proteins are required for efficient receptor transport and targeting (since both Sec8-Ct and NR2B-Ct impaired dendritic trafficking, the identity of these PDZ interactions cannot be determined from these experiments).

Morphological dissection of AMPA receptor delivery into the spine surface

In order to directly test a local role of the exocyst in AMPAR trafficking, we monitored the transport of AMPARs from the dendritic shaft into adjacent spines and their delivery into the spine plasma membrane. To this end, we coexpressed GFP-tagged GluR1 with a constitutively active, untagged form of α CaMKII (truncated α CaMKII, or tCaMKII) to trigger its synaptic delivery (see Supplementary data and Hayashi *et al*, 2000), with or without Exo70-N. Protein expression was allowed for 36 h, then slices were fixed and the surface distribution of the recombinant receptor in CA1 hippocampal neurons was monitored by immunostaining using an anti-GFP antibody under nonpermeabilized conditions (the GFP tag is placed at the extracellular N-terminus of the receptor). The immunostaining was visualized with an infrared fluorophore (Cy5). This experimental approach allows us to independently monitor the total amount of receptor (GFP channel), the fraction exposed to the cell surface (Cy5 channel) and the presence of the exocyst dominant negative construct (RFP channel). The distribution of surface and total receptor in spines and dendrites was quantified by line plots of fluorescence intensity across spine heads and adjacent dendritic shafts (see Figure 5A for representative examples of the quantification of GFP and surface immunostaining from dendrite–spine pairs with or without coexpressed Exo70-N). Surface ratios were then calculated as the ratio between the intensity of the Cy5 signal (surface receptor) and GFP signal (total receptor) after background subtraction (Gerges *et al*, 2004a; Brown *et al*, 2005). Importantly, this assay is internally normalized for immunostaining variability, because surface labeling is always acquired in pairs of spines and adjacent dendritic shafts. This also means that absolute changes in surface receptors at spines or dendrites cannot be evaluated, but only relative changes between the two compartments. Finally, spines are selected solely from their GFP image, so the analysis is done blind with respect to their surface immunostaining.

As shown in Figure 5B (first panel), surface expression of GluR1-GFP was slightly higher in spines as compared to their adjacent dendritic shafts when the exocyst function is intact. In contrast, Exo70-N led to a marked reduction in the amount of receptors that reach the spine surface, to the point that surface expression at spines became significantly lower than at dendrites (Figure 5B, second panel). This effect can also be evaluated from the leftward shift in the cumulative probability distribution of spine-to-dendrite ratios in the presence of Exo70-N (Figure 5B, third panel).

A reduction of surface expression at spines may be due to an impaired translocation of the receptor into the spine. This step can be evaluated from the total receptor distribution (GFP channel) at spines and adjacent dendrites. As shown in Figure 5C (first panel), GluR1-GFP partitions equally between dendrites and spines in the absence of exocyst dominant negative constructs. In marked contrast, and to our surprise, Exo70-N led to a significant increase (almost doubling) in the amount of total receptor in spines as compared to dendrites

(Figure 5C, second panel) (this effect can also be appreciated from the rightward shift in the cumulative probability distributions in Figure 5C, third panel). Therefore, the decrease in receptor surface expression produced by Exo70-N is accompanied by an enrichment of receptors at spines.

The next trafficking step after translocation into spines is receptor insertion at the spine plasma membrane. This process can be evaluated independently by calculating ratios of surface to total receptor (Cy5/GFP) at spines and dendrites. This approach normalizes receptor surface expression to the total number of receptors in each compartment, and therefore, it reflects the intrinsic efficiency of receptor delivery at the plasma membrane. As shown in Figure 5D (first panel), surface ratios were slightly higher at spines than at dendrites when GluR1-GFP was coexpressed only with tCaMKII. On the other hand, blockade of Exo70 function with Exo70-N strongly impaired receptor insertion at the spine surface, as judged from the drastic reduction in the surface ratio at spines versus dendrites (Figure 5D, second and third panel).

These results support our interpretation that Exo70 plays a late step in AMPAR trafficking, even within the local environment of the synapse. Specifically, Exo70 seems to be required for the fusion of AMPAR-containing vesicles with the spine plasma membrane.

Interestingly, the accumulation of GluR1-GFP at intracellular compartments within the spine when Exo70 function is impaired also suggests that this pool of AMPARs reaches the synaptic membrane from intracellular compartments contained within the spine. We then tested whether a similar intra-spine pathway is followed for the constitutive synaptic delivery of the cycling population of AMPARs. This pathway can be monitored by expressing GluR2-GFP, which has been shown to constitutively reach spine and synaptic compartments (Shi *et al*, 2001). Therefore, we expressed GluR2-GFP in CA1 hippocampal neurons with or without Exo70-N-RFP, Sec8-Ct-RFP or NR2B-Ct-RFP, as control. Total receptor (GFP channel) and surface (Cy5 channel) distribution of GluR2-GFP was evaluated as described above (see Figure 5A). As shown in Figure 6A–C (first and second panels), and similar to the results obtained with GluR1-GFP, interference with Exo70 function drastically impaired the surface delivery of GluR2 at the spine, and led to a concomitant enrichment of the receptor in the intracellular space of the spine. Interestingly, Sec8-Ct also strongly impaired surface receptor delivery (Figure 6A, third panel), but this was not accompanied by receptor accumulation within the spine. In fact, Sec8-Ct produced a significant decrease in the fraction of receptors present at spines (Figure 6B, third panel). As control, NR2B-Ct did not affect receptor presence at spines, and only modestly decreased surface delivery at spines (Figure 6, fourth panels; see also Supplementary Figure 5 for a comparative analysis of the effects of Sec8-Ct and NR2B-Ct at the spine–dendrite pairs).

Taken together, these results indicate that the exocyst plays specialized roles in AMPAR synaptic trafficking, with Sec8 assisting both receptor translocation into the spine and delivery into the spine surface, while Exo70 mediates exclusively a final membrane fusion step within the spine.

In addition, these results strongly suggest that the exocyst mediates both constitutive and regulated delivery of AMPARs at synapses. As a functional test for the role of the exocyst in these two pathways, we carried out electrophysiological experiments in which the presence of recombinant receptors




Figure 5 Role of Exo70 in local AMPAR trafficking at spines. (A) Representative confocal image of a spine and the adjacent dendritic shaft from a neuron transfected with GluR1-GFP and tCaMKII (left) or with GluR1-GFP, tCaMKII and Exo70-N-RFP (right), as indicated. GFP fluorescence signal (green) represents total GluR1 receptor distribution. Surface GluR1-GFP receptor (cyan) was assayed by immunostaining with an anti-GFP antibody coupled to Cy5 under nonpermeabilized conditions. Scale bar: 2 μ m. Lower panels: representative line plot profiles of GFP and Cy5 fluorescence intensities across dendrite–spine pairs. Total (GFP) and surface (Cy5) receptor accumulation are calculated from the corresponding peaks of fluorescence intensity at spines and adjacent dendritic shafts. (B) Surface analysis (background-subtracted Cy5 values) from pairs of spines and dendrites in GluR1 plus tCaMKII transfected neurons (first panel; $n = 59$ spine/dendrite pairs from five different cells) or in GluR1 plus tCaMKII and Exo70-N cotransfected neurons (second panel; $n = 40$ spine/dendrite pairs from five different cells). Surface values are normalized by the mean dendrite value. Cumulative probabilities (third panel) of surface expression at spine over dendrite show significant differences between surface distributions of GluR1 plus tCaMKII versus GluR1 plus tCaMKII and Exo70-N transfected neurons. (C) Total receptor amount (GFP signal) in spines and dendrites are calculated in GluR1 plus tCaMKII transfected neurons and in neurons cotransfected with GluR1 plus tCaMKII and Exo70-N, as indicated. GFP fluorescence is normalized to the mean dendrite value. The cumulative probability (right) indicates that Exo70-N leads to the accumulation of total receptor in spines as compared to dendrites. (D) Surface ratio from the same pairs of spines and dendrites. Surface ratios were calculated by dividing the corresponding background-subtracted Cy5 and GFP values. Surface ratios are normalized by the mean dendrite value. Cumulative probabilities (right) of the spine/dendrite ratios show a significant difference between the surface ratio distribution of GluR1 plus tCaMKII versus GluR1 plus tCaMKII and Exo70-N transfected neurons.

at synapses is detected from their inward rectification properties (Hayashi *et al*, 2000; Shi *et al*, 2001). These electrophysiological experiments reinforced our fluorescence imaging data, and confirmed that the exocyst is required for the functional insertion of all AMPAR populations at synapses (see Supplementary data and Supplementary Figure 6).

Distinct roles of Sec8 and Exo70 in targeting and insertion of AMPA receptors at the postsynaptic membrane

In yeast, different subsets of exocyst subunits have been proposed to associate with exocytic vesicles versus the target plasma membrane (Boyd *et al*, 2004). However, it is

unknown what exocyst subassemblies reside in each compartment in the case of synaptic trafficking. Furthermore, it is still uncertain whether the target membrane for AMPAR exocytic delivery is directly at the synaptic membrane contained within the postsynaptic density (PSD), or whether AMPARs are inserted in extrasynaptic membranes and then reach their synaptic targets by lateral diffusion (Passafaro *et al*, 2001; Tardin *et al*, 2003; Adesnik *et al*, 2005; Triller and Choquet, 2005). Since Exo70 is required for local surface delivery of AMPARs (Figures 5 and 6), we then tested whether interference with this exocyst subunit would alter the association of AMPARs with the PSD. To this end, we isolated PSD fractions (protocol based on Peng *et al* (2004);




Figure 6 Role of Exo70 and Sec8 in the constitutive AMPAR trafficking at spines. (A) Similar to Figure 5B, surface analysis from pairs of spines and dendrites in GluR2 (first panel; $n = 21$ spine/dendrite pairs from four different cells), GluR2 plus Exo70-N (second panel; $n = 32$ spine/dendrite pairs from five different cells), GluR2 plus Sec8-Ct (third panel; $n = 58$ spine/dendrite pairs from five different cells) or GluR2 plus NR2B-Ct cotransfected neurons (fourth panel; $n = 62$ spine/dendrite pairs from five different cells). (B) Similar to Figure 5C, total receptor amount from the same pairs of spines and dendrites in GluR2 or in GluR2 plus Exo70-N, Sec8-Ct or NR2B-Ct cotransfected neurons. (C) Similar to Figure 5D, surface ratio of the same pairs of spines and dendrites in GluR2 or in GluR2 plus Exo70-N, Sec8-Ct or NR2B-Ct cotransfected neurons.

see Supplementary data) from hippocampal extracts of control slices or slices overexpressing Exo70-N. To our surprise, and despite the fact that Exo70-N impairs receptor delivery into synapses (Figure 1D) and spine plasma membrane (Figures 5 and 6), Exo70-N overexpression led to a marked accumulation of Sec8, Rab8 and AMPARs at the PSD fraction (Figure 7A and B; to note, none of these proteins is particularly enriched at the PSD fraction under basal conditions). Importantly, endogenous Exo70 and stable components of the PSD, such as PSD95 and α CaMKII, were not affected by expression of Exo70-N (Figure 7A and B). Another exocytic Rab protein, Rab11, and the small GTPase Rho were undetectable at the PSD fraction under either condition. Therefore, these results together with those monitoring receptor delivery at the spine surface (Figures 5 and 6), strongly suggest that AMPARs are directly inserted at the postsynaptic membrane (or at membranes biochemically associated to the PSD) in an Exo70-dependent manner. Truncation of the Exo70 C-terminus would lead to formation of an abortive complex at the PSD, in which incoming vesicles containing AMPARs, Sec8 and Rab8 remained associated but not yet fused with the postsynaptic membrane.

Importantly, this interpretation also implies that AMPAR targeting towards the postsynaptic membrane does not require a fully functional Exo70. Since Sec8 is the exocyst subunit that interacts with synaptic scaffolding molecules (Riefler *et al*, 2003; Sans *et al*, 2003) and it contributes to the transport of AMPARs into spines (Figure 6), we then

tested whether Sec8 may be involved in the targeting aspects of AMPAR delivery into the postsynaptic membrane. To this end, we carried out similar PSD fractionation experiments with hippocampal extracts from uninfected slices, and from slices overexpressing Sec8-Ct. As shown in Figure 7C and D, Sec8-Ct led to a drastic reduction in the amount of AMPARs and Sec8 at the PSD fraction. Interestingly, the association of NMDARs and PSD95 with the PSD was also decreased, in agreement with the depression of NMDA responses produced by Sec8-Ct (Figure 1A). The same result was obtained for the actin-dependent motor protein myosin Va, which is enriched in dendritic spines (Naisbitt *et al*, 2000; Walikonis *et al*, 2000). However, Sec8-Ct did not produce a generalized disassembly of the postsynaptic complex since Exo70 and α CaMKII were not affected by Sec8-Ct overexpression. This result with Sec8-Ct is virtually opposite to the one observed with Exo70-N (compare quantifications in Figure 7B and D), and reinforces the interpretation that Sec8 mediates the targeting of AMPAR-containing vesicles towards the synapse, whereas Exo70 is required for the fusion of these vesicles with the postsynaptic membrane.

Ultrastructural analysis of the role of the exocyst in AMPA receptor insertion at the postsynaptic membrane

The results presented above strongly suggest that AMPARs are able to reach the PSD and accumulate within it when Exo70 function is blocked. In order to test anatomically this interpretation and map with high-spatial resolution the role




Figure 7 Opposite effects of Exo70 and Sec8 dominant negatives in the association of AMPA receptors with the postsynaptic density. **(A)** Total proteins in synaptosomes (8% of the total fraction) and PSD-associated proteins were analyzed by immunoblotting using the indicated antibodies (AMPA receptors were detected with a mix of GluR1 and GluR2/3 antibodies). The CA1 area was dissected from 30 to 45 control (unin.) hippocampal slices or slices infected with Exo70-N (inf.). Therefore, an 'n' of 1 in this experiment represents the average of 30–45 slices. Protein extracts were processed to isolate the synaptosomal and PSD fractions (see Supplementary data). **(B)** Quantification of multiple independent experiments as the one shown in (A). Relative PSD enrichment was calculated as the intensity of the PSD band in the infected extracts over that of the uninfected ones, after normalization by their relative band intensities at the homogenates (input). Inset shows that infection did not produce a systematic effect on protein abundance at the inputs. **(C)** Similar to (A), comparing uninfected and infected slices expressing Sec8-Ct (same construct as in Figure 1). Total proteins in synaptosomes (3% of the total fraction) and PSD-associated proteins were analyzed by immunoblotting using the indicated antibodies. NMDARs were detected with a combination of NR1 and NR2A/B antibodies. **(D)**. Quantification of PSD enrichment for the different proteins in the presence or absence of Sec8-Ct was calculated as described for (B).

of Sec8 and Exo70 in AMPAR synaptic insertion, we have carried out immunogold electron microscopy. Organotypic hippocampal slices overexpressing Exo70-N or Sec8-Ct, or control (uninfected) slices were processed for postembedding immunogold staining with anti-GluR2/3 antibodies (see Supplementary data). Electron micrographs were taken randomly from excitatory (asymmetric) synapses in the stratum radiatum (CA1) of these slices (see representative examples in Figure 8A). As shown in Figure 8A and B, AMPAR immunogold particles were significantly enriched (2.6-fold) within the PSD in Exo70-N-expressing slices. In contrast,

Sec8-Ct produced a reduction in the fraction of AMPAR labeling at the PSD, although the difference was not significant with respect to uninfected slices.

This result confirms our biochemical observations and strengthens the interpretation that AMPARs approach the postsynaptic membrane but fail to fuse with it in the absence of a functional Exo70. In order to precisely map the trafficking step mediated by Exo70, we measured the distance between each gold particle and the synaptic cleft from slices expressing Exo70-N, Sec8-Ct or uninfected slices. The results are presented as cumulative distributions in Figure 8C (only




Figure 8 Electron microscopic analysis of the role of the exocyst in AMPA receptor synaptic insertion. (A) Representative micrographs of AMPA receptor immunogold labeling in control slices (Uninf), slices infected with Exo70 dominant negative (Exo70-N) or with Sec8 dominant negative (Sec8-Ct; same construct as in Figure 1). Immunogold particles at the PSD are indicated with arrows. Note accumulation of AMPA receptors within the PSD in Exo70-N-expressing neurons. The opposite effect was observed in Sec8-Ct-infected slices. (B) Quantification of AMPA receptor presence at the PSD from electron micrographs as the ones shown in (A). Percentage of PSD labeling was calculated as the number of gold particles at the PSD versus the total number of gold particles in each synaptic terminal (only gold particles within 800 nm of the synaptic cleft were included for quantification). *n* represents the number of synapses analyzed in each condition. (C) Quantification of AMPA receptor local distribution with respect to the synaptic cleft. The distance to the synaptic cleft was calculated for each immunogold particle from uninfected slices (black line), slices expressing Exo70-N (red line) or Sec8-Ct (blue line), and plotted as cumulative distributions. In order to assess local synaptic distribution of AMPA receptors, only immunogold particles within 60 nm of the synaptic cleft were included in this quantification (the average thickness of the PSD from our micrographs was 18 nm-thick line on the *x*-axis). *n* represents the number of gold particles analyzed in each condition. (D) Quantification of AMPA receptor lateral distribution within the PSD and on extrasynaptic membranes from uninfected slices (white columns) and Exo70-N-expressing slices (red columns). The lateral distance to the closest edge of the PSD was calculated for each immunogold particle and divided by the total length of the PSD (the average length of the PSD from our micrographs was 290 nm). Distances within the PSD were computed as negative values (relative lateral distances within the PSD cannot exceed -0.5 , since the position of the immunogold particle is always calculated with respect to the closest PSD edge).

gold particles within 60 nm of the synaptic cleft—approximately three times the average PSD thickness—were considered for this quantification). As shown in Figure 8C, AMPARs were found to accumulate in close proximity to the synaptic cleft in slices expressing Exo70-N (red line; leftward shift of the cumulative distribution), even from the shortest measurable distances within the PSD. Sec8-Ct produced the opposite effect, that is, AMPARs were less abundant in the vicinity of the PSD (blue line; rightward shift in the cumulative distribution). These results indicate that, in the absence of functional Exo70, AMPARs approach the postsynaptic membrane as much as physically possible, within the resolution of the immunogold labeling technique.

Since Exo70-N seems to trap AMPARs at their final fusion step with the postsynaptic membrane, we then decided to map where this insertion takes place laterally with respect to the postsynaptic density. To this end, we measured the lateral distance to the edge of the PSD for all immunogold particles present within the PSD or on the lateral extrasynaptic membrane. As shown in Figure 8D, in the presence of Exo70-N, AMPARs predominantly accumulate within the PSD, but in close proximity to its edge (almost 40% of the labeling is found within the distal 10% of the PSD) (Sec8-Ct was not included in this analysis due to the low number of gold particles present at the PSD under those conditions). Once again, this analysis supports the interpretation that AMPARs are inserted into the synaptic membrane directly from the

PSD or close to its edge, in a process mediated by the Exo70 subunit of the exocyst.

Discussion

Delivery of AMPARs into synapses requires a precise and continuous coordination between sorting of AMPAR-containing vesicles towards the synapse, and fusion of these incoming vesicles with the postsynaptic membrane. Here, we show that the exocyst complex links these two distinct functions through the concerted actions of Sec8 and Exo70. This conclusion is based on four main experimental observations. First, both Sec8 and Exo70 are required for AMPAR function at synapses, as assayed electrophysiologically. Second, both Sec8 and Exo70 dominant negative mutants prevent receptor delivery to the spine surface, but in addition, only blockade of Exo70 function leads to accumulation of AMPARs at intracellular compartments within the spine. Third, exocytic vesicle proteins and AMPARs remain associated with the PSD upon interference with Exo70 function, whereas a Sec8 dominant negative produces the opposite effect. And fourth, AMPARs accumulate within the distal part of the PSD in very close proximity to the synaptic cleft when Exo70 function is blocked. These observations lead us to propose a model for the anatomical and molecular events that lead to the delivery of AMPARs into synapses (Figure 9). First, AMPARs are transported through intracellular membranes, probably

belonging to the trans-Golgi network and recycling endosomes (Horton and Ehlers, 2003, 2004; Gerges *et al*, 2004a; Park *et al*, 2004), to reach the spine compartment. This process is constitutive for GluR2/GluR3 receptors, but requires CaMKII activation for GluR1/GluR2 receptors (Piccini and Malinow, 2002). The translocation of AMPARs into the spine is assisted by PDZ-dependent interactions of Sec8, but is independent from the Exo70 subunit of the exocyst. Exocytic vesicles containing AMPARs bud from intra-spine compartments and are targeted towards the postsynaptic density, in a process probably directed by interactions between the Sec8 subunit of the exocyst and synaptic scaffolding molecules (Figure 9, left panel). These vesicles dock with the postsynaptic membrane within the PSD, leading to a transient association between AMPARs, vesicular proteins, such as Sec8 and Rab8, and the postsynaptic density (Figure 9, upper right panel). This step is blocked by Sec8-Ct, presumably by competing with the Sec8-PDZ-dependent interactions that direct the vesicles towards the PSD. Finally, this complex is resolved by exocytic membrane fusion mediated by the Exo70 subunit of the exocyst (Figure 9, lower right panel). The C-terminus of Exo70 is required for this step, since Exo70-N prevents AMPAR delivery into the synaptic surface.

Our results reinforce the current view that different exocyst subunits perform individual functions during the exocytic process (Boyd *et al*, 2004), with Sec8 being specialized in vesicle transport and targeting, and Exo70 in membrane fusion. More importantly, this model provides new information about the distinct organization of this exocytic machinery in neurons for the delivery of AMPARs at synapses. The exocyst was originally proposed to act as a spatial landmark for polarized exocytosis in yeast (Finger *et al*, 1998). We now know that the exocyst mediates surface delivery of a variety of membrane proteins in different cell types. Therefore, it is expected that cell- and structure-specific proteins will recruit the exocyst to localize delivery of different cargo to their

specific targets. In the case of the synaptic delivery of AMPA receptors, the PDZ-dependent interaction between Sec8 and synaptic scaffolding molecules, such as PSD95, seems to be the determining factor for membrane targeting. This interpretation is consistent with the observation that up- or down-regulation of PSD95 synaptic levels bidirectionally controls AMPAR delivery at synapses in hippocampus (El-Husseini *et al*, 2000; El-Husseini Ael *et al*, 2002; Colledge *et al*, 2003; Ehrlich and Malinow, 2004) and cerebral cortex (Beique and Andrade, 2003).

Finally, these results have offered unexpected anatomical information about the subcellular pathways followed by AMPARs to reach their synaptic targets. It has been previously proposed that the constitutive delivery of GluR2 receptors occurs by direct insertion at the postsynaptic membrane, whereas GluR1-containing receptors would reach the synaptic membrane by lateral translocation, after being inserted at extrasynaptic membranes (Passafaro *et al*, 2001). This model is compatible with the reported lateral diffusion of AMPARs on the dendritic plasma membrane (Tardin *et al*, 2003; Groc *et al*, 2004). However, our results do not support this model for synaptic delivery. We have found that impairment of AMPAR surface delivery with the Exo70 dominant negative leads to the accumulation of both recombinant GluR1 and GluR2 receptors within the spine, and not in the surrounding dendritic space. Using immunogold electron microscopy, we have shown that endogenous AMPARs accumulate within the PSD, in very close proximity to the synaptic membrane, when Exo70 function is blocked. And finally, AMPARs biochemically associate with PSD fractions when surface delivery is impaired. Taken together, these results strongly suggest that AMPARs are targeted directly into the PSD, but become locked in an abortive complex at its intracellular side when membrane fusion is prevented. We do not know the reason for this discrepancy in the spatial pattern of AMPAR delivery, although a potentially important experimental difference may be the use of organotypic slice cultures




Figure 9 Model for the exocyst-mediated targeting and delivery of AMPA receptors at the postsynaptic membrane. (Left) Exocytic vesicles containing AMPA receptors reach dendritic spines, bud from intracellular compartments within the spine and are directed towards the PSD by interactions between Sec8 and synaptic scaffolding molecules, such as PSD95. (Upper right): Incoming vesicles dock at the PSD forming a transient complex with it. (Lower right) This complex is resolved by membrane fusion with the postsynaptic membrane. This process is mediated by Exo70. See further explanation in the main text. TGN: *trans*-Golgi network; RE: recycling endosome; EV: exocytic vesicle; DV: docked vesicle.

in our experiments versus dissociated neuronal cultures for the reports on the lateral translocation of AMPARs (see also two recent reports supporting intra-spine delivery in slices (Kopeck *et al*, 2006) and lateral diffusion in dissociated neurons (Adesnik *et al*, 2005)). Future experiments will be needed to resolve the physiological role of these different delivery pathways.

In conclusion, our results shed light into the complex array of events that coordinate the transport, targeting and insertion of AMPARs at the postsynaptic membrane, and provide a molecular and anatomical map of the intracellular machinery that catalyzes this process along dendrites and within the dendritic spine.

Materials and methods

Electrophysiology

Simultaneous whole-cell recordings were carried out from nearby pairs of infected/transfected and uninfected/untransfected CA1 pyramidal neurons using fluorescence and transmitted light illumination. Synaptic responses were evoked with two bipolar electrodes (stimulation pulses of 200 μ s, up to 20 V) placed on the Schaffer collateral fibers between 300 and 500 μ m of the recorded cell. The responses obtained from the two stimulating electrodes were averaged for each cell and counted as an 'n' of 1. This experimental configuration ensures that the potential effects of the expressed recombinant proteins are postsynaptic, since expression is always in CA1 neurons and stimulation is applied on Schaffer collateral fibers from CA3 neurons. Synaptic responses mediated by AMPA or NMDA were acquired at -60 or $+40$ mV, respectively, in the presence of 100 μ M picrotoxin to block GABA_A receptors. GABA_A responses were acquired at 0 mV in the absence of receptor

antagonists; therefore, they should be considered as a combination of mono- and disynaptic IPSCs. All synaptic responses were averaged over 50–100 trials. Voltage-clamp whole-cell recordings were acquired with a Multiclamp 700 A amplifier (Axon Instruments). Composition of perfusion and internal solutions are described in the Supplementary data.

Imaging

Confocal and electron microscopy experiments are described in the Supplementary data.

Biochemistry

Biochemical techniques for membrane fractionation and immunoprecipitations from synaptosomal preparations are described in the Supplementary data.

Supplementary data

Supplementary data are available at *The EMBO Journal* Online.

Acknowledgements

We thank Alan Saltiel for the plasmids containing the Exo70 and Sec8 coding sequences, Shu-Chan Hsu for the monoclonal antibody against Exo70, Roger Tsien for the red fluorescence protein variant tdimer2(12), Dorothy Sorenson and Chris Edwards for their assistance with the immunogold electron microscopy, members of the Esteban lab and María Soengas for critical reading of the manuscript. This work was supported by grants from the National Institute of Mental Health (grant MH070417) and the National Alliance for Research on Schizophrenia and Depression (NARSAD) to JAE, and grant GM63032 from the National Institute of General Medical Sciences to MRS. We also thank Dr Thomas Coles for sponsoring JAE as an Investigator of the National Alliance for the Mentally Ill (NAMI).

References

- Adesnik H, Nicoll RA, England PM (2005) Photoinactivation of native AMPA receptors reveals their real-time trafficking. *Neuron* **48**: 977–985
- Barry MF, Ziff EB (2002) Receptor trafficking and the plasticity of excitatory synapses. *Curr Opin Neurobiol* **12**: 279–286
- Beique JC, Andrade R (2003) PSD-95 regulates synaptic transmission and plasticity in rat cerebral cortex. *J Physiol* **546**: 859–867
- Boyd C, Hughes T, Pypaert M, Novick P (2004) Vesicles carry most exocyst subunits to exocytic sites marked by the remaining two subunits, Sec3p and Exo70p. *J Cell Biol* **167**: 889–901
- Bredt DS, Nicoll RA (2003) AMPA receptor trafficking at excitatory synapses. *Neuron* **40**: 361–379
- Brown TC, Tran IC, Backos DS, Esteban JA (2005) NMDA receptor-dependent activation of the small GTPase Rab5 drives the removal of synaptic AMPA receptors during hippocampal LTD. *Neuron* **45**: 81–94
- Collodge M, Snyder EM, Crozier RA, Soderling JA, Jin Y, Langeberg LK, Lu H, Bear MF, Scott JD (2003) Ubiquitination regulates PSD-95 degradation and AMPA receptor surface expression. *Neuron* **40**: 595–607
- Ehrlich I, Malinow R (2004) Postsynaptic density 95 controls AMPA receptor incorporation during long-term potentiation and experience-driven synaptic plasticity. *J Neurosci* **24**: 916–927
- El-Husseini AE, Schnell E, Chetkovich DM, Nicoll RA, Bredt DS (2000) PSD-95 involvement in maturation of excitatory synapses. *Science* **290**: 1364–1368
- El-Husseini Ael D, Schnell E, Dakoji S, Sweeney N, Zhou Q, Prange O, Gauthier-Campbell C, Aguilera-Moreno A, Nicoll RA, Bredt DS (2002) Synaptic strength regulated by palmitate cycling on PSD-95. *Cell* **108**: 849–863
- Esteban JA (2003) AMPA receptor trafficking: a road map for synaptic plasticity. *Mol Interv* **3**: 375–385
- Finger FP, Hughes TE, Novick P (1998) Sec3p is a spatial landmark for polarized secretion in budding yeast. *Cell* **92**: 559–571
- Garner CC, Nash J, Hagan RL (2000) PDZ domains in synapse assembly and signalling. *Trends Cell Biol* **10**: 274–280
- Gerdes NZ, Backos DS, Esteban JA (2004a) Local control of AMPA receptor trafficking at the postsynaptic terminal by a small GTPase of the Rab family. *J Biol Chem* **279**: 43870–43878
- Gerdes NZ, Tran IC, Backos DS, Harrell JM, Chinkers M, Pratt WB, Esteban JA (2004b) Independent functions of hsp90 in neurotransmitter release and in the continuous synaptic cycling of AMPA receptors. *J Neurosci* **24**: 4758–4766
- Groc L, Heine M, Cognet L, Brickley K, Stephenson FA, Lounis B, Choquet D (2004) Differential activity-dependent regulation of the lateral mobilities of AMPA and NMDA receptors. *Nat Neurosci* **7**: 695–696
- Hayashi Y, Shi SH, Esteban JA, Piccini A, Poncer JC, Malinow R (2000) Driving AMPA receptors into synapses by LTP and CaMKII: requirement for GluR1 and PDZ domain interaction. *Science* **287**: 2262–2267
- Hazuka CD, Foletti DL, Hsu SC, Kee Y, Hopf FW, Scheller RH (1999) The sec6/8 complex is located at neurite outgrowth and axonal synapse-assembly domains. *J Neurosci* **19**: 1324–1334
- Henley JM (2003) Protein interactions implicated in AMPA receptor trafficking: a clear destination and an improving route map. *Neurosci Res* **45**: 243–254
- Horton AC, Ehlers MD (2003) Dual modes of endoplasmic reticulum-to-Golgi transport in dendrites revealed by live-cell imaging. *J Neurosci* **23**: 6188–6199
- Horton AC, Ehlers MD (2004) Secretory trafficking in neuronal dendrites. *Nat Cell Biol* **6**: 585–591
- Hsu SC, Hazuka CD, Roth R, Foletti DL, Heuser J, Scheller RH (1998) Subunit composition, protein interactions, and structures of the mammalian brain sec6/8 complex and septin filaments. *Neuron* **20**: 1111–1122
- Hsu SC, Ting AE, Hazuka CD, Davanger S, Kenny JW, Kee Y, Scheller RH (1996) The mammalian brain rsec6/8 complex. *Neuron* **17**: 1209–1219
- Inoue M, Chang L, Hwang J, Chiang SH, Saltiel AR (2003) The exocyst complex is required for targeting of Glut4 to the plasma membrane by insulin. *Nature* **422**: 629–633

- Kee Y, Yoo JS, Hazuka CD, Peterson KE, Hsu SC, Scheller RH (1997) Subunit structure of the mammalian exocyst complex. *Proc Natl Acad Sci USA* **94**: 14438–14443
- Kim E, Sheng M (2004) PDZ domain proteins of synapses. *Nat Rev Neurosci* **5**: 771–781
- Kopec CD, Li B, Wei W, Boehm J, Malinow R (2006) Glutamate receptor exocytosis and spine enlargement during chemically induced long-term potentiation. *J Neurosci* **26**: 2000–2009
- Kornau HC, Schenker LT, Kennedy MB, Seeburg PH (1995) Domain interaction between NMDA receptor subunits and the postsynaptic density protein PSD-95. *Science* **269**: 1737–1740
- Luscher C, Xia H, Beattie EC, Carroll RC, von Zastrow M, Malenka RC, Nicoll RA (1999) Role of AMPA receptor cycling in synaptic transmission and plasticity. *Neuron* **24**: 649–658
- Malinow R, Malenka RC (2002) AMPA receptor trafficking and synaptic plasticity. *Annu Rev Neurosci* **25**: 103–126
- McGee AW, Brecht DS (2003) Assembly and plasticity of the glutamatergic postsynaptic specialization. *Curr Opin Neurobiol* **13**: 111–118
- Naisbitt S, Valtschanoff J, Allison DW, Sala C, Kim E, Craig AM, Weinberg RJ, Sheng M (2000) Interaction of the postsynaptic density-95/guanylate kinase domain-associated protein complex with a light chain of myosin-V and dynein. *J Neurosci* **20**: 4524–4534
- Niethammer M, Kim E, Sheng M (1996) Interaction between the C terminus of NMDA receptor subunits and multiple members of the PSD-95 family of membrane-associated guanylate kinases. *J Neurosci* **16**: 2157–2163
- Nishimune A, Isaac JT, Molnar E, Noel J, Nash SR, Tagaya M, Collingridge GL, Nakanishi S, Henley JM (1998) NSF binding to GluR2 regulates synaptic transmission. *Neuron* **21**: 87–97
- Nong Y, Huang YQ, Salter MW (2004) NMDA receptors are moving in. *Curr Opin Neurobiol* **14**: 353–361
- Park M, Penick EC, Edwards JG, Kauer JA, Ehlers MD (2004) Recycling endosomes supply AMPA receptors for LTP. *Science* **305**: 1972–1975
- Passafaro M, Piech V, Sheng M (2001) Subunit-specific temporal and spatial patterns of AMPA receptor exocytosis in hippocampal neurons. *Nat Neurosci* **4**: 917–926
- Peng J, Kim MJ, Cheng D, Duong DM, Gygi SP, Sheng M (2004) Semi-quantitative proteomic analysis of rat forebrain postsynaptic density fractions by mass spectrometry. *J Biol Chem* **279**: 21003–21011
- Perez-Otano I, Ehlers MD (2005) Homeostatic plasticity and NMDA receptor trafficking. *Trends Neurosci* **28**: 229–238
- Piccini A, Malinow R (2002) Critical postsynaptic density 95/disc large/zonula occludens-1 interactions by glutamate receptor 1 (GluR1) and GluR2 required at different subcellular sites. *J Neurosci* **22**: 5387–5392
- Prybylowski K, Chang K, Sans N, Kan L, Vicini S, Wenthold RJ (2005) The synaptic localization of NR2B-containing NMDA receptors is controlled by interactions with PDZ proteins and AP-2. *Neuron* **47**: 845–857
- Riefler GM, Balasingam G, Lucas KG, Wang S, Hsu SC, Firestein BL (2003) Exocyst complex subunit sec8 binds to postsynaptic density protein-95 (PSD-95): a novel interaction regulated by cyprin (cytosolic PSD-95 interactor). *Biochem J* **373**: 49–55
- Sans N, Petralia RS, Wang YX, Blahos II J, Hell JW, Wenthold RJ (2000) A developmental change in NMDA receptor-associated proteins at hippocampal synapses. *J Neurosci* **20**: 1260–1271
- Sans N, Prybylowski K, Petralia RS, Chang K, Wang YX, Racca C, Vicini S, Wenthold RJ (2003) NMDA receptor trafficking through an interaction between PDZ proteins and the exocyst complex. *Nat Cell Biol* **5**: 520–530
- Scannevin RH, Hagan RL (2000) Postsynaptic organization and regulation of excitatory synapses. *Nat Rev Neurosci* **1**: 133–141
- Sheng M (2001) Molecular organization of the postsynaptic specialization. *Proc Natl Acad Sci USA* **98**: 7058–7061
- Sheng M, Lee SH (2001) AMPA receptor trafficking and the control of synaptic transmission. *Cell* **105**: 825–828
- Shi S, Hayashi Y, Esteban JA, Malinow R (2001) Subunit-specific rules governing AMPA receptor trafficking to synapses in hippocampal pyramidal neurons. *Cell* **105**: 331–343
- Shin DM, Zhao XS, Zeng W, Mozhayeva M, Muallem S (2000) The mammalian Sec6/8 complex interacts with Ca(2+) signaling complexes and regulates their activity. *J Cell Biol* **150**: 1101–1112
- Song I, Kamboj S, Xia J, Dong H, Liao D, Hagan RL (1998) Interaction of the N-ethylmaleimide-sensitive factor with AMPA receptors. *Neuron* **21**: 393–400
- Steiner P, Alberi S, Kulangara K, Yersin A, Sarria JC, Regulier E, Kasas S, Dietler G, Muller D, Catsicas S, Hirling H (2005) Interactions between NEEP21, GRIP1 and GluR2 regulate sorting and recycling of the glutamate receptor subunit GluR2. *EMBO J* **24**: 2873–2884
- Tardin C, Cognet L, Bats C, Lounis B, Choquet D (2003) Direct imaging of lateral movements of AMPA receptors inside synapses. *EMBO J* **22**: 4656–4665
- TerBush DR, Maurice T, Roth D, Novick P (1996) The Exocyst is a multiprotein complex required for exocytosis in *Saccharomyces cerevisiae*. *EMBO J* **15**: 6483–6494
- Ting AE, Hazuka CD, Hsu SC, Kirk MD, Bean AJ, Scheller RH (1995) rSec6 and rSec8, mammalian homologs of yeast proteins essential for secretion. *Proc Natl Acad Sci USA* **92**: 9613–9617
- Triller A, Choquet D (2005) Surface trafficking of receptors between synaptic and extrasynaptic membranes: and yet they do move!. *Trends Neurosci* **28**: 133–139
- Vega IE, Hsu SC (2001) The exocyst complex associates with microtubules to mediate vesicle targeting and neurite outgrowth. *J Neurosci* **21**: 3839–3848
- Vik-Mo EO, Olteidal L, Hoivik EA, Kleivdal H, Eidet J, Davanger S (2003) Sec6 is localized to the plasma membrane of mature synaptic terminals and is transported with secretogranin II-containing vesicles. *Neuroscience* **119**: 73–85
- Walikonis RS, Jensen ON, Mann M, Provance Jr DW, Mercer JA, Kennedy MB (2000) Identification of proteins in the postsynaptic density fraction by mass spectrometry. *J Neurosci* **20**: 4069–4080
- Wang S, Liu Y, Adamson CL, Valdez G, Guo W, Hsu SC (2004) The mammalian exocyst, a complex required for exocytosis, inhibits tubulin polymerization. *J Biol Chem* **279**: 35958–35966
- Wyszynski M, Valtschanoff JG, Naisbitt S, Dunah AW, Kim E, Standaert DG, Weinberg R, Sheng M (1999) Association of AMPA receptors with a subset of glutamate receptor-interacting protein *in vivo*. *J Neurosci* **19**: 6528–6537
- Yamagata M, Sanes JR, Weiner JA (2003) Synaptic adhesion molecules. *Curr Opin Cell Biol* **15**: 621–632
- Yeaman C, Grindstaff KK, Wright JR, Nelson WJ (2001) Sec6/8 complexes on trans-Golgi network and plasma membrane regulate late stages of exocytosis in mammalian cells. *J Cell Biol* **155**: 593–604